

A Graduate Course in Probability

Liviu I. Nicolaescu

University of Notre Dame

Last revised [2026-05-25, 09:53:27](#) .

Copyright © 2019, All rights reserved
Liviu I. Nicolaescu¹

¹Started December 11, 2018. Completed October 19, 2021 . Last modified on 2026-05-25.

Introduction

In no other branch of mathematics is it so easy for experts to blunder as in probability theory.

Martin Gardner

I have to confess that my mathematical formation is not that of a probabilist. I am a geometer/analyst by training. About fifteen years ago I stumbled on some probabilistic geometry questions. The ad-hoc methods I used were producing encouraging but unsatisfactory answers. A chance encounter with a trained probabilist led me to a pretty advanced monograph dealing with related problems from a probabilistic view point. I spent a sabbatical year learning probability so I could understand that book.

I eventually did understand that book, I was able to phrase the original questions in a better language and I even offered answers to questions I could not conceive before. A “side effect” of this effort was that I got a taste of probability.

To the geometer in me, the probabilistic thinking looked (and still looks) like mathematics with a bit more, somewhat similar to classical mechanics, that is mathematics with a sprinkle of physical intuition. I find this subject fresh, full of interesting and enticing questions. This is how my probabilistic journey began and I have been enjoying it since. In the meantime I matured a bit more by teaching probability, both at undergraduate and graduate level. This book partially reflects this personal journey.

Probability theory has grown out of many concrete examples and questions and I firmly believe that probabilistic thinking can only be grasped through examples. Compared to other mathematical areas I am familiar with, probability contains an unusually large number of counterintuitive results. To me, these represent one of the attractive features of the subject. So a substantial part of this book is devoted to examples, some truly fundamental and quite a few more esoteric, but which are aesthetically very pleasing and pedagogically very revealing. Some of these examples are recurring, appearing in many places in the text and, as we develop more and more sophisticated technology, we dig a deeper and deeper into them.

While teaching probability I discovered that probabilistic simulations enhance the understanding of probabilistic thinking. That is why I have included a brief introduction to R and a few of the simple codes that allows one to do basic Monte-Carlo simulations. I hope I can tempt the reader to try a few

of these and be amazed, like myself and my students, of the remarkable agreement between practice and theory.

I have divided the book into five chapters. The first one concentrates on the measure theoretic foundations of probability and its theoretical part. It is essentially the content of Kolmogorov's foundational monograph. I assume that the reader is familiar with the measure theory and integration. I survey this subject and I present complete proofs only of results that have important probabilistic applications or significance.

The first genuinely probabilistic concept is that of independence and I prove early on Kolmogorov's zero-one theorem. It is a striking all-or-nothing result and its deeper implications are gradually revealed in the later parts of the book. The ubiquitous concept of random variable and its numerical characteristics are discussed in detail. Along the way I discuss the various modes of convergence of random variables. I made sure the reader has the opportunity to see these ideas at work so I present many classical random variables and some of their probabilistic occurrences. Among the classical problems/themes I discuss I should mention, the inclusion-exclusion principle, sieves and Poissonization, Poisson processes, the coupon collector problem, the longest common subsequence problem.

Section 4, one of the largest of this chapter, is devoted to the concept of conditional expectation, a central probabilistic concept that takes some getting used to. Analytically, the existence of conditional expectation is a simple consequence of the Radon-Nicodym theorem. This however hides its probabilistic significance. I opted for the more involved approach that reveals the meaning of this object as the best predictor given certain information.

To get to the heart of the rather subtle concept of conditional expectation I tried to present many examples, from simple computations to more sophisticated applications to stochastic optimization problems such as the classical secretary problem. I spend considerable time on the concept of kernels a.k.a. random measures, regular conditional distributions and disintegration of measure describing the various connections between them. I opted to only sketch the proof of the existence of regular conditional distributions since I felt that the missing details add little to the understanding of this important concept. Instead, I have included a large and varied number of concrete examples to give the reader a better feel of this concept.

The last section of this chapter is an introduction to stochastic processes. The central result of this section is Kolmogorov's existence/consistency theorem that guarantees that various objects discussed in the previous sections do indeed have a mathematical existence. I decided to present a complete proof of this result so the reader can see the source of this existence, namely Tikhonov's compactness theorem, a result that is deeply rooted in the foundations of mathematics.

Chapter 2 is devoted to a major theme in probability, the law of large numbers and its relatives. To quote Gnedenko and Kolmogorov, [91], "*all epistemological value of the theory of probability is based on this: that large-scale random phenomena in their collective action create strict, nonrandom regularity*".

The first section is devoted to the Strong Law of Large Numbers. I present Kolmogorov's proof that reduces this result to the convergence of random series with independent summands. I find the Law of Large Numbers philosophically surprising since it extracts order out of chaos. The Monte Carlo method is one convincing manifestation of the order-out-chaos phenomenon. I could not pass the opportunity to introduce the concept of entropy and its application via the law of large numbers to coding/compression of data.

The second section is devoted to the CLT, Central Limit Theorem. I discuss in some detail the concept of weak convergence of probability measures on Euclidean spaces and present two proofs of the CLT, the classical one based on the Fourier transform, and a less familiar one, based on the theory of convolution semigroups.

The third section is devoted to concentration inequalities. We describe the basics of Chernoff's estimates and produce a few fundamental concentration inequalities. As an application we discuss Lindenstrauss-Johnson lemma stating that the geometry of a cloud of points in a high-dimensional vector space is, with high confidence, little disturbed by an orthogonal projection onto a random subspace of much smaller dimension.

Section 4 is devoted to more modern considerations, namely uniform limits of empirical processes. The Glivenko-Cantelli is the pioneering result in this direction. I also discuss more recent results showing how this uniform convergence can be obtained by combining the concentration results in the previous section and the concept of VC-families/dimension. I briefly describe the significance of such results to PAC-learning, a concept central in machine learning.

The last section of this chapter is a brief introduction to the theory of Brownian motion. I used it as an opportunity to discuss more concepts and results involving stochastic processes such as Gaussian processes and Kolmogorov's continuity theorem.

Chapter 3 is devoted to the castle that J. L. Doob built, namely the theory of martingales, discrete and continuous. I present in detail the theoretical pillars of this edifice: stopping/sampling, asymptotic behavior, maximal inequalities and I discuss a large and diverse collection of examples: occurrence of patterns, Galston-Watson processes, optimal gambling strategies, Azuma and McDiarmid inequalities and their application to combinatorial optimization problems, backwards martingales, exchangeable sequences, de Finetti's theorem, and asymptotics in Polya's urn problem, Brownian motion.

Chapter 4 is an introduction to Markov chains. This beautiful and rich subject is still actual, growing, and has many applications and ramifications. The first three sections are devoted to the "classical" part of this subject and culminates with the law of large numbers for such stochastic processes. Section 4 is devoted to a more recent (1950's) point of view, namely the connection between reversible Markov chains and electrical networks. I adopt a more geometric approach based on the old observations of H. Weyl and R. Bott (see [23]) that Kirchhoff's laws have a Hodge theoretic description. The last section is devoted to finite Markov chains I describe various ways of estimating the rate of convergence of irreducible recurrent Markov chains. The chapter ends with brief discussion of the Markov Chain Monte Carlo methods.

The last chapter of the book is the shortest and is devoted to the classical ergodic theorems. I have included it because I felt I owed it to the reader to highlight a principle that unifies and clarifies the main limit theorems in Chapters 2 and 4.

As the title indicates, this book is meant as an introduction to the modern, i.e., post Kolmogorov's axiomatization, theory of probability. The reader is assumed to have some familiarity with measure theory and integration and be comfortable with the basic objects and concepts of modern analysis: metric/topological spaces, convergence, compactness. In a few places, familiarity with basic properties of Banach spaces is assumed.

This book could serve as a textbook for a year-long basic graduate course in probability. With this purpose in mind I have included a relatively large number of exercises, many of them nontrivial and highlighting aspects I did not include in the main body of the text.

The book grew up from notes for a one-semester graduate course in probability that I taught at the University of Notre Dame. That course covered Chapter 1, the classical limit theorems (Sec.2.1-2.3) and discrete time martingales (Sec. 3.1-3.2). Some of the proofs appear in fine print as a suggestion to the potential student/instructor that they can be skipped at a first encounter with this subject.

Work on this book has been my constant happy companion during these improbable pandemic times. I hope I was able to convey my curiosity, fascination and enthusiasm about probability and convince some readers to dig deeper into this intellectually rewarding subject.

Notre Dame, May 2022

Notation and conventions

- We set $\mathbb{N} := \mathbb{Z}_{>0}$, $\mathbb{N}_0 := \mathbb{Z}_{\geq 0}$.
- We set $\mathbb{R}_+ := [0, \infty)$.
- For $a, b \in \mathbb{Z}$, $a \leq b$ we set $\llbracket a, b \rrbracket := [a, b] \cap \mathbb{Z}$.
- For $n \in \mathbb{N}$ we set $\mathbb{I}_n := \llbracket 1, n \rrbracket$.
- For $x \in \mathbb{R}$ we set $\lfloor x \rfloor := \max \mathbb{Z} \cap (-\infty, x]$, $\lceil x \rceil := \min \mathbb{Z} \cap [x, \infty)$.
- For $n \in \mathbb{N}$ we denote by $\mathfrak{S}_n$ the group of permutations of $\mathbb{I}_n$.
- $x \wedge y := \min(x, y)$, $x \vee y := \max(x, y)$.
- $i := \sqrt{-1}$
- Given a subset A of a set X we denote by A^c its complement (in X).
- For any set X we denote by 2^X the collection of all the subsets of X .
- For any set X we denote by 2_0^X the collection of all the *finite* subsets of X .
- We will denote by $|S|$ or $\#S$ the cardinality of a set S .
- For natural numbers $n \geq k$ we denote by $(n)_k$ the falling factorial,

$$(n)_k := n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}.$$

- If T is a topological space, then $\mathcal{B}_T$ denotes the σ -algebra of Borel subsets of T .
- We denote by λ the standard Lebesgue measure on $\mathbb{R}$ and by λ_n the standard Lebesgue measure on $\mathbb{R}^n$.
- If $(\Omega, \mathcal{F})$ is a measurable space and $(\mathcal{A}_i)_{i \in I}$ is a collection of subsets of $\mathcal{F}$, then $\sigma(\mathcal{A}_i, i \in I)$ denotes the smallest sub- σ -algebra of $\mathcal{F}$ containing all the collections $\mathcal{A}_i$. Sometimes I will use the alternate notation

$$\bigvee_{i \in I} \mathcal{A}_i := \sigma(\mathcal{A}_i, i \in I).$$

- For a collection $(X_i)_{i \in I}$ of random variables defined on the same probability space we denote by $\sigma(X_i; i \in I)$ the sub- σ -algebra generated by these variables.

- Given an ambient set Ω and a subset $A \subset \Omega$ I denote by $I_A : \Omega \rightarrow \{0, 1\}$ the *indicator function* of A ,

$$I_A(\omega) = \begin{cases} 1, & \omega \in A, \\ 0, & \omega \notin A. \end{cases}$$

- We denote by ω_n the volume of the unit ball in $\mathbb{R}^n$ and by σ_{n-1} the “area” of the unit $(n - 1)$ -dimensional sphere in $\mathbb{R}^n$.

$$\omega_n = \frac{1}{n} \sigma_{n-1}, \quad \sigma_{n-1} = \frac{2\Gamma(1/2)^n}{\Gamma(n/2)}.$$

Contents

Introduction	i
Notation and conventions	v
	Page
Chapter 1. Foundations	1
§1.1. Measurable spaces	2
1.1.1. Sigma-algebras	2
1.1.2. Measurable maps	5
§1.2. Measures and integration	12
1.2.1. Measures	12
1.2.2. Independence and conditional probability	21
1.2.3. Integration of measurable functions	30
1.2.4. L^p spaces	36
1.2.5. Measures on compact metric spaces	38
§1.3. Invariants of random variables	40
1.3.1. The distribution and the expectation of a random variable	40
1.3.2. Higher order integral invariants of random variables	47
1.3.3. Classical examples of discrete random variables	52
1.3.4. Classical examples of continuous probability distributions	62
1.3.5. Product probability spaces and independence	66
1.3.6. Convolution of Borel measures on the real axis	74
1.3.7. Poisson processes	76
1.3.8. Modes of convergence of random variables	79
§1.4. Conditional expectation	87
1.4.1. Conditioning on a sigma-subalgebra	88
1.4.2. Some applications of conditioning	97
1.4.3. Conditional independence	106
1.4.4. Kernels and regular conditional distributions	107
1.4.5. Disintegration of measures	115

§1.5. What are stochastic processes?	120
1.5.1. Definition and examples	120
1.5.2. Kolmogorov's existence theorem	124
§1.6. Exercises	129
Chapter 2. Limit theorems	147
§2.1. The Law of Large Numbers	148
2.1.1. Random series	148
2.1.2. The Law of Large Numbers	153
2.1.3. Entropy and compression	160
§2.2. The Central Limit Theorem	165
2.2.1. Weak and vague convergence	166
2.2.2. The characteristic function	180
2.2.3. The Central Limit Theorem	188
2.2.4. Semigroup approach to CLT	190
§2.3. Concentration inequalities	196
2.3.1. The Chernoff bound	197
2.3.2. Some applications	202
§2.4. Uniform laws of large numbers	207
2.4.1. The Glivenko-Cantelli theorem	208
2.4.2. More general uniform limit laws	211
2.4.3. PAC learning	218
§2.5. The Brownian motion	221
2.5.1. Heuristics	221
2.5.2. Gaussian measures and processes	224
2.5.3. The Brownian motion	232
§2.6. Exercises	243
Chapter 3. Martingales	269
§3.1. Basic facts about martingales	269
3.1.1. Definition and examples	270
3.1.2. Doob decomposition	275
3.1.3. Discrete stochastic integrals	277
3.1.4. Stopping and sampling: discrete time	278
3.1.5. Applications of the optional sampling theorem	283
3.1.6. Concentration inequalities: martingale techniques	288
3.1.7. Uniform laws of large numbers revisited	294
§3.2. Limit theorems: discrete time	297
3.2.1. Almost sure convergence	297
3.2.2. Uniform integrability	305
3.2.3. Uniformly integrable martingales	311
3.2.4. Applications of the optional sampling theorem	316
3.2.5. Uniformly integrable submartingales	322
3.2.6. Maximal inequalities and L^p -convergence	329
3.2.7. Backwards martingales	334

3.2.8.	Exchangeable sequences of random variables	336
§3.3.	Generalities about filtered processes	344
3.3.1.	Continuous parameter filtrations	344
3.3.2.	Stopping and optional times	346
§3.4.	The Brownian motion as a filtered process	348
3.4.1.	Some fundamental properties of the Brownian motion	348
3.4.2.	Some applications	355
3.4.3.	The support theorem	357
§3.5.	Continuous-time martingales	362
3.5.1.	Definition and examples of continuous-time martingales	362
3.5.2.	Limit theorems	368
3.5.3.	Sampling and stopping	371
§3.6.	Some applications	375
3.6.1.	The invariance theorem and applications	376
3.6.2.	The Kolmogorov-Smirnov statistics	381
§3.7.	Exercises	388
Chapter 4.	Markov chains	399
§4.1.	Markov chains	400
4.1.1.	Definition and basic concepts	400
4.1.2.	Examples	406
§4.2.	The dynamics of homogeneous Markov chains	409
4.2.1.	Classification of states	409
4.2.2.	The strong Markov property	415
4.2.3.	Transience and recurrence	416
4.2.4.	Invariant measures	422
§4.3.	Martingale techniques	430
4.3.1.	Recurrence, transience and Lyapunov functions	430
4.3.2.	Optimal stopping times	439
§4.4.	Asymptotic behavior	453
4.4.1.	The ergodic theorem	453
4.4.2.	Aperiodic chains	455
4.4.3.	The coupling technique	458
§4.5.	Electric networks	460
4.5.1.	Reversible Markov chains as electric networks	460
4.5.2.	Sources, currents and chains	461
4.5.3.	Kirkhoff's laws and Hodge theory	463
4.5.4.	A probabilistic perspective of Kirchoff's laws	468
4.5.5.	Degenerations	471
4.5.6.	Applications	478
§4.6.	Finite Markov chains	483
4.6.1.	The Perron-Frobenius theory	484
4.6.2.	Variational methods	495
4.6.3.	Markov Chain Monte Carlo	503

§4.7. Exercises	510
Chapter 5. Elements of Ergodic Theory	523
§5.1. The ergodic theorem	523
5.1.1. Measure preserving maps and invariant sets	523
5.1.2. Ergodic theorems	531
§5.2. Applications	539
5.2.1. Limit theorems	539
5.2.2. Mixing	541
§5.3. Exercises	548
Chapter 6. An introduction to stochastic calculus	553
§6.1. Itô's integral with respect to the Brownian motion	555
6.1.1. Itô integrals of progressive processes	555
6.1.2. Itô's integral as a stochastic process	560
6.1.3. Localization	564
6.1.4. Local martingales	570
6.1.5. Finite variation processes	571
6.1.6. Itô calculus	575
6.1.7. Some computational examples	583
§6.2. A more general stochastic integral	591
6.2.1. Quadratic variation of a local martingale	592
6.2.2. The bracket of two local martingales	600
6.2.3. Continuous semimartingales	601
6.2.4. The stochastic integral with respect to an L^2 -bounded martingale	602
6.2.5. Stochastic integration with respect to local martingales	608
6.2.6. Stochastic integration with respect to semimartingales	610
6.2.7. Itô's formula revisited	613
§6.3. Some applications	615
6.3.1. The Doléans exponential	615
6.3.2. Lévy's characterization of Brownian motion	618
6.3.3. Martingale representation theorem	619
§6.4. Exercises	622
Appendix A. A few useful facts	625
§A.1. The Gamma function	625
§A.2. Basic invariants of frequently used probability distributions	626
§A.3. A glimpse at R	627
Bibliography	643
Index	651

Foundations

At the beginning of the twentieth century probability was in a fluid state. There was no clear mathematical concept of probability, and ad-hoc methods were used to rigorously formulate classical questions. Probability at that stage was a collection of interesting problems in search of a coherent setup. According to Jean Ville, a PhD student of M. Fréchet, in Paris probability was viewed among mathematicians as “*an honorable pastime for those who distinguished themselves in pure mathematics*”.

The whole enterprise seemed to be concerned with concepts that lie outside mathematics. Henri Poincaré himself wrote that “*one can hardly give a satisfactory definition of probability*”. As Richard von Mises pointed out in 1928, the German word for probability, “*wahrscheinlich*”, translates literally as “*truth resembling*”; see [218]. Bertrand Russell was quoted as saying in 1929 that “*Probability is the most important concept in modern science, especially as nobody has the slightest notion of what it means*”. The philosophical underpinnings of this concept are discussed even today. For more on this aspect we refer to the recent delightful book [56].

In his influential 1900 International Congress address in Paris D. Hilbert recognized this state of affairs and the importance of the subject. In the sixth problem of his famous list of 23 he asked, among other things, for rigorous foundations of probability. These were laid by A. N. Kolmogorov in his famous 1933 monograph [119]. According to Kolmogorov himself, this was not a research work, but a work of synthesis. A brilliant synthesis I might add. His point of view was universally adopted and modern probability theory was born. The theory of probability can now be informally divided into two eras: before and after Kolmogorov.

The present chapter is devoted to this foundational work of Kolmogorov. The pillars of probability theory are the concept of probability or sample space, random variables, independence, conditional expectations, and consistency, i.e., the existence of random variables or processes with prescribed statistics.

So efficient is his axiomatization that to the untrained eye, probability, as envisaged by Kolmogorov, may seem like a slice of measure theory. In a 1963 interview Kolmogorov complained that his axioms have been so successful on the theoretic side that many mathematicians lost interest in the problems and applications that were and are the main engines of growth of this subject. I understand his criticism since I too was one of those mathematicians that was not interested in these applications. Now I know better.

In this chapter I present these pillars of probability theory and prove their main properties. I have included a large number of detailed examples meant to convey the subtleties, depth, power and richness of these concepts. No abstract theorem can capture this richness.

I want to close with a personal anecdote that I find revealing. A few years ago, at a conference, I had a conversation with J. M. Bismut, a known probabilist whose mathematical interests were becoming more and more geometric. He noticed that I was in the middle of a mathematical transition in the opposite direction, from geometry towards probability, and he asked me what prompted it. I explained my motivation, how I discovered that probability is not just a glorious part of measure theory and how much I struggled to truly understand the concept of conditional expectation, a concept eminently probabilistic. He smiled and said: “*Probability theory is measure theory plus conditional expectation*”. I know it is an oversimplification, but it contains a lot of truth.

1.1. Measurable spaces

1.1.1. Sigma-algebras. Fix a nonempty set Ω .

Definition 1.1.1. (a) A collection $\mathcal{A}$ of subsets of Ω is called an *algebra* of Ω if it satisfies the following conditions

- (i) $\emptyset, \Omega \in \mathcal{A}$.
- (ii) $\forall A, B \in \mathcal{A}, A \cup B \in \mathcal{A}$.
- (iii) $\forall A \in \mathcal{A}, A^c \in \mathcal{A}$.

(b) A collection $\mathcal{S}$ of subsets of Ω is called a σ -*algebra* (or *sigma-algebra*) of Ω if it is an algebra of Ω and the union of any countable subfamily of $\mathcal{S}$ is a set in $\mathcal{S}$, i.e.,

$$\forall (A_n)_{n \in \mathbb{N}} \in \mathcal{S}^{\mathbb{N}}, \bigcup_{n \geq 1} A_n \in \mathcal{S}. \quad (1.1.1)$$

(c) A *measurable space* is a pair $(\Omega, \mathcal{S})$, where $\mathcal{S}$ is a sigma-algebra of subsets of Ω . The subsets $S \in \mathcal{S}$ are called $(\mathcal{S})$ -*measurable*. $\square$

Remark 1.1.2. To prove that an algebra $\mathcal{S}$ is a σ -algebra it suffices to verify (1.1.1) *only for increasing* sequence of subsets $B_n \in \mathcal{S}$. Indeed, if $(A_n)_{n \in \mathbb{N}}$ is an arbitrary family in $\mathcal{S}$ the new family of sets in $\mathcal{S}$

$$B_n = \bigcup_{k=1}^n A_k, \quad n \in \mathbb{N},$$

is increasing and its union coincides with the union of the family $(A_n)_{n \in \mathbb{N}}$. $\square$

Example 1.1.3. (a) The collection 2^Ω of all subsets of Ω is obviously a σ -algebra.

(b) Suppose that $\mathcal{S}$ is a (σ) -algebra of a set Ω and $F : \widehat{\Omega} \rightarrow \Omega$ is a map. Then the preimage

$$F^{-1}(\mathcal{S}) = \{ F^{-1}(S); S \in \mathcal{S} \}$$

is a (σ) -algebra of subsets of $\widehat{\Omega}$. The σ -algebra $F^{-1}(\mathcal{S})$ is denoted by $\sigma(F)$ and it is called the σ -*algebra generated by F* or the *pullback of $\mathcal{S}$ via F* . We will often use the more suggestive notation

$$\{F \in S\} := F^{-1}(S) = \{ \hat{\omega} \in \widehat{\Omega}; F(\hat{\omega}) \in S \}.$$

(c) Given $A \in \Omega$ we denote by $\mathcal{S}_A$ the σ -algebra generated by A , i.e.,

$$\mathcal{S}_A = \{ \emptyset, A, A^c, \Omega \}.$$

We will refer to it as the *Bernoulli algebra* with success A . Note that $\mathcal{S}_A$ is the pullback of $\mathcal{P}^{\{0,1\}}$ via the indicator function $\mathbf{I}_A : \Omega \rightarrow \{0, 1\}$.

(d) If $\mathcal{C} \subset \mathcal{P}^\Omega$ is a family of subsets of Ω , then we denote by $\sigma(\mathcal{C})$ the σ -algebra generated by $\mathcal{C}$, i.e., the intersection of all σ -algebras that contain $\mathcal{C}$. In particular, if $\mathcal{S}_1, \mathcal{S}_2$ are σ -algebras of Ω , then we set

$$\mathcal{S}_1 \vee \mathcal{S}_2 := \sigma(\mathcal{S}_1 \cup \mathcal{S}_2).$$

More generally, for any family $(\mathcal{S}_i)_{i \in I}$ of σ -algebras we set

$$\bigvee_{i \in I} \mathcal{S}_i := \sigma \left(\bigcup_{i \in I} \mathcal{S}_i \right).$$

(e) Suppose that we are given a countable partition $\{A_n\}_{n \in \mathbb{N}}$ of Ω

$$\Omega = \bigsqcup_{n \in \mathbb{N}} A_n.$$

The sets A_n are called the *chambers* of the partition. The σ -algebra generated by this partition is the σ -algebra consisting of all the subsets of Ω who are unions of chambers. This σ -algebra can be viewed as the σ -algebra generated by the map

$$X : \Omega \rightarrow \mathbb{N}, \quad X = \sum_{n \in \mathbb{N}} n \mathbf{I}_{A_n},$$

so that $A_n = X^{-1}(\{n\})$.

(f) If $(\mathcal{S}_i)_{i \in I}$ is a family of (σ) -algebras of Ω , then their intersection

$$\bigcap_{i \in I} \mathcal{S}_i \subset \mathcal{P}^\Omega$$

is a (σ) -algebra of Ω .

(g) If $(\Omega_1, \mathcal{S}_1)$ and $(\Omega_2, \mathcal{S}_2)$ are two measurable spaces, then we denote by $\mathcal{S}_1 \otimes \mathcal{S}_2$ the sigma algebra of $\Omega_1 \times \Omega_2$ generated by the collection

$$\{S_1 \times S_2 : S_1 \in \mathcal{S}_1, S_2 \in \mathcal{S}_2\} \subset \mathcal{P}^{\Omega_1 \times \Omega_2}.$$

(h) If X is a topological space and $\mathcal{T}_X \subset \mathcal{P}^X$ denotes the family of open subsets, then the *Borel* σ -algebra of X , denoted by $\mathcal{B}_X$, is the σ -algebra generated by $\mathcal{T}_X$. The sets in $\mathcal{B}_X$ are called the *Borel subsets* of X . Note that since any open set in $\mathbb{R}^n$ is a countable union of open cubes we have

$$\mathcal{B}_{\mathbb{R}^n} = \mathcal{B}_{\mathbb{R}}^{\otimes n}. \quad (1.1.2)$$

Any *finite dimensional* real vector space V can be equipped with a topology by choosing a linear isomorphism $L : V \rightarrow \mathbb{R}^{\dim V}$. This topology is independent of the choice of the isomorphism L . It can be alternatively identified as the smallest topology on V such that all the linear maps $V \rightarrow \mathbb{R}$ are continuous. We denote by $\mathcal{B}_V$ the sigma-algebra of Borel subsets determined by this topology.

We set $\bar{\mathbb{R}} = [-\infty, \infty]$. As a topological space it is homeomorphic to $[-1, 1]$. For simplicity we will refer to the Borel subsets of $\bar{\mathbb{R}}$ simply as *Borel sets*.

(i) If $(\Omega, \mathcal{S})$ is a measurable space and $X \subset \Omega$, then the collection

$$\mathcal{S}|_X := \{ S \cap X : S \in \mathcal{S} \} \subset 2^X$$

is a σ -algebra of X called *the trace of $\mathcal{S}$ on X* . $\square$

Remark 1.1.4 (Nedoma's pathology). Suppose that $(\Omega, \mathcal{S})$ is a measurable space. The product $\Omega \times \Omega$ contains a distinguished set, the diagonal

$$\Delta = \{ (\omega, \omega) : \omega \in \Omega \} \subset \Omega \times \Omega.$$

Then Δ is *not measurable* with respect to the product sigma-algebra $\mathcal{S} \otimes \mathcal{S}$ if

$$\text{Card } \Omega > \aleph_c = \text{Card } \mathbb{R}.$$

For a proof we refer to [184, Sec. 21.8].

Suppose that Ω is a *Hausdorff* topological vector space, and $\mathcal{S} = \mathcal{B}_X$ is its associated Borel sigma-algebra. The diagonal Δ is closed with respect to the product topology. In particular it belongs to the Borel sigma-algebra defined by the product topology. However, if $\text{Card } \Omega > \aleph_c$, then the diagonal it is not measurable with respect to the the product $\mathcal{S} \otimes \mathcal{S}$! In other words the product of Borel sigma-algebras is strictly smaller than the Borel sigma-algebra $\mathcal{B}_{X \times X}$ determined by the product topology! This phenomenon is referred to as the *Nedoma's pathology*. $\square$

Definition 1.1.5. Let $\mathcal{C}$ be a collection of subsets of a set Ω . We say that $\mathcal{C}$ is a π -system if it is closed under finite intersections, i.e.,

$$\forall A, B \in \mathcal{C} : A \cap B \in \mathcal{C}.$$

The collection $\mathcal{C}$ is called a λ -system if it satisfies the following conditions.

- (i) $\emptyset, \Omega \in \mathcal{C}$.
- (ii) if $A, B \in \mathcal{C}$ and $A \subset B$, then $B \setminus A \in \mathcal{C}$.
- (iii) If $A_1 \subset A_2 \subset \dots$ belong to $\mathcal{C}$, then so does their union.

$\square$

Note that a collection $\mathcal{C}$ is a σ -algebra if it is simultaneously a π and a λ -system.¹ Since the intersection of any family of λ -systems is a λ -system we deduce that for any collection $\mathcal{C} \subset 2^\Omega$ there exists a smallest λ -system containing $\mathcal{C}$. We denote this system by $\Lambda(\mathcal{C})$ and we will refer to it as the λ -system generated by $\mathcal{C}$.

Example 1.1.6. Suppose that $\mathcal{H}$ is the collection of half-infinite intervals

$$(-\infty, x], \quad x \in \mathbb{R}.$$

Then $\mathcal{H}$ is π -system of $\mathbb{R}$. The λ -system generated by $\mathcal{H}$ contains all the open intervals. Since any open subset of $\mathbb{R}$ is a countable union of open intervals we deduce that $\Lambda(\mathcal{H})$ coincides with the Borel σ -algebra $\mathcal{B}_{\mathbb{R}}$.

If X is a topological space and $\mathcal{T}_X$ is the collection of open subsets, then $\mathcal{T}_X$ is a π -system. $\square$

¹Check this.

Theorem 1.1.7 (Dynkin's $\pi - \lambda$ theorem). *Suppose that $\mathcal{P}$ is a π -system. Then*

$$\Lambda(\mathcal{P}) = \sigma(\mathcal{P}).$$

In other words, any λ -system that contains $\mathcal{P}$, also contains the σ -algebra generated by $\mathcal{P}$.

Proof. Since any σ -algebra is a λ -system we deduce $\Lambda(\mathcal{P}) \subset \sigma(\mathcal{P})$. Thus it suffices to show that

$$\sigma(\mathcal{P}) \subset \Lambda(\mathcal{P}). \quad (1.1.3)$$

Equivalently, it suffices to show that $\Lambda(\mathcal{P})$ is a σ -algebra. This happens if and only if the λ -system $\Lambda(\mathcal{P})$ is also a π -system. Hence it suffices to show that $\Lambda(\mathcal{P})$ is closed under (finite) intersections.

For any subset $A \subset \Omega$ we define

$$\mathcal{L}_A := \{ B \in \mathcal{Z}^\Omega : A \cap B \in \Lambda(\mathcal{P}) \}.$$

It suffices to show that

$$\Lambda(\mathcal{P}) \subset \mathcal{L}_A, \quad \forall A \in \Lambda(\mathcal{P}). \quad (1.1.4)$$

Observe that $\mathcal{L}_A$ is a λ -system if $A \in \Lambda(\mathcal{P})$. Indeed, $\Omega \in \mathcal{L}_A$ since $A \in \Lambda(\mathcal{P})$. The properties (ii) and (iii) in the definition of a λ -system are clearly satisfied since $\Lambda(\mathcal{P})$ is a λ -system. Thus, to prove (1.1.4), it suffices to show that

$$\mathcal{P} \subset \mathcal{L}_A, \quad \forall A \in \Lambda(\mathcal{P}). \quad (1.1.5)$$

Note that since $\mathcal{P}$ is a π -system

$$\mathcal{P} \subset \mathcal{L}_B, \quad \forall B \in \mathcal{P}.$$

In particular, since $\mathcal{L}_B$ is a λ -system, we deduce

$$\Lambda(\mathcal{P}) \subset \mathcal{L}_B, \quad \forall B \in \mathcal{P}.$$

Thus, if $A \in \Lambda(\mathcal{P})$ and $B \in \mathcal{P}$, then $A \cap B \in \Lambda(\mathcal{P})$. In other words, $B \in \mathcal{L}_A, \forall B \in \mathcal{P}, \forall A \in \Lambda(\mathcal{P})$, i.e.,

$$\mathcal{P} \subset \mathcal{L}_A, \quad \forall A \in \Lambda(\mathcal{P}).$$

This proves (1.1.5) and completes the proof of the $\pi - \lambda$ -theorem. $\square$

1.1.2. Measurable maps.

Definition 1.1.8. A map $F : \Omega_1 \rightarrow \Omega_2$ called *measurable* with respect to the σ -algebras $\mathcal{S}_i$ on Ω_i , $i = 1, 2$ or $(\mathcal{S}_1, \mathcal{S}_2)$ -*measurable* if $F^{-1}(\mathcal{S}_2) \subset \mathcal{S}_1$, i.e.,

$$F^{-1}(S_2) \in \mathcal{S}_1, \quad \forall S_2 \in \mathcal{S}_2.$$

Two measurable spaces $(\Omega_i, \mathcal{S}_i), i = 1, 2$, are called *isomorphic* if there exists a bijection $F : \Omega_1 \rightarrow \Omega_2$ such that $F^{-1}(\mathcal{S}_2) = \mathcal{S}_1$ or, equivalently, both F and its inverse F^{-1} are measurable. $\square$

Definition 1.1.9. Suppose that $(\Omega, \mathcal{S})$ is a measurable space. A function $f : \Omega \rightarrow \bar{\mathbb{R}}$ is called *$\mathcal{S}$ -measurable* if, for any Borel subset $B \subset \bar{\mathbb{R}}$ we have $f^{-1}(B) \in \mathcal{S}$. $\square$

Example 1.1.10. (a) The composition of two measurable maps is a measurable map.

(b) A subset $S \subset \Omega$ is $\mathcal{S}$ -measurable if and only if the indicator function $\mathbf{I}_S$ is a measurable function.

(c) If $\mathcal{A}$ is the σ -algebra generated by a finite or countable partition

$$\Omega = \bigsqcup_{i \in I} A_i, \quad I \subset \mathbb{N},$$

then a function $f : \Omega \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ is $\mathcal{A}$ -measurable if and only if it is constant in the chambers A_i of this partition. $\square$

The measurability of a map $F : (\Omega_1, \mathcal{S}_1) \rightarrow (\Omega_2, \mathcal{S}_2)$ imposes infinitely many constraints on F , one constraint for each measurable set $S_2 \in \mathcal{S}_2$. It is very impractical to decide the measurability of such a map since very often $\mathcal{S}_2$ has a very complicated description. The next result is extremely useful in practice since it shows that often the measurability of a map is decided by a lot fewer and more transparent constraints.

Proposition 1.1.11. *Consider a map $F : (\Omega_1, \mathcal{S}_1) \rightarrow (\Omega_2, \mathcal{S}_2)$ between two measurable spaces. Suppose that $\mathcal{C}_2$ is a π -system of Ω_2 such that $\sigma(\mathcal{C}_2) = \mathcal{S}_2$. Then the following statements are equivalent.*

- (i) *The map F is measurable.*
- (ii) $F^{-1}(C) \in \mathcal{S}_1, \forall C \in \mathcal{C}_2$.

Proof. Clearly (i) $\Rightarrow$ (ii). The opposite implication follows from the $\pi - \lambda$ theorem since the set

$$\{ C \in \mathcal{S}_2; F^{-1}(C) \in \mathcal{S}_1 \}$$

is a λ -system containing the π -system $\mathcal{C}_2$ that generates $\mathcal{S}_2$. $\square$

Corollary 1.1.12. *If $F : X \rightarrow Y$ is a continuous map between topological spaces, then it is $(\mathcal{B}_X, \mathcal{B}_Y)$ measurable.*

Proof. Denote by $\mathcal{T}_Y$ the collection of open subsets of Y . Then $\mathcal{T}_Y$ is a π -system and, by definition, it generates $\mathcal{B}_Y$. Since F is continuous, for any $U \in \mathcal{T}_Y$ the set $F^{-1}(U)$ is open in X and thus belongs to $\mathcal{B}_X$. $\square$

Corollary 1.1.13. *Let $(\Omega, \mathcal{S})$ be a measurable space. A function $X : \Omega \rightarrow \mathbb{R}$ is $(\mathcal{S}, \mathcal{B}_{\mathbb{R}})$ -measurable if and only if the sets $X^{-1}((-\infty, x])$ are $\mathcal{S}$ -measurable for any $x \in \mathbb{R}$.*

Proof. It follows from the previous corollary by observing that the collection

$$\{ (-\infty, x]; x \in \mathbb{R} \} \subset \mathcal{B}_{\mathbb{R}}$$

is a π -system and the σ -algebra it generates is $\mathcal{B}_{\mathbb{R}}$. $\square$

Remark 1.1.14. In measure theory and analysis, sigma-algebras lie in the background and rarely come to the forefront. In probability they play a more important role having to do with how they are perceived.

One should think of Ω as the collection of all the possible outcomes of a random experiment. A σ -algebra of Ω can be viewed as the totality of information we can collect using certain measurements about the outcomes $\omega \in \Omega$. Let us explain this vague statement on a simple example.

For example, suppose that the set of possible outcomes is $[0, 1)$, but our measuring devices detect with certainty only the first digit of the decimal expansion of a number in $[0, 1)$. We say that a subset S of $[0, 1)$ is measurable if using our device we can conclude with absolute certainty that an outcome ω belongs or not to S . In this case the only measurable subsets are unions of the intervals $[\frac{k-1}{10}, \frac{k}{10})$, $k = 1, \dots, 10$.

Suppose now we are given a measurable space $(\Omega, \mathcal{S})$ and a function $X : \Omega \rightarrow \mathbb{R}$. Can we measure the value of X at an outcome ω using the same measurements that determine $\mathcal{S}$?

Suppose that we can absolutely confirm about the outcome ω of an experiment is whether $X(\omega) \leq x$ for any given $x \in \mathbb{R}$. In other words, we can detect by measurements the collection of sets

$$\{X \leq x\} := X^{-1}((-\infty, x]), \quad x \in \mathbb{R}.$$

In particular, we can detect whether $X(\omega) > x$, i.e., we can detect the sets $\{X > x\} = \{X \leq x\}^c$. More generally, we can determine the sets

$$\{a < X \leq b\} = \{X > a\} \cap \{X \leq b\}.$$

Indeed, we can do this using two measurements: one measurement to decide if $X \leq a$ and one to decide if $X \leq b$. Moreover, we are allowed to perform countably many measurements. In particular, we can decide if

$$\omega \in \bigcap_{n \in \mathbb{N}} \{x - 1/n < X(\omega) \leq x + 1/n\},$$

or, equivalently, if $X(\omega) = x$.

We say that a set S is X -measurable if given $\omega \in \Omega$ we can decide by doing countably many measurements on X whether $\omega \in S$. If $S_1, \dots, S_n, \dots \subset \Omega$ are known to be X -measurable, then their union is X -measurable. Indeed,

$$\omega \in \bigcup_{n \in \mathbb{N}} S_n \iff \exists n \in \mathbb{N} : \omega \in S_n.$$

Let us observe that the set theoretic conditions imposed on a sigma-algebra have logical/linguistic counterparts. Thus, the statement

$$\omega \in \bigcap_{i \in I} S_i$$

translates into the formula $\forall i \in I, \omega \in S_i$, while the statement

$$\omega \in \bigcup_{i \in I} S_i$$

translates into the formula $\exists i \in I, \omega \in S_i$.

Conversely, statements involving the quantifiers $\exists, \forall$ can be translated into set theoretic statements.

The information we can collect by doing such measurements of the function X is collected into the sigma-algebra $\sigma(X) = X^{-1}(\mathcal{B}_{\mathbb{R}})$ generated by the map $X : \Omega \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$. $\square$

Corollary 1.1.15. *Consider a pair of maps between measurable spaces*

$$F_i : (\Omega, \mathcal{S}) \rightarrow (\Omega_i, \mathcal{S}_i), \quad i = 1, 2.$$

Then the following statements are equivalent.

- (i) *The maps F_i are measurable.*
- (ii) *The map*

$$F_1 \times F_2 : \Omega \rightarrow \Omega_1 \times \Omega_2, \quad \omega \mapsto (F_1(\omega), F_2(\omega))$$

is $(\mathcal{S}, \mathcal{S}_1 \otimes \mathcal{S}_2)$ -measurable.

Proof. (i) $\Rightarrow$ (ii) Observe that if the maps F_1, F_2 are measurable then

$$\begin{aligned} F_1^{-1}(S_1), F_2^{-1}(S_2) &\in \mathcal{S}, \quad \forall S_1 \in \mathcal{S}_1, S_2 \in \mathcal{S}_2 \\ \Rightarrow (F_1 \times F_2)^{-1}(S_1 \times S_2) &= F_1^{-1}(S_1) \cap F_2^{-1}(S_2) \in \mathcal{S}, \quad \forall S_1 \in \mathcal{S}_1, S_2 \in \mathcal{S}_2. \end{aligned}$$

Since the collection $S_1 \times S_2, S_i \in \mathcal{S}_i, i = 1, 2$, is a π -system that, by definition, generates $\mathcal{S}_1 \otimes \mathcal{S}_2$ we see that the last statement is equivalent with the measurability of $F_1 \times F_2$.

(ii) $\Rightarrow$ (i) For $i = 1, 2$ we denote by π the natural projection $\Omega_1 \times \Omega_2 \rightarrow \Omega_i, (\omega_1, \omega_2) \mapsto \omega_i$. The maps π_i are $(\mathcal{S}_1 \otimes \mathcal{S}_2, \mathcal{S}_i)$ measurable and $F_i = \pi_i \circ (F_1 \times F_2)$. $\square$

Definition 1.1.16. For any measurable space $(\Omega, \mathcal{S})$ we denote by $\mathcal{L}^0(\mathcal{S}) = \mathcal{L}^0(\Omega, \mathcal{S})$ the space of $\mathcal{S}$ -measurable random variables, i.e., $(\mathcal{S}, \mathcal{B}_{\bar{\mathbb{R}}})$ -measurable functions $\Omega \rightarrow \bar{\mathbb{R}}$.

The subset of $\mathcal{L}^0(\Omega, \mathcal{S})$ consisting of nonnegative functions is denoted by $\mathcal{L}_+^0(\Omega, \mathcal{S})$, while the subspace of $\mathcal{L}^0(\Omega, \mathcal{S})$ consisting of bounded measurable functions is denoted $\mathcal{L}^\infty(\Omega, \mathcal{S})$. $\square$

Remark 1.1.17. The algebraic operations on $\mathbb{R}$ admit (partial) extensions to $\bar{\mathbb{R}}$.

$$c + \pm\infty = \pm\infty, \infty + \infty = \infty, \quad c \cdot \infty = \infty, \quad \forall c > 0.$$

As we know, there are a few “illegal” operations

$$\infty - \infty, \quad 0 \cdot \infty, \quad \frac{0}{0} \text{ etc.} \quad \square$$

Proposition 1.1.18. Fix a measurable space $(\Omega, \mathcal{S})$. Then the following hold.

(i) For any $X, Y \in \mathcal{L}^0(\Omega, \mathcal{S})$ and any $c \in \mathbb{R}$ we have

$$X + Y, XY, cX \in \mathcal{L}^0(\Omega, \mathcal{S}),$$

whenever these functions are well defined.

(ii) If $(X_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}^0(\Omega, \mathcal{S})$ such that, for any $\omega \in \Omega$ the limit

$$X_\infty(\omega) = \lim_{n \rightarrow \infty} X_n(\omega)$$

exists. Then $X_\infty : \Omega \rightarrow \bar{\mathbb{R}}$ is also $\mathcal{S}$ -measurable.

(iii) If $(X_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}^0(\Omega, \mathcal{S})$. For any $\omega \in \Omega$ we set

$$Y_\infty(\omega) = \inf_{n \in \mathbb{N}} X_n(\omega), \quad Z_\infty(\omega) = \sup_{n \in \mathbb{N}} X_n(\omega).$$

Then $Y_\infty, Z_\infty \in \mathcal{L}^0(\Omega, \mathcal{S})$.

Proof. (i) Denote by $\mathcal{D}$ the subset of $\bar{\mathbb{R}}^2$ consisting of the pairs (x, y) for which $x + y$ is well defined,

$$\mathcal{D} = \bar{\mathbb{R}}^2 \setminus \{(\infty, -\infty), (-\infty, \infty)\}.$$

The set $\mathcal{D}$ is obviously a Borel subset of $\bar{\mathbb{R}}^2$ since it is open. Observe that $X + Y$ is the composition of two measurable maps

$$\Omega \rightarrow \mathcal{D} \subset \bar{\mathbb{R}}^2, \quad \omega \mapsto (X(\omega), Y(\omega)), \quad \mathcal{D} \rightarrow \bar{\mathbb{R}}, \quad (x, y) \mapsto x + y.$$

Above, the first map is measurable according to Corollary 1.1.15 and the second map is Borel measurable since it is continuous. The measurability of XY and cX is established in a similar fashion.

(ii) Observe first that the set $\{X_\infty > -\infty\}$ is measurable because

$$X_\infty(\omega) > -\infty \iff \exists M \in \mathbb{Z}, \exists N \in \mathbb{N}, \forall n > N, X_n(\omega) > M.$$

We will show next that for any $x \in \mathbb{R}$ the set $\{X_\infty(\omega) > x\}$ is $\mathcal{S}$ -measurable. Note that

$$X_\infty(\omega) > x \iff \exists \nu \in \mathbb{N}, \exists N = N(\omega) \in \mathbb{N} : \forall n \geq N : X_n(\omega) > x + 1/\nu.$$

Equivalently

$$\{X_\infty(\omega) > x\} = \bigcup_{\nu \in \mathbb{N}} \bigcup_{N \in \mathbb{N}} \bigcap_{n \geq N} \{X_n > x + 1/\nu\} \in \mathcal{S}.$$

(iii) The proof is very similar to the proof of (ii) so we leave the details to the reader. $\square$

Corollary 1.1.19. *For any measurable function $f \in \mathcal{L}^0(\Omega, \mathcal{S})$, its positive and negative parts,*

$$f^+ := \max(f, 0), \quad f^- := \max(-f, 0)$$

are also measurable.

Proof. The function f^+ is the composition of the continuous function $x^+ = \max(x, 0)$ with f . $\square$

Definition 1.1.20. A function $f \in \mathcal{L}^0(\Omega, \mathcal{S})$ is called *elementary* or *step function* if its range is a *finite* subset of $\mathbb{R}$. We denote by $\text{Elem}(\Omega, \mathcal{S})$ the set of elementary functions. $\square$

More concretely, a function $f : \Omega \rightarrow \mathbb{R}$ is elementary if there exist *finitely many disjoint measurable sets* $A_1, \dots, A_N \in \mathcal{S}$, and constants $c_1, \dots, c_N \in \mathbb{R}$ such that

$$f(\omega) = \sum_{k=1}^N c_k \mathbf{I}_{A_k}(\omega), \quad \forall \omega \in \Omega. \quad (1.1.6)$$

The decomposition (1.1.6) of an elementary function f is not unique. Among the various decompositions there is a canonical one

$$f = \sum_{r \in \mathbb{R}} r \mathbf{I}_{f^{-1}(r)}.$$

The above sum is finite since $f^{-1}(r)$ is empty for all but finitely many r 's.

Let us also observe that $\text{Elem}(\Omega, \mathcal{S})$ is a vector space. Indeed if f_0, f_1 are elementary functions with ranges R_0 and respectively R_1 , then their sum is measurable and its range is contained in $R_0 + R_1$. This is a finite set since R_0, R_1 are finite. Clearly the multiplication of an elementary function by a scalar also produces an elementary function.

Any nonnegative measurable function is the pointwise limit of an increasing sequence of elementary functions. To see this, for each $n \in \mathbb{N}$ we define

$$D_n : [0, \infty) \rightarrow [0, \infty), \quad D_n(r) := \sum_{k=1}^{n2^n} \frac{k-1}{2^n} \mathbf{I}_{[(k-1)2^{-n}, k2^{-n})}(r).$$

Let us observe that if $r \in [0, n]$, then $D_n(r)$ truncates the binary expansion of r after n digits. E.g., if $r \in [0, 1)$ and

$$r = 0.\epsilon_1\epsilon_2\dots\epsilon_n\dots := \sum_{k=1}^{\infty} \frac{\epsilon_k}{2^k}, \quad \epsilon_k \in \{0, 1\},$$

then

$$D_n(r) = 0.\epsilon_1\dots\epsilon_n.$$

This shows that $(D_n)_{n \in \mathbb{N}}$ is a nondecreasing sequence of functions and

$$\lim_{n \rightarrow \infty} D_n(r) = r, \quad \forall r \geq 0.$$

For $f \in \mathcal{L}_+^0(\Omega, \mathcal{S})$ and $n \in \mathbb{N}$ we define $D_n[f] : (\Omega, \mathcal{S}) \rightarrow [0, \infty)$

$$D_n[f](\omega) := D_n(f(\omega)) = \sum_{k=1}^{n2^n} \frac{k-1}{2^n} \mathbf{I}_{[(k-1)2^{-n}, k2^{-n})}(f(\omega)) + n \mathbf{I}_{[n, \infty)}(f(\omega)). \quad (1.1.7)$$

We deduce that the sequence of nonnegative elementary functions $D_n[f]$ converges increasingly to f .

Definition 1.1.21. Let $(\Omega, \mathcal{S})$ be a measurable. A collection $\mathcal{M}$ of $\mathcal{S}$ -measurable functions is called a *monotone class* of $(\Omega, \mathcal{S})$ if it satisfies the following conditions.

- (i) $\mathbf{I}_\Omega \in \mathcal{M}$.
- (ii) If $f, g \in \mathcal{M}$ are *bounded* and $a, b \in \mathbb{R}$, then $af + bg \in \mathcal{M}$.
- (iii) If (f_n) is an increasing sequence of nonnegative random variables in $\mathcal{M}$ with finite pointwise limit f_∞ , then $f_\infty \in \mathcal{M}$.

□

Theorem 1.1.22 (Monotone Class Theorem). *Suppose that $\mathcal{M}$ is a monotone class of the measurable space $(\Omega, \mathcal{S})$ and $\mathcal{C}$ is a π -system that generates $\mathcal{S}$ and such that $\mathbf{I}_C \in \mathcal{M}$, $\forall C \in \mathcal{C}$. Then $\mathcal{M}$ contains $\mathcal{L}^\infty(\Omega, \mathcal{S})$ and all the nonnegative $\mathcal{S}$ -measurable functions.*

Proof. Observe that the collection

$$\mathcal{A} := \{A \in \mathcal{S} : \mathbf{I}_A \in \mathcal{M}\}$$

is a λ -system containing the π -system $\mathcal{C}$ so $\mathcal{A} = \sigma(\mathcal{C}) = \mathcal{S}$, by the $\pi - \lambda$ theorem. Thus $\mathcal{M}$ contains all the elementary functions. Since any nonnegative measurable function is an increasing pointwise limit of elementary functions we deduce that $\mathcal{M}$ contains all the nonnegative measurable functions. Finally, if f is a bounded measurable function, then f^+, f^- are nonnegative and bounded measurable functions so $f^+, f^- \in \mathcal{M}$ and thus

$$f = f^+ - f^- \in \mathcal{M}.$$

□

Definition 1.1.23. The σ -algebra generated by a collection $(X_i)_{i \in I}$ of real-valued functions on a set Ω is

$$\sigma(X_i, i \in I) := \bigvee_{i \in I} X_i^{-1}(\mathcal{B}_{\mathbb{R}}).$$

□

The next result provides an interpretation of the concept of measurability along the lines of Remark 1.1.14.

Theorem 1.1.24 (Dynkin). *Suppose that $F : (\Omega, \mathcal{S}) \rightarrow (\Omega', \mathcal{S}')$ is a measurable map. Let $X : \Omega \rightarrow \mathbb{R}$ be an $\mathcal{S}$ -measurable function. Recall that $\sigma(F) = F^{-1}(\mathcal{S}')$. Then the following are equivalent.*

- (i) *The function X is $(\sigma(F), \mathcal{B}_{\mathbb{R}})$ -measurable.*
- (ii) *There exists an $(\mathcal{S}', \mathcal{B}_{\mathbb{R}})$ -measurable function $X' : \Omega' \rightarrow \mathbb{R}$ such that $X = X' \circ F$.*

Proof. Clearly, (ii) $\Rightarrow$ (i). To prove that (i) $\Rightarrow$ (ii) consider the family $\mathcal{M}$ of $\sigma(F)$ -measurable functions of the form $X' \circ F$, $X' \in \mathcal{L}^0(\Omega', \mathcal{S}')$. We will prove that $\mathcal{M} = \mathcal{L}^0(\Omega, \sigma(F))$. We will achieve using the monotone class theorem.

Step 1. $I_\Omega \in \mathcal{M}$.

Step 2. $\mathcal{M}$ is a vector space. Indeed if $X, Y \in \mathcal{M}$ and $a, b \in \mathbb{R}$, then there exist $\mathcal{S}'$ -measurable functions X', Y' such that

$$X = X' \circ F, \quad Y = Y' \circ F, \quad aX + bY = (aX' + bY') \circ F.$$

Hence $aX + bY \in \mathcal{M}$.

Step 3. $I_A \in \mathcal{M}$, $\forall A \in \sigma(F)$. Indeed, since $A \in \sigma(F)$ there exists $A' \in \mathcal{S}'$ such that

$$A = F^{-1}(A')$$

so $I_A = I_{A'} \circ F$. Hence $\mathcal{M}$ contains all the $\sigma(F)$ -measurable elementary functions.

Step 4. Suppose now that $X \in \mathcal{L}^0(\Omega, \sigma(F))$ is nonnegative. Then there exists an increasing sequence $(X_n)_{n \in \mathbb{N}}$ of $\sigma(F)$ -measurable nonnegative elementary functions that converges pointwise to X . For every $n \in \mathbb{N}$ there exists an $\mathcal{S}$ -measurable elementary function $X'_n : \Omega' \rightarrow \mathbb{R}$ such that

$$X_n(\omega) = X'_n(F(\omega)), \quad \forall \omega \in \Omega$$

Define

$$\Omega'_0 := \{ \omega' \in \Omega'; \text{ the limit } \lim_{n \rightarrow \infty} X'_n(\omega') \text{ exists and it is finite} \}$$

Let us observe that Ω'_0 is $\mathcal{S}'$ -measurable because

$$\omega' \in \Omega'_0 \iff \forall \nu \geq 1, \exists N \geq 1, \forall m, n \geq N : |X'_n(\omega') - X'_m(\omega')| < 1/\nu,$$

i.e.,

$$\Omega'_0 = \bigcap_{\nu \in \mathbb{N}} \bigcup_{N \geq 1} \bigcap_{m, n \geq N} \{ |X'_n(\omega') - X'_m(\omega')| < 1/\nu \}.$$

Clearly, $F(\Omega) \subset \Omega'_0$. For any $\omega' \in \Omega'$ we set

$$X'_\infty(\omega') := \begin{cases} \lim_{n \rightarrow \infty} X'_n(\omega'), & \omega' \in \Omega'_0, \\ 0, & \omega' \in \Omega' \setminus \Omega'_0. \end{cases}$$

Arguing as in the proof of Proposition 1.1.18(ii) we deduce that X'_∞ is $\mathcal{S}'$ -measurable. For any $\omega \in \Omega$ the sequence $X'_n(F(\omega)) = X_n(\omega)$ is increasing and the limit

$$\lim_{n \rightarrow \infty} X'_n(F(\omega))$$

exists and it is finite. Hence

$$X'_\infty(F(\omega)) = X(\omega), \quad \forall \omega \in \Omega.$$

This proves that $\mathcal{M}$ is a monotone class in $\mathcal{L}^0(\Omega, \sigma(F))$ that is also a vector space so it coincides with $\mathcal{L}^0(\Omega, \sigma(F))$. $\square$

Corollary 1.1.25. Suppose that $X_1, \dots, X_n : (\Omega, \mathcal{S}) \rightarrow \mathbb{R}$ are $\mathcal{S}$ -measurable random variables. The function $X : \Omega \rightarrow \mathbb{R}$ is $\sigma(X_1, \dots, X_n)$ -measurable if and only if there exists an $\mathcal{B}_{\mathbb{R}^n}$ -measurable function $u : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$X = u(X_1, \dots, X_n).$$

Proof. Apply the above theorem with $(\Omega', \mathcal{S}') = (\mathbb{R}^n, \mathcal{B}_{\mathbb{R}^n})$ and

$$F(\omega) = (X_1(\omega), \dots, X_n(\omega)).$$

□

Remark 1.1.26. We see that, in its simplest form, Corollary 1.1.25 describes a measure theoretic form of functional dependence. Thus, if in a given experiment we can measure the quantities $X_1, \dots, X_n$ and we know that the information $X \leq c$ can be decided only by measuring the quantities $X_1, \dots, X_n$, then X is in fact a (measurable) function of $X_1, \dots, X_n$. In plain English this sounds tautological. In particular, this justifies the choice of term “measurable”. □

1.2. Measures and integration

1.2.1. Measures. Throughout this section $(\Omega, \mathcal{S})$ will denote a measurable space. Given a function $f : X \rightarrow \mathbb{R}$ we will use the notation $\{f \leq c\}$ to denote the subset $f^{-1}((-\infty, c])$. The sets $\{a \leq f \leq b\}$ etc. are defined in a similar fashion.

Definition 1.2.1. A *measure* on $(\Omega, \mathcal{S})$ is a function $\mu : \mathcal{S} \rightarrow [0, \infty]$, $S \mapsto \mu[S]$ such that the following hold.

- $\mu[\emptyset] = 0$, and
- it is σ -additive, i.e., for any sequence of pairwise disjoint $\mathcal{S}$ -measurable sets $(A_n)_{n \in \mathbb{N}}$ we have

$$\mu \left[\bigcup_{n \in \mathbb{N}} A_n \right] = \sum_{n \geq 1} \mu[A_n]. \quad (1.2.1)$$

The measure is called σ -finite if there exists an increasing sequence of $\mathcal{S}$ -measurable sets

$$A_1 \subset A_2 \subset \dots$$

such that

$$\bigcup_{n \in \mathbb{N}} A_n = \Omega \text{ and } \mu[A_n] < \infty, \quad \forall n \in \mathbb{N}.$$

The measure is called *finite* if $\mu[\Omega] < \infty$. A *probability measure* is a measure $\mathbb{P}$ such that $\mathbb{P}[\Omega] = 1$. We will denote by $\text{Prob}(\Omega, \mathcal{S})$ the set of probability measures on $(\Omega, \mathcal{S})$. □

Remark 1.2.2. The σ -additivity condition (1.2.1) is equivalent to a pair of conditions that are more convenient to verify in concrete situations.

- (i) μ is *finitely additive*, i.e., for any finite collection of $\mathcal{S}$ -measurable sets $A_1, \dots, A_n$ we have

$$\mu \left[\bigcup_{k=1}^n A_k \right] = \sum_{k=1}^n \mu[A_k].$$

- (ii) μ is *increasingly continuous* i.e., for any increasing sequence of $\mathcal{S}$ -measurable sets $A_1 \subset A_2 \subset \dots$

$$\mu \left[\bigcup_{n \in \mathbb{N}} A_n \right] = \lim_{n \rightarrow \infty} \mu[A_n]. \quad (1.2.2)$$

If $\mu[\Omega] < \infty$ and μ is finitely additive, then the increasing continuity condition (ii) is equivalent with the *decreasing continuity* condition, i.e., for any decreasing sequence of $\mathcal{S}$ -measurable sets $B_1 \supset B_2 \supset \dots$

$$\mu \left[\bigcap_{n \in \mathbb{N}} B_n \right] = \lim_{n \rightarrow \infty} \mu[B_n]. \quad (1.2.3)$$

Indeed, the sequence $B_n^c = \Omega \setminus B_n$ is increasing and $\mu[B_n^c] = \mu[\Omega] - \mu[B_n]$. This last equality could be meaningless if $\mu[\Omega] = \infty$ $\square$

Definition 1.2.3. (a) A *measured space* is a triplet $(\Omega, \mathcal{S}, \mu)$, where $(\Omega, \mathcal{S})$ is a measurable space and $\mu : \mathcal{S} \rightarrow [0, \infty]$ is a measure. $\square$

Our next result shows that a *finite* measure is uniquely determined by its restriction to an algebra generating the sigma-algebra where it is defined.

Proposition 1.2.4. Consider a measurable space $(\Omega, \mathcal{S})$ and two finite measures $\mu_1, \mu_2 : \mathcal{S} \rightarrow [0, \infty]$ such that $\mu_1[\Omega] = \mu_2[\Omega] < \infty$, then the collection

$$\mathcal{E} := \{ S \in \mathcal{S}; \mu_1[S] = \mu_2[S] \}$$

is a λ -system. In particular, if $\mu_1[C] = \mu_2[C]$ for any set C that belongs to a π -system $\mathcal{C}$, then μ_1 and μ_2 coincide on the σ -algebra generated by $\mathcal{C}$.

Proof. Clearly $\emptyset, \Omega \in \mathcal{E}$. If $A, B \in \mathcal{E}$ and $A \subset B$, then

$$\mu_1[A] = \mu_2[A] < \infty, \quad \mu_1[B] = \mu_2[B] < \infty$$

so

$$\mu_1[B \setminus A] = \mu_1[B] - \mu_1[A] = \mu_2[B] - \mu_2[A] = \mu_2[B \setminus A],$$

so $B \setminus A \in \mathcal{E}$. The condition (iii) in the Definition 1.1.5 of a λ -system follows from the σ -additivity of the measures μ_1, μ_2 . $\square$

Definition 1.2.5. A *probability space*, or *sample space*, is a measured space $(\Omega, \mathcal{S}, \mathbb{P})$, where $\mathbb{P}$ is a probability measure. In this case we use the following terminology.

- The subsets $S \in \mathcal{S}$ are called the *events* of the sample space.
- An event $S \in \mathcal{S}$ is called *almost sure* (or a.s.) if $\mathbb{P}[S] = 1$. An event S is called *improbable* if $\mathbb{P}[S] = 0$.
- The measurable functions $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \bar{\mathbb{R}}$ are called *random variables*.
- A random variable $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \bar{\mathbb{R}}$ is called a.s. *finite* if

$$\mathbb{P}[|X| < \infty] = 1.$$

- A random variable on $(\Omega, \mathcal{S}, \mathbb{P})$ is called *deterministic* if there exists $c \in \mathbb{R}$ such that $X = c$ a.s..

$\square$

 Traditionally the random variables have capitalized names X, Y, Z etc to distinguish them from deterministic quantities that are indicated in small caps. We will try to adhere to this convention throughout this book

Example 1.2.6. (a) If $(\Omega, \mathcal{S})$ is a measurable space, then for any $\omega_0 \in \Omega$, the *Dirac measure* concentrated at ω_0 is the probability measure

$$\delta_{\omega_0} : \mathcal{S} \rightarrow [0, \infty), \quad \delta_{\omega_0}[S] = \begin{cases} 1, & \omega_0 \in S, \\ 0, & \omega_0 \notin S. \end{cases}$$

(b) Suppose that S is a finite or countable set. A measure on $(S, 2^S)$ is uniquely determined by the function

$$w : S \rightarrow [0, \infty], \quad w(s) = \mu[\{s\}].$$

We say that $\mu[\{s\}]$ is the *mass* of s with respect to μ . The function w is referred to as the *weight function* of the measure. Often, for simplicity, we will write

$$\mu[s] := \mu[\{s\}].$$

The associated measure μ_w is a probability measure if

$$\sum_{s \in S} w(s) = 1.$$

When S is finite and

$$w(s) = \frac{1}{|S|}, \quad \forall s \in S,$$

then the associated probability measure μ_w is called the *uniform probability measure* on the finite set S .

(c) Suppose that $F : (\Omega, \mathcal{S}) \rightarrow (\Omega', \mathcal{S}')$ is a measurable map between measurable spaces. Then any measure μ on Ω induces a measure $F_{\#}\mu$ on Ω' according to the rule

$$F_{\#}\mu[S'] := \mu[F^{-1}(S')].$$

The measure $F_{\#}\mu$ is called the *pushforward of μ via F* .

(d) Fix a set T with two elements, $T = \{0, 1\}$. For any $p \in (0, 1)$ the probability measure $\beta_p : 2^T \rightarrow [0, \infty)$ defined by

$$\beta_p[1] = p, \quad \beta_p[0] = q := 1 - p$$

is called the *Bernoulli distribution* with success probability p . We abbreviate it by $\text{Ber}(p)$.

(e) Given finite or countable sets $\Omega_1, \dots, \Omega_n$, and probability measures $\mu_i : 2^{\Omega_i} \rightarrow [0, 1]$, we obtain a probability measure

$$\mu := \mu_1 \otimes \dots \otimes \mu_n : 2^{\Omega_1 \times \dots \times \Omega_n} \rightarrow [0, 1]$$

by setting

$$\mu[(\omega_1, \dots, \omega_n)] = \mu_1[\omega_1] \cdots \mu_n[\omega_n], \quad \forall (\omega_1, \dots, \omega_n) \in \Omega_1 \times \dots \times \Omega_n.$$

In particular, there exists a probability measure $\beta_p^{\otimes n}$ on $\{0, 1\}^n$.

Note that we have a random variable

$$N : \{0, 1\}^n \rightarrow \mathbb{N}_0, \quad N((\epsilon_1, \dots, \epsilon_n)) = \epsilon_1 + \dots + \epsilon_n, \quad \forall \epsilon_1, \dots, \epsilon_n \in \{0, 1\}.$$

The push-forward $\mathbb{P} = \mathbb{P}_{n,p} := N_{\#}\beta_p^{\otimes n}$ is a probability measure on $\{0, 1, \dots, n\}$ called the *binomial distribution* corresponding to n independent trials with success probability p and failure probability

$q = 1 - p$. It is abbreviated $\text{Bin}(n, p)$. Note that $\text{Bin}(1, p) = \text{Ber}(p)$. For any $k \in \{0, 1, \dots, n\}$ we have

$$\mathbb{P} = \sum_{k=0}^n \mathbb{P}[k] \delta_k,$$

where

$$\begin{aligned} \mathbb{P}[k] &= \beta_p^{\otimes n}[N = k] = \sum_{\epsilon_1 + \dots + \epsilon_n = k} \beta_p^{\otimes n}[(\epsilon_1, \dots, \epsilon_n)] \\ &= \sum_{\epsilon_1 + \dots + \epsilon_n = k} p^k q^{n-k} = \binom{n}{k} p^k q^{n-k}. \end{aligned}$$

(f) The Lebesgue measure λ defines a measure on $\mathcal{B}_{\mathbb{R}}$. For any compact interval $[a, b]$ the *uniform probability measure* on $[a, b]$ is

$$\frac{1}{b-a} \mathbf{I}_{[a,b]} \lambda. \quad \square$$

Definition 1.2.7. Let X be a topological space. As usual $\mathcal{B}_X$ denotes the σ -algebra of Borel subsets of X . A measure on X is called *Borel* if it is defined on $\mathcal{B}_X$. $\square$

The Lebesgue measure on $\mathbb{R}$ is a Borel measure.

Definition 1.2.8. Suppose that $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$. Its distribution is the Borel probability measure $\mathbb{P}_X$ on $\mathbb{R}$ defined by

$$\mathbb{P}_X[B] = \mathbb{P}[X \in B], \quad \forall B \in \mathcal{B}_{\mathbb{R}}.$$

In other words, $\mathbb{P}_X$ is the pushforward of $\mathbb{P}$ by X , $\mathbb{P}_X = X_{\#}\mathbb{P}$. $\square$

Definition 1.2.9. Suppose that μ is a measure on the measurable space $(\Omega, \mathcal{S})$.

(i) A set $N \subset \Omega$ is called *μ -negligible* if there exists a set $S \in \mathcal{S}$ such that

$$N \subset S \text{ and } \mu[S] = 0.$$

We denote by $\mathcal{N}_{\mu}$ the collection of μ -negligible sets.

(ii) The σ -algebra $\mathcal{S}$ is said to be *complete* with respect to μ (or *μ -complete*) if it contains all the μ -negligible subsets.

(iii) The *μ -completion* of $\mathcal{S}$ is the σ -algebra $\mathcal{S}^{\mu} := \sigma(\mathcal{S}, \mathcal{N}_{\mu})$. $\square$

Remark 1.2.10. (a) It may be helpful to think of a sample space $(\Omega, \mathcal{S}, \mathbb{P})$ as the collection of all possible outcomes ω of an experiment with unpredictable results. The observer may not be able to distinguish through measurements all the possible outcomes, but she is able to distinguish some features or properties of various outcomes. An event can be understood as the collection of the all outcomes having an observable or measurable property. The probability $\mathbb{P}$ associates a likelihood of a certain property to be observed at the end of such a random experiment.

Take for example the experiment of flipping n times a coin with 0/1 faces. One natural sample space for this experiment is based on the set $\Omega = \{0, 1\}^n$.

If we assume that the coin is fair, then it is natural to conclude that each outcome $\omega \in \Omega$ is equally likely. Suppose that we can distinguish all the outcomes. In this case

$$\mathcal{S} = 2^\Omega.$$

Since there are 2^n outcomes that are equally likely to occur we obtain a probability measure $\mathbb{P}$ given by

$$\mathbb{P}[S] = \frac{|S|}{2^n}, \quad \forall S \in \mathcal{S}.$$

A random variable on a sample space is a numerical attribute X that we can assign to each outcome ω of a random experiment with the following feature: for any $c \in \mathbb{R}$ the property $X(\omega) \leq c$ is observable, i.e., the set $X^{-1}((-\infty, c])$ belongs to the collection $\mathcal{S}$ of observable properties. For example, in the situation of n fair coin tosses, the number N of 1's observed at the end of n tosses is a random variable.

(b) Often one speaks of *sampling a probability distribution* on $\mathbb{R}$. Modern computer systems can sample many distributions. More concretely, we say that a probability measure μ on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ can be sampled by a computer system if that computer can produce a random² experiment whose outcome is a random number X so that, when we run the experiment a *large* number of times n , it generates numbers $x_1, \dots, x_n$ and, for any $c \in \mathbb{R}$, the fraction of these numbers that is $\leq c$ is very close to $\mu[(-\infty, c]]$.

When we speak of sampling a random variable X , we really mean sampling its probability distribution $\mathbb{P}_X$. $\square$

Clearly $\mathcal{S}^\mu$ is the smallest μ -complete σ -algebra containing $\mathcal{S}$. The proof of the following result can be safely left to the reader.

Proposition 1.2.11. *Suppose that μ is a measure on the σ -algebra $\mathcal{S} \subset 2^\Omega$.*

(i) *The completion $\mathcal{S}^\mu$ has the alternate description*

$$\mathcal{S}^\mu = \{ S \cup N; \ S \in \mathcal{S}, \ N \in \mathcal{N}_\mu \} \subset 2^\Omega.$$

(ii) *The measure μ admits a unique extension to a probability measure $\bar{\mu} : \mathcal{S}^\mu \rightarrow [0, \infty)$. More precisely*

$$\forall S \in \mathcal{S}, \ N \in \mathcal{N}_\mu \quad \bar{\mu}[S \cup N] = \mu[S].$$

$\square$

Definition 1.2.12. A set $S \subset \mathbb{R}$ is called *Lebesgue measurable* if it belongs to the λ -completion of $\mathcal{B}_{\mathbb{R}}$. $\square$

The most versatile method of constructing measures is *Carathéodory Extension Theorem*. We need to introduce the appropriate concepts.

Definition 1.2.13. Fix a set Ω and an algebra $\mathcal{F} \subset 2^\Omega$

(i) A function $\mu : \mathcal{F} \rightarrow [0, \infty]$ is called a *premeasure* if it satisfies the following conditions.

(a) $\mu[\emptyset] = 0$

²The precise term is pseudo-random since one cannot algorithmically simulate randomness.

- (b) μ is *finitely additive*, i.e., for any finite collection of disjoint sets $A_1, \dots, A_n \in \mathcal{F}$ we have

$$\mu \left[\bigcup_{k=1}^n A_k \right] = \sum_{k=1}^n \mu[A_k].$$

- (c) μ is *conditional countably additive*, i.e., for any sequence $(A_n)_{n \in \mathbb{N}}$ of disjoint sets in $\mathcal{F}$ whose union is a set $A \in \mathcal{F}$ we have

$$\mu[A] = \sum_{n \geq 1} \mu[A_n].$$

- (ii) The premeasure μ is called *σ -finite* if there exists a sequence of sets $(\Omega_n)_{n \in \mathbb{N}}$ in $\mathcal{F}$ such that

$$\Omega = \bigcup_{n \in \mathbb{N}} \Omega_n, \quad \mu[\Omega_n] < \infty, \quad \forall n \in \mathbb{N}.$$

□

Remark 1.2.14. Suppose that $\mu : \mathcal{F} \rightarrow [0, \infty)$ is a finitely additive function on an algebra $\mathcal{F}$ of subsets of a set Ω such that $\mu[\Omega] < \infty$. Then μ is a premeasure if and only if, for any decreasing sequence in $\mathcal{F}$,

$$F_1 \supset F_2 \supset \dots \quad \text{such that} \quad \bigcap_{n \geq 1} F_n = \emptyset,$$

we have

$$\lim_{n \rightarrow \infty} \mu[F_n] = 0.$$

Indeed, if $(A_n)_{n \in \mathbb{N}}$ is a sequence of disjoint sets of $\mathcal{F}$ whose union A is also a set in $\mathcal{F}$, then the sequence

$$F_n = A \setminus \bigcup_{k=1}^n A_k$$

is a decreasing sequence in $\mathcal{F}$ with empty intersection and

$$\mu[F_n] = \mu[A] - \sum_{k=1}^n \mu[A_k].$$

□

The (conditional) countable additivity condition in the definition of a premeasure could be challenging to verify. The next result gives a simple sufficient condition guaranteeing this countable additivity.

Theorem 1.2.15 (Alexandrov). *Suppose that K is a compact topological space, $\mathcal{F}$ is an algebra of subsets of K and $\mu : \mathcal{F} \rightarrow [0, 1]$ is a finitely additive function satisfying the following regularity property: for any $F \in \mathcal{F}$ and any $\varepsilon > 0$ there exists a set $F_- \in \mathcal{F}$ such that*

$$\text{cl}(F_-) \subset F, \quad \mu[F \setminus F_-] < \varepsilon.$$

Then μ is a premeasure.

□

Proof. Let us introduce a convenient terminology. For $\varepsilon > 0$ we define an ε -squeeze of a set $F \in \mathcal{F}$ to be a set $G \in \mathcal{F}$ such that $\text{cl}(G) \subset F$ and $\mu[F \setminus G] < \varepsilon$.

Lemma 1.2.16. *Suppose that $F_1, F_2 \in \mathcal{F}$, $F_2 \subset F_1$, and for $i = 1, 2$, G_i is an ε_i -squeeze of F_i . Then $G_1 \cap G_2$ is an $(\varepsilon_1 + \varepsilon_2)$ -squeeze of F_2 .*

Proof of Lemma 1.2.16. Clearly

$$\mathbf{cl}(G_1 \cap G_2) \subset \mathbf{cl}(G_2) \subset F_2,$$

$$F_2 \setminus (G_1 \cap G_2) = F_2 \cap (G_1 \cap G_2)^c = F_2 \cap (G_1^c \cup G_2^c) = (F_2 \cap G_1^c) \cup (F_2 \cap G_2^c),$$

and

$$\begin{aligned} \mu[F_2 \setminus (G_1 \cap G_2)] &= \mu[(F_2 \setminus G_1) \cup (F_2 \setminus G_2)] \\ &\leq \mu[F_2 \setminus G_1] + \mu[F_2 \setminus G_2] \leq \mu[F_1 \setminus G_1] + \mu[F_2 \setminus G_2] \leq \varepsilon_1 + \varepsilon_2. \end{aligned}$$

□

To prove that μ is a pre-measure it suffices to show that if $(F_n)_{n \in \mathbb{N}}$ is a decreasing sequence in $\mathcal{F}$ with empty intersection, then

$$\lim_{n \rightarrow \infty} \mu[F_n] = 0.$$

Fix $\varepsilon > 0$. For $n \in \mathbb{N}$, fix an $\frac{\varepsilon}{2^n}$ -squeeze G_n of F_n . Define

$$H_n := \bigcap_{k=1}^n G_k, \quad \varepsilon_n := \sum_{k=1}^n \frac{\varepsilon}{2^k} = \varepsilon(1 - 2^{-n}).$$

Applying Lemma 1.2.16 iteratively we deduce that H_n is an ε_n -squeeze of F_n . By construction the sequence H_n is decreasing and thus the sequence of closures $\mathbf{cl}(H_n)$ is decreasing as well. Note that

$$\bigcap_n \mathbf{cl}(H_n) \subset \bigcap_n F_n = \emptyset.$$

Since K is compact we deduce that there exists $N = N(\varepsilon) \in \mathbb{N}$ such that $\mathbf{cl}(H_N) = \emptyset$. Hence $H_N = \emptyset$ and since H_N is an ε_N -squeeze we deduce that, $\forall n \geq N$

$$\mu[F_n] \leq \mu[F_N] = \mu[F_N \setminus H_N] \leq \varepsilon_N < \varepsilon.$$

□

I refer to Gian-Carlo Rota [182, p.36-37] for a nice anecdote about Alexandrov's theorem. For a generalization of the above result I refer to [156, Prop.I.6.2].

The importance of pre-measures is justified by the next central existence result. For a proof I refer to [6, Sec. 1.3], [64, Chap. 3] or [118, Thm.1.53, 1.65].

Theorem 1.2.17 (Carathéodory Extension Theorem). *Suppose that $\mathcal{F}$ is an algebra of subsets of Ω and $\mu : \mathcal{F} \rightarrow [0, \infty]$ is a σ -finite premeasure on $\mathcal{F}$. Then the following hold.*

- (i) *The premeasure μ admits a unique extension to a measure $\tilde{\mu} : \sigma(\mathcal{F}) \rightarrow [0, \infty]$.*
- (ii) *For any $A \in \sigma(\mathcal{F})$ and any $\varepsilon > 0$ there exist mutually disjoint sets $A_1, \dots, A_m \in \mathcal{F}$ and $B_1, \dots, B_n \in \mathcal{F}$ such that*

$$A \subset \bigcup_{j=1}^m A_j, \quad \tilde{\mu} \left[\bigcup_{j=1}^m A_j \setminus A \right] < \varepsilon,$$

and

$$\tilde{\mu} \left[A \Delta \bigcup_{k=1}^n B_k \right] < \varepsilon.$$

□

Example 1.2.18. For any interval $I \subset \mathbb{R}$ we denote by $\lambda[I]$ its length. Let $\mathcal{F}$ denote the collection of subsets of $\mathbb{R}$ that are union of intervals of the type $(a, b]$, $-\infty \leq a < b < \infty$. This is an algebra of sets. Any F can be written in a (non)unique way as a union

$$F = \bigcup_{j=1}^n (a_i, b_i], \quad a_i < b_i \leq a_{i+1} < b_{i+1}, \quad \forall i = 1, \dots, n-1.$$

While this decomposition is not unique the sum

$$\lambda[F] = \sum_{i=1}^n \lambda[(a_i, b_i)],$$

depends only on F and not on the decomposition. It is not very hard to show that the correspondence

$$\mathcal{F} \ni F \mapsto \lambda[F] \in [0, \infty]$$

is finitely additive. The fact that λ is a premeasure, i.e., it is (conditionally) sigma-additive, is much more subtle, and it is ultimately rooted in the compactness of the closed and bounded intervals of $\mathbb{R}$. More precisely if we denote by $\mathcal{F}_n$ the trace of $\mathcal{F}$ to $[-n, n]$ and by λ_n the restriction of λ to $\mathcal{F}_n$, then Alexandrov's Theorem 1.2.15 implies that λ_n is a premeasure for any $n \in \mathbb{N}$. A simple argument then implies that λ itself is a premeasure. For details we refer to [6, Sec. 1.4] or [64, Chap. 3]. The resulting measure on $\mathcal{B}_{\mathbb{R}}$ is called the *Lebesgue measure* on $\mathbb{R}$ and we continue to denote it by λ . □

Definition 1.2.19. A *distribution function* is a right-continuous nondecreasing function

$$F : \bar{\mathbb{R}} \rightarrow [0, 1]$$

such that $F(-\infty) = 0$ and $F(\infty) = 1$. □

Example 1.2.20. Suppose that X is a random variable defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. The function

$$F_X : \mathbb{R} \rightarrow [0, 1], \quad F_X(x) = \mathbb{P}[X \leq x]$$

is a distribution function called the *cumulative distribution function* or *cdf* of the random variable X . □

Example 1.2.21 (Lebesgue-Stieltjes measures). Suppose that $F : \bar{\mathbb{R}} \rightarrow [0, 1]$ is a *distribution function*. Then there exists a unique Borel probability measure $\mu = \mu_F$ on $\mathcal{B}_{\mathbb{R}}$ such that

$$\mu[(x, y)] = F(y) - F(x), \quad \forall x \leq y \in \mathbb{R}. \quad (1.2.4)$$

The uniqueness follows from the fact the collection of intervals $(-\infty, x]$ is a π -system that generates the Borel algebra of $\mathbb{R}$. The existence follows from Caratheodory's extension theorem; see [6, Sec. 1.4] or [64, Chap 3.]. Below we will describe another existence proof that relies only the existence of the usual Lebesgue measure.

The above measure μ_F is called the *Stieltjes probability measure* associated to the distribution function F . Its extension to the completion $\mathcal{B}_{\mathbb{R}}^{\mu}$ is called the Lebesgue-Stieltjes measure associated to the distribution function F .

Conversely, if μ is a Borel probability, measure on $\mathbb{R}$, then μ is the Stieltjes measure associated to its *cumulative distribution function* (cdf) $F : \mathbb{R} \rightarrow [0, 1]$, $F(x) = \mu[(-\infty, x]]$. $\square$

Example 1.2.22 (Quantiles). Here is an alternate description of this measure based on a construction frequently used in statistics. Suppose that $F : \mathbb{R} \rightarrow [0, 1]$ is a cumulative distribution function. The *quantile function* of F is a generalized inverse of the nondecreasing function F . Here is a geometric description of Q .




Figure 1.1. Visualising a cdf and its quantile.

The non-decreasing function F has at most countable many discontinuities, all of jump type. Graph F in the xy plane and the fill in the gaps at its discontinuities by vertical segments; see Figure 1.1. The result is the completed graph of F . It is a "continuous" curve in the plane that may contain vertical segments. Given $p \in (0, 1)$, the horizontal line $y = p$ intersects this curve at a point or along a closed, horizontal, possibly degenerate segment. The quantile $Q(p)$ is the leftmost/smallest abscissa of a point on this intersection. For example, for F as in Figure 1.1 we have $Q(0.3) = 1$.

Formally

$$\begin{aligned} Q : [0, 1] &\rightarrow \mathbb{R}, \quad Q(p) := \inf \{x : p \leq F(x)\} \\ &= \inf F^{-1}([p, 1]). \end{aligned} \tag{1.2.5}$$

Since F is right-continuous the above definition is equivalent to

$$F^{-1}([p, 1]) = [Q(p), \infty].$$

Suppose that x_0 is a point of discontinuity of F and we set

$$p_0^- := \lim_{x \nearrow x_0} F(x) < F(x_0) =: p_0.$$

Note that $Q(p_0) = x_0$ and if $p \in (p_0^-, p_0]$, then $Q(p) = x_0$.

Note that for any $x \in \mathbb{R}$ we have

$$0 \leq y \leq F(x) \iff Q(y) \leq x, \tag{1.2.6}$$

$$Q^{-1}([-\infty, x]) = [0, F(x)]. \tag{1.2.7}$$

Indeed, $\ell \in Q^{-1}([-\infty, x])$ if and only if $Q(\ell) \leq x$, i.e., $\ell \leq F(x)$. In particular,

$$Q^{-1}((x, y]) = (F(x), F(y)], \quad \forall -\infty \leq x \leq y \leq \infty.$$

The quantile is *left* continuous. Indeed, let $p_n \nearrow p_0$. We will show that

$$\lim_n Q(p_n) = Q(p_0).$$

Note that $\lim_n Q(p_n) \leq Q(p_0)$ since Q is nondecreasing. To prove that we have equality we argue by contradiction. Set $x_n := Q(p_n)$, $x_0 = Q(p_0)$. Suppose

$$\lim_n x_n = x_\infty < x_0 = \inf \{ x; F(x) \geq p_0 \}.$$

From the definition of inf as the *greatest* lower bound we deduce that there exists $x_* \in (x_\infty, x_0)$ such that $F(x_*) < p_0$. Thus $F(x_n) \leq F(x_*)$. Since $p_n \nearrow p_0$ we deduce $p_n > F(x_*)$ for all n sufficiently large. This implies

$$x_* \notin \{ x; F(x) \geq p_n \} = [Q(p_n), \infty)$$

i.e., $x_n = Q(p_n) > x_*$, for all n sufficiently large. This contradicts the fact that $x_n \rightarrow x_\infty < x_*$.

If $\lambda_{[0,1]}$ denotes the Lebesgue measure³ on $[0, 1]$, then

$$Q_{\#} \lambda_{[0,1]}[(x, y)] = \lambda[Q^{-1}((x, y))] = F(y) - F(x).$$

Hence the pushforward measure $Q_{\#} \lambda_{[0,1]}$ satisfies (1.2.4) since it coincides with μ_F on the π -system consisting of the intervals of the form $(a, b]$ it coincides with μ_F on the sigma-algebra of Borel sets.

When F is the cumulative distribution function of a random variable, the associated quantile function is called the quantile of the random variable X and it is denoted by Q_X . $\square$

Definition 1.2.23 (Median). Suppose that X is a random variable, $F : \mathbb{R} \rightarrow [0, 1]$ is its cumulative distribution function, and $Q : [0, 1] \rightarrow \mathbb{R}$ is its quantile function. The interval $[Q_F(\frac{1}{2}), Q_F(\frac{1}{2}+0)]$ is called the *median interval* of X . A median of X is a point in the median interval. In other words, m is a median if

$$\mathbb{P}[X \leq m] \geq 1/2, \quad \mathbb{P}[X \geq m] \geq \frac{1}{2}.$$

$\square$

Let us observe that if $[m_-(X), m_+(X)]$ is the median interval of X , then

$$m_-(X) = \sup \{ x; \mathbb{P}[X \leq x] < \frac{1}{2} \}, \quad m_+(X) = \inf \{ x; \mathbb{P}[X \leq x] > \frac{1}{2} \}. \quad (1.2.8)$$

1.2.2. Independence and conditional probability. The next concepts are purely probabilistic in nature. They have no natural counterpart in the traditional measure theory.

Definition 1.2.24. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a sample space.

- (i) The events $A_1, A_2, \dots, A_n \in \mathcal{S}$ are called *independent* if, for any nonempty subset $\{i_1, \dots, i_k\} \subset \{1, \dots, n\}$ we have

$$\mathbb{P}[A_{i_1} \cap \dots \cap A_{i_k}] = \mathbb{P}[A_{i_1}] \cdots \mathbb{P}[A_{i_k}].$$

- (ii) The families of events $\mathcal{A}_1, \dots, \mathcal{A}_n \subset \mathcal{S}$ are called *independent* if for any $A_i \in \mathcal{A}_i$, $i = 1, \dots, n$, the events $A_1, \dots, A_n$ are independent.

- (iii) The (possibly infinite) collection of families of events $(\mathcal{A}_i)_{i \in I}$ is called *independent* if for any $i_1, \dots, i_n \in I$ the finite collection $\mathcal{A}_{i_1}, \dots, \mathcal{A}_{i_n}$ is independent.

- (iv) An *independency* is an independent collection $(\mathcal{S}_i)_{i \in I}$ of sigma-subalgebras of $\mathcal{S}$.

³The proof of the existence of the Lebesgue measure is based on Caratheodory's extension theorem.

- (v) A random variable X is independent of a sigma-subalgebra $\mathcal{F} \subset \mathcal{S}$ if the sigma subalgebras $\sigma(X)$ and $\mathcal{F}$ are independent
- (vi) The collection of random variables $X_i \in \mathcal{L}^0(\Omega, \mathcal{S})$, $i \in I$, is called *independent* if the collection of σ -algebras $(\sigma(X_i))_{i \in I}$ is independent.

□

📖 We will use the notation $X \perp\!\!\!\perp Y$ to indicate that the random variables X, Y are independent.

Remark 1.2.25. (a) We want to emphasize that the independence condition is sensitive to the choice of probability measure involved in this definition.

(b) It is possible that $n + 1$ events be dependent although any n of them are independent. Here is one such instance, [201, Ex. 3.5]. Suppose we flip a fair coin n times. In this case a natural sample space is

$$\Omega = \mathbb{2}^n = \{0, 1\}^n,$$

with the uniform probability measure. (Above, 1= Heads.) For $k = 1, \dots, n$ we denote by k the event “Heads at the k -th flip”, i.e.,

$$E_k = \{ \omega = (\omega_1, \dots, \omega_n) \in \Omega; \omega_k = 1 \}.$$

Denote by E_0 the event “the number of heads in these n flips is even”, i.e.,

$$E_0 = \{ \omega \in \Omega; \omega_1 + \dots + \omega_n \in 2\mathbb{Z} \}$$

Clearly

$$\mathbb{P}[E_k] = \frac{1}{2}, \quad \forall k = 1, \dots, n.$$

Since the probability of flipping an even number of Heads is equal to the probability of flipping an odd number of Heads, we deduce that

$$\mathbb{P}[E_0] = \frac{1}{2}.$$

For any subset $I \subset \{0, 1, \dots, n\}$ we set

$$E_I := \bigcap_{i \in I} E_i.$$

The events $E_1, \dots, E_n$ are independent. Observe that for any subset $I \subset \mathbb{I}_n$, $|I| = k < n$, we have

$$\begin{aligned} \mathbb{P}[E_0 \cap E_I] &= \mathbb{P}\left[\left\{ \omega \in \Omega; \omega_i = 1 \quad \forall i \in I, \sum_{j \notin I} \omega_j \equiv |I| \pmod{2} \right\} \right] \\ &= \underbrace{\mathbb{P}\left[\left\{ \omega \in \Omega; \omega_i = 1 \quad \forall i \in I \right\} \right]}_{\frac{1}{2^k}} \cdot \underbrace{\mathbb{P}\left[\left\{ \omega \in \Omega; \sum_{j \notin I} \omega_j \equiv |I| \pmod{2} \right\} \right]}_{\frac{1}{2}} \\ &= \frac{1}{2^{k+1}} = \mathbb{P}[E_0] \cdot \prod_{i \in I} \mathbb{P}[E_i]. \end{aligned}$$

Thus, any n of the events $E_0, E_1, \dots, E_n$ are independent. Finally, note that

$$\prod_{i=0}^n \mathbb{P}[E_i] = \frac{1}{2^{n+1}} \quad \text{and} \quad \mathbb{P}[E_0 \cap E_1 \cap \dots \cap E_n] = \begin{cases} 0, & n \text{ odd,} \\ \frac{1}{2^n}, & n \text{ even.} \end{cases}$$

This shows the events $E_0, E_1, \dots, E_n$ are dependent.

(c) If Ω is contained in each of the families of events $\mathcal{A}_1, \dots, \mathcal{A}_n$, then these families are independent if and only if

$$\mathbb{P}[A_1 \cap \dots \cap A_n] = \mathbb{P}[A_1] \cdots \mathbb{P}[A_n], \quad \forall A_k \in \mathcal{A}_k, \quad k = 1, \dots, n. \quad \square$$

Proposition 1.2.26. *Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a sample space and that $\mathcal{P}_1, \dots, \mathcal{P}_n \subset \mathcal{S}$ are π -systems each containing Ω . The following statements are equivalent*

- (i) *The families $\mathcal{P}_1, \dots, \mathcal{P}_n$ are independent*
- (ii) *The collection of σ -algebras $\sigma(\mathcal{P}_1), \dots, \sigma(\mathcal{P}_n)$ is independent.*

Proof. Clearly it suffices to prove only (i) $\Rightarrow$ (ii). Fix $S_i \in \mathcal{P}_i, i = 2, \dots, n$. Let

$$\mathcal{J} := \{S \in \mathcal{S} : \mathbb{P}[S \cap S_2 \cap \dots \cap S_n] = \mathbb{P}[S] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n]\}.$$

Note that $\mathcal{P}_1 \subset \mathcal{J}$. Next let us observe that $\mathcal{J}$ is a λ -system. Indeed if $A, B \in \mathcal{J}$ and $A \subset B$ then

$$\begin{aligned} \mathbb{P}[(B \setminus A) \cap S_2 \cap \dots \cap S_n] &= \mathbb{P}[(B \cap S_2 \cap \dots \cap S_n) \setminus (A \cap S_2 \cap \dots \cap S_n)] \\ &= \mathbb{P}[B] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n] - \mathbb{P}[A] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n] = \mathbb{P}[B \setminus A] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n]. \end{aligned}$$

If $A_1 \subset A_2 \subset \dots \subset A_\nu \subset \dots$ is an increasing sequence of events in $\mathcal{J}$ and

$$A = \lim_{\nu \rightarrow \infty} A_\nu = \bigcup_{\nu \geq 1} A_\nu$$

then

$$\begin{aligned} \mathbb{P}[A \cap S_2 \cap \dots \cap S_n] &= \lim_{\nu \rightarrow \infty} \mathbb{P}[A_\nu \cap S_2 \cap \dots \cap S_n] \\ &= \lim_{\nu \rightarrow \infty} \mathbb{P}[A_\nu] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n] = \mathbb{P}[A] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n]. \end{aligned}$$

The $\pi - \lambda$ theorem implies that $\sigma(\mathcal{P}_1) \subset \mathcal{J}$ so that

$$\mathbb{P}[A_1 \cap S_2 \cap \dots \cap S_n] = \mathbb{P}[A_1] \mathbb{P}[S_2] \cdots \mathbb{P}[S_n],$$

for all $A_1 \in \sigma(\mathcal{P}_1), S_i \in \mathcal{P}_i, i = 2, \dots, n$. Repeating the above argument we deduce

$$\mathbb{P}[A_1 \cap A_2 \cap \dots \cap A_n] = \mathbb{P}[A_1] \mathbb{P}[A_2] \cdots \mathbb{P}[A_n], \quad \forall A_k \in \sigma(\mathcal{P}_k), \quad k = 1, \dots, n.$$

Remark 1.2.25 shows that the σ -algebras $\sigma(\mathcal{P}_1), \dots, \sigma(\mathcal{P}_n)$ are independent. $\square$

Corollary 1.2.27. *Consider the random variables $X_1, \dots, X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$. The following statements are equivalent.*

- (i) *The random variables $X_1, \dots, X_n$ are independent.*
- (ii) *For any $x_1, \dots, x_n \in \mathbb{R}$*

$$\mathbb{P}[X_1 \leq x_1, \dots, X_n \leq x_n] = \mathbb{P}[X_1 \leq x_1] \cdots \mathbb{P}[X_n \leq x_n]$$

Proof. It follows from Proposition 1.2.26 applied to the π -systems

$$\mathcal{P}_k := \left\{ \{X_k \leq x_k\} : x_k \in (-\infty, \infty] \right\}, \quad k = 1, \dots, n.$$

$\square$

Corollary 1.2.28 (Partition of independencies). *Suppose that $(\mathcal{S}_i)_{i \in I}$ is an independency of $(\Omega, \mathcal{S}, \mathbb{P})$. For any partition $(I_\alpha)_{\alpha \in A}$ of I we set*

$$\mathcal{F}_\alpha := \bigvee_{i \in I_\alpha} \mathcal{S}_i, \quad \alpha \in A.$$

Then the collection $(\mathcal{F}_\alpha)_{\alpha \in A}$ is also an independency.

Proof. Denote by $\mathcal{C}_\alpha$ the π -system obtained by taking intersections of finitely many events from $\bigcup_{i \in I_\alpha} \mathcal{S}_i$. Note that

$$\bigcup_{i \in I_\alpha} \mathcal{F}_i \subset \Lambda(\mathcal{C}_\alpha)$$

The $\pi - \lambda$ theorem implies

$$\mathcal{F}_\alpha = \sigma(\mathcal{C}_\alpha), \quad \forall \alpha \in A.$$

The family $(\mathcal{C}_\alpha)_{\alpha \in A}$ is clearly independent. The conclusion now follows from Proposition 1.2.26. $\square$

Corollary 1.2.29. *Suppose that the random variables $X_1, \dots, X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ are independent. Then for any $1 < k < n$ and any Borel measurable functions $f : \mathbb{R}^k \rightarrow \mathbb{R}$, $g : \mathbb{R}^{n-k} \rightarrow \mathbb{R}$ the random variables*

$$f(X_1, \dots, X_k), \quad g(X_{k+1}, \dots, X_n)$$

are independent. $\square$

Definition 1.2.30 (Tail algebra). Consider a sequence $(\mathcal{S}_n)_{n \in \mathbb{N}}$ of sub- σ -algebras of $(\Omega, \mathcal{S}, \mathbb{P})$. The *tail algebra* of this sequence is σ -algebra

$$\mathcal{T} = \mathcal{T}(\mathcal{S}_n) := \bigcap_{m \in \mathbb{N}} \mathcal{T}_m, \quad \mathcal{T}_m := \bigvee_{n > m} \mathcal{S}_n. \quad (1.2.9)$$

The events in $\mathcal{T}$ are called *tail events*. $\square$

Remark 1.2.31. (a) An event S is a tail event of the sequence $(\mathcal{S}_n)_{n \in \mathbb{N}}$ if

$$\forall m \in \mathbb{N}; S \in \bigvee_{n > m} \mathcal{S}_n.$$

The sequence of σ -algebras $(\mathcal{S}_n)_{n \in \mathbb{N}}$ can be viewed as an information stream. The tail events are described by a stream of information and are characterized by the fact that their occurrence is unaffected by information at finitely moments of time in the stream.

(b) To a sequence of random variables $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ we associate the sequence of σ -algebras $\mathcal{S}_n = \sigma(X_n)$ and the event $C :=$ "the sequence $(X_n(\omega))_{n \geq 1}$ converges" is a tail event since it belongs to $\mathcal{T}_m$ for any m . Indeed, for any m we have

$$C := \left\{ \omega \in \Omega; \quad \forall \nu \in \mathbb{N}, \exists N > m, \quad \forall k_1, k_2 > N \mid X_{k_1}(\omega) - X_{k_2}(\omega) < \frac{1}{\nu} \right\}$$

$$\underbrace{\bigcap_{\nu} \bigcup_{N > m} \bigcap_{k_1, k_2 > N} \left\{ |X_{k_1}(\omega) - X_{k_2}(\omega)| < \frac{1}{\nu} \right\}}_{=: C_{m, \nu}}.$$

Next, observe that for $k_1, k_2 > N > m$ and $r > 0$ the event

$$\left\{ |X_{k_1}(\omega) - X_{k_2}(\omega)| < r \right\}$$

is $\mathcal{T}_m$ -measurable since X_{k_1} and X_{k_2} are $\mathcal{T}_m$ -measurable and so is their difference. Hence $C_{M,\nu} \in \mathcal{T}_m$ and thus

$$C = \bigcap_{\nu} C_{m,\nu} \in \mathcal{T}_m, \quad \forall m.$$

Hence $C \in \mathcal{T}$. □

Theorem 1.2.32 (Kolmogorov's zero-one law). *If A is a tail event of the independency $(\mathcal{S}_n)_{n \in \mathbb{N}}$, then $\mathbb{P}[A] = 0$ or $\mathbb{P}[A] = 1$.*

Proof. Let $\mathcal{T}_m$ as in (1.2.9). According to the principle of partition of independencies the collection $\mathcal{S}_1, \dots, \mathcal{S}_m, \mathcal{T}_m$ is an independency and, since $\mathcal{T} \subset \mathcal{T}_m$, the collection $\mathcal{S}_1, \dots, \mathcal{S}_m, \mathcal{T}$ is also an independency, $\forall m \in \mathbb{N}$. We deduce that the σ -algebras

$$\mathcal{T}, \mathcal{S}_1, \mathcal{S}_2, \dots$$

are independent. The principle of partition of independencies implies so $\{\mathcal{T}, \mathcal{T}_0\}$ is an independency. Hence, for any $A \in \mathcal{T}$, and any $B \in \mathcal{T}_0$, we have

$$\mathbb{P}[A \cap B] = \mathbb{P}[A]\mathbb{P}[B].$$

If above we choose $B = A \in \mathcal{T} \subset \mathcal{T}_0$ we deduce

$$\mathbb{P}[A] = \mathbb{P}[A]^2, \quad \forall A \in \mathcal{T} \Rightarrow \mathbb{P}[A] \in \{0, 1\}, \quad \forall A \in \mathcal{T}.$$

□

Definition 1.2.33. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space. A *zero-one event* is an event $S \in \mathcal{S}$ such that $\mathbb{P}[S] \in \{0, 1\}$. A *zero-one algebra* is a sigma-subalgebra $\mathcal{F} \subset \mathcal{S}$ consisting of zero-one events. □

Corollary 1.2.34. *Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent random variables on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Then the series*

$$\sum_{n \in \mathbb{N}} X_n$$

is either almost surely convergent, or almost surely divergent. In other words, the almost sure convergence is a zero-one event. □

Definition 1.2.35. Suppose that A, B are events in the sample space $(\Omega, \mathbb{P}, \mathcal{S})$ such that $\mathbb{P}[B] \neq 0$. The *conditional probability of A given B* is the number

$$\mathbb{P}[A|B] := \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]}.$$

□

Note that we have the useful *product formula*

$$\mathbb{P}[A \cap B] = \mathbb{P}[A|B]\mathbb{P}[B]. \tag{1.2.10}$$

In particular, we deduce that A, B are independent if and only if $\mathbb{P}[A] = \mathbb{P}[A|B]$. Note that the map

$$\mathbb{P}[-|B] : \mathcal{S} \rightarrow [0, 1], \quad S \mapsto \mathbb{P}[S|B]$$

is also a probability measure on $\mathcal{S}$. We say that it is *the probability measure obtained by conditioning on B* .

Remark 1.2.36. Observe that n events $A_1, \dots, A_n$, $n \geq 2$, are independent if and only if, for any nonempty subset $I \subset \{1, \dots, n\}$ of cardinality $< n$, and any $j \notin I$ we have

$$\mathbb{P}[A_j | A_I] = \mathbb{P}[A_j], \text{ where } A_I := \bigcap_{i \in I} A_i. \quad \square$$

Suppose we are given a finite or countable measurable partition of $(\Omega, \mathcal{S}, \mathbb{P})$

$$\Omega = \bigsqcup_{i \in I} A_i, \quad I \subset \mathbb{N}, \quad \mathbb{P}[A_i] \neq 0, \quad \forall i.$$

The law of total probability states that

$$\mathbb{P}[S] = \sum_{i \in I} \mathbb{P}[S | A_i] \mathbb{P}[A_i], \quad \forall S \in \mathcal{S}. \quad (1.2.11)$$

Indeed,

$$\mathbb{P}[S] = \sum_{i \in I} \mathbb{P}[S \cap A_i] \stackrel{(1.2.10)}{=} \sum_{i \in I} \mathbb{P}[S | A_i] \mathbb{P}[A_i].$$

In Exercise 1.11 I ask the reader to prove a conditional version of the law of total probability. More precisely given a finite or countable measurable partition $(A_i)_{i \in I}$ of $(\Omega, \mathcal{S}, \mathbb{P})$ and an event B such that $\mathbb{P}(B \cap A_i) > 0, \forall i$, then

$$\mathbb{P}[S | B] = \sum_{i=1}^n \mathbb{P}[S | A_i \cap B] \mathbb{P}[A_i | B]. \quad (1.2.12)$$

Note that (1.2.11) is a special case of (1.2.12) corresponding to the choice $B = \Omega$.

Example 1.2.37. Suppose that we have an urn containing b black balls and r red balls. A ball is drawn from the urn and discarded. Without knowing its color, what is the probability that a second ball drawn is black?

For $k = 1, 2$ denote by B_k the event “the k -th drawn ball is black”. We are asked to find $\mathbb{P}[B_2]$. The first drawn ball is either black (B_1) or not black (B_1^c). From the law of total probability we deduce

$$\mathbb{P}[B_2] = \mathbb{P}[B_2 | B_1] \mathbb{P}[B_1] + \mathbb{P}[B_2 | B_1^c] \mathbb{P}[B_1^c].$$

Observing that

$$\mathbb{P}[B_1] = \frac{b}{b+r} \quad \text{and} \quad \mathbb{P}[B_1^c] = \frac{r}{b+r},$$

we conclude

$$\begin{aligned} \mathbb{P}[B_2] &= \frac{b-1}{b+r-1} \cdot \frac{b}{b+r} + \frac{b}{b+r-1} \cdot \frac{r}{b+r} = \frac{b(b-1) + br}{(b+r)(b+r-1)} \\ &= \frac{b(b+r-1)}{(b+r)(b+r-1)} = \frac{b}{b+r} = \mathbb{P}[B_1]. \end{aligned}$$

Thus, the probability that the second extracted ball is black is equal to the probability that the first extracted ball is black. This seems to contradict our intuition because when we extract the second ball the composition of available balls at that time is different from the initial composition.

This is a special case of a more general result, due to S. Poisson, [39, Sec. 5.3].

Suppose in an urn containing b black and r red balls, n balls have been drawn first and discarded without their colors being noted. If another ball is drawn drawn next, the probability that it is black is the same as if we had drawn this ball at the outset, without having discarded the n balls previously drawn.

To quote John Maynard Keynes, [116, p.394],

This is an exceedingly good example of the failure to perceive that a probability cannot be influenced by the *occurrence* of a material event but only by such *knowledge* as we may have, respecting the occurrence of the event.

This example hides an even subtler phenomenon, namely *exchangeability*. We discuss this phenomenon in greater detail in Subsection 3.2.8. $\square$

Example 1.2.38 (The ballot problem). This is one of the oldest problems in probability.

Suppose that during an election between candidate H and T votes were cast successively. In the end H obtained a votes and T obtain votes and $a > b$ so H won. Denote by B the event that H was always in the lead while the voting took place. The ballot problem asks to find the probability of B given a and n .

We will visualize the voting process as an unbiased random walk on $\mathbb{Z}$. At the initial moment we are at the origin. A vote for H would move us a unit size step to the right, while a vote for T would move us one step to the left. The event B occurs if during this walk we always stay to the right of the origin,

Think of voting as flipping a *fair* coin with two sides, Heads and Tails. We denote by S_n its location after n coin flips. The sequence of random variables $(S_n)_{n \in \mathbb{N}}$ is called the standard (or unbiased) random walk on $\mathbb{Z}$.

Formally, we have a sequence of independent random variables $(X_n)_{n \in \mathbb{N}}$ such that

$$\mathbb{P}[X_n = 1] = \mathbb{P}[X_n = -1] = \frac{1}{2}, \quad \forall n \in \mathbb{N}.$$

The random variables with this distribution are called *Rademacher random variables*. Then

$$S_0 = 0, \quad S_n = S_0 + X_1 + \cdots + X_n.$$

Define a *zig-zag* to be a finite sequence of points or *nodes* in $\mathbb{Z}^2$

$$\omega = \{ (n_0, y_0), (n_0 + 1, y_1), \dots, (n_0 + k) \}$$

such that $|y_j - y_{j-1}| = 1$. We can visualize a zig-zag as the graph of a piecewise linear function obtained by connecting by line segments the successive nodes of the zig-zag; see Figure 1.2.

The node (n_0, y_0) is called the *initial node* of the zig-zag and $(n_0 + k, y_k)$ is called the *final node* of the zig-zag. We will refer to the natural number k as the length of the zig-zag. We will denote by $\Omega_{(n_0, y_0), (n_1, y_1)}$ the set of zig-zags that start at (n_0, y_0) and end at (n_1, y_1) . We set $\Delta n = n_1 - n_0$, $\Delta y = y_1 - y_0$. Then $\Delta n > 0$ and

$$\Delta y \equiv \Delta n \pmod{2}, \quad \#\Omega_{(n_0, y_0), (n_1, y_1)} = \binom{\Delta n}{h}, \quad h = \frac{\Delta n + \Delta y}{2}.$$

Each zig-zag in $\Omega_{(n_0, y_0), (n_1, y_1)}$ consists of Δn steps of which $\Delta_+ n$ go up, and the remaining $\Delta_- n$ go down so

$$\Delta n = \Delta_+ n + \Delta_- n, \quad \Delta y = \Delta_+ n - \Delta_- n.$$

It is convenient to visualize the random walk as a polygonal line obtained by successively connecting by a line segment the point $(n-1, S_{n-1})$ to the point (n, S_n) , $n \in \mathbb{N}$. The connecting line segment has slope X_n ;




Figure 1.2. A zig-zag describing a random walk started at $S_0 = 0$

The sample space in this problems is the space $\Omega_{n,y} = \Omega_{(0,0),(n,y)}$ of zig-zags ω that start at the origin and end at (n, y) . There are

$$\#\Omega_{n,y} = \binom{n}{a} = \binom{a+b}{b}, \quad a+b=n, \quad a-b=y,$$

equally likely zig-zags in this sample space. More precisely, there are $a+b$ steps of which a go up, and the remaining b go down.

The event B occurs iff $S_k > 0, \forall k = 1, \dots, n = a+b$.

Denote by $\Omega_{n,y}^+$ the set of zig-zags, $\omega \in \Omega_{n,y}$ that touch the horizontal axis only at $(0,0)$. Thus

$$\mathbb{P}[B] = \frac{\#\Omega_{n,y}^+}{\#\Omega_{n,y}}.$$

Note that all but the first node of a zig-zag in $\Omega_{n,y}^+$ is strictly above the horizontal axis. Thus, the first step of such a zig-zag is upwards, so $\Omega_{n,y}^+$ is in bijection with the space $\Omega_{(1,1),(n,y)}^+$ consisting of zig-zags from $(1,1)$ to (n,y) that do not touch the horizontal axis so that

$$\#\Omega_{n,y}^+ = \#\Omega_{(1,1),(n,y)}^+ = \#\Omega_{(1,1),(n,y)} - \#\Omega_{(1,1),(n,y)}^0.$$

where $\Omega_{(1,1),(n,y)}^0 \subset \Omega_{(1,1),(n,y)}$ consists of zig-zags that touch the horizontal axis.

For each zig-zag $\omega \in \Omega_{(1,1),(n,y)}^0$ denote by $k(\omega)$ the first moment it touches the horizontal axis. Denote by ω^r the zig-zag obtained from ω by reflecting in the horizontal axis the part of ω from $k(\omega)$ to n ; see Figure 1.3. The end point of ω^r is $(n, -(a-b))$.

The transformation $\omega \rightarrow \omega^r$ produces a bijection

$$\Omega_{(1,1),(n,-y)}^0 \rightarrow \Omega_{(1,1),(n,-y)}.$$

Indeed, any zig-zag $\omega' : (1,1) \rightarrow (n, b-a)$ must cross the horizontal axis. After the first touch we reflect it in this axis and obtain a zigzag $(1,1) \rightarrow (n, a-b)$ such that $\omega^r = \omega'$. Clearly ω touches the horizontal axis. The reflection trick used to produce this isomorphism is known as *André's reflection trick*. Hence

$$\#\Omega_{(1,1),(n,-y)}^0 = \binom{n-1}{b-1} = \binom{a+b-1}{a}, \quad (1.2.13)$$




Figure 1.3. The zigzag ω^r traces ω until ω hits the horizontal axis. At this moment the zigzag ω^r follows the opposite motion of ω (dashed line).

$$\#\Omega_{(1,1),(n,y)}^+ = \#\Omega_{(1,1),(n,y)} - \#\Omega_{(1,1),(n,-y)}^0 = \binom{a+b-1}{a-1} - \binom{a+b-1}{a}. \quad (1.2.14)$$

Note that

$$\begin{aligned} & \binom{a+b-1}{a-1} - \binom{a+b-1}{a} \\ &= \frac{(a+b-1)!}{(a-1)!(b-1)!} \left(\frac{1}{b} - \frac{1}{a} \right) = \frac{a-b}{a+b} \cdot \frac{(a+b)!}{a!b!} = \frac{a-b}{a+b} \binom{a+b}{a}, \\ & \mathbb{P}[B] = \frac{\#\Omega_{n,y}^+}{\#\Omega_{n,y}} = \frac{\binom{a+b-1}{b-1} - \binom{a+b-1}{a}}{\binom{a+b}{b}} = \frac{a-b}{a+b} = \frac{y}{n}. \end{aligned} \quad (1.2.15)$$

□

Proposition 1.2.39 (Bayes' formula). *Suppose we are given a finite or countable measurable partition of $(\Omega, \mathcal{S}, \mathbb{P})$*

$$\Omega = \bigsqcup_{i \in I} A_i, \quad I \subset \mathbb{N}, \quad \mathbb{P}[A_i] \neq 0, \quad \forall i.$$

Then, for any $S \in \mathcal{S}$ such that $\mathbb{P}[S] \neq 0$ and $i_0 \in I$ we have

$$\mathbb{P}[A_{i_0}|S] = \frac{\mathbb{P}[S|A_{i_0}]\mathbb{P}[A_{i_0}]}{\sum_{i \in I} \mathbb{P}[S|A_i]\mathbb{P}[A_i]}. \quad (1.2.16)$$

Proof. According to the law of total probability, the denominator in the right-hand-side of (1.2.16) equals $\mathbb{P}[S]$. Thus, the equality (1.2.16) is equivalent to

$$\mathbb{P}[A_{i_0}|S]\mathbb{P}[S] = \mathbb{P}[S|A_{i_0}]\mathbb{P}[A_{i_0}].$$

The product formula shows that both sides of the above equality are equal to $\mathbb{P}[A_{i_0} \cap S]$.

□

Remark 1.2.40. We should mention here a terminology favored by statisticians.

- The events A_k are called *hypotheses*.
- The probability $\mathbb{P}[A_k]$ is called *prior* (probability).
- The probability $\mathbb{P}[A_k|S]$ is called *posterior* (probability).

- The probability $\mathbb{P}[S|A_k]$ is called *likelihood*.

Here is one frequent application of Bayes' principle. Suppose that we observed a random event S we know that it can be caused only by one of the random events A_i . To decide which of the events A_i is more likely to have caused S we need to find the largest of the posteriors $\mathbb{P}[A_i|S]$. Bayes' formula shows that the most likely cause maximizes the numerator $\mathbb{P}[S|A_i]\mathbb{P}[A_i]$. $\square$

Example 1.2.41 (Biased coins). We say that a coin has bias $\theta \in (0, 1)$ if the probability of showing Heads when flipped is θ . Suppose that we have an urn containing c_1 coins with bias θ_1 and c_2 coins with bias θ_2 . Let $n := c_1 + c_2$ denote the total number of coins and set $p_i := \frac{c_i}{n}$, $i = 1, 2$. We assume that

$$c_1 < c_2 \quad \text{and} \quad \theta_1 > \theta_2, \quad (1.2.17)$$

i.e., there are fewer coins with higher bias. We draw a coin at random we flip it twice and we get Heads both times. What is the probability that the coin we have drawn has higher bias.

If θ denotes the (unknown) bias of the coin drawn at random, then we can think of θ as a random variable that takes two values θ_1, θ_2 with probabilities

$$\mathbb{P}[\theta_i] := \mathbb{P}[\theta = \theta_i] = p_i, \quad i = 1, 2.$$

Denote by E the event that two successive flips produce Heads. Then

$$\mathbb{P}[E|\theta_i] := \mathbb{P}[E|\theta = \theta_i] = \theta_i^2.$$

Bayes' formula shows that

$$\mathbb{P}[\theta_1|E] = \frac{\mathbb{P}[E|\theta_1]\mathbb{P}[\theta_1]}{\mathbb{P}[E|\theta_1]\mathbb{P}[\theta_1] + \mathbb{P}[E|\theta_2]\mathbb{P}[\theta_2]} = \frac{p_1\theta_1^2}{p_1\theta_1^2 + p_2\theta_2^2} = \frac{1}{1 + \frac{p_2}{p_1} \left(\frac{\theta_2}{\theta_1}\right)^2}.$$

Our assumption (1.2.17) shows that

$$\frac{c_2}{c_1} = \frac{p_2}{p_1} > 1 > \frac{\theta_2}{\theta_1}.$$

Observe that if $c_2\theta_2^2 > c_1\theta_1^2$, then

$$\mathbb{P}[\theta_1|E] < \frac{1}{2}.$$

Thus, in this case, if we observe two Heads, then the coin we randomly drew from the urn is less likely to be the one with bigger bias. For example if $\theta_1 = \frac{2}{3}$ and $\theta_2 = \frac{1}{3}$ and $c_2 > 8c_1$, then

$$\mathbb{P}[\theta_1|E] < \frac{1}{3},$$

so the randomly drawn coin is less likely to be the one heavily biased towards Heads. $\square$

1.2.3. Integration of measurable functions. We outline below, mostly without proofs, the construction and the basic facts about integration of measurable functions. For details we refer to [64, 131, 207].

Fix a measured space $(\Omega, \mathcal{S}, \mu)$. Recall that $\text{Elem}(\Omega, \mathcal{S})$ denotes the vector space of elementary $\mathcal{S}$ -measurable functions (see Definition 1.1.20). We denote by $\text{Elem}_+(\Omega, \mathcal{S})$ the convex cone of $\text{Elem}(\Omega, \mathcal{S})$ consisting of nonnegative elementary functions. Define

$$\mu : \text{Elem}_+(\Omega, \mathcal{S}) \rightarrow [0, \infty], \quad f \mapsto \mu[f] = \int_{\Omega} f(\omega) \mu[d\omega],$$

as follows. If

$$f = \sum_{k=1}^M a_k \mathbf{I}_{A_k}, \quad A_1, \dots, A_M \text{ disjoint},$$

then

$$\mu[f] = \int_{\Omega} f(\omega) \mu[d\omega] := \sum_{i=1}^M a_i \mu[A_i].$$

Note that if

$$f = \sum_{j=1}^N b_j \mathbf{I}_{B_j}, \quad B_1, \dots, B_n \text{ disjoint},$$

then $a_i = b_j$ if $A_i \cap B_j \neq \emptyset$. Hence

$$\sum_i a_i \mu[A_i] = \sum_i \sum_j a_i \mu[A_i \cap B_j] = \sum_j \sum_i b_j \mu[A_i \cap B_j] = \sum_j b_j \mu[B_j].$$

This shows that the value of $\int_{\Omega} f(\omega) \mu(d\omega)$ is independent of the decomposition of f as a linear combination of indicators of pairwise disjoint measurable sets.

The above integration map satisfies the following elementary properties.

$$\forall f, g \in \text{Elem}_+(\Omega, \mathcal{S}) \quad f \leq g \Rightarrow \mu[f] \leq \mu[g]. \quad (1.2.18a)$$

$$\forall a, b \geq 0, \quad f, g \in \text{Elem}_+(\Omega, \mathcal{S}) : \quad \mu[af + bg] = a\mu[f] + b\mu[g]. \quad (1.2.18b)$$

For $f \in \mathcal{L}_+^0(\Omega, \mathcal{S})$ we set

$$\mathcal{E}_+^f := \{ g \in \text{Elem}_+(\Omega, \mathcal{S}) ; \quad g \leq f \}.$$

The set $\mathcal{E}_+^f$ is nonempty since $0 \in \mathcal{E}_+^f$. Define

$$\boxed{\mu[f] = \int_{\Omega} f d\mu = \int_{\Omega} f(\omega) \mu[d\omega] := \sup_{g \in \mathcal{E}_+^f} \int_{\Omega} g(\omega) \mu[d\omega] \in [0, \infty)}. \quad (1.2.19)$$

Definition 1.2.42. A measurable function $f \in \mathcal{L}^0(\Omega, \mathcal{S})$ is called μ -integrable if

$$\mu[f^+], \quad \mu[f^-] < \infty.$$

In this case we define its *Lebesgue integral* to be

$$\int_{\Omega} f d\mu = \int_{\Omega} f(\omega) \mu[d\omega] = \mu[f] := \mu[f_+] - \mu[f_-].$$

We denote by $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ the set of μ -integrable functions and by $\mathcal{L}_+^1(\Omega, \mathcal{S}, \mu)$ the set of μ -integrable nonnegative functions. $\square$

Note that

$$\forall f, g \in \mathcal{L}_+^0(\Omega, \mathcal{S}) \quad f \leq g \Rightarrow \mu[f] \leq \mu[g]. \quad (1.2.20)$$

Moreover,

$$\forall f \in \mathcal{L}_+^0(\Omega, \mathcal{S}) : \quad \mu[f > 0] = 0 \iff \int_{\Omega} f d\mu = 0. \quad (1.2.21)$$

The integral $\mathcal{L}_+^0 \ni f \mapsto \mu[f] \in [0, \infty]$ enjoys the following key continuity property which is the “workhorse” of the Lebesgue integration theory.

Theorem 1.2.43 (Monotone Convergence theorem). *Suppose that $(f_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}_+^0(\Omega, \mathcal{S})$ that converges increasingly to $f \in \mathcal{L}_+^0(\Omega, \mathcal{S})$. Then*

$$\mu[f_n] \nearrow \mu[f] \text{ as } n \rightarrow \infty.$$

□

Proof. The sequence $\mu[f_n]$ is nondecreasing and is bounded above by $\mu[f]$. Hence it has a, possibly infinite, limit and

$$\lim_{n \rightarrow \infty} \mu[f_n] \leq \mu[f].$$

The proof of the opposite inequality

$$\lim_{n \rightarrow \infty} \mu[f_n] \geq \mu[f].$$

relies on a clever a clever trick. Fix $g \in \mathcal{E}_+^f$, $c \in (0, 1)$, and set

$$S_n := \{\omega \in \Omega; f_n(\omega) \geq cg(\omega)\}.$$

Since $f = \lim f_n$ and (f_n) is a nondecreasing sequence of functions we deduce that S_n is a nondecreasing sequence of measurable sets whose union is Ω . For any elementary function h the product $\mathbf{I}_{S_n} h$ is also elementary. For any $n \in \mathbb{N}$ we have $f_n \geq f_n \mathbf{I}_{S_n} \geq cg \mathbf{I}_{S_n}$ so that

$$\mu[f_n] \geq \mu[\mathbf{I}_{S_n} f_n] \geq c\mu[g \mathbf{I}_{S_n}].$$

If we write g as a finite linear combination

$$g = \sum_j g_j \mathbf{I}_{A_j}$$

with A_j pairwise disjoint, then we deduce

$$\mu[f_n] \geq c\mu[g \mathbf{I}_{S_n}] = c \sum_j g_j \mu[A_j \cap S_n].$$

The sequence of sets $(A_j \cap S_n)_{n \in \mathbb{N}}$ is nondecreasing and its union is A_j so that

$$\lim_{n \rightarrow \infty} \mu[f_n] \geq c \sum_j g_j \lim_{n \rightarrow \infty} \mu[A_j \cap S_n] = c \sum_j g_j \mu[A_j] = c\mu[g].$$

Hence

$$\lim_{n \rightarrow \infty} \mu[f_n] \geq c\mu[g], \quad \forall g \in \mathcal{E}_+^f, \quad \forall c \in (0, 1),$$

so that

$$\lim_{n \rightarrow \infty} \mu[f_n] \geq c\mu[f], \quad \forall c \in (0, 1).$$

Letting $c \nearrow 1$ we deduce $\lim_{n \rightarrow \infty} \mu[f_n] \geq \mu[f]$. □

Corollary 1.2.44. *For any $f \in \mathcal{L}_+^0(\Omega, \mathcal{S})$ we have*

$$\mu[f] = \lim_{n \rightarrow \infty} \mu[D_n[f]].$$

□

Corollary 1.2.45. *For any $f, g \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and $a, b \in \mathbb{R}$ such that $af + bg$ is well defined we have $af + bg \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and*

$$\int_{\Omega} (af + bg) d\mu = a \int_{\Omega} f d\mu + b \int_{\Omega} g d\mu. \quad (1.2.22)$$

Moreover, if $f, g \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and $f(\omega) \leq g(\omega), \forall \omega \in \Omega$ then

$$\int_{\Omega} f d\mu \leq \int_{\Omega} g d\mu.$$

□

Since $|f| = f^+ + f^-$ we deduce the following result.

Corollary 1.2.46. *Let $f \in \mathcal{L}^0(\Omega, \mathcal{S})$. Then*

$$f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \iff |f| \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu).$$

□

Corollary 1.2.47 (Markov's Inequality). *Suppose that $f \in \mathcal{L}_+^1(\Omega, \mathcal{S}, \mu)$. Then, for any $C > 0$, we have*

$$\mu[\{f \geq C\}] \leq \frac{1}{C} \int_{\Omega} f d\mu. \quad (1.2.23)$$

In particular, $f < \infty$, μ -a.e..

Proof. Note that

$$C\mathbf{I}_{\{f \geq C\}} \leq f \Rightarrow C\mu[\{f \geq C\}] = \int_{\Omega} C\mathbf{I}_{\{f \geq C\}} d\mu \leq \int_{\Omega} f d\mu.$$

□

Corollary 1.2.48. *If $f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$, then $\mu[\{|f| = \infty\}] = 0$.*

Proof. Note that

$$\mu[\{|f| = \infty\}] = \bigcap_{n \in \mathbb{N}} \mu[\{f > n\}].$$

On the other hand, Markov's inequality implies

$$\mu[\{f > n\}] \leq \frac{\mu[|f|]}{n} \rightarrow 0.$$

□

Proposition 1.2.49. *Suppose $f, g \in \mathcal{L}^0(\Omega, \mathcal{S})$ and $f = g$, μ -a.e.. Then*

$$f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \iff g \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu).$$

Moreover, if one of the above equivalent conditions hold, then $\mu[f] = \mu[g]$.

□

Remark 1.2.50. The presentation so far had to tread carefully around a nagging problem: given f, g in $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$, then $f(\omega) + g(\omega)$ may not be well defined for some ω . For example, it could happen that $f(\omega) = \infty$, $g(\omega) = -\infty$. Fortunately, Corollary 1.2.48 shows that the set of such ω 's is negligible. Moreover, if we redefine f and g to be equal to zero on the set where they had infinite values, then their integrals do not change. For this reason we alter the definition of $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ as follows.

$$\mathcal{L}^1(\Omega, \mathcal{S}, \mu) := \left\{ f : (\Omega, \mathcal{S}) \rightarrow \mathbb{R}; \ f \text{ measurable } \int_{\Omega} |f| d\mu < \infty \right\}.$$

Thus, in the sequel the integrable functions will be assumed to be everywhere finite.

With this convention, the space $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ is a vector space and the Lebesgue integral is a linear functional

$$\mu : \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \rightarrow \mathbb{R}, \quad f \mapsto \mu[f].$$

□

Remark 1.2.51 (Daniell-Stone integral). A *Daniell-Stone integral* is a triplet $(\Omega, \mathcal{E}, L)$, where Ω is a set, $\mathcal{E}$ is a vector space of bounded functions $\Omega \rightarrow \mathbb{R}$ and $L : \mathcal{E} \rightarrow \mathbb{R}$ is a linear map satisfying the following properties.

- (i) $1 \in \mathcal{E}$.
- (ii) $\forall f, g \in \mathcal{E}, \max(f, g), \min(f, g) \in \mathcal{E}$.
- (iii) If $f, g \in \mathcal{E}$ and $f \leq g$ then $L[f] \leq L[g]$.
- (iv) If $(f_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{E}$ such that $f_n \searrow 0$ as $n \rightarrow \infty$, then $L[f_n] \searrow 0$.

For example, if $(\Omega, \mathcal{S}, \mu)$ is a finite measured space and $\mathcal{E} = \mathcal{E}(\Omega, \mathcal{S}, \mu)$ is the subspace of elementary functions spanned by indicators of sets of finite measure, then the triplet $(\Omega, \mathcal{E}, \mu[-])$ is a Daniell-Stone integral.

The *Daniell-Stone theorem* states that there is only one way of producing Daniell-Stone integrals. More precisely, if $\mathcal{S}$ denotes the sigma-algebra of subsets of Ω generated by the functions $f \in \mathcal{E}$, then there exists a unique measure μ on $\mathcal{S}$ such that

$$\mathcal{E} \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \text{ and } \mu[f] = L[f], \quad \forall f \in \mathcal{E}.$$

For a proof we refer to [45, Thm. 7.7.4], [64, Sec.4.5] or [139, Chap.III]. $\square$

Recall that for any sequence $(x_n)_{n \in \mathbb{N}}$ of real numbers we have

$$\liminf_{n \rightarrow \infty} x_n = \lim_{k \rightarrow \infty} x_k^* := \inf_{n \geq k} x_n.$$

The sequence (x_k^*) is nondecreasing. The Monotone Convergence Theorem has the following useful immediate consequence.

Theorem 1.2.52 (Fatou's Lemma). *Suppose that $(f_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}_+^0(\Omega, \mathcal{S})$. Then*

$$\int_{\Omega} \liminf_{n \rightarrow \infty} f_n(\omega) \mu[d\omega] \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu.$$

$\square$

Proof. Set

$$g_k := \inf_{n \geq k} f_n.$$

Proposition 1.1.18(iii) implies that $g_k \in \mathcal{L}_+^0(\Omega, \mathcal{S})$. The sequence (g_k) is nondecreasing and

$$\liminf_{n \rightarrow \infty} f_n = \lim_{k \rightarrow \infty} g_k.$$

The Monotone Convergence Theorem implies that

$$\int_{\Omega} \liminf_{n \rightarrow \infty} f_n(\omega) \mu[d\omega] = \lim_{k \rightarrow \infty} \int_{\Omega} g_k d\mu.$$

Note that $g_k \leq f_n, \forall n \geq k$, and thus

$$\int_{\Omega} g_k d\mu \leq \int_{\Omega} f_n d\mu, \quad \forall n \geq k,$$

i.e.,

$$\int_{\Omega} g_k d\mu \leq \inf_{n \geq k} \int_{\Omega} f_n d\mu.$$

Letting $k \rightarrow \infty$ we deduce

$$\lim_{k \rightarrow \infty} \int_{\Omega} g_k d\mu \leq \lim_{k \rightarrow \infty} \inf_{n \geq k} \int_{\Omega} f_n d\mu = \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu.$$

$\square$

The next result illustrates one of the advantages of the Lebesgue integral over the Riemann integral: one needs less restrictive conditions to pass to the limit under the Lebesgue integral.

Theorem 1.2.53 (Dominated Convergence). *Suppose $(f_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ satisfying the following properties*

(i) *There exists $f \in \mathcal{L}^0(\Omega, \mathcal{S})$ such that*

$$\lim_{n \rightarrow \infty} f_n(\omega) = f(\omega), \quad \forall \omega \in \Omega.$$

(ii) There exists $g \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ such that

$$|f_n(\omega)| \leq g(\omega), \quad \forall \omega \in \Omega, \quad n \in \mathbb{N}.$$

Then $f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and

$$\lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu = \int_{\Omega} f d\mu, \quad (1.2.24a)$$

$$\lim_{n \rightarrow \infty} \int_{\Omega} |f_n(\omega) - f(\omega)| d\mu = 0. \quad (1.2.24b)$$

Proof. Set $g_n = |f| - f_n$. Then $g_n \geq 0$ and $\lim g_n = |f| - f$. Fatou's Lemma implies

$$\int_{\Omega} (|f| - f) d\mu \leq \liminf \int_{\Omega} (|f| - f_n) d\mu = \int_{\Omega} |f| d\mu - \limsup \int_{\Omega} f_n d\mu.$$

We deduce

$$\limsup \int_{\Omega} f_n d\mu \leq \int_{\Omega} f d\mu.$$

Arguing in the same fashion using the sequence $f_n - |f|$ we deduce

$$\int_{\Omega} f d\mu \leq \liminf \int_{\Omega} f_n d\mu.$$

Hence

$$\int_{\Omega} f d\mu \leq \liminf \int_{\Omega} f_n d\mu \leq \limsup \int_{\Omega} f_n d\mu \leq \int_{\Omega} f d\mu.$$

This proves (1.2.24a). The equality (1.2.24b) follows by applying (1.2.24a) to the sequence $g_n = |f_n - f|$. $\square$

Theorem 1.2.54 (Change in variables). Suppose that $(\Omega_0, \mathcal{S}_0)$, $(\Omega_1, \mathcal{S}_1)$ are measurable spaces and

$$\Phi : (\Omega_0, \mathcal{S}_0) \rightarrow (\Omega_1, \mathcal{S}_1)$$

is a measurable map. Fix a measure $\mu_0 : \mathcal{S}_0 \rightarrow [0, \infty]$ and a measurable function $f \in \mathcal{L}^0(\Omega_1, \mathcal{S}_1)$. Then

$$f \in \mathcal{L}^1(\Omega_1, \mathcal{S}_1, \Phi_{\#}\mu_0) \iff \Phi^*(f) := f \circ \Phi \in \mathcal{L}^1(\Omega_0, \mathcal{S}_0, \mu_0)$$

and

$$\int_{\Omega_0} \Phi^*(f) d\mu_0 = \int_{\Omega_1} f d\Phi_{\#}\mu_0. \quad (1.2.25)$$

Proof. Note that it suffices to prove the theorem in the case $f \geq 0$. The result is obviously true if $f \in \text{Elem}_+(\Omega_1, \mathcal{S}_1)$. The general case follows from the Monotone Convergence Theorem using the increasing approximation $D_n[f] \nearrow f$ of f by elementary functions; see (1.1.7). This has the property that $D_n[\Phi^*(f)] = \Phi^*(D_n[f])$. $\square$

Remark 1.2.55. Unlike the well known change-in-variables formula, the map T in (1.2.25) need not be bijective, only measurable.

If T is bijective with measurable inverse, then for any measure μ_1 on $(\Omega_1, \mathcal{S}_1)$ then (1.2.25) applied to the map T^{-1} reads

$$\int_{\Omega_1} f(\omega_1) \mu_1[d\omega_1] = \int_{\Omega_0} f(T\omega_0) T_{\#}^{-1} \mu_1[d\omega_0], \quad (1.2.26)$$

$\forall f \in \mathcal{L}^1(\Omega_1, \mathcal{S}_1, \mu_1)$.

If Ω_i are open subsets of $\mathbb{R}^n$, $\Phi : \Omega_0 \rightarrow \Omega_1$ is a C^1 -diffeomorphism onto,

$$x \mapsto y = \Phi(x),$$

then the classical change in variables formula for Lebesgue integral reads,

$$\int_{\Omega_1} f(y) \lambda[dy] = \int_{\Omega_0} f(\Phi(x)) |\det J_T(x)| \lambda[dx], \quad (1.2.27)$$

where $J_\Phi(x)$ is the Jacobian of the C^1 map $x \rightarrow \Phi x$.

In particular, if μ_0 is a Borel measure on Ω_0 that absolutely continuous with respect to the Lebesgue measure,

$$\mu_0[dx] = \rho_0(x) \lambda[dx],$$

then

$$\Phi_{\#} \mu_0[dy] = \rho_1(y) \lambda[dy], \quad \rho_1(y) = |\det J_{\Phi^{-1}}(y)| \rho_0(\Phi^{-1}y). \quad (1.2.28)$$

This follows from (1.2.25) and (1.2.27) with $f(y) = |\det J_{\Phi^{-1}}(y)| \rho_0(\Phi^{-1}y)$. $\square$

Proposition 1.2.56. Let $f \in \mathcal{L}_+^0(\Omega, \mathcal{S})$. Suppose that $\mu : \mathcal{S} \rightarrow [0, \infty]$ is a sigma-finite measure. Define

$$\mu_f : \mathcal{S} \rightarrow [0, \infty], \quad \mu_f[S] = \int_S f d\mu := \int_{\Omega} \mathbf{I}_S f d\mu.$$

Then, μ_f is a measure. Moreover

$$\mu_{f_0} = \mu_{f_1} \iff f_0 = f_1, \quad \mu - \text{almost everywhere.} \quad \square$$

The above result has an important converse. To state it we need to introduce the concept of *absolute continuity*.

Definition 1.2.57. Suppose that μ, ν are two measures on the measurable space $(\Omega, \mathcal{S})$. We say that ν is *absolutely continuous* with respect to μ , and we write this $\nu \ll \mu$ if

$$\forall S \in \mathcal{S} : \mu[S] = 0 \Rightarrow \nu[S] = 0. \quad \square$$

For a proof of the next result we refer to [21, 43, 207].

Theorem 1.2.58 (Radon–Nikodym). Suppose that μ, ν are two σ -finite measures on the measurable space $(\Omega, \mathcal{S})$. The following statements are equivalent.

- (i) $\nu \ll \mu$.
- (ii) There exists $\rho \in \mathcal{L}_+^0(\Omega, \mathcal{S})$ such that $\nu = \mu_\rho$, i.e.,

$$\nu[S] = \int_S \rho d\mu, \quad \forall S \in \mathcal{S}.$$

The function ρ is not unique, but it defines a unique element in $L_+^0(\Omega, \mathcal{S}, \mu)$ which we denote by $\frac{d\nu}{d\mu}$ and we will refer to it as the density of ν relative to μ . $\square$

1.2.4. L^p spaces. We recall here an important class of Banach spaces. For proofs and many more details we refer to [64, 131, 207]. We define an equivalence relation $\sim_\mu$ on $\mathcal{L}^0(\Omega, \mathcal{S})$ by declaring $f \sim_\mu g$ iff $\mu[f \neq g] = 0$. Note that

$$f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \text{ and } g \sim_\mu f \Rightarrow g \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \text{ and } \int_{\Omega} g d\mu = \int_{\Omega} f d\mu.$$

We set

$$L^0(\Omega, \mathcal{S}, \mu) := \mathcal{L}^0(\Omega, \mathcal{S}, \mu) / \sim_\mu, \quad L^1(\Omega, \mathcal{S}, \mu) := \mathcal{L}^1(\Omega, \mathcal{S}, \mu) / \sim_\mu.$$

For $p \in [1, \infty)$ we set

$$\mathcal{L}^p(\Omega, \mathcal{S}, \mu) := \left\{ f \in \mathcal{L}^0(\Omega, \mathcal{S}, \mu); \quad |f|^p \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu) \right\},$$

$$L^p(\Omega, \mathcal{S}, \mu) := \mathcal{L}^p(\Omega, \mathcal{S}, \mu) / \sim_\mu.$$

We will refer to the functions in $\mathcal{L}^p(\Omega, \mathcal{S}, \mu)$ as p -integrable functions. For $p \in [1, \infty)$ and $f \in \mathcal{L}^p(\Omega, \mathcal{S}, \mu)$ we set

$$\|f\|_p := \left(\int_\Omega |f|^p d\mu \right)^{\frac{1}{p}}.$$

Define

$$L^\infty(\Omega, \mathcal{S}, \mu) := \{ [f] \in L^0(\Omega, \mathcal{S}, \mu); \quad \exists g \in \mathcal{L}^\infty(\Omega, \mathcal{S}), \quad g \sim_\mu f \}.$$

For $f \in \mathcal{L}^0(\Omega, \mathcal{S})$ we define

$$\|f\|_\infty = \text{ess sup } |f| := \inf \{ a \geq 0; \quad \mu[|f| > a] = 0 \}.$$

Note that this quantity only depends on the $\sim_\mu$ -equivalence class of f and

$$L^\infty(\Omega, \mathcal{S}, \mu) = \{ f \in L^1(\Omega, \mathcal{S}, \mu); \quad \|f\|_\infty < \infty \}.$$

In this fashion we obtain for every $p \in [1, \infty]$ maps

$$\| \cdot \|_p : L^p(\Omega, \mathcal{S}, \mu) \rightarrow [0, \infty).$$

Theorem 1.2.59 (Hölder inequality). *Let $p, q \in [1, \infty]$ such that*

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Then for any $f \in \mathcal{L}^p(\Omega, \mathcal{S}, \mu)$ and $g \in \mathcal{L}^q(\Omega, \mathcal{S}, \mu)$ we have $fg \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and

$$\int_\Omega |fg| d\mu \leq \|f\|_p \cdot \|g\|_q. \quad (1.2.29)$$

□

Theorem 1.2.60 (Minkowski's inequality). *Let $p \in [1, \infty]$, Then,*

$$\forall f, g \in L^p(\Omega, \mathcal{S}, \mu) : \quad \|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

□

Theorem 1.2.61. *Fix a sigma-finite measured space $(\Omega, \mathcal{S}, \mu)$.*

- (i) *For any $p \in [1, \infty]$, the pair $(L^p(\Omega, \mathcal{S}, \mu), \| \cdot \|_p)$ is a Banach space.*
- (ii) *If $p \in [1, \infty)$, the vector subspace of p -integrable elementary functions is dense in $L^p(\Omega, \mathcal{S}, \mu)$. In particular, if $\mathcal{S}$ is generated as a sigma-algebra by a countable collection of sets, then $L^p(\Omega, \mathcal{S}, \mu)$ is separable.* □

The above density result follows from a combined application of the Monotone Class Theorem and the Monotone Convergence Theorem; see Exercise 1.9.

Suppose that $(\Omega, \mathcal{S}, \mu)$ is a measured space and $p \in [1, \infty]$. Denote by q the exponent conjugate to p , i.e.,

$$\frac{1}{p} + \frac{1}{q} = 1 \iff q = \frac{p}{p-1}.$$

If $g \in L^q(\Omega, \mathcal{S}, \mu)$, then Hölder's inequality shows that $fg \in L^1, \forall f \in L^p(\Omega, \mathcal{S}, \mu)$ and the resulting linear map

$$L^p(\Omega, \mathcal{S}, \mu) \ni f \mapsto \xi_g(f) := \int_{\Omega} gf d\mu \in \mathbb{R}$$

is continuous.

Theorem 1.2.62. *Suppose that $(\Omega, \mathcal{S}, \mu)$ is a sigma-finite measured space and $p \in (1, \infty)$. Then the map*

$$L^q(\Omega, \mathcal{S}, \mu) \ni g \mapsto \xi_g \in L^p(\Omega, \mathcal{S}, \mu)^* = \text{the dual of the Banach space } L^p(\Omega, \mathcal{S}, \mu)$$

is a bijective isometry of Banach spaces. □

1.2.5. Measures on compact metric spaces. Up to this point we have indicated how one can use a measure to define an integral. The integral is a linear functional on an appropriate space of measurable spaces.

On certain measurable spaces one can invert this process. Suppose that X is a topological space and $\mathcal{B} = \mathcal{B}_X$ is the sigma algebra of Borel sets. We denote by $C_b(X)$ the vector space of bounded continuous functions on X . This is equipped with the sup-norm

$$\|f\|_{\infty} = \sup_{x \in X} |f(x)|.$$

Any finite Borel measure μ on $\mathcal{B}$ defines via integration a continuous linear functional

$$I_{\mu} : C_b(X) \rightarrow \mathbb{R}, \quad I_{\mu}[f] = \int_X f(x) \mu[dx].$$

This linear functional satisfies the positivity condition

$$I_{\mu}[f] \geq 0, \quad \forall f \in C_b(X), \quad f \geq 0. \quad (\text{Pos})$$

On metric spaces the measure μ is uniquely determined by the associated functional μ . More precisely, we have the following fact.

Proposition 1.2.63. *If X is a metric space and μ, ν are two finite Borel measures such that*

$$I_{\mu}[f] = I_{\nu}[f], \quad \forall f \in C_b(X),$$

then $\mu[B] = \nu[B]$ for any subset $B \subset X$.

Proof. Since the Borel sigma-algebra of X is generated by the π -system $\mathcal{C}_X$ of closed subsets it suffices to show that

$$\mu[C] = \nu[C], \quad \forall C \in \mathcal{C}_X.$$

To see that this indeed the case fix $C \in \mathcal{C}_X$ and, for any $n \in \mathbb{N}$ denote by D_n the closed set

$$D_n := \{x \in X; \text{dist}(x, C) \geq 1/n\}.$$

Define $f_n \in C_b(X)$

$$f_n(x) := \frac{\text{dist}(x, D_n)}{\text{dist}(x, D_n) + \text{dist}(x, C)}.$$

The function f_n is identically 1 on C and identically 0 on D_n . Moreover

$$\lim_{n \rightarrow \infty} f_n(x) = \mathbf{I}_C(x), \quad \forall x \in X.$$

Using the Dominated Convergence Theorem we deduce

$$\mu[C] = \lim_{n \rightarrow \infty} I_\mu[f_n] = \lim_{n \rightarrow \infty} I_\nu[f_n] = \nu[C].$$

□

We want to include a useful consequence of the above proof.

Corollary 1.2.64. *Suppose that X is a metric space and μ is a finite Borel measure on X . Then the space $C_b(X)$ is dense in $L^1(X, \mathcal{B}_X, \mu)$.* □

We have the following remarkable result.

Theorem 1.2.65 (Riesz Representation). *Suppose that X is a compact metric space and L is a linear functional on $C(X)$ satisfying the positivity condition (Pos). Then there exists a unique finite Borel measure μ on X such that*

$$L[f] = I_\mu[f], \quad \forall f \in C(X).$$

Idea of proof. Observe that the triplet $(K, C(K), L)$ is a Daniell-Stone integral; see Remark 1.2.51. Indeed, observe that L is continuous since

$$|L[f]| \leq L[1] \cdot \|f\|_\infty, \quad \forall f \in C(K).$$

If $(f_n)_{n \geq 0}$ is a sequence of continuous functions converging decreasingly to 0, then Dini's theorem implies that f_n converge *uniformly* to 0, so $L[f_n] \searrow 0$. Moreover, the sigma-algebra generated by the continuous functions on K coincides with the Borel sigma-algebra since any closed set $S \subset K$ is the zero set of the continuous function $x \mapsto \text{dist}(x, C)$. Theorem 1.2.65 is now obviously a special case of the Daniell-Stone theorem; see Remark 1.2.51. □

For a details we refer to [66, Sec. IV.6, Thm.3] or [207, Thm. 13.5].

Example 1.2.66. We can use the above result to construct probability measures on a smooth compact manifold M of dimension m . As shown in e.g. [159, Sec. 3.4.1] a Riemann metric g on M , defines a continuous linear functional

$$C(M) \ni f \mapsto \int_M f dV_g \in \mathbb{R},$$

usually referred to as the integral with respect to the volume element determined by g . The Riesz Representation Theorem shows that this corresponds to the integral with respect to a finite Borel measure Vol_g on M called the *metric measure*. The metric volume of M is then

$$\text{Vol}_g[M] = \int_M \mathbf{1}_M dV_g.$$

We can associate to it the metric probability measure $\mathbb{P}_g$

$$\mathbb{P}_g[B] := \frac{1}{\text{Vol}_g[M]} \text{Vol}_g[B],$$

for any Borel subset $B \subset M$.

In particular, if M is a compact submanifold of an Euclidean space $\mathbb{R}^N$, then it comes equipped with an induced metric and as such, with a finite metric measure μ_M and thus with a probability measure $\mathbb{P}_M$. We will refer to this probability measure as the *Euclidean probability measure*.

Suppose for example that $M = S^m$ is the unit sphere in $\mathbb{R}^m$

$$S^m := \{ (x_0, x_1, \dots, x_m) \in \mathbb{R}^{m+1}; x_0^2 + \dots + x_m^2 = 1 \}.$$

The Euclidean volume of S^m is (see e.g. [159, Eq. (9.1.10)])

$$\sigma_m := \frac{2\pi^{(m+1)/2}}{\Gamma(\frac{m+1}{2})}$$

and the Euclidean probability measure is

$$\mathbb{P}_{S^m} = \frac{1}{\sigma_m} \mu_{S^m}.$$

For example, if $m = 1$, then μ_{S^1} is expressed traditionally as $d\theta$, where θ is the angular coordinate. Hence

$$\mathbb{P}_{S^1}[d\theta] = \frac{1}{2\pi} d\theta. \quad (1.2.30)$$

If we use spherical coordinates (φ, θ) on S^2 , where φ denotes the Latitude and θ the Longitude, then

$$\mathbb{P}_{S^2}[d\varphi d\theta] = \frac{1}{4\pi} \sin \varphi d\varphi d\theta. \quad (1.2.31)$$

□

1.3. Invariants of random variables

We have defined the random variables as measurable functions on a probability space. In concrete examples this probability space is not specifically mentioned. In fact there could be different looking random variables describing essentially the same random quantity.

Consider for example the simplest example of rolling a fair die and observing the number N that shows up. The possible values of N are $\{1, \dots, 6\}$. We equip $\mathbb{I}_6$ with the uniform probability measure and then we can view N as the map

$$N : \mathbb{I}_6 \rightarrow \mathbb{R}, \quad N(k) = k, \quad \forall k \in \mathbb{I}_6.$$

Consider now a different experiment. Pick a point x uniformly random in $(0, 1]$. We receive a reward $R(x) = k \in \mathbb{I}_6$ if $\lceil 6x \rceil = k$. The functions N and R are obviously different but the random quantities they described are very similar and they should have many things in common.

This is analogous to the situation we encounter in geometry or physics when the same physical or geometric object can be given different descriptions using different coordinates. The laws of physics or geometry are however independent of coordinates. Technically, this means they are described in terms of tensors.

In this section we explain a few basic techniques for describing the behavior of random variables that capture the similarities we observe intuitively.

1.3.1. The distribution and the expectation of a random variable. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For any random variable $X \in \mathcal{L}^0(\Omega, \mathcal{S})$ the most basic invariant is its *probability distribution* or the *law* of X , i.e., the pushforward

$$\mathbb{P}_X := X_{\#} \mathbb{P}. \quad (1.3.1)$$

Thus $\mathbb{P}_X$ is a Borel probability measure on $\mathbb{R}$ and, as such, it is uniquely determined by the *cumulative distribution function* (cdf)

$$F(x) = F_X(x) := \mathbb{P}[X \leq x].$$

More precisely, $\mathbb{P}_X$ can be identified with the associated Lebesgue-Stieltjes measure,

$$\mathbb{P}_X = dF_X.$$

When the random variable X is *discrete*, i.e., the range of X is a finite or countable discrete subset $\mathfrak{X} \subset \mathbb{R}$, then $\mathbb{P}_X$ is completely determined by the “mass” of each $x \in \mathfrak{X}$,

$$\mathbb{P}_X[\{x\}] = \mathbb{P}[X = x].$$

For this reason in this case the probability distribution of X is often referred as the *probability mass function* (or pmf) of X .

The *quantile* of X is the quantile of its cdf; see Example 1.2.22. More precisely, the quantile is the function

$$Q_X : [0, 1] \rightarrow \bar{\mathbb{R}}, \quad Q_X(p) = \inf \left\{ x \in \bar{\mathbb{R}}; \quad \mathbb{P}[X \leq x] \geq p \right\}. \quad (1.3.2)$$

Given a Borel probability measure μ on $\bar{\mathbb{R}}$, we will use the notation $X \sim \mu$ to indicate that the probability distribution of X is μ , i.e., $\mathbb{P}_X = \mu$.

Any probability measure μ on $(\bar{\mathbb{R}}, \mathcal{B}_{\bar{\mathbb{R}}})$ tautologically defines a random variable with probability distribution μ . If we denote by $\mathbb{1}_{\bar{\mathbb{R}}}$ the identity map $\bar{\mathbb{R}} \rightarrow \bar{\mathbb{R}}$, then the random variable

$$X = \mathbb{1}_{\bar{\mathbb{R}}} : (\bar{\mathbb{R}}, \mathcal{B}_{\bar{\mathbb{R}}}, \mu) \rightarrow \bar{\mathbb{R}}$$

has probability distribution $\mathbb{P}_X = \mu$. Because of this fact random variables are often identified with their probability distributions. We will use the notations

$$X \stackrel{d}{=} Y \quad \text{or} \quad X \sim Y$$

to indicate that X and Y have the same distribution.

Definition 1.3.1 (Expectation). The *expectation* or the *mean* of the integrable random variable $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ is the quantity

$$\mathbb{E}[X] = \mathbb{E}_{\mathbb{P}}[X] := \int_{\Omega} X(\omega) \mathbb{P}[d\omega]. \quad \square$$

We deduce from the Change in Variables Theorem 1.2.54 that

$$\int_{\mathbb{R}} x \mathbb{P}_X[dx] = \int_{\mathbb{R}} \mathbb{1}_{\mathbb{R}}(x) X_{\#} \mathbb{P}[dx] = \int_{\Omega} \mathbb{1}_{\mathbb{R}}(X(\omega)) \mathbb{P}[d\omega] = \mathbb{E}[X]$$

so that obtain the useful formula

$$\mathbb{E}[X] = \int_{\mathbb{R}} x \mathbb{P}_X[dx]. \quad (1.3.3)$$

If $F(x) = F_X(x)$ is the cdf of X , $F(x) = \mathbb{P}[X \leq x]$, then the distribution $\mathbb{P}_X$ is the Lebesgue-Stieltjes measure dF determined by F and (1.3.3) takes the classical form

$$\mathbb{E}[X] = \int_{\mathbb{R}} x dF(x). \quad (1.3.4)$$

The above equality shows that

$$X \stackrel{d}{=} Y \Rightarrow \mathbb{E}[X] = \mathbb{E}[Y].$$

More generally, for any Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(X)$ is integrable or nonnegative we have⁴

$$\mathbb{E}[f(X)] = \int_{\mathbb{R}} f(x) \mathbb{P}_X[dx]. \quad (1.3.5)$$

In other words, the expectation of a random variable is determined by its probability distribution alone, and not on the precise nature of the sample space on which it is defined.

For example, the random variables N and R described at the beginning of this section have the same distribution and thus they have the same mean

$$\mathbb{E}[N] = \mathbb{E}[R] = \frac{1 + \dots + 6}{6} = \frac{7}{2}.$$

Remark 1.3.2 (Bertrand's paradox). More often than not, in concrete problems the sample space where a random variable is defined is not explicitly mentioned. Sometimes this can create a problem. Consider the following classical example.

Pick a chord at random on a unit circle. What is the probability that its length is at least $\sqrt{3}$, the length of the edge of an equilateral triangle inscribed in that unit circle?

The answer depends on the concept of “*at random*” we utilize.

For example, we can think that a chord is determined by two points θ_1, θ_2 on the circle or, equivalently, by a pair of numbers in $[0, 2\pi]$. The corresponding chord has length $\leq \sqrt{3}$ if and only if $|\theta_1 - \theta_2| \geq \frac{2\pi}{3}$. The region in the square $[0, 2\pi]$ occupied by pairs (θ_1, θ_2) satisfying $|\theta_1 - \theta_2| \geq \frac{2\pi}{3}$ consists of two isosceles right triangles with legs of size $\frac{2\pi}{3}$ with vertices $(0, 2\pi)$ and $(2\pi, 0)$. By gluing these triangles along their hypotenuses we get a square one third the size of $[0, 2\pi]$. Assuming that the point (θ_1, θ_2) is chosen uniformly inside the square $[0, 2\pi]$ we deduce that the probability that the chord has length at most $\sqrt{3}$ is $\frac{1}{9}$.

On the other hand, a chord is uniquely determined by the location of its midpoint inside the unit circle. The chord has length at least $\sqrt{3}$ if and only if the midpoint is at distance at least $\frac{1}{2}$ from the center. Assuming that the midpoint is chosen uniformly inside this circle, we deduce that the probability that the chord is at least $\sqrt{3}$ is $\frac{3}{4}$ since the disk of radius $\frac{1}{2}$ occupies $\frac{1}{4}$ of the unit disk.

We can try to decide empirically which is correct answer, but any simulation/experiment must adopt a certain model of randomness. Things are even more complex. The set of chords has a natural symmetry given by the group of rotations about the origin. Any “reasonable” model of randomness ought to be compatible with this symmetry. In mathematical terms this means that the underlying probability measure ought to be invariant with respect to this symmetry.

As a set, we can identify the set of chords with the unit disk: we can describe a chord by indicating the location of its midpoint. The problem boils down to choosing a rotation invariant Borel measure on the unit disk. The quotient of the disk with respect to the group of rotation is a segment. In particular, any probability measure μ on the unit interval defines a rotation invariant probability measure $\mathbb{P}_\mu$ defined on the unit disk, determined by the requirements

$$\mathbb{P}_\mu[0 \leq r \leq r_1, \theta_0 \leq \theta \leq \theta_1] = \frac{\theta_1 - \theta_0}{2\pi} \mu[0, r_1].$$

Hence, there are infinitely many geometric randomness models. In our first model of randomness, the measure μ is the distribution the Lebesgue measure on $[0, 1]$ and $\mathbb{P}_\mu = dr d\theta$. In the second model of

⁴In undergraduate probability classes this formula is often referred as LOTUS: the Law Of The Unconscious Statistician.

randomness the measure μ is $2rdr$ and $\mathbb{P}_\mu = \frac{1}{\pi}rdrd\theta$, the normalized Lebesgue measure on the unit disk. $\square$

If $X, Y \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ and $a, b \in \mathbb{R}$, then $aX + bY \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ and

$$\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y] \quad (1.3.6)$$

The above linearity of the expectation is a very powerful tool. Here is a simple illustration.

Example 1.3.3. Suppose that $n \geq 3$ birds are arranged along a circle looking towards the center. At a given moment each bird randomly and independently turns his head to the left or to the right, with equal probabilities. After they turn their heads, some birds will be visible by one of their neighbors, and some not. Denote by X_n the number of birds that are invisible to their neighbors. We want to compute $\mathbb{E}[X_n]$, the expected number of invisible birds. We leave the reader to convince herself/himself that X_n is indeed a well defined mathematical object.

For $k = 1, \dots, n$ we denote by B_k the event that the k -th bird is invisible to its neighbors. Then

$$X_n = \sum_{k=1}^n \mathbf{I}_{B_k} \text{ and } \mathbb{E}[X_n] = \sum_{k=1}^n \mathbb{E}[\mathbf{I}_{B_k}] = \sum_{k=1}^n \mathbb{P}[B_k] = n\mathbb{P}[B_1].$$

The probability that the first bird is invisible to its neighbors is computed by observing that this happens iff its right neighbor turns his head right and its left neighbor turns his head left. Since they do this independently with probabilities $\frac{1}{2}$ we deduce

$$\mathbb{P}[B_1] = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}.$$

Hence

$$\mathbb{E}[X_n] = \frac{n}{4}.$$

To appreciate how efficient this computation is we present an alternate method.

We will determine the expectation by determining the probability distribution of X_n , or equivalently its *probability generating function* (pgf)

$$G_{X_n}(t) = \mathbb{E}[t^{X_n}] = \sum_{k \geq 0} \mathbb{P}[X_n = k] t^k.$$

I learned the argument below from *Luke Whitmer*, a student in one of my undergraduate probability courses.

Assume the birds sit on the edges of a convex n -gon $\mathcal{P}_n$. Orienting an edge corresponds to describing in which direction the corresponding bird is looking. We will refer to a choice of orientations of the edges of $\mathcal{P}_n$ as an *orientation* of $\mathcal{P}_n$. We denote by Ω_n the collection of orientations of $\mathcal{P}_n$. Note that $|\Omega_n| = 2^n$.

Fix a cyclic clockwise labelling of the vertices of n -gon, $v_1, v_2, \dots, v_n$ and define v_m for $m \in \mathbb{N}$ by requiring $v_i = v_j$ if $i \equiv j \pmod{n}$. The i -th bird sits on the edge $E_i := [v_i, v_{i+1}]$. The i -th bird, or equivalently the edge E_i , is invisible to its neighbors if E_{i-1} is oriented from v_i to v_{i-1} and E_{i+1} is oriented from v_{i+1} to v_{i+2} . Given an orientation ω of $\mathcal{P}_n$ we denote by $x_n(\omega)$ the number of invisible edges in this orientation. Thus

$$\mathbb{P}[X_n = j] = \frac{\#\{\omega \in \Omega_n; x_n(\omega) = j\}}{2^n}.$$

We distinguish two cases.

1. $n = 2k$. Denote by $\mathcal{P}_n^+$ the polygon obtained from $\mathcal{P}_n$ by collapsing the edges $E_1, E_3, E_5, \dots$. As vertices of the new polygon we can take the collapsed edges. The edges of the new polygon are

$$E_1^+ = E_2, E_2^+ = E_4, \dots, E_k^+ = E_{2k}.$$

Similarly, we denote by $\mathcal{P}_n^-$ the polygon obtained from $\mathcal{P}_n$ by collapsing the edges $E_2, E_4, \dots$. We can take the collapsed edges as vertices of the new polygon. Its edges are

$$E_1^- = E_1, E_2^- = E_3, \dots, E_k^- = E_{2k-1}.$$

Note that an orientation of $\mathcal{P}_n$ induces orientations of $\mathcal{P}_n^\pm$ and conversely, orientations $\mathcal{P}_n^\pm$ determine an orientation of $\mathcal{P}_n$. We denote by $\Omega_n^\pm$ the set of orientations of $\mathcal{P}_n^\pm$. We thus have a bijection

$$\Omega_n \ni \omega \mapsto (\omega_+, \omega_-) \in \Omega_n^+ \times \Omega_n^-.$$

Suppose now that we have an oriented m -gon $\mathcal{Q}_m$. If $q_1, \dots, q_m$ are the vertices $\mathcal{Q}_m$ we say that v_i is an *out-vertex* if both edges at v_i are oriented away from v_i and it is an *in-vertex*, if both edges at v_i are oriented towards v_i . A *neutral* vertex is a vertex with an incoming edge and one outgoing edge. For an orientation ω of $\mathcal{Q}_m$ we denote by $y_m(\omega)$ the number of out-vertices.

Fix an orientation on $\mathcal{P}_n$. An edge E_i is invisible in this orientation if and only if the corresponding vertex in $\mathcal{P}_n^{(-1)^i}$ is an out-vertex. More explicitly, if i is even/odd, then the corresponding vertex in $\mathcal{P}_n^\pm$ is an out-vertex. Note that,

$$x_{2k}(\omega) = y_k(\omega_+) + y_k(\omega_-). \quad (1.3.7)$$

We denote by $x_{n,j}$ the number of oriented n -gons with j invisible edges and we set

$$P_n(t) = \sum_{j \geq 0} x_{n,j} t^j = \sum_{\omega \in \Omega_n} t^{x_n(\omega)}.$$

Note that

$$G_{X_n}(t) = \frac{1}{2^n} P_n(t).$$

We denote by $y_{m,j}$ the number of oriented m -gons with j out-vertices and we set

$$Q_m(t) := \sum_{j \geq 0} y_{m,j} t^j = \sum_{\omega \in \Omega_m} t^{y_m(\omega)}.$$

From (1.3.7) we deduce

$$P_{2k}(t) = Q_k(t)^2. \quad (1.3.8)$$

2. $n = 2k + 1$. Fix an orientation of $\mathcal{P}_n$. Consider a new oriented n -gon $\mathcal{Q}_n$ with edges, in clockwise order

$$E'_1, E'_2, \dots, E'_n,$$

where E'_i carries the orientation of the edge $E_{(2i-1) \bmod n}$ of $\mathcal{P}_n$. Denote the vertices of $\mathcal{Q}_n$ by $q_1, q_2, \dots, q_n$, so the two edges that meet at q_i are E'_{i-1} and E'_i .

Imagine stepping in a clockwise fashion on the edges of $\mathcal{P}_n$ and skipping every other edge and labelling by E'_i the i -th edge we stepped on. Observe that the edge $E_{2i \bmod n}$ of $\mathcal{P}_n$ is invisible iff the vertex q_{i+1} (where $E'_i \leftrightarrow E_{2i-1}$ and $E'_{i+1} \leftrightarrow E_{2i+1}$ meet) is an out-vertex. Thus, the number of invisible edges of $\mathcal{P}_n$ is equal to the number of out-vertices of $\mathcal{Q}_n$. Hence

$$P_{2k+1}(t) = Q_{2k+1}(t). \quad (1.3.9)$$

To determine $Q_m(t)$ fix an orientation ω of an m -gon $\mathcal{Q}_m$. As we travel clockwise from one vertex to the next, the out- and in-vertices alternate: once we leave an out-vertex, the first non-neutral vertex we meet is an in-vertex and similarly once we leave an in-vertex the first non-neutral vertex we encounter is an out-vertex. In particular this shows that there is an equal number of in and out-vertices. Fix a cyclic labelling $\{1, 2, \dots, m\}$ of the vertices of $\mathcal{Q}_m$. If $y_m(\omega) = j$ then $z_m(\omega) = j$ so the set S of locations of in-/out-vertices has cardinality $2j$,

$$S = \{1 \leq \ell_1 < \ell_2 < \dots < \ell_{2j} \leq m, \}.$$

The above discussion shows that if ℓ_1 is an out/in-vertex, then all vertices $\ell_3, \ell_5, \dots$ are out/in-vertices while the even vertices $\ell_2, \ell_4, \dots$ are in/out-vertices. This shows that

$$y_{m,j} = 2 \binom{n}{2j}, \quad Q_m(t) = \sum_{j \geq 0} \binom{n}{2j} t^j,$$

$$Q_m(t^2) = (1+t)^m + (1-t)^m.$$

Hence

$$\begin{aligned} P_{2k}(t^2) &= ((1+t)^k + (1-t)^k)^2 = (1+t)^{2k} + (1-t)^{2k} + 2(1-t^2)^k, \\ P_{2k+1}(t^2) &= (1-t)^{2k+1} + (1+t)^{2k+1}. \end{aligned}$$

We conclude that

$$G_{X_n}(t) = \frac{1}{2^n} \times \begin{cases} (1-\sqrt{t})^{2k+1} + (1+\sqrt{t})^{2k+1}, & n = 2k+1, \\ (1+\sqrt{t})^{2k} + (1-\sqrt{t})^{2k} + 2(1-t)^k, & n = 2k. \end{cases}$$

The mean of X_n is

$$\mathbb{E}[X_n] = G'_{X_n}(1).$$

□

Example 1.3.4. Consider the unit circle $S^1 := \{z \in \mathbb{C}; |z| = 1\}$ equipped with its natural arclength measure $\ell[d\theta]$; see Example 1.2.66. In particular $\ell[S^1] = 2\pi$.

Choose a Borel subset $R \subset S^1$ of arclength $\frac{2\pi}{3}$ so R occupies two thirds of the circle. Set $G = S^1 \setminus R$, paint R in red, and paint G in green. We want to show that there is an inscribed square with at least three red vertices. We will do so using a clever probabilistic method we learned from [204, Ex. 1.1.15].

Let $\mathbb{P}$ denote the uniform probability measure on S^1 , $\mathbb{P}[d\theta] = \frac{1}{2\pi}\ell[d\theta]$ so $\mathbb{P}[R] = \frac{2}{3}$. Note that the vertices of an inscribed square have the form $z, iz, i^2z, i^3z, z \in S^1$. Define,

$$X_0, X_1, X_2, X_3 : S^1 \rightarrow \{0, 1\}, \quad X_k(z) = \mathbf{I}_R(i^k z), \quad S := X_0 + X_1 + X_2 + X_3.$$

There exists an inscribed square with at least 3 red vertices iff there exists $z \in S^1$ such that $S(z) > 2$.

Clearly, each of the random variables X_k is a Bernoulli random variable with success probability $\frac{2}{3}$. Hence

$$\mathbb{E}[S] = 4\mathbb{E}[X_0] = \frac{8}{3} > 2.$$

Hence there exists z such that $S(z) > 2$. □

Theorem 1.3.5. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ and $\mathcal{F}, \mathcal{G} \subset \mathcal{S}$ are two independent sigma-subalgebras. If $X \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$, $Y \in \mathcal{L}^1(\Omega, \mathcal{G}, \mathbb{P})$, then $XY \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ and

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]. \quad (1.3.10)$$

Proof. Observe that the equality (1.3.10) is bilinear in X and Y . The equality holds for $X = \mathbf{I}_F$, $F \in \mathcal{F}$ and $Y = \mathbf{I}_G$, $G \in \mathcal{G}$ and thus, by bilinearity, it holds for $X \in \text{Elem}(\Omega, \mathcal{F})$ and $Y \in \text{Elem}(\Omega, \mathcal{G})$.

If X, Y are nonnegative, then $D_n[X]D_n[Y] \nearrow XY$ and the Monotone Convergence Theorem shows that (1.3.10) holds for $X, Y \geq 0$. □

Corollary 1.3.6. Suppose that $X, Y \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ are independent random variables such that $XY \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$. Then

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]. \quad (1.3.11)$$

Proof. Use Theorem 1.3.5 with $\mathcal{F} = \sigma(X)$ and $\mathcal{G} = \sigma(Y)$. □

Corollary 1.3.7. Suppose that the random variables $X_1, \dots, X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ are independent. Then, for any Borel measurable functions $f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f_i(X_i) \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$$

we have $f_1(X_1) \cdots f_n(X_n) \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ and

$$\mathbb{E}[f_1(X_1) \cdots f_n(X_n)] = \mathbb{E}[f_1(X_1)] \cdots \mathbb{E}[f_n(X_n)].$$

Proof. Follows inductively from Corollary 1.3.6 by observing that for any $k = 2, \dots, n$ the random variables $f_1(X_1) \cdots f_{k-1}(X_{k-1})$ and $f_k(X_k)$ are independent. □

Corollary 1.3.8. Let $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ and suppose that $\mathcal{F} \subset \mathcal{S}$ is sigma-subalgebra. Then the following are equivalent.

(i) For any Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(X) \in \mathcal{L}^1$ and any $F \in \mathcal{F}$

$$\mathbb{E}[f(X)\mathbf{I}_F] = \mathbb{P}[F]\mathbb{E}[f(X)].$$

(ii) *The random variable X is independent of $\mathcal{F}$.*

Proof. The implication (i) $\Rightarrow$ (ii) follows by using $f = I_{(-\infty, x]}$, $x \in \mathbb{R}$. The converse follows from Theorem 1.3.6. $\square$

The following is not the usual definition of a convex function (see Exercise 1.31) but it has the advantage that it is better suited for the applications we have in mind.

Definition 1.3.9. Let I be an interval of the real axis. A continuous function $\varphi : I \rightarrow \mathbb{R}$ is called *convex* if for any $x_0 \in I$ there exists a linear function $\ell(x)$ such that⁵

$$\ell(x_0) = \varphi(x_0), \quad \ell(x) \leq \varphi(x), \quad \forall x \in I.$$

The convex function is called *strictly convex* if for any $x_0 \in I$ there exists a linear function $\ell(x)$ such that

$$\ell(x_0) = \varphi(x_0), \quad \ell(x) < \varphi(x), \quad \forall x \in I \setminus \{x_0\}. \quad \square$$

For example, if $\varphi : I \rightarrow \mathbb{R}$ is C^2 , then φ is convex (resp. strictly convex) if $\varphi''(x) \geq 0$ (resp. $\varphi'(x) > 0$), $\forall x \in I$.

Theorem 1.3.10 (Jensen's Inequality). *Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space, $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$, and $\varphi : I \rightarrow \mathbb{R}$ is a convex function defined on an interval I that contains the range of X . Then $\mathbb{E}[\varphi(X)]$ is well defined (possibly infinite) and*

$$\varphi(\mathbb{E}[X]) \leq \mathbb{E}[\varphi(X)]. \quad (1.3.12)$$

Moreover, if φ is strictly convex, then $\varphi(\mathbb{E}[X]) = \mathbb{E}[\varphi(X)]$ iff X is a.s. constant.

Proof. Observe that when φ is linear theorem is valid in the stronger form

$$\varphi(\mathbb{E}[X]) = \mathbb{E}[\varphi(X)].$$

We can find a linear function $\ell : \mathbb{R} \rightarrow \mathbb{R}$ such that $\varphi(x) \geq \ell(x)$, $\forall x \in I$ and it is clear that if the theorem is valid for the nonnegative convex function $g := \varphi - \ell$, then it is also valid for φ . Note that $\mathbb{E}[g(X)] \in [0, \infty]$ and thus the addition $\mathbb{E}[g(X)] + \ell(\mathbb{E}[X])$ is well defined and yields a well defined $\mathbb{E}[\varphi(X)]$, when $\varphi(X)$ is integrable or nonnegative. Moreover $\varphi(X)$ is integrable if and only if $g(X)$ is so. Because of this, we set

$$\mathbb{E}[\varphi(X)] := \infty \quad \text{if } \varphi(X) \text{ is not integrable.}$$

Set $\mu := \mathbb{E}[X]$ and observe that $\mu \in I$ since $X \in I$ a.s.. Choose a linear function $\ell : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\ell(x) \leq \varphi(x), \quad \forall x \in I \quad \text{and} \quad \ell(\mu) = \varphi(\mu).$$

Then

$$\varphi(\mathbb{E}[X]) = \varphi(\mu) = \ell(\mu) = \mathbb{E}[\ell(X)] \leq \mathbb{E}[\varphi(X)].$$

If φ is strictly convex, then we can choose $\ell(x)$ such that

$$\ell(x) < \varphi(x), \quad \forall x \in I \setminus \{\mu\} \quad \text{and} \quad \ell(\mu) = \varphi(\mu).$$

If X is not a.s. constant neither is the nonnegative random variable $\varphi(X) - \ell(X)$ so

$$\mathbb{E}[\varphi(X) - \varphi(\mu)] = \mathbb{E}[\varphi(X) - \ell(X)] > 0.$$

$\square$

⁵The graph of such an ℓ is tangent to the graph of φ at x_0 .

For any convex function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ we define the φ -entropy of an integrable random variable X to be the quantity

$$\mathbb{H}_\varphi[X] := \mathbb{E}[\varphi(X)] - \varphi(\mathbb{E}[X]). \quad (1.3.13)$$

Jensen's inequality shows that $\mathbb{H}_\varphi[X] \geq 0$.

1.3.2. Higher order integral invariants of random variables. On a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ we have the inclusions

$$L^{p_1}(\Omega, \mathcal{S}, \mathbb{P}) \subset L^{p_0}(\Omega, \mathcal{S}, \mathbb{P}), \quad \forall 1 \leq p_0 < p_1 \leq \infty.$$

Indeed, let $X \in \mathcal{L}^{p_1}(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$p := \frac{p_1}{p_0}, \quad \varphi(x) = x^p, \quad x \geq 0, \quad Y = |X|^{p_0}.$$

Since $p_1 > p_0$ the function φ is convex and we have

$$(\|X\|_{p_0})^{p_1} = \mathbb{E}[|X|_0^p]^p = \varphi(\mathbb{E}[Y]) \stackrel{(1.3.12)}{\leq} \mathbb{E}[\varphi(Y)] = (\|X\|_{p_1})^{p_1}.$$

In particular, if $p_0 = 1 \leq p$ we deduce

$$\mathbb{E}[|X|]^p \leq \mathbb{E}[|X|^p]. \quad (1.3.14)$$

Given $k \in \mathbb{N}$ and $X \in \mathcal{L}^k(\Omega, \mathcal{S}, \mathbb{P})$ we define the k -th momentum of X to be the quantity

$$\mu_k[X] := \mathbb{E}[X^k].$$

Note that $\mu_1[X] = \mathbb{E}[X]$.

Definition 1.3.11 (Variance). Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space. Suppose that $X \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$ is a random variable with mean $\mu := \mathbb{E}[X]$. The *variance* of X is the real number

$$\text{Var}[X] = \mathbb{E}[(X - \mu)^2].$$

The *standard deviation* of X is the quantity

$$\sigma[X] := \sqrt{\text{Var}[X]}. \quad \square$$

Observe that

$$\text{Var}[X] = 0 \iff X = \mathbb{E}[X] \text{ a.s..}$$

The quadratic function

$$q(t) = \mathbb{E}[(X - t)^2] = t^2 - 2\mu t + \mathbb{E}[X^2]$$

achieves its minimum at $t = \mu$ so that

$$\text{Var}[X] = \min_{t \in \mathbb{R}} \mathbb{E}[(X - t)^2].$$

Thus the standard deviation is the L^2 -distance from X to the 1-dimensional space of deterministic quantities. The variance can be given the alternate description

$$\text{Var}[X] = \mathbb{E}[X^2] - \mu^2 = \mu_2[X] - \mu_1[X]^2. \quad (1.3.15)$$

Indeed, if we set $\mu := \mathbb{E}[X]$, then

$$\text{Var}[X] = q(\mu) = \mathbb{E}[X^2] - \mu^2.$$

This shows that the variance is a special case of φ -entropy. More precisely,

$$\text{Var}[X] = \mathbb{H}_\varphi[X] = \mathbb{E}[\varphi(X)] - \varphi(\mathbb{E}[X]), \quad \varphi(x) = x^2.$$

Note that

$$\text{Var} [aX + b] = a^2 \text{Var} [X], \quad \forall a, b \in \mathbb{R}. \quad (1.3.16)$$

Indeed, set $\bar{X} := X - \mu$ and $Z := aX + b$. Then

$$\begin{aligned} \text{Var} [X] &= \mathbb{E} [\bar{X}^2], \quad Z - \mathbb{E}[Z] = a(X - \mathbb{E}[X]) = a\bar{X}, \\ \text{Var} [Z] &= \mathbb{E} [a^2 \bar{X}^2] = a^2 \text{Var} [X]. \end{aligned}$$

Theorem 1.3.12 (Chebyshev's inequality). *Let $X \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$ Set $\mu := \mathbb{E}[X]$ and $\sigma = \sigma[X]$. Then*

$$\mathbb{P} [|X - \mu| \geq c\sigma] \leq \frac{1}{c^2}, \quad \forall c > 0. \quad (1.3.17)$$

Equivalently

$$\mathbb{P} [|X - \mu| \geq r] \leq \frac{\text{Var}[X]}{r^2} = \frac{\sigma^2}{r^2}, \quad \forall r > 0. \quad (1.3.18)$$

Proof. Set $Y := |X - \mu|^2$. Then

$$\mathbb{P} [|X - \mu| > r] = \mathbb{P} [Y > r^2] \stackrel{(1.2.23)}{\leq} \frac{1}{r^2} \mathbb{E}[Y] = \frac{\text{Var}[X]}{r^2}.$$

Chebyshev's inequality (1.3.17) now follows from (1.3.18) by setting $r = c\sigma$. $\square$

Theorem 1.3.13 (Cantelli). *Let $X \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$ Set $\mu := \mathbb{E}[X]$ and $\sigma = \sigma[X]$. Then for any $x > \mu$ we have*

$$\mathbb{P} [X - \mu \geq x] \leq \frac{\sigma^2}{\sigma^2 + x^2} \quad (1.3.19)$$

This inequality is often referred to as the one-sided Chebyshev's inequality.

Proof. Without loss of generality we can assume that $\mu = 0$. For any $t \geq 0$ we set $Y = (X + t)^2$. Then

$$\mathbb{E}[Y] = \sigma^2 + t^2$$

and

$$\mathbb{P} [X \geq x] \leq \mathbb{P} [Y \geq (x + t)^2] \stackrel{(1.3.18)}{\leq} \frac{\sigma^2 + t^2}{(x + t)^2}, \quad \forall t \geq 0.$$

The inequality (1.3.19) is obtained by setting $t = \frac{\sigma^2}{x}$ in the above inequality. This is where the minimum of the right-hand side is attained. $\square$

Corollary 1.3.14. *Let $X \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$ Set $\mu := \mathbb{E}[X]$ and $\sigma = \sigma[X]$. If m is a median of X (see Definition 1.2.23), then $|\mu - m| \leq \sigma$.*

Proof. Since $m - \mu$ is also a median of $X - \mu$ we can assume $\mu = 0$. Suppose that $m \geq 0$. The inequality (1.3.19) implies that $\mathbb{P} [X \geq \sigma] \leq \frac{1}{2}$. Hence $\mathbb{P} [X \leq \sigma] > \frac{1}{2}$. Using (1.2.8) we deduce $m \leq m_+(X) \leq \sigma$.

—f $m \leq 0$, then $-m \geq 0$ is a median of $-X$ and we deduce from above that $-m \leq \sigma$. $\square$

Definition 1.3.15. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and $X, Y \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$. We set

$$\mu_X := \mathbb{E}[X], \quad \mu_Y := \mathbb{E}[Y].$$

(i) The *covariance* of X, Y is the quantity

$$\text{Cov} [X, Y] := \mathbb{E} [(X - \mu_X)(Y - \mu_Y)].$$

(ii) If X, Y are not deterministic we define the *correlation coefficient* of X and Y to be

$$\rho[X, Y] := \frac{\text{Cov}[X, Y]}{\sigma[X]\sigma[Y]}.$$

□

Proposition 1.3.16. *Let $X, Y \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$. Then the following hold.*

- (i) $\text{Cov}[X, Y] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$.
- (ii) *If X, Y are independent, then $\text{Cov}[X, Y] = 0$.*
- (iii) $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y] + 2\text{Cov}[X, Y]$.
- (iv) *If X, Y are independent, then $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y]$.*

Proof. Set

$$\mu_X := \mathbb{E}[X], \quad \bar{X} = X - \mu_X, \quad \mu_Y := \mathbb{E}[Y], \quad \bar{Y} = Y - \mu_Y.$$

(i) We have

$$\begin{aligned} \text{Cov}[X, Y] &= \mathbb{E}[\bar{X}\bar{Y}] = \mathbb{E}[XY] - \underbrace{\mathbb{E}[\mu_X Y]}_{\mu_X \mu_Y} - \underbrace{\mathbb{E}[\mu_Y X]}_{\mu_X \mu_Y} + \mu_X \mu_Y \\ &= \mathbb{E}[XY] - \mu_X \mu_Y. \end{aligned}$$

(ii) Corollary 1.3.6 shows that if X, Y are independent, then $\mathbb{E}[XY] = \mu_X \mu_Y$, i.e., $\text{Cov}[X, Y] = 0$.

(iii) Next

$$\begin{aligned} \text{Var}[X + Y] &= \mathbb{E}[(\bar{X} + \bar{Y})^2] = \mathbb{E}[\bar{X}^2] + \mathbb{E}[\bar{Y}^2] + 2\mathbb{E}[\bar{X}\bar{Y}] \\ &= \text{Var}[X] + \text{Var}[Y] + 2\text{Cov}[X, Y]. \end{aligned}$$

(iv) This follows from (ii) and (iii). □

Corollary 1.3.17. *If $X_1, \dots, X_n \in \mathcal{L}^2(\Omega, \mathcal{S}, \mathbb{P})$ are independent, then*

$$\text{Var}[X_1 + \dots + X_n] = \text{Var}[X_1] + \dots + \text{Var}[X_n]. \quad (1.3.20)$$

□

Example 1.3.18. Consider a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and two events $A, B \in \mathcal{S}$. We have

$$\text{Cov}[\mathbf{I}_A, \mathbf{I}_B] = \mathbb{P}[A \cap B] - \mathbb{P}[A]\mathbb{P}[B].$$

Thus A, B are independent iff $\text{Cov}[\mathbf{I}_A, \mathbf{I}_B] = 0$. □

Definition 1.3.19 (Moment generating function). Let X be a random variable defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that $e^{tX} \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ for all t in an open interval I containing 0. The *moment generating function* or *mgf* of X is the function

$$\mathbb{M}_X : I \rightarrow \mathbb{R}, \quad \mathbb{M}_X(t) = \mathbb{E}[e^{tX}].$$

□

The proof of following result is left to you as an exercise.

Proposition 1.3.20. *Let $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ be a random variable.*

- (i) If $\mathbb{M}_X(t) = \mathbb{E}[e^{tX}]$ is well defined for all $t \in (-t_0, t_0)$, then all the momenta of X are well defined and

$$\mathbb{M}_X(t) = \sum_{n=0}^{\infty} \mu_n[X] \frac{t^n}{n!}, \quad \forall |t| < t_0. \quad (1.3.21)$$

- (ii) If all the momenta of X are well defined and power series

$$\sum_{n=0}^{\infty} \mu_n[X] \frac{t^n}{n!},$$

converges $\forall t \in (-t_0, t_0)$, then its sum is $\mathbb{M}_X(t)$, $\forall |t| < t_0$.

□

Corollary 1.3.21. Suppose that $X_1, \dots, X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ are independent random variables such that $e^{tX_k} \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ for any $k = 1, \dots, n$ and any t in an open interval $I \subset \mathbb{R}$ that contains the origin. Then

$$\mathbb{M}_{X_1 + \dots + X_n}(t) = \mathbb{M}_{X_1}(t) \cdots \mathbb{M}_{X_n}(t), \quad \forall t \in I.$$

Proof. This is a special case of Corollary 1.3.7 corresponding to the choices

$$f_1(x) = \dots = f_n(x) = e^{tx}, \quad t \in I.$$

□

Remark 1.3.22 (The moment problem). Denote by Prob the set of Borel probability measures on the real axis and by $\text{Prob}^{\infty-}$ the subset of Prob consisting of probability measures $\mathbf{p}$ such that

$$\int_{\mathbb{R}} |x|^k \mathbf{p}[dx] < \infty, \quad \forall k \in \mathbb{N}.$$

For $\mathbf{p} \in \text{Prob}^{\infty-}$ and $k \in \mathbb{N}_0$ we set

$$\mu_k[\mathbf{p}] := \int_{\mathbb{R}} x^k \mathbf{p}[dx].$$

We denote by $\mathbb{R}^{\mathbb{N}_0}$ the set of sequences of real numbers $\underline{s} = (s_n)_{n \geq 0}$. We have a map

$$\boldsymbol{\mu} : \text{Prob}^{\infty-} \rightarrow \mathbb{R}^{\mathbb{N}_0}, \quad \boldsymbol{\mu}[\mathbf{p}] = (\mu_n[\mathbf{p}])_{n \geq 0}.$$

The *moment problem* asks the following.

- (i) Describe the range of $\boldsymbol{\mu}$, i.e., given a sequence of real numbers $\underline{s} = s_0, s_1, \dots$, decide if there exists $\mathbf{p} \in \text{Prob}^{\infty-}$ such that $\mu_n[\mathbf{p}] = s_n, \forall n \geq 0$.
- (ii) Is it true that the moments uniquely determine a probability measure, i.e., given $\underline{s}$ in the range of $\boldsymbol{\mu}$ is it true that there exists a *unique* $\mathbf{p} \in \text{Prob}^{\infty-}$ such that $\boldsymbol{\mu}[\mathbf{p}] = \underline{s}$?

Part (i) of the moment problem is completely understood in the sense that there are known several necessary and sufficient conditions for a sequence $\underline{s}$ to be the sequence of momenta of a probability measure on $\mathbb{R}$. We refer to [187, Chap. 3] for more details.

As for part (ii), it is known that a sequence $\underline{s}$ can be the sequence of momenta of *several* probability measures; see Exercise 1.39. On the other hand, there are known sufficient conditions on $\underline{s}$ guaranteeing the uniqueness of measure with that sequence of momenta; see [187, Chap. 4] for more

details. In particular, if X is a random variable such that e^{tX} is integrable for any t in an open interval containing 0, then $\mathbb{P}_X$ is uniquely determined by its moments, [187, Cor. 4.14]. $\square$

We formulate for the record the last uniqueness result mentioned above. In Exercise 2.53 we outline a proof of this special case.

Theorem 1.3.23. *Let $X, Y \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ such that there exist $r > 0$ with the property that*

$$\mathbb{E}[e^{tX}], \mathbb{E}[e^{tY}] < \infty, \quad \forall |t| < r.$$

Then

$$X \stackrel{d}{=} Y \iff \mathbb{M}_X(t) = \mathbb{M}_Y(t), \quad \forall |t| < r. \quad \square$$

Corollary 1.3.24. *Suppose that $\mathbb{P}_0, \mathbb{P}_1$ are Borel probability measures on $\mathbb{R}$ supported on $[0, 1]$, i.e.,*

$$\mathbb{P}_0[\mathbb{R} \setminus [0, 1]] = \mathbb{P}_1[\mathbb{R} \setminus [0, 1]] = 0.$$

Then

$$\mathbb{P}_0 = \mathbb{P}_1 \iff \int_{\mathbb{R}} x^n \mu_0[dx] = \int_{\mathbb{R}} x^n \mu_1[dx], \quad \forall n \in \mathbb{N}. \quad \square$$

Proof. Note that

$$\int_{\mathbb{R}} x^n \mathbb{P}_i[dx] \leq 1 \Rightarrow \int_{\mathbb{R}} e^{tx} \mathbb{P}_i[dx] < \infty, \quad \forall t \in \mathbb{R}$$

and

$$\begin{aligned} \int_{\mathbb{R}} e^{tx} \mathbb{P}_0[dx] &= \int_{\mathbb{R}} e^{tx} \mathbb{P}_1[dx], \quad \forall t \in \mathbb{R} \\ \iff \int_{\mathbb{R}} x^n \mathbb{P}_0[dx] &= \int_{\mathbb{R}} x^n \mathbb{P}_1[dx], \quad \forall n \in \mathbb{N}_0. \end{aligned} \quad \square$$

To a random variable X with range contained in $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ we can associate its *probability generating function* (or pgf)

$$G_X(t) := \sum_{n \geq 0} \mathbb{P}[X = n] t^n = \mathbb{E}[t^X].$$

Note that

$$G_X(1) = 1, \quad G'_X(1) = \mathbb{E}[X], \quad G''_X(1) = \mathbb{E}[X(X-1)]. \quad (1.3.22)$$

Similarly, if X, Y are two independent $\mathbb{N}_0$ -valued random variables, then

$$G_{X+Y}(t) = \mathbb{E}[t^{X+Y}] = \mathbb{E}[t^X] \mathbb{E}[t^Y] = G_X(t) G_Y(t).$$

1.3.3. Classical examples of discrete random variables. The theory of probability has grown mostly from concrete intriguing examples. In this process people encountered various frequently occurring patterns encoded by some ubiquitous random variables. We describe a few of them in the following subsections. These examples are part of the theory of probability and have many and varied uses. Their knowledge is absolutely necessary for a genuine understanding of probability.

Before Kolmogorov (and currently in most undergraduate probability courses), the world of random variables was divided into three categories: discrete, continuous and neither, or mixed. The discrete random variables are those whose ranges are discrete subsets of $\mathbb{R}$. A random variable X is called continuous if its probability distribution $\mathbb{P}_X$ is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$. We throw in the third category the random variables that do not fit in these two categories. We want to describe a few classical example of discrete and continuous random variables that play an important role in probability. Throughout our presentation we will frequently assume that given a sequence $(\mu_n)_{n \in \mathbb{N}}$ of Borel probability measures on $\mathbb{R}$ there exists a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and independent random variables $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ such that $\mathbb{P}_{X_n} = \mu_n, \forall n \in \mathbb{N}$. The fact that such a thing is possible is a consequence of Kolmogorov's existence theorem, Theorem 1.5.5.

We begin by introducing some frequently occurring discrete random variables by describing the random experiments where they appear.

Example 1.3.25 (Bernoulli random variables). Suppose we perform a random experiment aiming to observe the occurrence of a certain event S , $p := \mathbb{P}[S]$. When S has occurred we say that we have registered a success. Traditionally such an experiment is called a *Bernoulli trial* with success probability p . When the event S is not observed we say that the experiment was a failure. The failure probability is $q := 1 - p$. The Bernoulli trial is encoded by the random variable I_S which takes the value 1 when we register a success, and the value 0 otherwise. We also say that I_S is a *Bernoulli random variable*. Observe that

$$\mathbb{E}[I_S] = p, \quad \text{Var}[I_S] = \mathbb{E}[I_S^2] - (\mathbb{E}[I_S])^2 = p - p^2 = pq.$$

Note that any random variable with range $\{0, 1\}$ is a Bernoulli random variable since

$$X = I_{\{X=1\}}. \quad \square$$

Example 1.3.26 (Binomial random variables). Suppose that we perform the experiment in the above example n times, and the results of these experiments are independent of each other. We denote by N the number of successes observed during these n trials.⁶ We say that N is a *binomial random variable* corresponding to n trials with success probability p and we indicate this $N \sim \text{Bin}(n, p)$.

For $k = 1, \dots, n$ we denote by S_k the event “the k -th trial was a success”. Then

$$N = \sum_{k=1}^n I_{S_k} \quad \text{and} \quad \mathbb{E}[N] = \sum_{k=1}^n \mathbb{E}[I_{S_k}] = np.$$

Since the events $(S_k)_{1 \leq k \leq n}$ are independent we deduce from Corollary 1.3.7 that

$$\text{Var}[N] = \sum_{k=1}^n \text{Var}[I_{S_k}] = npq.$$

⁶Think for example that you roll a pair of dice 10 times and you aim to count how many times the sum of the numbers on the dice is 7. In this case success is when the sum is 7 and it is not hard to see that the probability of success is $\frac{1}{6}$.

Next observe that

$$G_N(s) = q + ps, \quad \mathbb{M}_N(t) = q + pe^t,$$

so

$$G_N(s) = G_{\mathbf{I}_{S_1}}(t)^n = (q + ps)^n, \quad \mathbb{M}_N(t) = \mathbb{M}_{\mathbf{I}_{S_1}}(t)^n = (q + pe^t)^n. \quad \square$$

This string of Bernoulli trials can be realized abstractly in the probability space

$$\left(\{0, 1\}^n, 2^{\{0, 1\}^n}, \beta_p^{\otimes n} \right)$$

described in Example 1.2.6(e). The events

$$S_k := \{ (\epsilon_1, \dots, \epsilon_n) \in \{0, 1\}^n; \quad \epsilon_k = 1 \}, \quad k = 1, \dots, n,$$

are independent and $\mathbb{P}[S_k] = p, \forall k = 1, \dots, n$. Then

$$\mathbf{I}_{S_k}(\epsilon) = \epsilon_k, \quad \forall \epsilon = (\epsilon_1, \dots, \epsilon_n) \in \{0, 1\}^n.$$

As explained in Example 1.2.6(e), the probability distribution of N is given by the equalities

$$\mathbb{P}[N = k] = \binom{n}{k} p^k q^{n-k}, \quad k = 0, 1, \dots, n.$$

Equivalently,

$$\mathbb{P}_N = \sum_{k=0}^n \binom{n}{k} p^k q^{n-k} \delta_k.$$

□

Example 1.3.27 (Waiting for successes). Suppose that we perform *independent* Bernoulli trials until we register the first success. We denote by T_1 the moment we observe the first success, $T_1 \in \mathbb{N} \cup \{\infty\}$. The random variable T_1 is a *geometric random variable* with success probability p . We write this $T_1 \sim \text{Geom}(p)$.

Observe that $T_1 = n$ iff the first $n - 1$ trials were failures and the n -th trial was a success. Thus

$$\mathbb{P}[T_1 = n] = q^{n-1}p.$$

In particular, $\mathbb{P}[T_1 = \infty] = 0$. We deduce that the probability distribution of T_1 is

$$\mathbb{P}_{T_1} = \sum_{n \geq 1} p q^{n-1} \delta_n.$$

Moreover

$$\mathbb{E}[T_1] = \sum_{n \geq 1} n p q^{n-1} = p \sum_{n \geq 1} n q^{n-1} = p \frac{d}{dq} \sum_{n \geq 0} q^n = \frac{p}{(1-q)^2} = \frac{1}{p}. \quad (1.3.23)$$

Here is a simple plausibility test for this result. Suppose we roll a die until we first roll a 1. The probability of rolling a 1 is $\frac{1}{6}$ so it is to be expected that we need 6 rolls until we roll our first 1.

We have

$$\begin{aligned} \mathbb{E}[T_1^2] - \mathbb{E}[T_1] &= \sum_{n=1}^{\infty} n(n-1) p q^{n-1} = \sum_{n=2}^{\infty} n(n-1) p q^{n-1} \\ &= p q \sum_{n=2}^{\infty} n(n-1) q^{n-2} = p q \frac{d^2}{dq^2} \left(\frac{1}{1-q} \right) = \frac{2pq}{(1-q)^3} = \frac{2q}{p^2}. \end{aligned}$$

We deduce that

$$\mathbb{E}[T_1^2] = \frac{2q}{p^2} + \frac{1}{p}, \quad \text{Var}[T_1] = \frac{2q}{p^2} + \frac{1}{p} - \frac{1}{p^2} = \frac{q}{p^2}.$$

Note that

$$\mathbb{M}_{T_1}(t) = \mathbb{E}[e^{tT_1}] = \sum_{n=1}^{\infty} pq^{n-1}e^{nt} = pe^t \sum_{m=0}^{\infty} (qe^t)^m = \frac{pe^t}{1-e^t}.$$

Consider now a more general situation. Fix $k \in \mathbb{N}$ and perform independent Bernoulli trials until we observe the k -th success. Denote by T_k the number trials until we record the k -th success. Note that

$$T_k = T_1 + (T_2 - T_1) + (T_3 - T_2) + \cdots + (T_k - T_{k-1}).$$

Due to the independence of the trials, once we observe the i -th success it is as if we start the experiment anew, so the waiting time $T_{i+1} - T_i$ until we observe the next success, the $(i+1)$ -th, is a random variable with the same distribution as T_1

$$T_{i+1} - T_i \stackrel{d}{=} T_1, \quad \forall i \in \mathbb{N}.$$

Hence $\mathbb{E}[T_{i+1} - T_i] = \mathbb{E}[T_1] = \frac{1}{p}$ so

$$\mathbb{E}[T_k] = k\mathbb{E}[T_1] = \frac{k}{p}. \quad (1.3.24)$$

The probability distribution of T_k is computed as follows. Note that $T_k = n$ if during the first $n-1$ trials we observed exactly $k-1$ successes, and at the n -th trial we observed another success. Hence

$$\mathbb{P}[T_k = n] = \binom{n-1}{k-1} p^{k-1} q^{n-k} \cdot p = \binom{n-1}{k-1} p^k q^{n-k}, \quad (1.3.25)$$

and

$$\mathbb{M}_{T_k}(t) = \left(\frac{pe^t}{1-e^t} \right)^k.$$

Since the waiting times between two consecutive successes are independent random variables we deduce

$$\text{Var}[T_k] = k \text{Var}[T_1] = \frac{kq}{p^2}.$$

The above probability measure on $\mathbb{R}$ is called the *negative binomial distribution* and T_k is called a *negative binomial random variable* corresponding to k successes with probability p . We write this $T_k \sim \text{NegBin}(k, p)$. $\square$

Let us describe a classical and less than obvious application of the geometric random variables.

Example 1.3.28 (The coupon collector problem). The coupon collector's problem arises from the following scenario. Suppose that each box of cereal contains one of m different coupons. Once you obtain one of every type of coupons, you can send in for a prize. Ann wants that prize and, for that reason, she buys one box of cereals everyday. Assuming that the coupon in each box is chosen independently and uniformly at random from the m possibilities and that Ann does not collaborate with others to collect coupons, how many boxes of cereal is she expected to buy before she obtain at least one of every type of coupon?

Let N denote the number of boxes bought until Ann has at least one of every coupon. We want to determine $\mathbb{E}[N]$. For $i = 1, \dots, n-1$ denote by N_i the number of boxes she bought while she had exactly i coupons. The first box she bought contained one coupon. Then she bought N_1 boxes

containing the coupon you already had. After $1 + N_1$ boxes she has two coupons. Next, she bought N_2 boxes containing one of the two coupons you already had etc. Hence⁷

$$N = 1 + N_1 + \cdots + N_{m-1}.$$

Let us observe first that for $i = 1, \dots, m-1$ we have

$$N_i \sim \text{Geom}(p_i), \quad p_i = \frac{m-i}{m}, \quad q_i = 1 - p_i = \frac{i}{m}.$$

Indeed, at the moment she has i coupons, a success occurs when she buys one of the remaining $m-i$ coupons. The probability of buying one such coupon is thus $\frac{m-i}{m}$. Think of buying a box at this time as a Bernoulli trial with success probability $\frac{m-i}{m}$. The number N_i is then equal to the number of trials until you register the first success. This argument also shows that the random variables N_i are independent. In particular,

$$\mathbb{E}[N_i] = \frac{1}{p_i} = \frac{m}{m-i}.$$

From the linearity of expectation we deduce

$$\begin{aligned} \mathbb{E}[N] &= 1 + \mathbb{E}[N_1] + \mathbb{E}[N_2] + \cdots + \mathbb{E}[N_{m-1}] \\ &= m \underbrace{\left(1 + \frac{1}{2} + \cdots + \frac{1}{m-1} + \frac{1}{m}\right)}_{=: H_m}. \end{aligned}$$

Asymptotically H_m differs from $\log m$ by the mysterious Euler-Mascheroni constant $\gamma \approx 0.5772$, i.e.,

$$\lim_{m \rightarrow \infty} (H_m - \log m) = \gamma.$$

Thus the expected number of boxes needed to collect all the m coupons is about $m \log m + m\gamma$. $\square$

Remark 1.3.29. We can ask a more general question. For $k \geq 1$ we denote by $X_k = X_{k,m}$ the number of boxes Ann has to buy until she has at least k of each of these m coupons. We have seen that that $\mathbb{E}[X_{1,m}] = mH_m$. One can show that as $m \rightarrow \infty$ we have

$$\mathbb{E}[X_{k,m}] = m(\log m + (k-1) \log \log m + \gamma - \log(k-1)! + o(1)),$$

where γ is the Euler-Mascheroni constant. For details we refer to [72, 157]. $\square$

Example 1.3.30 (The hypergeometric distribution). Suppose that we have a bin containing w white balls and b black balls. We select n balls at random from the bin and we denote by X the number of white balls among the selected ones. This is a random variable with range $0, 1, \dots, n$ called the *hypergeometric random variable* with parameters w, b, n . We will use the notation $X \sim \text{HGeom}(w, b; n)$ to indicate this and we will refer to its pmf as the *hypergeometric distribution*. For example, if A is the number of aces in a random poker hand, then $A \sim \text{HGeom}(4, 48; 5)$.

To compute $\mathbb{P}[X = k]$ when $X \sim \text{HGeom}(w, b; n)$ note that a favorable outcome for the event $X = k$ is determined by a choice of k white balls (out of w) and another independent choice of $n-k$ black balls (out of b) so that the number of favorable outcomes is

$$\binom{w}{k} \binom{b}{n-k}.$$

⁷Here we tacitly assume that we can describe quantities N_i as measurable functions defined on the same probability space. In Exercise 1.14 we ask the reader to do this. It is more challenging than it looks.

The number of possible outcomes of a random draw of n balls $\binom{w+b}{n}$. Hence

$$\mathbb{P}[X = k] = \frac{\binom{w}{k} \binom{b}{n-k}}{\binom{w+b}{n}}.$$

Its probability generating function is

$$G_X(s) = \frac{1}{\binom{N}{n}} \sum_{k=0}^n \binom{w}{k} \binom{b}{n-k} s^k, \quad N := w + b.$$

We can identify $G_X(s)$ as the coefficient of x^n in the polynomial

$$Q(s, x) = \frac{1}{\binom{N}{n}} (1 + sx)^w (1 + x)^b.$$

We have

$$\frac{\partial Q}{\partial s}(s, x) = \frac{wx(1+x)^b}{\binom{N}{n}} (1 + sx)^{w-1}.$$

The mean of X is $G'_X(1)$ and it is equal to the coefficient of x^n in

$$\frac{\partial Q}{\partial s}(1, x) = \frac{wx}{\binom{w+b}{n}} (1+x)^{N-1} = \frac{w \binom{N-1}{n-1}}{\binom{N}{n}} = \frac{wn}{N} = \frac{wn}{w+b}.$$

Hence

$$\mathbb{E}[\text{HGeom}(w, b; n)] = \frac{w}{w+b} \cdot n. \quad (1.3.26)$$

This equality has a very natural interpretation. Initially, the ratio of white balls in the bin was $r := \frac{w}{w+b}$. If we randomly draw n balls from the bin, we expect a ratio r of them to be white so we expect to draw rw white balls. $\square$

Example 1.3.31 (Poisson random variables). These random variables count the number N of random rare events that occur in a given unit of time. E.g., N could mean the number of computers in a large organization that die during one fiscal year. They depend on a parameter λ and we indicate this using the notation $N \sim \text{Poi}(\lambda)$. If $N \sim \text{Poi}(\lambda)$, then

$$\mathbb{P}[N = n] = e^{-\lambda} \frac{\lambda^n}{n!}, \quad \text{i.e., } \mathbb{P}_N = \sum_{n=0}^{\infty} e^{-\lambda} \frac{\lambda^n}{n!} \delta_n$$

Then

$$\mathbb{E}[N] = \sum_{n \geq 0} e^{-\lambda} \frac{n \lambda^n}{n!} = \lambda e^{-\lambda} \sum_{n \geq 1} \frac{\lambda^{n-1}}{(n-1)!} = \lambda.$$

The moment generating function of N is

$$\mathbb{M}_N(t) = \mathbb{E}[e^{tN}] = e^{-\lambda} \sum_{n \geq 0} \frac{(\lambda e^t)^n}{n!} = e^{\lambda(e^t - 1)}.$$

We have

$$\mathbb{M}'_N(t) = \lambda e^t e^{\lambda(e^t - 1)}, \quad \mathbb{M}''_N(t) = \lambda e^t e^{\lambda(e^t - 1)} + (\lambda e^t)^2 e^{\lambda(e^t - 1)}$$

so

$$\mathbb{E}[N^2] = \mathbb{M}''_N(0) = \lambda + \lambda^2, \quad \text{Var}[N] = \lambda.$$

$\square$

Example 1.3.32 (The inclusion-exclusion principle). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $A_1, \dots, A_n \in \mathcal{S}$. We want to compute the probability distribution of the random variable

$$N = \sum_{k=1}^n \mathbf{I}_{A_k}.$$

If the events A_k were independent and had identical probabilities, then $N \sim \text{Bin}(n, p)$. Set

$$\mathbb{I}_n := \{1, \dots, n\}.$$

For $m = 0, 1, \dots, n$ we denote by Ω_m set of points $\omega \in \Omega$ that belong to *exactly* m of the sets $A_1, \dots, A_n$. In other words $\Omega_m = \{N = m\}$. Note that

$$\Omega_0^c = A_1 \cup \dots \cup A_n.$$

For $I \subset \mathbb{I}_n$ we set

$$A_I := \begin{cases} \bigcap_{i \in I} A_i, & I \neq \emptyset, \\ \Omega, & I = \emptyset. \end{cases}$$

For $k \in \{0, 1, 2, \dots, n\}$ we define

$$s_k = s_k^n := \sum_{\substack{I \subset \mathbb{I}_n, \\ |I|=k}} \mathbb{P}[A_I]. \quad (1.3.27)$$

The *inclusion-exclusion principle* states that

$$\mathbb{P}[N = m] = \mathbb{P}[\Omega_m] = \sum_{k=0}^{n-m} (-1)^k \binom{m+k}{m} s_{m+k}, \quad \forall m = 0, 1, \dots, n. \quad (1.3.28)$$

Using the above equality with $m = 0$ we obtain the better known formula

$$\mathbb{P}[A_1 \cup \dots \cup A_n] = 1 - \mathbb{P}[\Omega_0] = \sum_{k=1}^n (-1)^{k-1} \sum_{\substack{I \subset \mathbb{I}_n, \\ |I|=k}} \mathbb{P}[A_I] = \sum_{k=1}^n (-1)^{k-1} s_k. \quad (1.3.29)$$

To prove (1.3.28) we set

$$S_k = S_k^n := \sum_{\substack{I \subset \mathbb{I}_n, \\ |I|=k}} \mathbf{I}_{A_I}. \quad (1.3.30)$$

Note that

$$s_k^n = \mathbb{E}[S_k^n].$$

We will prove that

$$\mathbf{I}_{\Omega_m} = \sum_{k=0}^{n-m} (-1)^k \binom{m+k}{m} S_{m+k}. \quad (1.3.31)$$

Indeed, using the equality $\mathbf{I}_{A \cap B} = \mathbf{I}_A \cdot \mathbf{I}_B$ we deduce

$$\begin{aligned} \mathbf{I}_{\Omega_m} &= \sum_{\substack{I \subset \mathbb{I}_n \\ |I|=m}} \left(\prod_{i \in I} \mathbf{I}_{A_i} \prod_{j \in \mathbb{I}_n \setminus I} \mathbf{I}_{A_j^c} \right) = \sum_{\substack{I \subset \mathbb{I}_n \\ |I|=m}} \left(\prod_{i \in I} \mathbf{I}_{A_i} \prod_{j \in \mathbb{I}_n \setminus I} (1 - \mathbf{I}_{A_j}) \right) \\ &= \sum_{k=0}^{n-m} (-1)^k \sum_{|J|=m+k} c(J) \mathbf{I}_{A_J}. \end{aligned}$$

Now observe that for any subset $J \subset \mathbb{I}_n$ of cardinality $m+k$ there are $\binom{m+k}{m}$ different way of writing $\mathbf{I}_{A_J}$ as a product

$$\mathbf{I}_{A_J} = \mathbf{I}_{A_I} \mathbf{I}_{A_{J \setminus I}}, \quad |I| = m.$$

Thus $c(J) = \binom{m+k}{m}$ for $|J| = m+k$. We deduce

$$\sum_{|J|=m+k} c(J) \mathbf{I}_{A_J} = \binom{m+k}{m} S_{m+k}.$$

Using the linearity of expectation we deduce from (1.3.31) that

$$\mathbb{P}[\Omega_m] = \mathbb{E}[\mathbf{I}_{\Omega_m}] = \sum_{k=0}^{n-m} (-1)^k \binom{m+k}{m} \mathbb{E}[S_{m+k}],$$

where $\mathbb{E}[S_{m+k}] = s_{m+k}$.

Associated to the equality (1.3.28) there is a sequence of inequalities called the *Bonferroni inequalities*. For $\ell \in \mathbb{N}$ and $\frac{n-m}{2} \geq \ell$

$$\sum_{k=0}^{2\ell-1} (-1)^k \binom{m+k}{m} s_{m+k} \leq \mathbb{P}[\Omega_m] \leq \sum_{k=0}^{2\ell} (-1)^k \binom{m+k}{m} s_{m+k}. \quad (1.3.32)$$

The above inequalities follow from the “*motivic*” *Bonferroni inequalities*

$$\begin{aligned} \sum_{k=0}^{2\ell-1} (-1)^k \binom{m+k}{m} S_{m+k} &\leq \mathbf{I}_{\Omega_m} \\ &\leq \sum_{k=0}^{2\ell} (-1)^k \binom{m+k}{m} S_{m+k}, \quad 1 \leq \ell \leq \frac{n-m}{2}. \end{aligned} \quad (1.3.33)$$

To prove this we fix $\omega \in \Omega$. We have to show that

$$\sum_{k=0}^{2\ell-1} (-1)^k \binom{m+k}{m} S_{m+k}(\omega) \leq \mathbf{I}_{\Omega_m}(\omega) \leq \sum_{k=0}^{2\ell} (-1)^k \binom{m+k}{m} S_{m+k}(\omega) \quad (1.3.34)$$

for $k \leq \frac{n-m}{2}$. Define

$$I_\omega := \{i \in \mathbb{I}_n; \omega \in A_i\}, \quad r(\omega) := |I_\omega| = \sum_{k=1}^n \mathbf{I}_{A_k}(\omega).$$

Note that $\mathbf{I}_{A_I}(\omega) = 0$ if $|I| > r(\omega)$. In particular, this shows that all the terms in the inequality (1.3.33) are equal to zero if $r(\omega) < m$.

Suppose that $r(\omega) \geq m$. Then, for any $k \leq r$, we have

$$S_k(\omega) = \sum_{\substack{I \subset I_\omega \\ |I|=k}} \mathbf{I}_{A_I}(\omega) = \binom{r}{k}.$$

Thus, the inequality (1.3.34) evaluated at ω is equivalent to

$$\sum_{k=0}^{2\ell-1} (-1)^k \binom{m+k}{m} \binom{r}{m+k} \leq \mathbf{I}_{\Omega_m}(\omega) \leq \sum_{k=0}^{2\ell} (-1)^k \binom{m+k}{m} \binom{r}{m+k}. \quad (1.3.35)$$

The inclusion-exclusion identity (1.3.28) shows that the inequalities become equalities for $2\ell > r - m$ so we assume $2\ell \leq r - m$.

For $r = m$ the inequality (1.3.35) is obvious since the sums in the left and right-hand sides consist of a single term equal to $1 = \mathbf{I}_{\Omega_m}(\omega)$. Assume $r > m$. In this case (1.3.35) is equivalent to

$$\sum_{k=0}^{2\ell-1} (-1)^k a_k \leq 0 \leq \sum_{k=0}^{2\ell} (-1)^k a_k, \quad a_k := \binom{m+k}{m} \binom{r}{m+k}. \quad (1.3.36)$$

Observe that

$$a_k = \binom{r}{m} \binom{p}{k}, \quad p = r - m.$$

The inequality (1.3.36) reduces to

$$\begin{aligned} & \binom{p}{0} - \binom{p}{1} + \cdots + \binom{p}{2\ell-2} - \binom{p}{2\ell-1} \leq 0 \\ & 0 \leq \binom{p}{0} - \binom{p}{1} + \binom{p}{2} + \cdots - \binom{p}{2\ell-1} + \binom{p}{2\ell}, \end{aligned}$$

where $2\ell \leq p$. These inequalities are immediate consequences of two well known properties of the binomial coefficients, namely their symmetry

$$\binom{p}{k} = \binom{p}{p-k},$$

and their *unimodality*

$$\binom{p}{0} \leq \binom{p}{1} \leq \cdots \leq \binom{p}{\lfloor p/2 \rfloor} = \binom{p}{\lfloor (p+1)/2 \rfloor} \geq \binom{p}{\lfloor p/2 \rfloor + 1} \geq \cdots \geq \binom{p}{p}.$$

For $m = 0$ we obtain the inequalities

$$\sum_{k=1}^n \mathbb{P}[A_k] - \sum_{1 \leq i < j \leq n} \mathbb{P}[A_i \cap A_j] \leq \mathbb{P}[A_1 \cup \cdots \cup A_n] \leq \sum_{k=1}^n \mathbb{P}[A_k].$$

The right-hand-side inequality is referred to as the *union bound*. □

Remark 1.3.33 (Binomial inversion). Consider the upper triangular matrices

$$A = (a_{\ell m})_{0 \leq \ell, m}, \quad a_{\ell m} = \begin{cases} (-1)^{m+\ell} \binom{m}{\ell}, & \ell \leq m, \\ 0, & \ell > m, \end{cases}$$

and

$$B = (b_{kn})_{0 \leq k \leq n}, \quad b_{kn} = \begin{cases} \binom{n}{k}, & k \leq n, \\ 0, & k > n. \end{cases}$$

The collections $((x-1)^m)_{m \geq 0}$ and $(x^n)_{n \geq 0}$ are bases of the space $\mathbb{R}[x]$ of polynomials with real coefficients. Newton's binomial formula implies

$$(x-1)^m = \sum_{\ell} a_{\ell m} x^{\ell}, \quad x^n = \sum_k b_{kn} (x-1)^k.$$

Hence $A^{-1} = B$, $B^{-1} = A$. This fact is known as *binomial inversion*. Note that (1.3.28) reads

$$\mathbb{P}[\Omega_{\ell}] = \sum_{m \geq \ell} a_{\ell m} s_m,$$

We deduce that

$$s_k = \sum_{m \geq k} b_{km} \mathbb{P}[\Omega_m].$$

We set $X = I_{A_1} + \cdots + I_{A_n}$. In Exercise 1.24 we ask the reader to prove that

$$s_k = \mathbb{E} \left[\binom{X}{k} \right].$$

□

Example 1.3.34 (Sieves and poissonization). Suppose now that we have an upper triangular array of measurable sets $(A_{n,i})_{i \in \mathbb{I}_n}$, $n \in \mathbb{N}$.

$$\begin{array}{ccccccc} & & A_{1,1} & & & & \\ & & A_{2,1}, & A_{2,2} & & & \\ & & \vdots & \vdots & & & \\ & & A_{n,1}, & A_{n,2}, & A_{n,3} & \cdots & A_{n,n} \\ & & \vdots & \vdots & \vdots & \vdots & \vdots \end{array}$$

For $n \geq q$ we denote by Ω_m^n the set of points in Ω that belong to exactly m of the sets $A_{n,1} \dots, A_{n,n}$, i.e., $\Omega_m^n = \{X_n = m\}$. Using Bonferroni's inequalities we deduce that for fixed ℓ and $n > 2\ell + m$ we have

$$\sum_{k=0}^{2\ell-1} (-1)^k \binom{m+k}{m} s_{m+k}^n \leq \mathbb{P}[\Omega_m^n] \leq \sum_{k=0}^{2\ell} (-1)^k \binom{m+k}{m} s_{m+k}^n. \quad (1.3.37)$$

Suppose now that there exists $\lambda > 0$ such that, for any $q \in \mathbb{N}$ we have

$$\lim_{n \rightarrow \infty} s_q^n = \frac{\lambda^q}{q!}. \quad (1.3.38)$$

If we let $n \rightarrow \infty$ in (1.3.37) we obtain

$$\frac{1}{m!} \sum_{k=0}^{2\ell-1} (-1)^k \frac{\lambda^k}{k!} \leq \liminf_{n \rightarrow \infty} \mathbb{P}[\Omega_m^n] \leq \limsup_{n \rightarrow \infty} \mathbb{P}[\Omega_m^n] \leq \frac{1}{m!} \sum_{k=0}^{2\ell} (-1)^k \frac{\lambda^k}{k!}.$$

If we now let $\ell \rightarrow \infty$ we deduce

$$\lim_{n \rightarrow \infty} \mathbb{P}[\Omega_m^n] = \frac{e^{-\lambda} \lambda^m}{m!}.$$

We can rephrase this in an equivalent way. Set

$$X_n := \sum_{k=1}^n I_{A_{n,k}}.$$

Then $\Omega_m^n = \{X_n = m\}$ and thus we showed that if (1.3.37) holds, then

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_n = m] = \mathbb{P}[\text{Poi}(\lambda) = m],$$

where we recall that $\text{Poi}(\lambda)$ denotes a Poisson random variable with parameter λ .

The phenomenon depicted above is referred under the generic name of *poissonization* or *Poisson approximation*. Let us observe that if the events $A_{n,k}$ are independent and $\mathbb{P}[A_{n,i}] = \frac{\lambda}{n}$, then

$$s_k^n = \binom{n}{k} \left(\frac{\lambda}{n} \right)^k \sim \frac{\lambda^k}{k!} \text{ as } n \rightarrow \infty.$$

In this case $X_n = \text{Bin}(n, \lambda/n)$. The success probability $\frac{\lambda}{n}$ is small for large n and for this reason the Poisson distribution is sometimes referred as the *law of rare events*.

The estimation techniques based on various versions of the inclusion-exclusion principle are called *sieves*. We refer to [197, Chap. 2, 3] for a more detailed description of far reaching generalizations of the inclusion-exclusion principle and associated sieves. $\square$

Example 1.3.35 (Fixed points of random permutations). Let us show how the above arguments work on the classical *derangements problem*. Denote by $\mathfrak{S}_n$ the group of permutations of $\mathbb{I}_n$. We equip it with the uniform probability measure so each permutation σ has probability $\frac{1}{n!}$. For each $\sigma \in \mathfrak{S}_n$ we denote by $F(\sigma) = F_n(\sigma)$ its number of fixed points, i.e.,

$$F(\sigma) = \#\{k \in \mathbb{I}_n; \sigma(k) = k\}.$$

Thus $F : \mathfrak{S}_n \rightarrow \{0, 1, \dots, n\}$ can be viewed as a random variable.

A *derangement* is a permutation σ with no fixed points, i.e., $F(\sigma) = 0$. A concrete occurrence of a derangement can be observed when a group of n , slightly inebriated, passengers board a plane and pick seats at random. A derangement occurs when none of them sits on his/her preassigned seat.

We want to compute the probability distribution of F_n , i.e., the probabilities

$$\mathbb{P}[F_n = m], \quad k = 0, 1, \dots, n.$$

For $j \in \mathbb{I}_n$ we denote by E_j the event $\sigma(j) = j$. The set of permutations that fix j can be identified with the set of permutations of $\mathbb{I}_n \setminus \{j\}$ so

$$\mathbb{P}[E_j] = \frac{(n-1)!}{n!} = \frac{1}{n}.$$

Observe that

$$F_n = \sum_{k=1}^n \mathbf{I}_{E_k},$$

so

$$\mathbb{E}[F_n] = \sum_{k=1}^n \mathbb{E}[\mathbf{I}_{E_k}] = \sum_{k=1}^n \mathbb{P}[E_k] = 1. \quad (1.3.39)$$

Thus the expected number of fixed points is rather low: a random permutation has, on average, one fixed point.

Let us compute the probability distribution of F . For each $I \subset \mathbb{I}_n$ we set

$$E_I = \bigcup_{i \in I} E_i.$$

Thus $\sigma \in E_I$ if and only if the permutation σ fixes all the points in I . We deduce that if $|I| = k$, then

$$\mathbb{P}[E_I] = \frac{(n-k)!}{n!} \text{ and } s_k = s_k^n := \sum_{|I|=k} \mathbb{P}[E_I] = \binom{n}{k} \frac{(n-k)!}{n!} = \frac{1}{k!}.$$

Note that if $F_n(\sigma) = m$, then σ fixes exactly k points and (1.3.28) yields

$$\mathbb{P}[F_n = m] = \sum_{k=0}^{n-m} (-1)^k \binom{m+k}{m} s_{m+k} = \frac{1}{m!} \sum_{k=0}^{n-m} (-1)^k \frac{1}{k!}.$$

In particular, the number of derangements is

$$\mathbb{P}[F_n = 0] = \sum_{k=0}^n (-1)^k \frac{1}{k!}.$$

The equality $\mathbb{E}[F_n] = 1$ yields an interesting identity

$$1 = \sum_{m=1}^n m \mathbb{P}[F_n = m] = \sum_{m=1}^n \frac{1}{(m-1)!} \left(\sum_{k=0}^{n-m} (-1)^k \frac{1}{k!} \right).$$

Note that

$$\lim_{n \rightarrow \infty} \mathbb{P}[F_n = m] = \frac{e^{-1}}{m!}. \quad (1.3.40)$$

The sequence $\frac{e^{-1}}{m!}$, $m \geq 0$ describes the Poisson distribution $\text{Poi}(1)$. $\square$

1.3.4. Classical examples of continuous probability distributions. We want to describe a few example of random variables whose probability distributions are absolutely continuous with respect to the Lebesgue measure on the real axis. They all have the form

$$\mathbb{P}[dx] = p(x) \lambda[dx], \quad p \in \mathcal{L}_+^1(\mathbb{R}, \mathcal{B}_{\mathbb{R}}, \lambda), \quad \int_{\mathbb{R}} p(x) \lambda[dx] = 1.$$

The function p is called the *probability density* of the Borel probability measure on $\mathbb{R}$. To ease the notational burden we will use the simpler notation

$$p(x)dx := p(x) \lambda[dx]$$

Such distributions are classically known as *continuous probability distributions*. The probabilistic significance of the examples discussed in this section will gradually be revealed in the book.

Example 1.3.36 (Uniform distribution). A random variable X is said to be *uniformly distributed* or *uniform* in the interval $[a, b]$, and we write this $X \sim \text{Unif}(a, b)$, if

$$\mathbb{P}_X[dx] = \frac{1}{b-a} \mathbf{I}_{[a,b]} dx.$$

When $X \sim \text{Unif}(a, b)$ we have

$$\mathbb{M}_X(t) = \frac{1}{b-a} \int_a^b e^{tx} dx = \frac{e^{tb} - e^{ta}}{t(b-a)} = \sum_{n \geq 1} \frac{b^n - a^n}{n(b-a)} \frac{t^{n-1}}{(n-1)!}.$$

In particular we deduce

$$\mu_n[X] = \frac{1}{n+1} \frac{b^{n+1} - a^{n+1}}{b-a}. \quad (1.3.41)$$

$\square$

Example 1.3.37 (Gaussian random variables). The *Gaussian* or *normal* random variables form a 2-parameter family $N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ where $X \sim N(\mu, \sigma^2)$ iff

$$\mathbb{P}_X[dx] = \gamma_{\mu, \sigma^2}(x) dx, \quad \gamma_{\mu, \sigma^2}(x) := \frac{1}{\sqrt{2\pi} \sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

We will use the simpler notation $\gamma_{\sigma^2}(x) := \gamma_{0, \sigma^2}$. The measure

$$\mathbf{\Gamma}_{\mu, \sigma^2}[dx] := \gamma_{\mu, \sigma^2}(x) dx$$




Figure 1.4. The graph of the Gauss bell $\gamma_{0,\sigma}$ for $\sigma = 1$ (dotted red curve) and $\sigma = 0.1$ (continuous blue curve). Note that sharper profiles correspond to smaller standard deviations.

is called the *Gaussian measure* on $\mathbb{R}$ with mean μ and variance σ^2 . Let us observe

$$X \sim N(\mu, \sigma^2) \iff \frac{1}{\sigma}(X - \mu) \sim N(0, 1).$$

Indeed if we set

$$Y := \frac{1}{\sigma}(X - \mu),$$

then

$$\begin{aligned} \mathbb{P}[Y \leq y] &= \mathbb{P}[(x - \mu)/\sigma \leq y] = \mathbb{P}[x \leq \sigma y + \mu] = \int_{-\infty}^{\sigma y + \mu} \gamma_{\mu, \sigma^2}(x) dx \\ &= \sigma \int_{-\infty}^y \gamma_{\mu, \sigma^2}(\sigma t + \mu) dt = \int_{-\infty}^y \gamma_{0,1}(t) dt. \end{aligned}$$

Thus

$$\mathbb{E}[X] = \mathbb{E}[Y] + \mu, \quad \text{Var}[X] = \sigma^2 \text{Var}[Y].$$

We have

$$\mathbb{E}[Y] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} y e^{-y^2/2} dy = 0,$$

and

$$\text{Var}[Y] = \mathbb{E}[Y^2] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} y^2 e^{-y^2/2} dy = \frac{2}{\sqrt{2\pi}} \int_0^{\infty} y^2 e^{-y^2/2} dy$$

$$(s = y^2/2, y = \sqrt{2s})$$

$$= \frac{2}{\sqrt{\pi}} \int_0^{\infty} s^{1/2} e^{-s} ds = \frac{2}{\sqrt{\pi}} \Gamma(3/2) = \frac{2}{\sqrt{\pi}} \cdot \frac{1}{2} \Gamma(1/2) = 1,$$

where at the last two steps we used basic facts about the Gamma function recalled in Proposition A.1.2. We deduce that

$$X \sim N(\mu, \sigma^2) \Rightarrow \mathbb{E}[X] = \mu, \quad \text{Var}[Y] = \sigma^2. \quad (1.3.42)$$

A variable $X \sim N(0, 1)$ is called a *standard normal* random variable. Its cdf is

$$\Phi(x) := \mathbb{P}[X \leq x] = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-x^2/2} dx, \quad (1.3.43)$$

plays an important role in probability and statistics. The quantity

$$\frac{\mathbb{P}[X > x]}{\gamma_1(x)}$$

is called the *Mills ratio* of the standard normal random variable. It satisfies the inequalities

$$\frac{x}{x^2 + 1} \gamma_1(x) \leq \mathbb{P}[X > x] \leq \frac{1}{x} \gamma_1(x). \quad (1.3.44)$$

In Exercise 1.34 we outline a proof of this inequality.

Observe that if $X \sim N(0, 1)$, and $\sigma \in \mathbb{R}$, then $\sigma X \in N(0, \sigma^2)$ and

$$\mathbb{M}_{\sigma X}(t) = \mathbb{E}[e^{t\sigma X}] = \mathbb{M}_X[\sigma t].$$

On the other hand, if $X \sim N(0, 1)$, then

$$\begin{aligned} \mathbb{M}_X(t) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{tx - x^2/2} dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{(2tx - x^2 - t^2)/2} e^{t^2/2} dx \\ &= e^{t^2/2} \cdot \underbrace{\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-(x-t)^2/2} dx}_{=1} = e^{t^2/2}. \end{aligned} \quad (1.3.45)$$

Thus

$$\mu_{2m}[X] = \frac{(2m)!}{2^m m!} = (2m-1)!!, \quad \mu_{2m-1}[X] = 0, \quad \forall m \in \mathbb{N}. \quad \square$$

Example 1.3.38 (Gamma distributions). The *Gamma distributions* with parameters ν, λ are defined by

$$\Gamma_{\nu, \lambda}[dx] = g_{\nu}(x; \lambda) dx.$$

where the densities $g_{\nu}(x; \lambda)$, $\lambda, \nu > 0$ are given by

$$g_{\nu}(x; \lambda) = \frac{\lambda^{\nu}}{\Gamma(\nu)} x^{\nu-1} e^{-\lambda x} \mathbf{I}_{(0, \infty)}. \quad (1.3.46)$$

From the definition of the Gamma function we deduce that $g_{\nu}(x; \lambda)$ is indeed a probability density, i.e.,

$$\int_0^{\infty} g_{\nu}(x; \lambda) dx = 1.$$

We will use the notation $X \sim \text{Gamma}(\nu, \lambda)$ to indicate that $\mathbb{P}_X = \Gamma_{\nu, \lambda}$.

The $\text{Gamma}(1, \lambda)$ -random variables play a special role in probability. They are called *exponential random variables* with parameter λ . We will use the notation $X \sim \text{Exp}(\lambda)$ to indicate that X is such a random variable. The distribution of $\text{Exp}(\lambda)$ is

$$\text{Exp}(\lambda) \sim \lambda e^{-\lambda x} \mathbf{I}_{(0, \infty)} dx.$$

We will have more to say about exponential variables in the next subsection.

The parameter ν is sometimes referred to as the *shape* parameter. Figure 1.5 may explain the reason for this terminology.

For $n = 1, 2, 3, \dots$ the distribution $\text{Gamma}(n, \lambda)$ is also known as an *Erlang distribution* and has a simple probabilistic interpretation. If the waiting time T for a certain event is exponentially distributed with rate λ , e.g., the waiting time for a bus to arrive, then the waiting time for n of these events to occur independently and in succession is a $\text{Gamma}(n, \lambda)$ random variable. We will prove this later.




Figure 1.5. The graphs of $g_\nu(x; \lambda)$ for $\nu > 1$ and $\nu < 1$. Note that the graph explodes at the origin if $\nu < 1$.

The distribution $g_{n/2}(x; 1/2)$, where $n = 1, 2, \dots$, plays an important role in statistics it also known as the *chi-squared distribution with n degrees of freedom* and it is traditionally denoted by $\chi^2(n)$. One can show that if $X_1, \dots, X_n$ are independent standard normal random variables, then the random variable

$$X_1^2 + \dots + X_n^2$$

has a chi-squared distribution of degree n .

If $X \sim \text{Gamma}(\nu, \lambda)$ is a Gamma distributed random variable, then X is s -integrable for any $s \geq 1$. Moreover, for any $k \in \{1, 2, \dots\}$ we have

$$\begin{aligned} \mu_k[X] &= \frac{\lambda^\nu}{\Gamma(\nu)} \int_0^\infty x^{k+\nu-1} e^{-\lambda x} dx \\ (x = \lambda^{-1}t, dx = \lambda^{-1}dt, \lambda x = t, x^{k+\nu-1} &= \lambda^{-(k+\nu-1)}t^{k+\nu-1}) \\ &= \frac{1}{\lambda^k \Gamma(\nu)} \int_0^\infty t^{k+\nu-1} e^{-t} dt = \frac{\Gamma(k+\nu)}{\lambda^k \Gamma(\nu)}. \end{aligned}$$

We deduce

$$\begin{aligned} \mathbb{E}[X] &= \mu_1[X] = \frac{\Gamma(\nu+1)}{\lambda \Gamma(\nu)} = \frac{\nu}{\lambda}, \\ \text{Var}[X] &= \mu_2[X] - \mu_1[X]^2 = \frac{\Gamma(\nu+2)}{\lambda^2 \Gamma(\nu)} - \frac{\nu^2}{\lambda^2} = \frac{\nu(\nu+1) - \nu^2}{\lambda^2} = \frac{\nu}{\lambda^2}. \end{aligned}$$

Finally, if $X \sim \text{Gamma}(\nu, \lambda)$, then for $t < \lambda$ we have

$$\begin{aligned} \mathbb{M}_X(t) &= \frac{\lambda^\nu}{\Gamma(\nu)} \int_0^\infty x^{\nu-1} e^{-(\lambda-t)x} dx \\ x = y/(\lambda-t) & \\ &= \frac{\lambda^\nu}{\Gamma(\nu)(\lambda-t)^\nu} \int_0^\infty y^{\nu-1} e^{-y} dy = \left(\frac{\lambda}{\lambda-t} \right)^\nu. \end{aligned}$$

□

Example 1.3.39 (Beta distributions). The *Beta distribution* with parameters $a, b > 0$ is defined by the probability density function

$$\beta_{a,b}(x) = \frac{1}{B(a,b)} x^{a-1} (1-x)^{b-1} \mathbf{I}_{(0,1)}.$$

The normalizing constant $B(a, b)$ is the *Beta function* (A.1.2),

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

We will use the notation $X \sim \text{Beta}(a, b)$ to indicate that the pdf of X is a Beta distribution with parameters a, b .

Suppose that $X \sim \text{Beta}(a, b)$. Then

$$\begin{aligned} \mathbb{E}[X] &= \frac{1}{B(a,b)} \int_0^1 x^a (1-x)^{b-1} dx = \frac{B(a+1, b)}{B(a, b)} \\ &\stackrel{(A.1.4)}{=} \frac{\Gamma(a+1)\Gamma(a+b)}{\Gamma(a)\Gamma(a+b+1)} = \frac{a}{a+b}, \\ \mathbb{E}[X^2] &= \frac{1}{B(a,b)} \int_0^1 x^{a+1} (1-x)^{b-1} dx = \frac{\Gamma(a+2)\Gamma(a+b)}{\Gamma(a)\Gamma(a+b+2)} = \frac{a(a+1)}{(a+b)(a+b+1)}. \end{aligned}$$

Hence

$$\begin{aligned} \text{Var}[X] &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{a}{a+b} \left(\frac{a+1}{a+b+1} - \frac{a}{a+b} \right) \\ &= \frac{a}{a+b} \cdot \frac{(a+1)(a+b) - a(a+b+1)}{(a+b)(a+b+1)} = \frac{ab}{(a+b)^2(a+b+1)}. \end{aligned}$$

Note that $\text{Beta}(1, 1) = \text{Unif}([0, 1])$. The distribution $\text{Beta}(1/2, 1/2)$ is called the *arcsine distribution*. In this case

$$\beta_{1/2, 1/2}(x) = \frac{1}{\pi} \frac{1}{\sqrt{x(1-x)}},$$

and

$$\int_0^x \beta_{1/2, 1/2}(s) ds = \frac{2}{\pi} \arcsin \sqrt{x}.$$

We refer to Exercise 1.45 for an alternate interpretation of $\text{Beta}(1/2, 1/2)$. □

In Appendix A.2 we have listed the basic integral invariants of several frequently occurring probability distributions.

1.3.5. Product probability spaces and independence. Suppose $(\Omega_i, \mathcal{S}_i)$, $i = 0, 1$, are two measurable spaces. Recall that $\mathcal{S}_0 \otimes \mathcal{S}_1$ is the sigma-algebra of subsets of $\Omega_0 \times \Omega_1$ generated by the collection $\mathcal{R}$ of “rectangles” of the form $S_0 \times S_1$, $S_i \in \mathcal{S}_i$, $i = 0, 1$.

The goal of this subsection is to show that two sigma-finite measures μ_i on $\mathcal{S}_i$, $i = 0, 1$ induce in a canonical way a measure $\mu_0 \otimes \mu_1$ uniquely determined by the condition

$$\mu_0 \otimes \mu_1[S_0 \times S_1] = \mu_0[S_0] \mu_1[S_1], \quad \forall S_i \in \mathcal{S}_i, \quad i = 0, 1.$$

The collection $\mathcal{A}$ of subsets of $\Omega_0 \times \Omega_1$ that are finite disjoint unions of rectangles is an algebra. This suggests using Carathéodory’s existence theorem to prove this claim.

We choose a different route that bypasses Carathéodory’s existence theorem. This alternate, more efficient approach, is driven by the Monotone Class Theorem and simultaneously proves a

central result in integration theory, the Fubini-Tonelli Theorem. For every measurable space $(\Omega, \mathcal{S})$ we denote by $\mathcal{L}^0(\Omega, \mathcal{S})_*$ the space of $\mathcal{S}$ measurable functions $f : \Omega \rightarrow \mathbb{R}$.

Lemma 1.3.40. *Suppose that*

$$f \in \mathcal{L}^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)_* \cup \mathcal{L}_+^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1).$$

Then, for any $\omega_1 \in \Omega_1$ the function $f_{\omega_1}^0 : \Omega_0 \rightarrow \mathbb{R}$,

$$f_{\omega_1}^0(\omega_0) = f(\omega_0, \omega_1)$$

is $\mathcal{S}_0$ -measurable and, for any $\omega_0 \in \Omega_0$, the function $f_{\omega_0}^1 : (\Omega_1, \mathcal{S}_1) \rightarrow \mathbb{R}$,

$$f_{\omega_0}^1(\omega_1) = f(\omega_0, \omega_1)$$

is $\mathcal{S}_1$ -measurable.

Proof. We prove only the statement concerning $f_{\omega_1}^0$. For simplicity will write f_{ω_1} instead of $f_{\omega_1}^0$. We will use the Monotone Class Theorem 1.1.22.

Denote by $\mathcal{M}$ the collection of functions $f \in \mathcal{L}^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \times \mathcal{S}_1)_*$ such that f_{ω_1} is $\mathcal{S}_0$ -measurable, $\forall \omega_1 \in \Omega_1$. Clearly is $f, g \in \mathcal{M}$ are bounded then $af + bg \in \mathcal{M}$, $\forall a, b \in \mathbb{R}$

The collection $\mathcal{R}$ of rectangles is a π -system. Note that for any rectangle $R = S_0 \times S_1$ the function $f = \mathbf{I}_R$ belongs to $\mathcal{M}$. Indeed, for any $\omega_1 \in \Omega_1$ we have

$$f_{\omega_1} = \begin{cases} \mathbf{I}_{S_0}, & \omega_1 \in S_1, \\ 0, & \omega_1 \in \Omega_1 \setminus S_1. \end{cases}$$

If (f_n) is an increasing sequence of functions in $\mathcal{M}$ so is the sequence of slices f_{n, ω_1} so the limit f is also in $\mathcal{M}$. By the Monotone Class Theorem the collection $\mathcal{M}$ contains all the nonnegative measurable functions. Since $\mathcal{M}$ is a vector space, it must coincide with $\mathcal{L}^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)_*$.

When $f \in \mathcal{L}_+^0$, but f is allowed to have infinite values, the function f is the increasing limit of a sequence in $\mathcal{M}$. Hence this situation is also included in the conclusions of the lemma. $\square$

Theorem 1.3.41 (Fubini-Tonelli). *Let $(\Omega_i, \mathcal{S}_i, \mu_i)$, $i = 0, 1$ be two sigma-finite measured spaces.*

(i) *There exists a measure μ on $\mathcal{S}_0 \otimes \mathcal{S}_1$ uniquely determined by the equalities*

$$\mu[S_0 \times S_1] = \mu_0[S_0] \mu_1[S_1], \quad \forall S_0 \in \mathcal{S}_0, S_1 \in \mathcal{S}_1.$$

We will denote this measure by $\mu_0 \otimes \mu_1$.

(ii) *For each nonnegative function $f \in \mathcal{L}_+^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$ the functions*

$$\omega_0 \mapsto \mathbf{I}_1[f](\omega_0) := \int_{\Omega_1} f(\omega_0, \omega_1) \mu_1[d\omega_1] \in [0, \infty],$$

$$\omega_1 \mapsto \mathbf{I}_0[f](\omega_1) := \int_{\Omega_0} f(\omega_0, \omega_1) \mu_0[d\omega_0] \in [0, \infty]$$

are measurable and

$$\begin{aligned} & \int_{\Omega_0} \left(\int_{\Omega_1} f(\omega_0, \omega_1) \mu_1[d\omega_1] \right) \mu_0[d\omega_0] \\ &= \int_{\Omega_0 \times \Omega_1} f(\omega_0, \omega_1) \mu_0 \otimes \mu_1[d\omega_0 d\omega_1] \\ &= \int_{\Omega_1} \left(\int_{\Omega_0} f(\omega_0, \omega_1) \mu_0[d\omega_0] \right) \mu_1[d\omega_1]. \end{aligned} \tag{1.3.47}$$

In particular, if only one of the three terms above is finite, then all three are finite and equal.

(iii) *Let $f \in \mathcal{L}^1(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1, \mu_0 \otimes \mu_1)$. Then each of the three terms in (1.3.47) is well defined, finite and the equalities (1.3.47) hold.*

Proof. We will carry the proof in several steps.

Step 1. We will prove that for every positive function $f \in \mathcal{L}^0(\Omega_0 \times \Omega_1, \mathcal{S}_0 \times \mathcal{S}_1)$ the nonnegative function

$$\omega_0 \mapsto \mathbf{I}_1[f](\omega_0) = \int_{\Omega_1} f(\omega_0, \omega_1) \mu_1[d\omega_1]$$

is measurable so the integral

$$I_{1,0}[f] := \int_{\Omega_0} \left(\int_{\Omega_1} f(\omega_0, \omega_1) \mu_1[d\omega_1] \right) \mu_0[d\omega_0] \in [0, \infty]$$

is well defined.

This follows from Dynkin's $\pi - \lambda$ Theorem arguing exactly as in the proof of Lemma 1.3.40. For $S \in \mathcal{S}_0 \otimes \mathcal{S}_1$ we set

$$\mu_{1,0}[S] = I_{1,0}[\mathbf{I}_S].$$

Note that

$$\mathbf{I}_1[\mathbf{I}_{S_0 \times S_1}] = \int_{\Omega_1} \mathbf{I}_{\Omega_0 \times \Omega_1}(\omega_0, \omega_1) \mu_1[d\omega_1].$$

If $\omega_0 \in \Omega_0 \setminus S_0$ the integral is 0. If $\omega_0 \in S_0$ the integral is

$$\int_{\Omega_1} \mathbf{I}_{S_1} d\mu_1 = \mu_1[S_1].$$

Hence

$$\mathbf{I}_1[\mathbf{I}_{S_0 \times S_1}] = \mu_1[S_1] \mathbf{I}_{S_0}.$$

We deduce

$$I_{1,0}[S_0 \times S_1] = \mu_1[S_1] \int_{\Omega_0} \mathbf{I}_{S_0} d\mu_0 = \mu_0[S_0] \cdot \mu_1[S_1].$$

Clearly if $A, A' \in \mathcal{S}$ are disjoint, then $\mathbf{I}_{A \cup A'} = \mathbf{I}_A + \mathbf{I}_{A'}$ so that

$$I_{1,0}[\mathbf{I}_{A \cup A'}] = I_{1,0}[\mathbf{I}_A] + I_{1,0}[\mathbf{I}_{A'}]$$

and

$$\mu_{1,0}[A \cup A'] = \mu_{1,0}[A] + \mu_{1,0}[A'].$$

If

$$A_1 \subset A_2 \subset \dots$$

is an increasing sequence of sets in $\mathcal{S}$ and

$$A = \bigcup_{n \geq 1} A_n,$$

then invoking the Monotone Convergence Theorem we first deduce that $I_{1,0}[\mathbf{I}_{A_n}]$ is a nondecreasing sequence of measurable functions converging to $I_{1,0}[\mathbf{I}_A]$ and then we conclude that $\mu_{1,0}[A_n]$ converges to $\mu_{1,0}[A]$. Hence $\mu_{1,0}$ is a measure on $\mathcal{S} = \mathcal{S}_0 \otimes \mathcal{S}_1$.

Step 2. A similar argument shows that

$$\mu_{0,1}[S] = \int_{\Omega_1} \left(\int_{\Omega_0} \mathbf{I}_S(\omega_0, \omega_1) \mu_0[d\omega_0] \right) \mu_1[d\omega_1]$$

is also a sigma-finite measure on $\mathcal{S} = \mathcal{S}_0 \otimes \mathcal{S}_1$. Note that

$$\mu_{1,0}[S_0 \times S_1] = \mu_{0,1}[S_0 \times S_1], \quad \forall S_0 \in \mathcal{S}_0, \quad S_1 \in \mathcal{S}_1.$$

Thus $\mu_{1,0}[R] = \mu_{0,1}[R], \forall R \in \mathcal{R}$.

We want to show that if ν is another measure on $\mathcal{S}$ such that $\nu[R] = \mu_{1,0}[R]$ for any $R \in \mathcal{R}$, then $\nu[A] = \mu_{1,0}[A], \forall A \in \mathcal{S}$.

To see this assume first that μ_0 and μ_1 are finite measures. Then $\Omega_0 \times \Omega_1 \in \mathcal{R}$

$$\mu_{1,0}[\Omega_0 \times \Omega_1] = \nu[\Omega_0 \times \Omega_1] < \infty$$

and since $\mathcal{R}$ is a π -system we deduce from Proposition 1.2.4 that $\mu_{1,0} = \nu$ on $\mathcal{S}$.

To deal with the general case choose two increasing sequences $E_n^i \in \mathcal{S}_i, i = 0, 1$ such that

$$\mu_i[E_n^i] < \infty, \quad \forall n \quad \text{and} \quad \Omega_i = \bigcup_{n \geq 1} E_n^i, \quad i = 0, 1.$$

Define

$$E_n := E_n^0 \times E_n^1, \quad \mu_i^n[S_i] := \mu_i[S_i \cap E_n^i], \quad S_i \in \mathcal{S}_i, \quad i = 0, 1, \\ \nu^n[A] := \nu[A \cap E_n], \quad \forall A \in \mathcal{S}.$$

Using the measures μ_i^n we form as above the measures $\mu_{1,0}^n$ and we observe that

$$\mu_{1,0}^n[A] = \mu_{0,1}[A \cap E_n], \quad \forall n, \quad \forall A \in \mathcal{S}.$$

For any rectangle R , the intersection $R \cap E_n$ is a rectangle and

$$\mu_{1,0}^n[R] = \nu^n[R], \quad \forall n.$$

Thus

$$\mu_{1,0}^n[A] = \mu^n[A], \quad \forall n \in \mathbb{N}, \quad A \in \mathcal{S}.$$

If we let $n \rightarrow \infty$ in the above equality we deduce that $\mu_{1,0} = \nu$ on $\mathcal{S}$.

We deduce that $\mu_{0,1} = \mu_{1,0}$. Thus the measures $\mu_{0,1}$ and $\mu_{1,0}$ coincide on the algebra of sets generated by the rectangles and thus they must coincide on the $\mathcal{S}_0 \otimes \mathcal{S}_1$. This common measure is denoted by $\mu_0 \otimes \mu_1$ and it clearly satisfies statement (i) in the theorem

Step 3. From **Step 2** we deduce that (1.3.47) is true for $f = I_S, \forall S \in \mathcal{S}_0 \otimes \mathcal{S}_1$. From this, using the Monotone Class Theorem exactly as in the proof of Lemma 1.3.40 we deduce (1.3.47) in its entire generality. The claim in (iii) follows from the fact that any integrable function f is the difference of two nonnegative integrable functions $f = f^+ - f^-$ and the claim is true for $f^\pm$. $\square$

The above construction can be iterated. More precisely, given sigma-finite measured spaces $(\Omega_k, \mathcal{S}_k, \mu_k), k = 1, \dots, n$, we have a measure $\mu = \mu_1 \otimes \dots \otimes \mu_n$ uniquely determined by the condition

$$\mu[S_1 \times S_2 \times \dots \times S_n] = \mu_1[S_1] \mu_2[S_2] \dots \mu_n[S_n], \quad \forall S_k \in \mathcal{S}_k, \quad k = 1, \dots, n.$$

Remark 1.3.42. Recall that λ denotes the Lebesgue measure on $\mathbb{R}$. The measure $\lambda^{\otimes n}$ on $\mathcal{B}_{\mathbb{R}^n}$ is called the n -dimensional Lebesgue measure and will be denoted by λ_n or simply λ , when no confusion is possible. A subset of $\mathbb{R}^n$ is called *Lebesgue measurable* if it belongs to the completion of the Borel sigma-algebra with respect to the Lebesgue measure.

One can prove that if a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is absolutely Riemann integrable (see [158, Chap.15]), then it is also Lebesgue integrable with respect to the Lebesgue measure on $\mathbb{R}^n$ and, moreover

$$\int_{\mathbb{R}^n} f(x) |dx| = \int_{\mathbb{R}^n} f(x) \lambda[dx],$$

where the left-hand-side integral is the (improper) Riemann integral.

We recommend the reader to try to prove this fact or at least to try to understand why a Riemann integrable function defined on a cube is Lebesgue measurable. This is not obvious because there exist Riemann integrable functions that are not Borel measurable.

For example, if $C \subset [0, 1]$ is the Cantor set, then there exists a subset A of C that are not Borel because the cardinality of the set 2^C is bigger than the cardinality of the family of Borel subsets of C . The subset A is Lebesgue measurable since C is Lebesgue negligible. The indicator function I_A is Riemann integrable but not Borel measurable.

The change in variables for the Riemann integral shows that if U, V are open subsets of $\mathbb{R}^n$ and $F : U \rightarrow V$ is a C^1 -diffeomorphism onto V , then

$$F_{\#}^{-1} \lambda_V[dx] = |\det J_F(x)| \lambda_U[dx].$$

$\square$

Let us present a few useful consequences of Fubini's theorem.

Proposition 1.3.43. Suppose that X is a nonnegative random variable defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For any $p \in [1, \infty)$ we have

$$\mathbb{E}[X^p] = p \int_0^\infty x^{p-1} \mathbb{P}[X > x] dx. \quad (1.3.48)$$

In particular,

$$\mathbb{E}[X] = \int_0^\infty \mathbb{P}[X > x] dx. \quad (1.3.49)$$

Proof. We have

$$\begin{aligned} p \int_0^\infty x^{p-1} \mathbb{P}[X > x] dx &= \int_0^\infty \left(\int_\Omega \mathbf{I}_{\{X > x\}}(\omega) \mathbb{P}[d\omega] \right) p x^{p-1} dx \\ &= \int_{\substack{(\omega, x) \in \Omega \times [0, \infty) \\ 0 \leq x < X(\omega)}} p x^{p-1} \mathbb{P} \otimes \lambda[d\omega dx] \end{aligned}$$

(use Fubini-Tonelli)

$$= \int_\Omega \left(\int_0^{X(\omega)} p x^{p-1} dx \right) \mathbb{P}[d\omega] = \int_\Omega X^p(\omega) \mathbb{P}[d\omega] = \mathbb{E}[X^p].$$

□

We want to point out that when $p = 1$ the equality

$$\int_{\substack{(\omega, x) \in \Omega \times [0, \infty) \\ 0 \leq x < X(\omega)}} \mathbb{P} \otimes \lambda[d\omega dx] = \mathbb{E}[X]$$

simply says that $\mathbb{E}[X]$ is equal to the “area” below the graph of the function $X : \Omega \rightarrow [0, \infty)$.

Example 1.3.44. Suppose that X is a random variable that takes only nonnegative *integral* values. Then

$$\mathbb{P}_X = \sum_{n \geq 0} \mathbb{P}[X = n] \delta_n,$$

and

$$\begin{aligned} \mathbb{E}[X] &\stackrel{(1.3.48)}{=} \int_0^\infty \mathbb{P}[X > x] dx \\ &= \sum_{n \geq 0} \int_n^{n+1} \mathbb{P}[X > x] dx = \sum_{n \geq 0} \mathbb{P}[X > n]. \end{aligned} \quad (1.3.50)$$

Let us apply this identity to a geometric random variable with success probability p , $T \sim \text{Geom}(p)$. Note that $\mathbb{P}[T > n]$ is the probability that the waiting time for a success is $> n$ or, equivalently, the probability that the first n trials are failures. Hence

$$\mathbb{P}[T > n] = q^n \text{ so } \mathbb{E}[T] = \sum_{n \geq 0} q^n = \frac{1}{1 - q} = \frac{1}{p}.$$

Similarly

$$\begin{aligned} \mu_2[T] &= \mathbb{E}[T^2] = 2 \sum_{n \geq 0} n \mathbb{P}[T > n] \\ &= 2 \sum_{n \geq 1} n q^n = 2q \sum_{n \geq 1} n q^{n-1} = \frac{2q}{(1 - q)^2} = \frac{2q}{p^2}. \end{aligned}$$

In particular

$$\text{Var}[T] = \mathbb{E}[T^2] - \mathbb{E}[T]^2 = \frac{q}{p^2}.$$

□

Example 1.3.45. Suppose that T is an *exponential random variable with parameter λ* , i.e., a random variable with the exponential probability distribution

$$\mathbb{P}_T[dt] = \lambda e^{-\lambda t} \mathbf{I}_{(0,\infty)} dt$$

This random variable describes the waiting time for an event to happen, e.g., the waiting time for a laptop to crash, or the waiting time for a bus to arrive at a bus station. The quantity $\lambda e^{-\lambda t} dt$ is the probability that the waiting time is in the interval $(t, t + dt]$. Then

$$\mathbb{P}[T > t] = \int_t^\infty \lambda e^{-\lambda \tau} d\tau = e^{-\lambda t}, \quad \mathbb{E}[T] = \int_0^\infty e^{-\lambda t} dt = \frac{1}{\lambda}.$$

We see that $\frac{1}{\lambda}$ is measured in units of time. For this reason λ is called the *rate* and describes how many rare events take place per unit of time.

Similarly

$$\begin{aligned} \mu_2[T] = \mathbb{E}[T^2] &= 2 \int_0^\infty t \mathbb{P}[T > t] dt = 2 \int_0^\infty t e^{-\lambda t} dt = \frac{2}{\lambda^2} \int_0^\infty s e^{-s} ds \\ &= \frac{2}{\lambda^2} \Gamma(2) = \frac{2}{\lambda^2}. \end{aligned}$$

The function $S(t) := \mathbb{P}[T > t]$ is called the *survival function*. For example, if T denotes the life span of a laptop, then $S(t)$ is the probability that a laptop survives more than t units of time.

The exponential distribution enjoys the so called *memoryless property*

$$\mathbb{P}[T > t + s | T > s] = \mathbb{P}[T > t]. \quad (1.3.51)$$

For example, if T is the waiting time for a bus to arrive then, given that you've waited more than s units of time, the probability that you will have to wait at least t extra is the same as if you have not waited at all. The proof of (1.3.51) is immediate.

$$\mathbb{P}[T > t + s | T > s] = \frac{\mathbb{P}[T > t + s]}{\mathbb{P}[T > s]} = \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} = e^{-\lambda t} = \mathbb{P}[T > t]. \quad \square$$

Example 1.3.46 (Integration by parts). Suppose that μ_0, μ_1 are two Borel probability measures on $\mathbb{R}$ supported on $[0, \infty)$, i.e.

$$\mu_k[(-\infty, 0)] = 0, \quad k = 0, 1.$$

We set

$$F_k(x) = \mu_k[(-\infty, x)], \quad k = 0, 1,$$

so that μ_k is the Lebesgue-Stieltjes measure determined by F_k . Note that

$$F_k(0) = \mu_k[\{0\}].$$

Classically, the integral

$$\int_{[0,a]} u(x) \mu_k[dx]$$

was denoted by

$$\int_0^a u(x) dF_k(x).$$

This classical notation is a bit ambiguous due to the following simple fact

$$\int_{[0,a]} u(x) \mu_k[dx] = u(0)F_k(0) + \int_{(0,a]} u(x) \mu_k[dx].$$

We want to prove a version of the integration-by-parts formula. Namely, we will show that if one of the functions F_0, F_1 is continuous, then

$$\int_0^a F_0(x) dF_1(x) = F_0(a)F_1(a) - F_0(0)F_1(0) - \int_0^a F_1(x) dF_0(x). \quad (1.3.52)$$

Assume for simplicity that F_1 is continuous so $F_1(0) = 0$. Set $\mu := \mu_0 \otimes \mu_1$. Observe that since

$$F_0(a)F_1(a) - F_0(0)F_1(0) = F_0(a)F_1(a) = \mu \left[\underbrace{[0, a] \times [0, a]}_{S_a} \right].$$

Using the Fubini-Tonelli theorem we deduce

$$\int_0^a F_1(x) dF_1(x) = \int_{[0, a]} \left(\int_{\mathbb{R}} \mathbf{I}_{(-\infty, x]}(y) \mu_1[dy] \right) \mu_0[dx]$$

(F_1 is continuous)

$$= \int_{[0, a]} \left(\int_{[0, a]} \mathbf{I}_{[0, x]}(y) \mu_1[dy] \right) \mu_0[dx] = \mu[R_0],$$

where

$$R_0 := \{ (x, y) \in \mathbb{R}^2; \ 0 \leq y < x \leq a, \ y < x \}.$$

Similarly

$$\int_0^a F_0(y) dF_1(y) = \int_{[0, a]} \left(\int_{[0, a]} \mathbf{I}_{[0, y]} \mu_0[dx] \right) \mu_1[dy] = \mu[R_1],$$

$$R_1 := \{ (x, y) \in \mathbb{R}^2; \ 0 \leq x \leq y \leq a \},$$

Observe that the regions R_0, R_1 are disjoint.

The region R_0 is the part of the square $S_a = [0, a] \times [0, a]$ strictly below the diagonal $y = x$, while R_1 is the part of this square above or this diagonal. Hence $S_a = R_0 \cup R_1$ and thus

$$\mu[R_0] + \mu[R_1] = \mu[S_a].$$

Let us observe that the integration by parts formula is not true if both F_0, F_1 are discontinuous. Take for example the case $\mu_0 = \mu_1 = \frac{1}{2}(\delta_1 + \delta_3)$. Then

$$F_0(x) = F_1(x) = F(x) = \begin{cases} 0, & x < 1. \\ \frac{1}{2}, & 1 \leq x < 3, \\ 1, & x \geq 3. \end{cases}$$

In this case we have

$$\int_0^2 F(x) dF(x) = \int_{[0, 2]} F(x) \mu_0[dx] = \frac{1}{2}F(1) = \frac{1}{4}, \quad F(2)^2 = \frac{1}{4}.$$

so

$$2 \int_0^2 F(x) dF(x) \neq F(2)^2.$$

The reason for this failure has a simple geometric origin: the diagonal $\{y = x\}$ may not be $\mu_0 \otimes \mu_1$ -negligible. The continuity assumption allowed us to discard the diagonal of the square because in this case it is indeed negligible. $\square$

Definition 1.3.47. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$.

- (i) Suppose that V is a finite dimensional vector space. We denote by $\mathcal{B}_V$ the sigma-algebra of Borel subsets of V . A V -valued *random vector* is a measurable map

$$\mathbf{X} : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (V, \mathcal{B}_V).$$

Its *probability distribution* is the pushforward measure $\mathbb{P}_{\mathbf{X}} := \mathbf{X}_{\#}\mathbb{P}$. By definition, $\mathbb{P}_{\mathbf{X}}$ is a Borel probability measure on V .

- (ii) The *joint probability distribution* of the random variables

$$X_1, \dots, X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$$

is the probability distribution of the random vector

$$\mathbf{X} := (X_1, \dots, X_n) : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}^n.$$

We will denote by $\mathbb{P}_{X_1, \dots, X_n}$ the joint distribution.

□

Observe that joint probability distribution $\mathbb{P}_{X_1, \dots, X_n}$ is uniquely determined by the probabilities

$$\mathbb{P}[X_1 \leq x_1, \dots, X_n \leq x_n], \quad x_1, \dots, x_n \in \mathbb{R}.$$

Note also that if $\pi_i : \mathbb{R}^n \rightarrow \mathbb{R}$ denotes the natural projection $(x_1, \dots, x_n) \mapsto x_i, i = 1, 2, \dots, n$, then

$$\mathbb{P}_{X_i} = (\pi_i)_{\#}\mathbb{P}_{X_1, \dots, X_n}.$$

The probability distributions $\mathbb{P}_{X_i}$ are often referred as the *marginals* (or *marginal distributions*) of the joint probability distribution $\mathbb{P}_{X_1, \dots, X_n}$.

Proposition 1.3.48. *Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and*

$$X_1, \dots, X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$$

are random variables with probability distributions $\mathbb{P}_{X_1}, \dots, \mathbb{P}_{X_n}$. The following statements are equivalent.

- (i) *The random variables $X_1, \dots, X_n$ are independent.*
- (ii) $\mathbb{P}_{X_1, \dots, X_n} = \mathbb{P}_{X_1} \otimes \dots \otimes \mathbb{P}_{X_n}$.

Proof. The random variables $X_1, \dots, X_n$ are independent iff for any Borel sets $B_1, \dots, B_n \subset \mathbb{R}$ we have

$$\mathbb{P}[X_1 \in B_1, \dots, X_n \in B_n] = \mathbb{P}[X_1 \in B_1] \cdots \mathbb{P}[X_n \in B_n]$$

$$\iff \mathbb{P}_{X_1, \dots, X_n}[B_1 \times \dots \times B_n] = \mathbb{P}_{X_1} \otimes \dots \otimes \mathbb{P}_{X_n}[B_1 \times \dots \times B_n].$$

Thus the random variables $X_1, \dots, X_n$ are independent iff the measures $\mathbb{P}_{X_1, \dots, X_n}$ and $\mathbb{P}_{X_1} \otimes \dots \otimes \mathbb{P}_{X_n}$ coincide on the set of rectangles $B_1 \times \dots \times B_n$, i.e.,

$$\mathbb{P}_{X_1, \dots, X_n} = \mathbb{P}_{X_1} \otimes \dots \otimes \mathbb{P}_{X_n}.$$

This set of rectangles is a π -system that generates the Borel algebra of $\mathbb{R}^n$. The conclusion follows from Proposition 1.2.4. □

1.3.6. Convolution of Borel measures on the real axis.

Definition 1.3.49. Let μ, ν be two finite Borel measures on $(\mathbb{R}^k, \mathcal{B}_{\mathbb{R}^k})$. The *convolution* of μ with ν is the Borel measure $\mu * \nu$ on $(\mathbb{R}^k, \mathcal{B}_{\mathbb{R}^k})$ defined by

$$\mu * \nu[B] = \int_{\mathbb{R}^k} \mu[B - y] \nu[dy], \quad \forall B \in \mathcal{B}_{\mathbb{R}^k}. \quad (1.3.53)$$

□

For $y \in \mathbb{R}^k$ we denote by S_y the shift $S_y : \mathbb{R}^k \rightarrow \mathbb{R}^k$, $S_y(x) = x + y$ and set $\mu_y := (S_y)_\# \mu$. Note that for any Borel set $B \subset \mathbb{R}^k$ we have

$$\mu_y[B] = \mu[S^{-1}(B)] = \mu[B - y],$$

so we can rewrite (1.3.53) in the form

$$\mu * \nu[-] = \int_{\mathbb{R}^k} \mu_y[-] \nu[dy].$$

A simple argument based on the Monotone Convergence Theorem shows that $\mu * \nu$ is indeed a Borel measure on $\mathbb{R}$. By letting $B = \mathbb{R}$ in (1.3.53) we see that $\mu * \nu$ is indeed a finite measure. It is a probability measure if both μ and ν are.

Note that $\mu * \nu$ is a *mixture* in the sense that it is obtained by averaging of the family of probability measures $(\mu_y)_{y \in \mathbb{R}}$ with respect to the probability measure $\nu[dy]$. For example, if

$$\nu = \sum_{i=1}^n \frac{1}{n} \delta_{x_i},$$

then

$$\mu * \nu = \frac{1}{n} \sum_{i=1}^n \mu_{x_i}.$$

In the remainder of this section I will concentrate exclusively on the one-dimensional case, $k = 1$.

Proposition 1.3.50. Let μ, ν be probability measures on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ and

$$\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad \Phi(x, y) = x + y.$$

Then $\mu * \nu = \Phi_\#(\mu \otimes \nu) = \nu * \mu$.

Proof. Let $B \in \mathcal{B}_{\mathbb{R}}$ and set $\hat{B} = \Phi^{-1}(B)$. Set

$$\hat{B}_y := \{x; (x, y) \in \hat{B}\} = B - y.$$

Then

$$\Phi_\#(\mu \otimes \nu)[B] = \int_{\mathbb{R}^2} \mathbf{I}_{\hat{B}} \mu \otimes \nu[dx dy]$$

(use Fubini-Tonelli)

$$= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} \mathbf{I}_{\hat{B}_y} \mu[dx] \right) \nu[dy] = \int_{\mathbb{R}} \mu[B - y] \nu[dy] = \mu * \nu[B].$$

The equality $\mu * \nu = \nu * \mu$ follows by changing the order of integration in the Fubini-Tonelli theorem.

□

Corollary 1.3.51. Let $X, Y \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ be two independent random variables with distributions $\mathbb{P}_X$ and $\mathbb{P}_Y$. Then

$$\mathbb{P}_{X+Y} = \mathbb{P}_X * \mathbb{P}_Y.$$

Proof. Since X, Y are independent we have $\mathbb{P}_{X,Y} = \mathbb{P}_X \otimes \mathbb{P}_Y$. Note that $\mathbb{P}_{X+Y} = \Phi_{\#} \mathbb{P}_{X,Y}$. The conclusion now follows from Proposition 1.3.50. $\square$

Remark 1.3.52. (a) Suppose that F_{μ} is the cdf of the probability measure μ , i.e.,

$$F_{\mu}(c) = \mu[(-\infty, c]], \quad \forall c \in \mathbb{R}.$$

Then the cdf $F_{\mu*\nu}$ of $\mu * \nu$ satisfies

$$F_{\mu*\nu}(c) = \int_{\mathbb{R}} F_{\mu}(c-x) \nu[dx], \quad \forall c \in \mathbb{R}.$$

We write this equality as

$$F_{\mu*\nu} = F_{\mu} * \nu. \quad (1.3.54)$$

If μ is absolutely continuous with respect to the Lebesgue measure λ on $\mathbb{R}$ so

$$\mu[dx] = \rho_{\mu}(x)dx, \quad \nu[dx] = \rho_{\nu}(x)dx, \quad \rho_{\mu}, \rho_{\nu} \in L^1(\mathbb{R}, \lambda),$$

then $\mu * \nu \ll \lambda$ and

$$\mu * \nu[dx] = \rho_{\mu*\nu}(x)dx, \quad \rho_{\mu*\nu}(x) = \rho_{\mu} * \rho_{\nu}(x) := \int_{\mathbb{R}} \rho_{\mu}(x-y) \nu[dy].$$

To see this it suffices to check that for any $c \in \mathbb{R}$ we have

$$\mu * \nu[(-\infty, c]] = \int_{-\infty}^c \rho_{\mu*\nu}(x)dx.$$

We have

$$\begin{aligned} \mu * \nu[(-\infty, c]] &= \int_{\mathbb{R}} \mu[(-\infty, c-y)] \nu[dy] = \int_{\mathbb{R}} \left(\int_{-\infty}^{c-y} \rho_{\mu}(x)dx \right) \nu[dy] \\ &= \int_{\mathbb{R}} \left(\int_{-\infty}^{c-y} \rho_{\mu}(x)dx \right) \nu[dy] = \int_{\mathbb{R}} \left(\int_{-\infty}^c \rho_{\mu}(z-y)dz \right) \nu[dy] \end{aligned}$$

(use Fubini)

$$= \int_{-\infty}^c \left(\int_{\mathbb{R}} \rho_{\mu}(z-y) \nu[dy] \right) dz = \int_{-\infty}^c \rho_{\mu*\nu}(z) [dz].$$

(b) Any Borel probability measure μ on $\mathbb{R}$ is the probability distribution of the random variable

$$\mathbb{1}_{\mathbb{R}} : (\mathbb{R}, \mathcal{B}_{\mathbb{R}}, \mu) \rightarrow \mathbb{R}, \quad \mathbb{1}_{\mathbb{R}}(x) = x.$$

If μ_1, μ_2, μ_3 are different Borel probability measures on $\mathbb{R}$, then we can define three independent random variables

$$\begin{aligned} X_1, X_2, X_3 &: (\mathbb{R}^3, \mathcal{B}_{\mathbb{R}^3}, \mu_1 \otimes \mu_2 \otimes \mu_3) \rightarrow \mathbb{R}, \\ X_k(x_1, x_2, x_3) &= x_k, \quad k = 1, 2, 3. \end{aligned}$$

Note that $\mathbb{P}_{X_k} = \mu_k, \forall k = 1, 2, 3$. Since $(X_1 + X_2) \perp X_3$ and $X_1 \perp (X_2 + X_3)$ we deduce

$$(\mu_1 * \mu_2) * \mu_3 = \mathbb{P}_{(X_1+X_2)+X_3} = \mathbb{P}_{X_1+(X_2+X_3)} = \mu_1 * (\mu_2 * \mu_3).$$

Similarly

$$\mu_1 * \mu_2 = \mathbb{P}_{X_1+X_2} = \mathbb{P}_{X_2+X_1} = \mu_2 * \mu_1.$$

Note that $\mu * \nu[\mathbb{R}] = \mu[\mathbb{R}] \cdot \nu[\mathbb{R}]$. In particular, the space $\text{Prob}(\mathbb{R})$ of Borel probability measures on $\mathbb{R}$ has a structure of commutative semigroup with respect to the convolution. The Dirac measure δ_0 is the identity element of this semigroup. $\square$

1.3.7. Poisson processes. Suppose that we have a stream of events occurring in succession at random times $S_1 \leq S_2 \leq S_3 \leq \dots$ such that the waiting times between two successive occurrences

$$T_1 = S_1, \quad T_2 = S_2 - S_1, \dots, T_n = S_n - S_{n-1}, \dots$$

are i.i.d. exponential random variables $T_n \sim \text{Exp}(\lambda)$, $n = 1, 2, \dots$. We set $S_0 := 0$.

It may help to think of the sequence (T_n) as inter-arrival times for a bus. The first bus arrives at the station at time $S_1 = T_1$. Once the n -th bus has left the station, the waiting time for the next bus to arrive is an exponential random variable T_{n+1} , independent of the preceding waiting times. From this point of view, S_n is the arrival time of the n -th bus.

For $t > 0$ we denote by $N(t)$ the number events that of occurred during the time interval $[0, t]$. In terms of streams of busses, $N(t)$ would count the number of buses that have arrived at the station in the interval $[0, t]$. In other words

$$N(t) = \max \{ n \geq 1; S_n \leq t \} = \# \{ n \geq 1; S_n \leq t \}.$$

This is a discrete random variable with range $\{0, 1, 2, 3, \dots\}$. The collection of random variables $\{N(t), t \geq 0\}$ is called the *Poisson process* with intensity λ . Note that

$$N(t) = \sum_{n=1}^{\infty} I_{[0,t]}(S_n).$$

Let us find the distribution (pmf) of $N(t)$. We have

$$\mathbb{P}[N(t) = 0] = \mathbb{P}[T_1 > t] = e^{-\lambda t} = \text{the survival function of } \text{Exp}(\lambda).$$

If $n > 0$, then $N(t) = n$ if and only if the n -th bus arrived sometime during the interval $[0, t]$, i.e., $S_n \leq t$, but the $(n+1)$ -th bus has not arrived in this time interval. We deduce

$$\mathbb{P}[N(t) = n] = \mathbb{P}[\{S_n \leq t\} \setminus \{S_{n+1} \leq t\}] = \mathbb{P}[S_n \leq t] - \mathbb{P}[S_{n+1} \leq t].$$

If we denote by $F_n(t)$ the cdf of S_n , then we can rewrite the above equality in the form

$$\mathbb{P}[N(t) = n] = F_n(t) - F_{n+1}(t).$$

We have

$$\begin{aligned} \mathbb{P}_{S_n} &= \underbrace{\text{Exp}(\lambda) * \dots * \text{Exp}(\lambda)}_n \\ &= \underbrace{\text{Gamma}(\lambda, 1) * \dots * \text{Gamma}(\lambda, 1)}_n \stackrel{(1.6.8a)}{=} \text{Gamma}(\lambda, n). \end{aligned}$$

Hence, for $n > 0$

$$F_{n+1}(t) = \frac{\lambda^{n+1}}{\Gamma(n+1)} \int_0^t s^n e^{-\lambda s} ds = \frac{\lambda^{n+1}}{n!} \int_0^t s^n e^{-\lambda s} ds.$$

For $n > 0$, we integrate by parts to obtain

$$F_{n+1}(t) = - \left(\frac{\lambda^n}{n!} s^n e^{-\lambda s} \right) \Big|_{s=0}^{s=t} + \frac{\lambda^n}{(n-1)!} \int_0^t s^{n-1} e^{-\lambda s} ds = - \frac{(t\lambda)^n}{n!} e^{-\lambda t} + F_n(t).$$

Hence

$$\mathbb{P}[N(t) = n] = F_n(t) - F_{n+1}(t) = \frac{(t\lambda)^n}{n!} e^{-\lambda t}, \quad n > 0. \quad (1.3.55)$$

This shows that $N(t)$ is a Poisson random variable, $N(t) \sim \text{Poi}(\lambda t)$.

The family of random variables $(N(t))$ is nondecreasing and thus there exist right and left limits

$$N(t-0) = \lim_{s \nearrow t} N(s), \quad N(t+0) = \lim_{s \searrow t} N(s).$$

It is not difficult to see that

$$\forall t \geq 0, \quad N(t) = N(t+0), \quad N(t) - N(t-0) \in \{0, 1\} \quad \text{a.s.} \quad (1.3.56)$$

The Poisson process plays an important role in probability since it appears in many situations and displays many surprising phenomena. One such interesting phenomenon is the *waiting time paradox*, [76, I.4]. To better appreciate this paradox we consider two separate situations.

Suppose first that buses arrive at a bus station following a Poisson stream with frequency λ . Bob arrives at the bus station at a time $t \geq 0$, the bus is not there and he is waiting for the next one. His *waiting time* is

$$W_t := S_{N(t)+1} - t$$

We want to compute its expectation $w_t := \mathbb{E}[W_t]$. There are two possible heuristic arguments.

- (i) The memoryless property of the exponential distribution shows that w_t should be independent of t so $w_t = w_0 = \frac{1}{\lambda}$.
- (ii) Bob's arrival time t is uniformly distributed in the inter-arrivals interval $(S_{N(t)}, S_{N(t)+1})$ of expected length $\frac{1}{\lambda}$ and, as in the earlier deterministic computation, the expectation should be half its length, $\frac{1}{2\lambda}$.

We will show that (i) provides the correct answer. However, even the reasoning (ii) holds a bit of truth. To see what is happening we compute the expectations of $S_{N(t)}$ and $S_{N(t)+1}$. We have

$$\mathbb{E}[S_{N(t)}] = \int_0^t \mathbb{P}[S_{N(t)} > x] dx.$$

Note that

$$\mathbb{P}[S_{N(t)} > x] = \sum_{n \geq 0} \mathbb{P}[S_{N(t)} > x, N(t) = n].$$

On the other hand,

$$\mathbb{P}[S_{N(t)} > x, N(t) = n] = \mathbb{P}[x < S_n \leq t, S_n + T_{n+1} > t].$$

The random variables S_n and T_{n+1} are independent and the joint distribution of (S_n, T_{n+1}) is

$$\mathbb{P}_{S_n, T_{n+1}}[ds d\tau] = \underbrace{\frac{\lambda^n}{(n-1)!} s^{n-1} e^{-\lambda s} \lambda e^{-\lambda \tau}}_{\rho(s, \tau)} ds d\tau$$

so

$$\begin{aligned} \mathbb{P}[x < S_n < t, S_n + T_{n+1} > t] &= \int_{\substack{x < s \leq t \\ s + \tau > t}} \rho(s, \tau) ds d\tau \\ &= \int_x^t \left(\int_{t-s}^\infty \rho(s, \tau) d\tau \right) ds = \int_x^t \mathbb{P}[T_{n+1} > t-s] \frac{\lambda^n}{(n-1)!} s^{n-1} e^{-\lambda s} ds \end{aligned}$$

$$= \int_x^t e^{-\lambda(t-s)} \frac{\lambda^n}{(n-1)!} s^{n-1} e^{-\lambda s} ds = \frac{e^{-\lambda t} \lambda^n}{(n-1)!} \int_x^t s^{n-1} ds = \frac{e^{-\lambda t} \lambda^n}{n!} (t^n - x^n).$$

We deduce

$$\mathbb{P}[S_{N(t)} > x] = \sum_{n \geq 0} \frac{e^{-\lambda t} \lambda^n}{n!} (t^n - x^n) = 1 - e^{-\lambda(t-x)},$$

$$\mathbb{E}[S_{N(t)}] = \int_0^t (1 - e^{-\lambda(t-x)}) dx = t - e^{-\lambda t} \int_0^t e^{\lambda x} dx = t - \frac{e^{-t}}{\lambda} (e^{\lambda t} - 1).$$

Hence

$$\mathbb{E}[S_{N(t)}] = t - \frac{1}{\lambda} + \frac{e^{-\lambda t}}{\lambda} = \frac{1}{\lambda} \mathbb{E}[N(t) - 1 + e^{-\lambda t}]. \quad (1.3.57)$$

Let us compute $\mathbb{E}[S_{N(t)+1}]$. Again, we have

$$\mathbb{P}[S_{N(t)+1} > x] = \sum_{n \geq 0} \mathbb{P}[S_{N(t)+1} > x, N(t) = n],$$

and

$$\begin{aligned} \mathbb{P}[S_{N(t)+1} > x, N(t) = n] &= \mathbb{P}[S_n \leq t, S_{n+1} \geq \max(t, x)] \\ &= \begin{cases} \mathbb{P}[S_n \leq t, S_n + T_{n+1} \geq t], & x \leq t, \\ \mathbb{P}[S_n \leq t, S_n + T_{n+1} \geq x], & x > t. \end{cases} \end{aligned}$$

For any $c \geq t$ we have

$$\begin{aligned} \mathbb{P}[S_n \leq t, S_n + T_{n+1} \geq c] &= \int_{\substack{s \leq t, \\ s+\tau \geq c}} \rho(s, t) ds d\tau \\ &= \int_0^t \left(\int_{c-s}^\infty \rho(s, \tau) d\tau \right) ds = \frac{\lambda^n}{(n-1)!} \int_0^t e^{-\lambda(c-s)} s^{n-1} e^{-\lambda s} ds = \frac{e^{-\lambda c} (\lambda t)^n}{n!}. \end{aligned}$$

Observing that

$$\sum_{n \geq 0} \frac{e^{-\lambda c} (\lambda t)^n}{n!} = e^{-\lambda(c-t)}$$

we deduce that

$$\mathbb{P}[S_{N(t)} > x] = \begin{cases} 1, & x \leq t, \\ e^{-\lambda(x-t)}, & x > t. \end{cases}$$

Hence

$$\mathbb{E}[S_{N(t)+1}] = \int_0^t dx + e^{\lambda t} \int_t^\infty e^{-\lambda x} dx = t + \frac{1}{\lambda} = \frac{1}{\lambda} \mathbb{E}[N(t) + 1], \quad (1.3.58)$$

and

$$w_t = \mathbb{E}[S_{N(t)+1}] - t = \frac{1}{\lambda}.$$

In fact much more is true. One can show (see [175, Sec. 3.6]) that the waiting time W_t is an exponential random variable, $W_t \sim \text{Exp}(\lambda)$, in agreement with the conclusion of the argument (i).

The above computation show that the expectation of $L_t = S_{N(t)+1} - S_{N(t)}$ is

$$\mathbb{E}[L_t] = \frac{2}{\lambda} - \frac{e^{-\lambda t}}{\lambda} \approx \frac{2}{\lambda} \text{ for } t \text{ large.}$$

We have reached counterintuitive conclusions. The expected waiting time from the moment bus $N(t)$ left the station until bus $N(t)+1$ arrives in the station is *twice* the expected inter-arrival times $\mathbb{E}[T_n]$!

On the other hand, the actual expected time w_t from epoch t until the arrival of bus $N(t) + 1$ is the usual expected inter-arrival time. This shows that even the argument (ii) captures a bit of what is going on since w_t is close to half the expected length of the inter-arrival interval $(S_{N(t)}, S_{N(t)+1})$.

The number of busses arriving during a time interval $[0, t]$ is $N(t)$. The busses arrive with a frequency of $\frac{1}{\lambda}$ per unit of time, so we should expect to wait $t = \frac{1}{\lambda} \mathbb{E}[N(t)]$ units of time for $N(t)$ busses to arrive. However, formula (1.3.57) shows that we should expect less than t units of time for $N(t)$ busses to arrive. On the other hand, formula (1.3.58) shows that we should expect $t + 1 = \frac{1}{\lambda} \mathbb{E}[N(t) + 1]$ units of time for $N(t) + 1$ busses to arrive! We refer to Remark 3.2.36 for another (technical) explanation for this paradoxical divergence of conclusions.

The Poisson processes are special cases of *renewal processes*. For an enjoyable and highly readable introduction to renewal processes we refer to [76] or [175, Chap. 3]. For a more in-depth presentation of these processes and some of their practical applications we refer to [7]. $\square$

1.3.8. Modes of convergence of random variables. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$.

Definition 1.3.53 (Almost sure convergence). We say that the sequence of random variables

$$X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P}), \quad n \in \mathbb{N},$$

converges *almost surely* (or a.s.) to $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ if there exist $\Omega_0 \in \mathcal{S}$ such that

$$\mathbb{P}[\Omega_0] = 1, \quad \lim_{n \rightarrow \infty} X_n(\omega) = X(\omega), \quad \forall \omega \in \Omega_0.$$

We will use the notation $X_n \xrightarrow{\text{a.s.}} X$ to indicate the a.s. convergence. $\square$

Tautologically, the a.s. convergence is well defined in L^0 . To describe a useful criterion for a.s. convergence we need to rely on a very versatile classical result.

Definition 1.3.54. For any sequence of events $(A_n)_{n \in \mathbb{N}} \subset \mathcal{S}$ we denote by A_n i.o. the event “ A_n occurs infinitely often”,

$$A_n \text{ i.o.} := \bigcap_{m \geq 1} \bigcup_{n \geq m} A_n.$$

Thus

$$\omega \in A_n \text{ i.o.} \iff \forall m \in \mathbb{N} \quad \exists n \geq m : \omega \in A_n. \quad \square$$

Theorem 1.3.55 (Borel-Cantelli Lemma). Consider a sequence of events $(A_n)_{n \in \mathbb{N}} \subset \mathcal{S}$.

(i) If

$$\sum_{n \geq 1} \mathbb{P}[A_n] < \infty.$$

Then $\mathbb{P}[A_n \text{ i.o.}] = 0$.

(ii) Conversely, if the events $(A_n)_{n \in \mathbb{N}}$ are independent then $\mathbb{P}[A_n \text{ i.o.}] \in \{0, 1\}$, and

$$\mathbb{P}[A_n \text{ i.o.}] = 0 \iff \sum_{n \geq 1} \mathbb{P}[A_n] < \infty. \quad (1.3.59)$$

Proof. (i) We set

$$N := \sum_{n \geq 1} I_{A_n}.$$

Note that $\{A_n \text{ i.o.}\} = \{N = \infty\}$. From the Monotone Convergence Theorem we deduce

$$\mathbb{E}[N] = \sum_{n \geq q} \mathbb{E}[\mathbf{I}_{A_n}] = \sum_{n \geq 1} \mathbb{P}[A_n] < \infty$$

so $\mathbb{P}[N = \infty] = 0$.

(ii) Kolmogorov's zero-one theorem shows that when the events $(A_n)_{n \geq 1}$ are independent we have $\mathbb{P}[A_n \text{ i.o.}] \in \{0, 1\}$.

To prove (1.3.59) we have to show that if

$$\sum_{n \geq 1} \mathbb{P}[A_n] = \infty,$$

then $\mathbb{P}[A_n \text{ i.o.}] = 1$. We have

$$\mathbb{P}\left[\bigcup_{n \geq m} A_n\right] = 1 - \mathbb{P}\left[\bigcap_{n \geq m} A_n^c\right]$$

(use the independence of A_n)

$$= 1 - \prod_{n \geq m} (1 - \mathbb{P}[A_n])$$

$$(1 - x \leq e^{-x}, \forall x \in \mathbb{R})$$

$$\geq 1 - e^{-\sum_{n \geq m} \mathbb{P}[A_n]} = 1.$$

Hence

$$\mathbb{P}[A_n \text{ i.o.}] = \lim_{m \rightarrow \infty} \mathbb{P}\left[\bigcup_{n \geq m} A_n\right] = 1.$$

□

Remark 1.3.56. Statement (i) in Theorem 1.3.55 is usually referred to as the *First Borel-Cantelli Lemma* while statement (ii) is usually referred to as the *Second Borel-Cantelli Lemma*. Exercises 3.12 and 3.20 present refinements of the Borel-Cantelli lemmas. □

Observe that $X_n \rightarrow X$ a.s. if and only if, for any $\nu \in \mathbb{N}$

$$\mathbb{P}[\{ |X_n - X| > 1/\nu \} \text{ i.o.}] = 0.$$

The Borel-Cantelli Lemma now implies the following result.

Corollary 1.3.57. Suppose that there exists $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ such that the sequence $X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ satisfies

$$\sum_{n \geq 1} \mathbb{P}[|X_n - X| > \varepsilon] < \infty, \quad \forall \varepsilon > 0.$$

Then $X_n \xrightarrow{\text{a.s.}} X$.

□

Proof. The Borel-Cantelli Lemma implies that

$$\mathbb{P}[|X_n - X| > \varepsilon \text{ i.o.}] = 0, \quad \forall \varepsilon > 0.$$

Hence, for any $\varepsilon > 0$ there exists a negligible set $S_\varepsilon \in \mathcal{S}$ such that, for any $\omega \in \Omega \setminus S_\varepsilon$ we have

$$\limsup_{n \rightarrow \infty} |X_n(\omega) - X(\omega)| \leq \varepsilon.$$

Set

$$S_\infty = \bigcup_{k \in \mathbb{N}} S_{1/k}.$$

We deduce that for any $\omega \in \Omega \setminus S_\infty$ we have

$$\limsup_{n \rightarrow \infty} |X_n(\omega) - X(\omega)| \leq 1/k, \quad \forall k \in \mathbb{N}.$$

□

Example 1.3.58. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent Bernoulli variables, $X_n \sim \text{Ber}(p_n)$. This sequence defines a random subset of $\mathbb{N}$,

$$S = \{n \in \mathbb{N}; X_n = 1\}.$$

The random set S is a.s. finite iff $\mathbb{P}[X_n = 1 \text{ i.o.}] = 0$. Since the random variables X_n are independent we deduce that $\mathbb{P}[X_n = 1 \text{ i.o.}] \in \{0, 1\}$. The Borel-Cantelli lemma implies

$$S \text{ is a.s. finite} \iff \sum_{n \geq 1} p_n < \infty.$$

Hence

$$S \text{ is a.s. infinite} \iff \sum_{n \geq 1} p_n = \infty.$$

□

Definition 1.3.59. We say that the sequence $X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ converges *in probability* to the random variable $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ if, $\forall \varepsilon > 0$, we have

$$\lim_{n \rightarrow \infty} \mathbb{P}[|X_n - X| > \varepsilon] = 0.$$

We will use the notation $X_n \xrightarrow{p} X$ to indicate convergence in probability. □

Observe that if $X_n \rightarrow X$ in probability and, for any $n \in \mathbb{N}$, we have $X_n = X'_n$ a.s., then $X'_n \rightarrow X$ in probability. Thus the convergence in probability is correctly defined in $L^0(\Omega, \mathcal{S}, \mathbb{P})$.

The convergence in probability is equivalent to the convergence defined by a metric on $L^0(\Omega, \mathcal{S}, \mathbb{P})$. For $X, Y \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ we set

$$\text{dist}(X, Y) := \mathbb{E}[\min(|X - Y|, 1)] \quad (1.3.60)$$

Clearly $\text{dist}(X, Y) = \text{dist}(Y, X)$ and

$$\text{dist}(X, Z) \leq \text{dist}(X, Y) + \text{dist}(Y, Z).$$

Note that $\text{dist}(X, Y) = 0$ iff $X = Y$ a.s. so “dist” is a metric on $L^0(\Omega, \mathcal{S}, \mathbb{P})$.

Proposition 1.3.60. Let $X, X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$. Then the following statements are equivalent.

- (i) $X_n \rightarrow X$ in probability as $n \rightarrow \infty$.
- (ii) $\text{dist}(X_n, X) \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Set

$$\rho(x) := \min(|x|, 1), \quad Y_n := X_n - X.$$

Using Markov's inequality we deduce that for any $n \geq 1$ and any $\varepsilon \in (0, 1)$ we have

$$\varepsilon \mathbb{P}[|Y_n| > \varepsilon] = \varepsilon \mathbb{P}[\rho(Y_n) > \varepsilon] \leq \mathbb{E}[\rho(Y_n)] = \text{dist}(Y_n, 0).$$

This shows that (ii) $\Rightarrow$ (i).

Conversely, observe that, for any $\varepsilon > 0$, we have

$$\mathbb{E}[\rho(Y_n)] = \int_{|Y_n| \leq \varepsilon} \rho(Y_n) d\mathbb{P} + \int_{|Y_n| > \varepsilon} \rho(Y_n) d\mathbb{P} \leq \varepsilon + \mathbb{P}[|Y_n| > \varepsilon].$$

This proves that $0 \leq \liminf \text{dist}(Y_n, 0) \leq \limsup \text{dist}(Y_n, 0) \leq \varepsilon, \forall \varepsilon > 0$. $\square$

The next result describes the relationships between a.s. convergence and convergence in probability.

Theorem 1.3.61. *Let $X, X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$. Then the following hold.*

- (i) *If $X_n \rightarrow X$ a.s., then $X_n \rightarrow X$ in probability.*
- (ii) *If $X_n \rightarrow X$ in probability, then (X_n) contains a subsequence that converges a.s. to X .*
- (iii) *The sequence X_n converges in probability to X if and only if any subsequence contains a further subsequence that is a.s. convergent to X .*

Proof. (i) Set $Y_n := X_n - X$. Since $Y_n \rightarrow 0$ a.s. we have $\min(|Y_n|, 1) \rightarrow 0$ a.s.. From the Dominated Convergence Theorem we deduce

$$\text{dist}(X_n, X) = \mathbb{E}[|Y_n|] \rightarrow 0,$$

so that $Y_n \xrightarrow{p} 0$.

(ii) Suppose that $Y_n \rightarrow 0$ in probability. We deduce that for any $k \in \mathbb{N}$ there exists $n_k \in \mathbb{N}$ such that

$$\forall n \geq n_k : \mathbb{P}[|Y_n| > 1/k] < \frac{1}{2^k}.$$

Now observe that for any $m > 0$, the series

$$\sum_{k \geq 1} \mathbb{P}[|Y_{n_k}| > 1/m]$$

is convergent since, for $k > m$ we have

$$\mathbb{P}[|Y_{n_k}| > 1/m] \leq \mathbb{P}[|Y_{n_k}| > 1/k] < \frac{1}{2^k}.$$

The desired conclusion now follows from Corollary 1.3.57.

(iii) Recall that a sequence in a metric space converges to a given point if and only if any subsequence contains a sub-subsequence converging to that point. The properties (i) and (ii) show that the sequence (X_n) satisfies this condition with respect to the metric dist defined by ρ . $\square$

Corollary 1.3.62. *If the sequence (X_n) in $L^0(\Omega, \mathcal{S}, \mathbb{P})$ converges in probability to X , then for any continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ the sequence $f(X_n)$ converges in probability to $f(X)$.*

Proof. The sequence (X_n) satisfies the necessary and sufficient conditions (iii) in Theorem 1.3.61. Since f is continuous, the sequence $f(X_n)$ satisfies these necessary and sufficient conditions as well. $\square$

The next result is also an immediate consequence of Theorem 1.3.61(iii).

Corollary 1.3.63. Suppose that (X_n) and (Y_n) are two sequences of a.s. finite random variables converging in probability to the a.s. finite variables X and respectively Y . Then $X_n + Y_n$ converges in probability to $X + Y$. $\square$

Definition 1.3.64. Let $p \in [1, \infty)$. We say that the sequence $(X_n)_{n \in \mathbb{N}} \subset L^0(\Omega, \mathcal{S}, \mathbb{P})$ converges in p -mean or in L^p to $X \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ if

$$X, X_n \in L^p(\Omega, \mathcal{S}, \mathbb{P}), \quad \forall n \in \mathbb{N},$$

and

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X_n - X|^p] = 0.$$

The convergence in the L^∞ -norm is referred to as a.s. uniform convergence. $\square$

Proposition 1.3.65. If $X_n \rightarrow X$ in p -mean, $p \in [1, \infty)$, then $X_n \rightarrow X$ in probability. In particular, X_n admits a subsequence that converges a.s. to X .

Proof. Set $Y_n := X_n - X$. Then

$$\mathbb{P}[|Y_n| > \varepsilon] = \mathbb{P}[|Y_n|^p > \varepsilon^p] \stackrel{(1.2.23)}{\leq} \frac{1}{\varepsilon^p} \mathbb{E}[|Y_n|^p] \rightarrow 0 \text{ as } n \rightarrow \infty.$$

$\square$

Corollary 1.3.66. Let $p \in [1, \infty)$. If

$$\sum_{n \geq 1} \|X_n - X\|_{L^p}^p < \infty.$$

Then $X_n \rightarrow X$, a.s. and in p -mean.

Proof. The convergence in p -mean is obvious. Markov's inequality implies that for any $\varepsilon > 0$

$$\sum_{n \geq 1} \mathbb{P}[|X_n - X| > \varepsilon] \leq \frac{1}{\varepsilon^p} \sum_{n \geq 1} \|X_n - X\|_{L^p}^p < \infty$$

The conclusion now follows from Corollary 1.3.57. $\square$

Remark 1.3.67. Of all the modes of convergence discussed in this section, I find the a.s. convergence to be the most “slippery” one. I should remark that this type of convergence is not a metric type convergence, i.e., there is no metric on the space of random variables such that a.s. convergence corresponds to the convergence with respect to this metric.

This becomes a hindrance when dealing with stochastic processes. Suppose that we are given a uncountable family $(X_t)_{t > 0}$ of random variables. Then the concept of a.s. limit as $t \searrow 0$ is ill defined and needs to be dealt with a lot of care.

On a practical level, there are few versatile tricks that can be used to establish a.s. convergence. The “cheapest” one is formulated in Corollary 1.3.57. It relies ingeniously on the Borel-Cantelli lemma.

We will discuss a few fundamental a.s. convergence results in this book: Kolmogorov's one-series theorem, the strong law of large numbers, Birkhoff's a.s. ergodic theorem and Doob's martingale convergence theorem. In the first three of the above examples the a.s. convergence is a consequence of certain very ingenious maximal inequalities. Doob's theorem is in some sense an outlier, because it relies on a completely different and again very ingenious principle. $\square$

Example 1.3.68. For each $n \in \mathbb{N}$ and each $1 \leq k \leq n$ we set

$$A_{k,n} = [(k-1)/n, k/n], \quad X_{k,n} = \mathbf{I}_{A_{k,n}} : [0, 1] \rightarrow \mathbb{R}.$$

Then the sequence of random variables

$$X_{1,1}, X_{1,2}, X_{2,2}, X_{1,3}, X_{2,3}, X_{3,3}, \dots$$

converges in mean and in probability to 0. It does not converge a.s. to 0 because for any $x \in [0, 1]$ infinitely many of these random variables are equal to 1 at x .

The related sequence $Y_{k,n} = nX_{k,n}$ converges in probability to 0 but not in mean since $\|Y_{k,n}\|_{L^1} = 1$. $\square$

Example 1.3.69 (Bernoulli). Suppose that $(X_n)_{n \geq 1}$ is a sequence of i.i.d. Bernoulli random variables with winning probability $p = \frac{1}{2}$. Set

$$S_n = X_1 + \dots + X_n \sim \text{Bin}(n, 1/2), \quad M_n = \frac{1}{n} S_n.$$

Then

$$\text{Var}[M_n] = \frac{1}{n^2} \text{Var}[S_n] = \frac{1}{n} \text{Var}[\text{Ber}(1/2)] = \frac{1}{4n}.$$

Hence

$$\|M_n - 1/2\|_{L^2} = \frac{1}{2\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

so that M_n converges in 2-mean to $\frac{1}{2}$ and thus, in probability to $\frac{1}{2}$. Intuitively, M_n is the fraction of Heads in a string of n independent fair coin flips. From Chebyshev's inequality we deduce that

$$\mathbb{P}[|M_n - 1/2| > \varepsilon] \leq \frac{\varepsilon^2}{4n}.$$

It turns out that this deviation probability is much smaller. In (2.3.13a) we will show that

$$\mathbb{P}[|M_n - 1/2| > \varepsilon] \leq 2e^{-2n\varepsilon^2}.$$

For example if $\varepsilon = 10^{-2}$, $n = 10^5$ then

$$\mathbb{P}[|M_{10^5} - 1/2| > 0.01] \leq 2e^{-20} \approx 4.2 \times 10^{-9}. \quad \square$$

Example 1.3.70 (Longest common subsequence). Consider a finite set $\mathcal{A}$, $|\mathcal{A}| = k$, called *alphabet*. A word of length n in the alphabet $\mathcal{A}$ is a finite sequence of the form

$$\underline{x} := (x_1, \dots, x_n) \in \mathcal{A}^n.$$

A *subsequence* of such a word is a word of the form

$$(x_{f(1)}, \dots, x_{f(\ell)}) \in \mathcal{A}^\ell,$$

where f an increasing function $f : \{1, \dots, \ell\} \rightarrow \{1, \dots, n\}$. The natural number ℓ is called the length of the subsequence.

A common subsequence of two words $\underline{x}, \underline{y} \in \mathcal{A}^n$ is a word $w \in \mathcal{A}^\ell$ that is a subsequence of both. For example, if $\mathcal{A} = \{H, T\}$, then H, T, H, T, T is a subsequence of both words

$$\underline{H, T, T, H, H, T, T} \text{ and } \underline{T, H, T, H, T, T, H}$$

We are interested in the length of the longest common subsequence of two *random* words of length n on the alphabet $\mathcal{A}$. Such a problem arises in genetics. In that case the alphabet is $\{A, C, T, G\}$. The DNA molecules are described by (very long) words in this alphabet. The existence a long common

subsequence of two such words is an indication of a common ancestor of two living organisms with those DNAs.

From a mathematical point of view, we fix a probability measure π on an alphabet $\mathcal{A}$ and we choose independent random variables

$$\{X_n, Y_n; n \in \mathbb{N}\}$$

where X_n, Y_n are $\mathcal{A}$ -valued and have common distribution π .

One can think that these random variables are obtained as follows. Two individuals independently roll identical "dice" with faces labeled by $\mathcal{A}$ and whose occurrences are governed by π . The first individual generates the sequence (X_n) while the second individual generates the sequence Y_n . We denote by L_n the length of the longest common subsequence of the words

$$(X_1, \dots, X_n) \text{ and } (Y_1, \dots, Y_n).$$

We want to prove at a.s. and L^1 we have

$$\lim_{n \rightarrow \infty} \frac{L_n}{n} = R(\pi) := \sup_{n \geq 1} \frac{L_n}{n} \quad (1.3.61)$$

In particular, this shows that

$$\lim_{n \rightarrow \infty} \frac{L_n}{n} > L_1 > 0.$$

Note that L_1 is a Bernoulli random variable with success probability

$$p = \sum_{a \in \mathcal{A}} \pi[a]^2.$$

The equality (1.3.61) is due to Chvátal and Sankoff [41], but we will follow the presentation in [198, Chap. 1].

The key observation is that the sequence $(\ell_n)_{n \in \mathbb{N}}$ is *superadditive*, i.e.,

$$\ell_n + \ell_m \leq \ell_{m+n}, \quad \forall m, n \in \mathbb{N}. \quad (1.3.62)$$

The proof is very simple. We set $Z_n = (X_n, Y_n)$ and we observe that the random variable L_n is an invariant of the sequence of pairs $(Z_1, \dots, Z_n)$, $L_n = L(Z_1, \dots, Z_n)$. Clearly

$$L_m = L(Z_{n+1}, \dots, Z_{n+m}), \quad \forall m, n \in \mathbb{N}.$$

If we concatenate the longest common subsequence of $(Z_1, \dots, Z_n)$ with the longest common subsequence of $(Z_{n+1}, \dots, Z_{n+m})$ we obtain a common subsequence of $(Z_1, \dots, Z_n, Z_{n+1}, \dots, Z_{n+m})$ of length

$$L(Z_1, \dots, Z_n) + L(Z_{n+1}, \dots, Z_{n+m})$$

showing that

$$L(Z_1, \dots, Z_n) + L(Z_{n+1}, \dots, Z_{n+m}) \leq L(Z_1, \dots, Z_n, Z_{n+1}, \dots, Z_{n+m}),$$

i.e.,

$$L_m + L_n \leq L_{m+n}, \quad \forall m, n \in \mathbb{N}. \quad (1.3.63)$$

Taking the expectations of both sides in the above inequality we obtain (1.3.62).

The conclusion (1.3.61) is now an immediate consequence of the following elementary result.

Lemma 1.3.71 (Fekete). Suppose that $(x_n)_{n \geq 1}$ is a subadditive sequence of real numbers, i.e.,

$$x_{m+n} \leq x_m + x_n, \quad \forall m, n \in \mathbb{N}.$$

Then

$$\lim_{n \rightarrow \infty} \frac{x_n}{n} = \mu := \inf_{n \geq 1} \frac{x_n}{n}$$

Proof. Then, for any $c > \mu$ we can find $k = k(c) > 0$ such that $x_k \leq c$. The subadditivity condition implies $x_{kn} \leq nx_k$, $\forall n \in \mathbb{N}$, so that

$$\mu \leq \frac{x_{nk}}{nk} < c, \quad \forall n \in \mathbb{N}.$$

Hence

$$\mu \leq \liminf_{n \rightarrow \infty} \frac{x_n}{n} \leq c, \quad \forall c > \mu,$$

i.e.,

$$\mu = \liminf_{n \rightarrow \infty} \frac{x_n}{n}.$$

Now observe that for any $n \geq k(c) > 0$, there exist $m \in \mathbb{N}$ and $r \in \{0, 1, \dots, k(c) - 1\}$ such that $n = mk(c) + r$. Hence

$$x_n \leq mx_{k(c)} + r_r < mc + x_r$$

so that

$$\frac{x_n}{n} < \frac{(n-r)c}{n} + \frac{M_c}{n}, \quad M_c = \sup \{ |x_1| + \dots + |x_{k(c)}| \}.$$

Hence

$$\limsup_{n \rightarrow \infty} \frac{x_n}{n} \leq \limsup_{n \rightarrow \infty} \frac{(n-r)c}{n} = c, \quad \forall c > \mu.$$

This completes the proof of the lemma. $\square$

The conclusion (1.3.61) follows from Fekete's Lemma applied to the sequence $x_n = -L_n$. The inequality (1.3.63) show that

$$\frac{L_n}{n} \rightarrow R := \sup_n \frac{L_n}{n}.$$

Set $r = r(\pi) := \mathbb{E}[R]$. We deduce from the Cauchy-Schwartz inequality that

$$r \geq \mathbb{E}[L_1] = \sum_{a \in \mathcal{A}} \pi[a]^2 \geq \frac{1}{k} \left(\sum_{a \in \mathcal{A}} \pi(a) \right)^2 = \frac{1}{k} > 0.$$

The Dominated Convergence Theorem implies that

$$r = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[L_n].$$

The exact value of $r(\pi)$ is not known in general. In Example 3.1.36, using more sophisticated techniques, we will show that the limit $R(\pi)$ is constant, $R(\pi) = r$ and $\frac{L_n}{n}$ is highly concentrated around its mean r_n . $\square$

The concept of convergence in probability is weaker than the concepts of convergence a.s. or in p -mean. In many applications it is useful to know sufficient additional assumptions that will guarantee that a sequence convergent in probability is also convergent in p -mean. The a.s. convergence does not guarantee convergence in mean. The next elementary example is typical of what can go wrong.

Example 1.3.72. Consider the interval $[-1, 1]$ equipped with the uniform probability measure $\frac{1}{2}dx$. Consider the sequence of nonnegative random variables

$$X_n = 2^n \mathbf{I}_{[-2^{-n}, 2^{-n}]}$$

Note that $X_n \rightarrow 0$ a.s. but

$$\mathbb{E}[X_n] = \frac{2^n}{2} \int_{-2^{-n}}^{2^{-n}} dx = 1, \quad \forall n.$$

As we will see later in Chapter 3, the reason why the convergence in mean fails is the high concentration of X_n on sets of smaller and smaller measures. $\square$

Our next result is an example of a sufficient condition for a sequence converging in probability to also converge in the mean. It is a stepping stone towards the more refined results that we will discuss in Chapter 3.

Theorem 1.3.73 (Bounded Convergence Theorem). *Suppose that (X_n) is a sequence in $L^1(\Omega, \mathcal{S}, \mathbb{P})$ that converges in probability to $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. If the sequence (X_n) is bounded in $L^\infty(\Omega, \mathcal{S}, \mathbb{P})$, i.e.,*

$$M := \sup_{n \in \mathbb{N}} \|X_n\|_\infty < \infty,$$

then $X_n \rightarrow X$ in L^1 and

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n] = \mathbb{E}[X] \quad (1.3.64)$$

Proof. I follow the approach in [216, Thm. 1.4]. Since

$$|\mathbb{E}[X_n] - \mathbb{E}[X]| \leq \mathbb{E}[|X_n - X|],$$

and $|X_n - X| \rightarrow 0$ in probability, it suffices to consider only the special case $X = 0$, and $X_n \geq 0$ a.s.. In such an instance it suffices to prove only the equality (1.3.64).

For any $\varepsilon > 0$ we have

$$\mathbb{E}[X_n] = \mathbb{E}[X_n \mathbf{I}_{\{X_n \leq \varepsilon\}}] + \mathbb{E}[X_n \mathbf{I}_{\{X_n > \varepsilon\}}] \leq \varepsilon + M \mathbb{P}[X_n > \varepsilon].$$

Letting $n \rightarrow \infty$ taking to account that $X_n \geq 0$ and $X_n \rightarrow 0$ in probability we deduce

$$0 \leq \liminf_{n \rightarrow \infty} \mathbb{E}[X_n] \leq \limsup_{n \rightarrow \infty} \mathbb{E}[X_n] \leq \varepsilon, \quad \forall \varepsilon > 0.$$

$\square$

Remark 1.3.74. The Bounded Convergence theorem does not follow immediately from the Dominated Convergence Theorem which involves a.s. convergence. However, using Theorem 1.3.61(iii) we can use the Dominated Convergence Theorem to provide an alternate proof of the Bounded Convergence Theorem. $\square$

1.4. Conditional expectation

The concept of *conditioning* is a central pillar of the theory of probability. It has a genuinely probabilistic origin and very rich and subtle ramifications. Also, it takes some time getting used to it. This concept is one important reason why in probability sigma-algebras play a much more important role than in analysis.

Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$.

1.4.1. Conditioning on a sigma-subalgebra. The main formal constructions of this section are best understood if we first consider a special but very useful example.

Example 1.4.1 (Conditioning on a partition). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ and $(F_\alpha)_{\alpha \in A}$, $A \subset \mathbb{N}$, is a finite or countable partition of Ω with measurable and nonnegligible chambers, i.e.,

$$F_\alpha \in \mathcal{S}, \quad \mathbb{P}[F_\alpha] > 0, \quad \forall \alpha \in A.$$

We denote by $\mathcal{F}$ the sigma-algebra generated by this partition. In other words, $F \in \mathcal{F}$ if and only if it is a union of chambers F_α . This means that $\exists B \subset A$ such that

$$F = \bigcup_{\beta \in B} F_\beta.$$

Observe that a function $Y : \Omega \rightarrow \mathbb{R}$ is $\mathcal{F}$ -measurable if and only there exist real numbers $(y_\alpha)_{\alpha \in A}$ such that

$$Y = \sum_{\alpha \in A} y_\alpha \mathbf{I}_\alpha, \quad \mathbf{I}_\alpha := \mathbf{I}_{F_\alpha}.$$

Moreover

$$Y \in \mathcal{L}^1 \iff \sum_{\alpha} |y_\alpha| \mathbb{P}[F_\alpha] < \infty.$$

Suppose now that $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$. We define the *expectation of X given the event F_α* to be the expectation of X with respect to the conditional probability $\mathbb{P}[- | F_\alpha]$, i.e., the *number*

$$\bar{x}_\alpha = \mathbb{E}[X | F_\alpha] := \frac{1}{\mathbb{P}[F_\alpha]} \mathbb{E}[X \mathbf{I}_\alpha] = \frac{1}{\mathbb{P}[F_\alpha]} \int_{F_\alpha} X(\omega) \mathbb{P}[d\omega]. \quad (1.4.1)$$

We obtain an $\mathcal{F}$ -measurable *random variable*

$$\bar{X} = \sum_{\alpha} \bar{x}_\alpha \mathbf{I}_\alpha.$$

Note that

$$|\bar{x}_\alpha| \leq \frac{1}{\mathbb{P}[F_\alpha]} \mathbb{E}[|X| \mathbf{I}_\alpha]$$

so

$$\mathbb{E}[|\bar{X}|] \leq \sum_{\alpha} \mathbb{E}[|X| \mathbf{I}_\alpha] = \mathbb{E}[|X|] < \infty.$$

Since

$$\mathbb{E}[X \mathbf{I}_\alpha] = \mathbb{E}[\bar{X} \mathbf{I}_\alpha], \quad \forall \alpha \in A,$$

we deduce

$$\mathbb{E}[X \mathbf{I}_F] = \mathbb{E}[\bar{X} \mathbf{I}_F], \quad \forall F \in \mathcal{F}. \quad (1.4.2)$$

Note that if

$$\hat{X} = \sum_{\alpha} \hat{x}_\alpha \mathbf{I}_\alpha$$

is another $\mathcal{F}$ -measurable, integrable random variable that satisfies (1.4.2), then

$$\mathbb{P}[F_\alpha] \hat{x}_\alpha = \mathbb{E}[X \mathbf{I}_\alpha] = \mathbb{E}[\hat{X} \mathbf{I}_\alpha] = \mathbb{P}[F_\alpha] \hat{x}_\alpha, \quad \forall \alpha \in A,$$

so that $\hat{x}_\alpha = \bar{x}_\alpha, \forall \alpha$, i.e., $\bar{X}$ is uniquely determined by (1.4.2).

As a special case, suppose that $Y \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ has finite or countable range $\mathcal{Y}$. We obtain a countable measurable partition of Ω $(F_y)_{y \in \mathcal{Y}}$, $F_y = \{Y = y\}$. In this case

$$\hat{x}_y = \frac{1}{\mathbb{P}[\{Y = y\}]} \int_{\{Y=y\}} X(\omega) \mathbb{P}[d\omega].$$

If, additionally, the range of X is also finite or countable, then $\bar{X}$ coincides with the random variable $\mathbb{E}[X \mid Y]$ defined in Exercise 1.17.

If in (1.4.2) we set $F = \Omega$ we deduce

$$\mathbb{E}[X] = \mathbb{E}[\bar{X}] = \sum_{\alpha} \bar{x}_{\alpha} \mathbb{P}[F_{\alpha}] = \sum_{\alpha} \mathbb{E}[X \mid F_{\alpha}] \mathbb{P}[F_{\alpha}]. \quad (1.4.3)$$

When $X = I_S$, then

$$\mathbb{E}[I_S \mid F_{\alpha}] = \frac{\mathbb{P}[S \cap F_{\alpha}]}{\mathbb{P}[F_{\alpha}]} = \mathbb{P}[S \mid F_{\alpha}].$$

In this special case the equality (1.4.3) becomes the *law of total probability*

$$\mathbb{P}[S] = \sum_{\alpha} \mathbb{P}[S \mid F_{\alpha}] \mathbb{P}[F_{\alpha}]. \quad (1.4.4)$$

□

The next result explains why the condition (1.4.2) is key to our further developments.

Proposition 1.4.2. *If $\mathcal{F} \subset \mathcal{S}$ is a sigma-subalgebra and $Y_0, Y_1 \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ are two $\mathcal{F}$ -measurable random variables such that*

$$\mathbb{E}[Y_0 I_F] = \mathbb{E}[Y_1 I_F], \quad \forall F \in \mathcal{F}, \quad (1.4.5)$$

then $Y_0 = Y_1$ a.s..

Proof. Set $Z = Y_0 - Y_1$. Then Z is $\mathcal{F}$ -measurable, integrable and satisfies

$$\mathbb{E}[Z I_F] = 0, \quad \forall F \in \mathcal{F}. \quad (1.4.6)$$

If we let $F = \{Z > 1/n\}$, $n \in \mathbb{N}$, we deduce that

$$\frac{1}{n} \mathbb{P}[Z > 1/n] \leq \mathbb{E}[Z I_{\{Z > 1/n\}}] = 0, \quad \forall n \in \mathbb{N}.$$

Thus

$$\mathbb{P}[Z > 1/n] = 0, \quad \forall n \in \mathbb{N} \Rightarrow \mathbb{P}[Z > 0] = 0.$$

A similar argument shows that $\mathbb{P}[Z < 0] = 0$. □

Definition 1.4.3. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space, $\mathcal{F} \subset \mathcal{S}$ a sigma subalgebra, and $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$. A version of the conditional expectation of X given $\mathcal{F}$ is an $\mathcal{F}$ -measurable random variable $\bar{X} \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ such that

$$\mathbb{E}[X I_F] = \mathbb{E}[\bar{X} I_F], \quad \forall F \in \mathcal{F}. \quad (1.4.7)$$

□

According to Proposition 1.4.2, any two random variables $\bar{X}_0, \bar{X}_1 \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ satisfying (1.4.7) are a.s. equal. Their equivalence class in $L^1(\Omega, \mathcal{F}, \mathbb{P})$ is denoted by $\mathbb{E}[X \parallel \mathcal{F}]$ and it is called *the conditional expectation* of X given $\mathcal{F}$. Also, if $X = Y$ a.s. and $\mathbb{E}[X \parallel \mathcal{F}]$ exists, then $\mathbb{E}[Y \parallel \mathcal{F}]$ also exists and $\mathbb{E}[X \parallel \mathcal{F}] = \mathbb{E}[Y \parallel \mathcal{F}]$ a.s..

🔗 **About the notation.** I am using different notations, one for the conditional expectation given an event, $\mathbb{E}[X \mid F]$, and another for the conditional expectation given a sigma-subalgebra, $\mathbb{E}[X \parallel \mathcal{F}]$. The reason is simple: I want to emphasize visually that the first is a *number* and the latter is a *function* (random variable).

Remark 1.4.4. Using the Monotone Convergence Theorem and the Monotone Class Theorem we deduce that the following are equivalent.

- (i) The random variable $\bar{X} \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ is a representative of $\mathbb{E}[X \parallel \mathcal{F}]$
- (ii) For any $Y \in \mathcal{L}^\infty(\Omega, \mathcal{F}, \mathbb{P})$

$$\mathbb{E}[XY] = \mathbb{E}[\bar{X}Y]. \quad (1.4.8)$$

- (iii) There exists a π -system $\mathcal{A} \subset \mathcal{F}$ that contains Ω , generates $\mathcal{F}$, and

$$\mathbb{E}[XI_A] = \mathbb{E}[\bar{X}I_A], \quad \forall A \in \mathcal{A}. \quad (1.4.9)$$

From Corollary 1.3.8 we deduce that

$$\bar{X} \text{ is } \mathcal{F} \text{ measurable, } \mathbb{E}[X - \bar{X}] = 0 \text{ and } (X - \bar{X}) \perp \mathcal{F} \Rightarrow \bar{X} = \mathbb{E}[X \parallel \mathcal{F}]. \quad (1.4.10)$$

□

We will soon prove (Theorem 1.4.8) that the conditional expectation of an integrable random variable given a sigma-subalgebra exists.

Definition 1.4.5. Given random variables $X \in L^0(\Omega, \mathcal{S}, \mathbb{P})$, $Y \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ we write

$$\boxed{\mathbb{E}[Y \parallel X] := \mathbb{E}[Y \parallel \sigma(X)]},$$

where $\sigma(X)$ denotes the sigma-subalgebra generated by X . This random variable is called the *conditional expectation of Y given X* . □

Remark 1.4.6. A function $\bar{Y} \in \mathcal{L}^1(\Omega, \sigma(X), \mathbb{P})$ represents $\mathbb{E}[Y \parallel X]$ if, for any $x \in \mathbb{R}$ we have

$$\int_{\{X \leq x\}} Y(\omega) \mathbb{P}[d\omega] = \int_{\{X \leq x\}} \bar{Y}(\omega) \mathbb{P}[d\omega].$$

Since $\mathbb{E}[Y \parallel X]$ is $\sigma(X)$ -measurable we deduce from Dynkin's Theorem 1.1.24 that there exists a Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(X) = \mathbb{E}[Y \parallel X] \quad \text{a.s.}$$

This is equivalent to the statement

$$\mathbb{E}[Y I_{\{X \leq x\}}] = \mathbb{E}[f(X) I_{\{X \leq x\}}], \quad \forall x \in \mathbb{R}. \quad (1.4.11)$$

“The value $f(x)$ of the function f at x ”⁸ is called the *conditional expectation of Y given $X = x$* and it is denoted by $\mathbb{E}[Y \mid X = x]$. Think of it as the conditional expectation of Y given the possible

⁸We used quotes since “the value at a point” is not a precise concept for a function unique up to almost everywhere equality.

negligible event $\{X = x\}$. The graph of $x \mapsto \mathbb{E}[Y | X = x]$ was classically referred to as *the regression curve*.

Note that

$$\mathbb{E}[Y] = \mathbb{E}[\bar{Y}] = \mathbb{E}[\mathbb{E}[Y | X]] = \mathbb{E}[f(X)].$$

Thus

$$\mathbb{E}[Y] = \mathbb{E}[f(X)] = \int_{\mathbb{R}} f(x) \mathbb{P}_X[dx].$$

We can rewrite the last equality as

$$\mathbb{E}[Y] = \int_{\mathbb{R}} \mathbb{E}[Y | X = x] \mathbb{P}_X[dx]. \quad (1.4.12)$$

This approach to computing the expectation of Y by relying on the above identity is referred to *computing the expectation of Y by conditioning on X* . This generalizes the elementary situation in Exercise 1.17. $\square$

Example 1.4.7. Suppose that $X, Y : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ are two random variables such that their joint probability distribution $\mathbb{P}_{X,Y} \in \text{Prob}(\mathbb{R}^2)$ is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^2$. This means that there exists a Lebesgue integrable function

$$p_{X,Y} : \mathbb{R}^2 \rightarrow [0, \infty)$$

such that

$$\mathbb{P}[(X, Y) \in B] = \int_B p_{X,Y}(x, y) dx dy, \quad \forall B \in \mathcal{B}_{\mathbb{R}^2}.$$

We denote by $\mathbb{P}_X$ and respectively $\mathbb{P}_Y$ the probability distributions of X and respectively Y . Note that the cumulative distribution function F_X of X is

$$F_X(c) = \mathbb{P}[X \leq c] = \int_{-\infty}^c \underbrace{\left(\int_{\mathbb{R}} p_{X,Y}(x, y) dy \right)}_{=: p_X(x)} dx = \int_{-\infty}^c p_X(x) dx.$$

This shows that $\mathbb{P}_X$ is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$ and

$$\mathbb{P}_X[dx] = p_X(x) dx.$$

Similarly

$$\mathbb{P}_Y[dy] = p_Y(y) dy = \int_{\mathbb{R}} p_{X,Y}(x, y) dx.$$

Classically, the probability distributions $\mathbb{P}_X$ and $\mathbb{P}_Y$ are called the *marginal distributions* of the random vector (X, Y) . We define

$$p_{Y|X=x}(y) := \begin{cases} \frac{p_{X,Y}(x, y)}{p_X(x)}, & p_X(x) \neq 0, \\ 0, & p_X(x) = 0. \end{cases}$$

Assume that Y is integrable. Define

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \int_{\mathbb{R}} y p_{Y|X=x}(y) dy$$

$$= \begin{cases} \frac{1}{p_X(x)} \int_{\mathbb{R}} y p_{X,Y}(x, y) dy, & p_X(x) \neq 0, \\ 0, & p_X(x) = 0. \end{cases}$$

Using the Fubini-Tonelli theorem and the integrability of Y we deduce that the above integrals are well defined and the resulting function f is Borel measurable. Note that

$$f(x)p_X(x) = \int_{\mathbb{R}} y p_{X,Y}(x, y) dy, \quad \forall x \in \mathbb{R}.$$

We want to show that $f(x) = \mathbb{E}[Y | X = x]$, i.e., $f(X)$ is a version of $\mathbb{E}[Y | X]$. We will show that it satisfies (1.4.11).

Let $c \in \mathbb{R}$. We have

$$\begin{aligned} \mathbb{E}[f(X)\mathbf{I}_{X \leq c}] &= \int_{-\infty}^c f(x)p_X(x)dx = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} y p_{X,Y}(x, y) dy \right) \mathbf{I}_{(-\infty, c]}(x) dx \\ &= \int_{\mathbb{R}^2} y \mathbf{I}_{(-\infty, c]}(x) p_{X,Y}(x, y) dx dy = \mathbb{E}[Y \mathbf{I}_{X \leq c}]. \end{aligned} \quad (1.4.13)$$

The function $f(x)$ is the conditional expectation $\mathbb{E}[Y | X = x]$ discussed in Remark 1.4.4.

Note that the event $\{X = x\}$ has probability zero so this nomenclature should be taken with a grain of sand since we cannot apply (1.4.1). Intuitively

$$\mathbb{E}[Y | X = x] = \lim_{\varepsilon \searrow 0} \mathbb{E}[Y | \{|X - x| < \varepsilon\}] \stackrel{(1.4.1)}{=} \lim_{\varepsilon \searrow 0} \frac{\mathbb{E}[Y \mathbf{I}_{|X-x|<\varepsilon}]}{\mathbb{P}[|X-x|<\varepsilon]}.$$

□

One issue we need to address is the existence of the conditional expectation. There is a fast proof based on the Radon–Nikodym theorem. We will use a more roundabout approach that sheds additional light on probabilistic the nature of conditional expectation. As an aside, let us mention that this approach leads to an alternate proof of the Radon–Nikodym theorem that does not rely on the concept of signed-measure.

Theorem 1.4.8. *For any $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and any sigma subalgebra $\mathcal{F} \subset \mathcal{S}$ there exists a conditional expectation $\mathbb{E}[X | \mathcal{F}] \in L^1(\Omega, \mathcal{F}, \mathbb{P})$.*

Proof. We follow the approach in [224]. We establish the existence gradually, first under more restrictive assumptions.

Step 1. Assume $X \in L^2(\Omega, \mathcal{S}, \mathbb{P})$. Then $L^2(\Omega, \mathcal{F}, \mathbb{P})$ is a closed subspace of $L^2(\Omega, \mathcal{S}, \mathbb{P})$. Denote by $P_{\mathcal{F}}X$ the orthogonal projection of X on this closed subspace. We claim that

$$P_{\mathcal{F}}X = \mathbb{E}[X | \mathcal{F}], \quad (1.4.14a)$$

$$X \geq 0 \Rightarrow \mathbb{E}[X | \mathcal{F}] \geq 0. \quad (1.4.14b)$$

Set $Y := P_{\mathcal{F}}X$. Since $X - Y \perp L^2(\Omega, \mathcal{F}, \mathbb{P})$ we deduce

$$\mathbb{E}[(X - Y)Z] = 0, \quad \forall Z \in L^2(\Omega, \mathcal{F}, \mathbb{P}).$$

In particular,

$$\mathbb{E}[(X - Y)\mathbf{I}_F] = 0, \quad \forall F \in \mathcal{F}.$$

This proves (1.4.14a). Now suppose that $X \geq 0$. For any $n \in \mathbb{N}$ we have

$$0 \leq \mathbb{E}[X \mathbf{I}_{\{Y \leq -1/n\}}] = \mathbb{E}[Y \mathbf{I}_{\{Y \leq -1/n\}}] \leq -\frac{1}{n} \mathbb{P}[Y \leq -1/n],$$

so

$$\mathbb{P}[Y \leq -1/n] = 0, \quad \forall n \in \mathbb{N}.$$

This proves (1.4.14b). Clearly, the resulting map

$$L^2(\Omega, \mathcal{S}, \mathbb{P}) \ni X \mapsto \mathbb{E}[X \mid \mathcal{F}] \in L^2(\Omega, \mathcal{F}, \mathbb{P})$$

is linear.

Step 2. Assume $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Decompose $X = X^+ - X^-$ and, for $n \in \mathbb{N}$, set

$$X_n^\pm = \min(X^\pm, n).$$

Note that $X_n^\pm \in L^\infty(\Omega, \mathcal{S}, \mathbb{P})$ and, as $n \rightarrow \infty$, $X_n^\pm \nearrow X^\pm$ a.s.. From Step 1 we deduce that the random variables $X_n^\pm$ have conditional expectations given $\mathcal{F}$. Choose versions

$$Y_n^\pm := \mathbb{E}[X_n^\pm \mid \mathcal{F}].$$

Since $X_n^\pm - X_m^\pm \geq 0$ a.s. if $m \leq n$ we deduce from (1.4.14b) that

$$0 \leq Y_m^\pm \leq Y_n^\pm, \quad \text{a.s., } \forall m \leq n.$$

We set

$$Y^\pm := \lim_{n \rightarrow \infty} Y_n^\pm.$$

From the Monotone Convergence Theorem we deduce that

$$\infty > \mathbb{E}[X^\pm] = \lim_{n \rightarrow \infty} \mathbb{E}[X_n^\pm] = \lim_{n \rightarrow \infty} \mathbb{E}[Y_n^\pm] = \mathbb{E}[Y^\pm].$$

This shows that the random variables $Y_\pm$ are integrable and in particular a.s. finite. We set

$$Y := Y^+ - Y^-.$$

We will show that Y is a version of the conditional expectation of X given $\mathcal{F}$. Let $F \in \mathcal{F}$. Then

$$\begin{aligned} \mathbb{E}[X \mathbf{I}_F] &= \mathbb{E}[X^+ \mathbf{I}_F] - \mathbb{E}[X^- \mathbf{I}_F] = \lim_{n \rightarrow \infty} \mathbb{E}[X_n^+ \mathbf{I}_F] - \lim_{n \rightarrow \infty} \mathbb{E}[X_n^- \mathbf{I}_F] \\ &= \lim_{n \rightarrow \infty} \mathbb{E}[Y_n^+ \mathbf{I}_F] - \lim_{n \rightarrow \infty} \mathbb{E}[Y_n^- \mathbf{I}_F] = \mathbb{E}[Y^+ \mathbf{I}_F] - \mathbb{E}[Y^- \mathbf{I}_F] = \mathbb{E}[Y \mathbf{I}_F]. \end{aligned}$$

This proves that Y is a version of $\mathbb{E}[X \mid \mathcal{F}]$. $\square$

Remark 1.4.9. (a) The sigma-subalgebra $\mathcal{F}$ should be viewed as encoding partial information that we have about a random experiment. Following a terminology frequently used in statistics, we refer to the $\mathcal{F}$ -measurable random variables as *predictors* determined by the information contained in $\mathcal{F}$.

Step 1 in the above proof shows that the conditional expectation $\bar{X}$ of a random variable X , given the partial information $\mathcal{F}$, should be viewed as the predictor that best approximates X given the information $\mathcal{F}$. The missing part $X - \bar{X}$ is independent of $\mathcal{F}$ so it is unknowable given only the information encoded by $\mathcal{F}$.

Intuitively, suppose we perform a random experiment with space of outcomes Ω . The result of one experiment is an outcome ω . We have at our disposal a “dæmon”⁹ who can only give yes or no answers to questions of the type: given $F \in \mathcal{F}$, does ω belong to F ? Then $\bar{X}(\omega)$ is the best guess about $X(\omega)$ using the “dæmonic information” available to us.

⁹Use the concept of dæmon in Socratic sense

Note that when $\mathcal{F} = \{\emptyset, \Omega\}$, then

$$\mathbb{E}[X] = \mathbb{E}[X] \mathbf{I}_\Omega.$$

To put it differently, if the only information we have about a random experiment is that there will be an outcome, then the most/best we can predict about a numerical characteristic of that outcome is its expectation.

(b) There is an alternate approach to proving the existence of conditional expectation. A random variable $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ defines a signed measure

$$\mu_X : \mathcal{F} \rightarrow [0, \infty), \quad \mu_X[F] = \int_F X(\omega) \mathbb{P}[d\omega], \quad \forall F \in \mathcal{F}.$$

This measure is absolutely continuous with $\mathbb{P}$ (restricted to $\mathcal{F}$). The Radon-Nikodym theorem implies that there exists an $\mathcal{F}$ -measurable integrable function $\rho_X \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mu_X[d\omega] = \rho_X(\omega) \mathbb{P}[d\omega]$, i.e.,

$$\int_F X(\omega) \mathbb{P}[d\omega] = \int_F \rho_X(\omega) \mathbb{P}[d\omega], \quad \forall F \in \mathcal{F}.$$

This shows that $\rho_X = \mathbb{E}[X \mid \mathcal{F}]$ a.s..

Conversely, one can show with considerable effort and ingenuity that the existence of conditional expectations implies the Radon-Nikodym Theorem. We refer to [224, Sec. 14.15] for details. $\square$

Definition 1.4.10. Given a sigma subalgebra $\mathcal{F} \subset \mathcal{S}$, and an event $S \in \mathcal{S}$, we define the *conditional probability* of S given $\mathcal{F}$ to be the random variable

$$\mathbb{P}[S \mid \mathcal{F}] := \mathbb{E}[\mathbf{I}_S \mid \mathcal{F}]. \quad \square$$

Example 1.4.11 (Conditioning on an event). Suppose that $S \in \mathcal{S}$ is an event such $0 < \mathbb{P}[S] < 1$. Let $Y \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then

$$\mathbb{E}[Y \mid \mathbf{I}_S] = \mathbb{E}[Y \mid S] \mathbf{I}_S + \mathbb{E}[Y \mid S^c] \mathbf{I}_{S^c},$$

where we recall that (see (1.4.1))

$$\mathbb{E}[Y \mid S] = \frac{1}{\mathbb{P}[S]} \mathbb{E}[Y \mathbf{I}_S]. \quad \square$$

Our next result lists the main properties of the conditional expectation.

Theorem 1.4.12. Suppose that $\mathcal{F} \subset \mathcal{S}$ is a sigma subalgebra. Then the following hold.

(i) Let $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. If Y is any version of $\mathbb{E}[X \mid \mathcal{F}]$, then $\mathbb{E}[Y] = \mathbb{E}[X]$. In other words

$$\mathbb{E}[\mathbb{E}[X \mid \mathcal{F}]] = \mathbb{E}[X]. \quad (1.4.15)$$

(ii) If $X, Y \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and $X \leq Y$ a.s., then $\mathbb{E}[X \mid \mathcal{F}] \leq \mathbb{E}[Y \mid \mathcal{F}]$ a.s..

(iii) The map

$$L^1(\Omega, \mathcal{S}, \mathbb{P}) \ni X \mapsto \mathbb{E}[X \mid \mathcal{F}] \in L^1(\Omega, \mathcal{F}, \mathbb{P})$$

is a linear contraction, i.e., it is linear and satisfies

$$\|\mathbb{E}[X \mid \mathcal{F}]\|_{L^1} \leq \|X\|_{L^1}, \quad \forall X \in L^1(\Omega, \mathcal{S}, \mathbb{P}).$$

(iv) If $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and $Y \in L^\infty(\Omega, \mathcal{F}, \mathbb{P})$, then

$$\mathbb{E}[XY \mid \mathcal{F}] = Y \mathbb{E}[X \mid \mathcal{F}].$$

(v) If $\mathcal{G} \subset \mathcal{F}$ is another sigma subalgebra, then for any $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ we have

$$\mathbb{E}[X \parallel \mathcal{G}] = \mathbb{E}[\mathbb{E}[X \parallel \mathcal{F}] \parallel \mathcal{G}].$$

(vi) If $0 \leq X_n \nearrow X$ a.s., $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$, then

$$\mathbb{E}[X_n \parallel \mathcal{F}] \nearrow \mathbb{E}[X \parallel \mathcal{F}], \text{ a.s. and } L^1.$$

(vii) If $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$, $n \in \mathbb{N}$, $X_n \geq 0$ a.s., $\liminf X_n \in L^1$ a.s., then

$$\mathbb{E}[\liminf X_n \parallel \mathcal{F}] \leq \liminf \mathbb{E}[X_n \parallel \mathcal{F}] \text{ a.s..}$$

(viii) If $X_n \rightarrow X$ a.s. and there exists $Y \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that $|X_n| \leq Y$ a.s., then

$$\mathbb{E}[X_n \parallel \mathcal{F}] \rightarrow \mathbb{E}[X \parallel \mathcal{F}] \text{ a.s..}$$

(ix) If $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function such that $\varphi(X)$ is integrable, then

$$\varphi(\mathbb{E}[X \parallel \mathcal{F}]) \leq \mathbb{E}[\varphi(X) \parallel \mathcal{F}] \text{ a.s..}$$

In particular, if we choose $\varphi(x) = |x|^p$, $p \geq 1$ we deduce that the conditional expectation defines a linear map

$$\mathbb{E}[\cdot \parallel \mathcal{F}] : L^p(\Omega, \mathcal{S}, \mathbb{P}) \rightarrow L^p(\Omega, \mathcal{F}, \mathbb{P})$$

that is linear contraction, i.e.,

$$\|\mathbb{E}[X \parallel \mathcal{F}]\|_{L^p} \leq \|X\|_{L^p}.$$

(x) If $\mathcal{G}$ is another sigma-algebra that is independent of $\sigma(X) \vee \mathcal{F}$, then

$$\mathbb{E}[X \parallel \mathcal{F} \vee \mathcal{G}] = \mathbb{E}[X \parallel \mathcal{F}].$$

In particular, if $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ is independent of $\mathcal{G}$, then

$$\mathbb{E}[X \parallel \mathcal{G}] = \mathbb{E}[X].$$

Proof. (i) Follows by choosing $F = \Omega$ in (1.4.7). (ii) Follows from the proof of Theorem 1.4.8.

(iii) The linearity follows from the fact that the defining condition (1.4.7) is linear in X . Now let $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. We have $X = X^+ - X^-$. Choose versions $Y^\pm$ of $\mathbb{E}[X^\pm \parallel \mathcal{F}]$. Then $Y_\pm \geq 0$ and

$$\begin{aligned} |\mathbb{E}[X \parallel \mathcal{F}]| &= |Y^+ - Y^-| \leq Y^+ + Y^- \\ &= \mathbb{E}[X^+ + X^- \parallel \mathcal{F}] = \mathbb{E}[|X| \parallel \mathcal{F}]. \end{aligned}$$

Hence

$$\|\mathbb{E}[X \parallel \mathcal{F}]\|_{L^1} \leq \mathbb{E}[\mathbb{E}[|X| \parallel \mathcal{F}]] = \mathbb{E}[|X|] = \|X\|_{L^1}.$$

(iv) Choose a version Z of $\mathbb{E}[X \parallel \mathcal{F}]$. Let $Y \in L^\infty(\Omega, \mathcal{F}, \mathbb{P})$. We have to show that YZ is a version of $\mathbb{E}[XY \parallel \mathcal{F}]$, i.e.,

$$\mathbb{E}[XY \mathbf{I}_F] = \mathbb{E}[ZY \mathbf{I}_F], \quad \forall F \in \mathcal{F}. \quad (1.4.16)$$

Let $F \in \mathcal{F}$. Since Z is a version of $\mathbb{E}[X \parallel \mathcal{F}]$ we deduce from (1.4.8) that

$$\mathbb{E}[XU] = \mathbb{E}[ZU], \quad \forall U \in L^\infty(\Omega, \mathcal{F}, \mathbb{P}).$$

In particular, $\forall F \in \mathcal{F}$ we have

$$\mathbb{E}[\underbrace{XY \mathbf{I}_F}_U] = \mathbb{E}[ZU] = \mathbb{E}[ZY \mathbf{I}_F].$$

Thus ZY satisfies (1.4.16).

(v) Choose a version Y of $\mathbb{E}[X \mid \mathcal{F}]$, and a version Z of $\mathbb{E}[Y \mid \mathcal{G}]$. We have to show that Z is also a version of $\mathbb{E}[X \mid \mathcal{G}]$. Let $G \in \mathcal{G}$. We have

$$\mathbb{E}[Y \mathbf{I}_B] = \mathbb{E}[Z \mathbf{I}_B],$$

$$\mathbb{E}[X \mathbf{I}_G] \stackrel{G \in \mathcal{F}}{=} \mathbb{E}[Y \mathbf{I}_G] = \mathbb{E}[Z \mathbf{I}_B].$$

(vi) Choose versions Y_n of $\mathbb{E}[X_n \mid \mathcal{F}]$ and Y of $\mathbb{E}[X \mid \mathcal{F}]$. Note that Y_n is increasing. The Monotone Convergence theorem implies that $\|X - X_n\|_{L^1} \rightarrow 0$. From (iii) we deduce

$$\|Y_n - Y\|_{L^1} \leq \|X_n - Y\|_{L^1}.$$

Proposition 1.3.65 implies that Y_n admits a subsequence that converges a.s. to Y . Since the sequence Y_n is increasing we deduce that the whole sequence converges a.s. to Y .

(vii) Set

$$Y_k = \inf_{n \geq k} X_n.$$

The sequence of random variables (Y_k) is increasing and converges a.s. to $X := \liminf X_n$. We deduce from (vi) that

$$\mathbb{E}[Y_k \mid \mathcal{F}] \nearrow \mathbb{E}[X \mid \mathcal{F}].$$

Note that since $Y_k \leq X_n, \forall n \geq k$, we have

$$\mathbb{E}[Y_k \mid \mathcal{F}] \leq Z_k := \inf_{n \geq k} \mathbb{E}[X_n \mid \mathcal{F}]$$

so

$$\mathbb{E}[X \mid \mathcal{F}] = \lim_k \mathbb{E}[Y_k \mid \mathcal{F}] \leq \lim_k Z_k = \liminf_k \mathbb{E}[X_n \mid \mathcal{F}].$$

(viii) Set $Y_n := X_n + Y$. Then $Y_n \geq 0$ and $Y_n \rightarrow X + Y$ a.s. We deduce from (vii) that

$$\mathbb{E}[X \mid \mathcal{F}] + \mathbb{E}[Y \mid \mathcal{F}] \leq \liminf \mathbb{E}[X_n \mid \mathcal{F}] + \mathbb{E}[Y \mid \mathcal{F}]$$

i.e.,

$$\mathbb{E}[X \mid \mathcal{F}] \leq \liminf \mathbb{E}[X_n \mid \mathcal{F}].$$

Similarly, we set $Z_n = Y - X_n$. Then $Z_n \geq 0$ and $Z_n \rightarrow Y - X$ a.s.. Applying (vii) to Z_n we deduce

$$\limsup \mathbb{E}[Z_n \mid \mathcal{F}] \leq \mathbb{E}[Z \mid \mathcal{F}].$$

(ix) We need to use a less familiar property of convex functions, [6, Thm.6.3.4]. More precisely, there exist *sequences* of real numbers $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ such that

$$\varphi(x) = \sup_{n \in \mathbb{N}} (a_n x + b_n), \quad \forall x \in \mathbb{R}.$$

Set $\ell_n(x) = a_n x + b_n$.¹⁰ Clearly

$$\ell_n(\mathbb{E}[X \mid \mathcal{F}]) = \mathbb{E}[\ell_n(X) \mid \mathcal{F}] \leq \mathbb{E}[\varphi(X) \mid \mathcal{F}].$$

Hence

$$\begin{aligned} \varphi(\mathbb{E}[X \mid \mathcal{F}]) &= \sup_{n \in \mathbb{N}} \ell_n(\mathbb{E}[X \mid \mathcal{F}]) \\ &= \sup_{n \in \mathbb{N}} \mathbb{E}[\ell_n(X) \mid \mathcal{F}] \leq \mathbb{E}[\varphi(X) \mid \mathcal{F}]. \end{aligned}$$

¹⁰When φ is C^1 the family ℓ_n coincides with the family of tangent lines $(\ell_q)_{q \in \mathbb{Q}}$, $\ell_q(x) = \varphi'(q)(x - q) + \varphi(q)$.

(x) Let $G \in \mathcal{G}$ and F in $\mathcal{F}$. Then, the random variables $\mathbf{I}_G$ and $X\mathbf{I}_F$ are independent so

$$\mathbb{E}[X\mathbf{I}_{F \cap G}] = \mathbb{E}[X\mathbf{I}_F\mathbf{I}_G] = \mathbb{E}[X\mathbf{I}_F]\mathbb{P}[G].$$

If Y is a version of $\mathbb{E}[X \mid \mathcal{F}]$, then Y is $\mathcal{F}$ -measurable and thus independent of G , so

$$\begin{aligned} \mathbb{E}[Y\mathbf{I}_{F \cap G}] &= \mathbb{E}[Y\mathbf{I}_F\mathbf{I}_G] = \mathbb{E}[Y\mathbf{I}_F]\mathbb{P}[G] \\ &= \mathbb{E}[X\mathbf{I}_F]\mathbb{P}[G] = \mathbb{E}[X\mathbf{I}_{F \cap G}], \quad \forall F \in \mathcal{F}, G \in \mathcal{G}. \end{aligned}$$

Since the collection

$$\{F \cap G; F \in \mathcal{F}, G \in \mathcal{G}\}$$

is a π -system generating $\mathcal{F} \vee \mathcal{G}$, we deduce from Dynkin's ($\pi - \lambda$) theorem that

$$\mathbb{E}[Y\mathbf{I}_S] = \mathbb{E}[X\mathbf{I}_S], \quad \forall S \in \mathcal{F} \vee \mathcal{G},$$

so that $\mathbb{E}[X \mid \mathcal{F} \vee \mathcal{G}] = Y$, i.e., $\mathbb{E}[X \mid \mathcal{F}] = \mathbb{E}[X \mid \mathcal{F} \vee \mathcal{G}]$. $\square$

1.4.2. Some applications of conditioning. To give the reader a taste of the power and uses of conditional expectation we describe some nontrivial and less advertised uses of conditional expectation.

Example 1.4.13. Suppose that a player rolls a die an indefinite amount of times. More formally, we are given a sequence independent random variables $(X_n)_{n \in \mathbb{N}}$, uniformly distributed on

$$\mathbb{I}_6 = \{1, 2, \dots, 6\}.$$

For $k \in \mathbb{N}$, we say that a k -run of length k occurred at time n if $n \geq k$ and

$$X_n = X_{n-1} = \dots = X_{n-k+1} = 6.$$

We set

$$R = R_k := \{n; \text{ a } k\text{-run occurred at time } n\} \subset \mathbb{N} \cup \{\infty\}, \quad T = T_k = \inf R_k,$$

where $\inf \emptyset := \infty$. Thus T is the moment when the first k -run is observed. We want to show that $\mathbb{E}[T] < \infty$.

Note that for each $n \in \mathbb{N}$ the event $\{T \leq n\}$ belongs to the sigma algebra $\mathcal{F}_n$ generated by $X_1, \dots, X_n$. The explanation is simple: if we know the results of the first n rolls of the die we can decide if a k -run was occurred. Consider the conditional probability

$$\mathbb{P}[\{T \leq n+k\} \mid \mathcal{F}_n] = \mathbb{E}[\mathbf{I}_{\{T \leq n+k\}} \mid \mathcal{F}_n].$$

This conditional probability is a *random variable*. Since the sigma-algebra $\mathcal{F}_n$ is defined by the partition

$$S_{i_1, \dots, i_n} := \{X_1 = i_1, \dots, X_n = i_n\}, \quad i_j \in \{1, \dots, 6\},$$

we see that $\mathbb{P}[T \leq n+k \mid \mathcal{F}_n]$ has the form

$$\mathbb{P}[T \leq n+k \mid \mathcal{F}_n] = \sum_{i_1, \dots, i_n=1}^6 p_{i_1, \dots, i_n|k} \mathbf{I}_{S_{i_1, \dots, i_n}},$$

where

$$p_{i_1, \dots, i_n|k} = \mathbb{P}[T \leq n+k \mid X_1 = i_1, \dots, X_n = i_n].$$

Note that, irrespective of the i_j -s, we have

$$p_{i_1, \dots, i_n|k} \geq \frac{1}{6^k} =: r.$$

Hence

$$\mathbb{P}[T \leq n + k \mid \mathcal{F}_n] \geq r, \quad \forall n.$$

In particular,

$$\mathbb{P}[T > n + k \mid \mathcal{F}_n] \leq (1 - r) < 1, \quad \forall n \in \mathbb{N}.$$

Now observe that for any $n \in \mathbb{N}$, $\ell \in \mathbb{N}_0$ we have $\{T > n + \ell k \mid \mathcal{F}_{n+\ell k}\}$. Hence

$$\begin{aligned} \mathbb{P}[T > n + (\ell + 1)k] &= \mathbb{E}[\mathbf{I}_{\{T > n + (\ell + 1)k\}} \mathbf{I}_{\{T > n + \ell k\}}] \\ &= \mathbb{E}[\mathbf{I}_{\{T > n + \ell k\}} \mathbb{E}[T > n + (\ell + 1)k \mid \mathcal{F}_{n+\ell k}]] \\ &\leq (1 - r) \mathbb{E}[\mathbf{I}_{\{T > n + \ell k\}}] = (1 - r) \mathbb{P}[T > n + \ell k]. \end{aligned}$$

Iterating, we deduce that for any $i \in \{1, \dots, k\}$ and any $\ell \in \mathbb{N}$ we have

$$\mathbb{P}[T > i + \ell k] < (1 - r)^\ell \mathbb{P}[T > i] \leq (1 - r)^\ell.$$

Now observe that

$$\mathbb{E}[T] = \sum_{n \in \mathbb{N}_0} \mathbb{P}[T > n] = \sum_{i=1}^k \sum_{\ell \in \mathbb{N}_0} \mathbb{P}[T > i + \ell k] < \sum_{i=1}^k \sum_{\ell \in \mathbb{N}_0} (1 - r)^\ell = \frac{k}{r} < \infty.$$

This proves that $\mathbb{E}[T]$ is finite. In Example 3.1.33 we will use martingale techniques to show that

$$\mathbb{E}[T] = \frac{6^{k+1} - 6}{5}.$$

□

Example 1.4.14 (Optimal stopping with finite horizon). Let us consider the following abstract situation.

Suppose we are given N random variables

$$X_1, \dots, X_N \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P}).$$

For $n \in \mathbb{I}_N := \{1, 2, \dots, N\}$ we denote by $\mathcal{F}_n$ the sigma-algebra generated by $X_1, \dots, X_n$. Suppose that we are also given a sequence of rewards

$$R_n \in \mathcal{L}^1(\Omega, \mathcal{F}_n, \mathbb{P}), \quad n \in \mathbb{I}_N.$$

A *stopping time* is a random variable

$$T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{I}_N$$

such that $\{T \leq n\} \in \mathcal{F}_n$, $\forall n \in \mathbb{I}_N$. Equivalently, T is a stopping time if and only if $\{T = n\} \in \mathcal{F}_n$, $\forall n$. Note that if T is a stopping time, then $\{T \geq n\} = \Omega \setminus \{T \leq n - 1\} \in \mathcal{F}_{n-1}$.

One should think of the collection $X_1, \dots, X_N$ as a finite stream of random quantities flowing in time, one quantity per unit of time. The reward R_n depends only on the observed values $X_1, \dots, X_n$, i.e., $R_n = R_n(X_1, \dots, X_n)$. A stopping time describes a decision when to stop the stream based only on the information accumulated up to the decision moment. After we observe the first quantity X_1 , we can decide if $T = 1$. If this not the case, we observe a second quantity and, using the information about X_1 , and X_2 we can decide to stop, i.e., if $T = 2$ or not. We continue until we either observe all the random quantities or at the first n such that $T = n$.

We set

$$R_T = \sum_{n \in \mathbb{I}_N} R_n \mathbf{I}_{\{T=n\}}.$$

In other words R_T is the reward at the random stopping time T . We denote by $\mathcal{T}$ the collection of all possible stopping times. Note that

$$\mathbb{E}[|R_T|] \leq \sum_{n=1}^N \mathbb{E}[|R_n|] < \infty, \quad \forall T \in \mathcal{T}.$$

We want to show that there exists $T_* \in \mathcal{T}$ such that

$$\mathbb{E}[R_{T_*}] = r := \sup_{T \in \mathcal{T}} \mathbb{E}[R_T]$$

Such a T_* is called an *optimal stopping time*. To prove the existence of an optimal time we establish a Fermat-like optimality condition that the optimal stopping times satisfy. We follow the dynamic programming strategy in [36, Chap. 3].

For $n \in \mathbb{I}_N$ we set

$$\mathcal{T}_n := \{ T \in \mathcal{T}; \quad T \geq n \}.$$

Note that

$$\mathcal{T} = \mathcal{T}_1 \supset \mathcal{T}_2 \supset \cdots \supset \mathcal{T}_N.$$

A stopping time T belongs to $\mathcal{T}_n$ if and only if the decision to stop comes only after we have observed the first n random variables in the stream, $X_1, \dots, X_n$.

We will detect an optimal stopping strategy using a process of “successive approximations”. The first approximation is the simplest strategy: pick the reward only at the end, after we have observed all the N variables in the stream. In this case the reward is $Y_N = R_N$. This may not give us the largest expected reward because some of the up-stream rewards could have been higher. We tweak this strategy a bit to produce a better outcome.

We wait to observe the first $N - 1$ variables in the stream, and then decide what to do. At this moment our reward is R_{N-1} . To decide what to do next we compare this reward with the expected reward R_N given that we observed $X_1, \dots, X_{N-1}$, i.e., with the conditional expectation $\mathbb{E}[R_N \mid \mathcal{F}_{N-1}]$. This is an $\mathcal{F}_{N-1}$ -measurable quantity, i.e., a quantity that is computable from the knowledge of $X_1, \dots, X_{N-1}$.

If the reward R_{N-1} that what we have in our hands is bigger than we expect to gain given our current information, we choose it and we stop. If not, we wait one more step to stop. More formally, we stop after $N - 1$ steps if $R_{N-1} \geq \mathbb{E}[R_N \mid \mathcal{F}_{N-1}]$ and we continue one more step otherwise. The decision is thus based on the random variable $Y_{N-1} = \max(R_{N-1}, \mathbb{E}[R_N \mid \mathcal{F}_{N-1}])$.

This heuristic suggests the following backwards induction.

$$Y_N := R_N, \quad Y_n := \max \{ R_n, \mathbb{E}[Y_{n+1} \mid \mathcal{F}_n] \}, \quad (1.4.17a)$$

$$T_n := \min \{ i \geq n; \quad R_i \geq Y_i \} = \min \{ i \geq n; \quad R_i = Y_i \}. \quad (1.4.17b)$$

Note that $T_n \geq n$ and, for any $k \geq n$,

$$\{ T_n > k \} = \{ R_k < \mathbb{E}[Y_{k+1} \mid \mathcal{F}_k] \} \in \mathcal{F}_k.$$

Hence $T_n \in \mathcal{T}_n$. We claim that for any $n = 1, \dots, N$ we have

$$Y_n \geq \mathbb{E}[R_T \mid \mathcal{F}_n], \quad \forall T \in \mathcal{T}_n. \quad (1.4.18a)$$

$$\mathbb{E}[R_{T_n} \mid \mathcal{F}_n] = Y_n. \quad (1.4.18b)$$

Hence

$$\mathbb{E}[R_{T_n} \mid \mathcal{F}_n] \geq \mathbb{E}[Y_n] = \mathbb{E}[R_T \mid \mathcal{F}_n], \quad \forall T \in \mathcal{T}_n.$$

By taking expectations we deduce

$$\mathbb{E}[R_{T_n}] = \sup_{T \in \mathcal{T}_n} \mathbb{E}[R_T]. \quad (1.4.19)$$

In particular, this shows that the stopping time T_1 is optimal.

The optimal stopping strategy T_1 has a natural description:

stop at the first moment when the reward at hand is greater or equal than the expected future reward, given the information we have at that moment.

The stopping strategy T_n is similar, but delayed for n units of times. We will prove (1.4.18a) and (1.4.18b) by backwards induction on n .

The inequality (1.4.18a) is clearly true for $n = N$. Assume it is true for n . Let $T \in \mathcal{T}_{n-1}$ and set $T' = \max\{T, n\}$. Then $T' \in \mathcal{T}_n$. For $A \in \mathcal{F}_{n-1}$ we have

$$\begin{aligned} \int_A R_T &= \int_{A \cap \{T=n-1\}} R_{n-1} + \int_{A \cap \{T \geq n\}} R_{T'} \\ (\{T \geq n\} \in \mathcal{F}_{n-1}) & \\ &= \int_{A \cap \{T=n-1\}} R_{n-1} + \int_{A \cap \{T \geq n\}} \mathbb{E}[R_{T'} \mid \mathcal{F}_{n-1}] \\ &= \int_{A \cap \{T=n-1\}} R_{n-1} + \int_{A \cap \{T \geq n\}} \mathbb{E}[\mathbb{E}[R_{T'} \mid \mathcal{F}_n] \mid \mathcal{F}_{n-1}] \\ (\text{use the induction assumption } \mathbb{E}[R_{T'} \mid \mathcal{F}_n] \leq Y_n) & \\ &\leq \int_{A \cap \{T=n-1\}} \underbrace{R_{n-1}}_{\leq Y_{n-1}} + \int_{A \cap \{T \geq n\}} \underbrace{\mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]}_{\leq Y_{n-1}} \leq \int_A Y_{n-1}. \end{aligned}$$

This proves the inequality (1.4.18a).

To prove the equality (1.4.18b), we run the above argument with $T = T_{n-1}$. Observe that in this case

$$\mathcal{U}_n := \{T = n-1\} = \{R_{n-1} \geq \mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]\} = \{Y_{n-1} = R_{n-1}\}, \quad (1.4.20a)$$

$$\begin{aligned} \mathcal{V}_n &:= \{T_{n-1} > n-1\} = \{R_{n-1} < \mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]\} \\ &= \{Y_{n-1} = \mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]\}. \end{aligned} \quad (1.4.20b)$$

We have $T_{n-1} = n-1$ on $\mathcal{U}_n$ and $T_{n-1} = T_n$ on $\mathcal{V}_n$ so that

$$\begin{aligned} \int_A R_{T_{n-1}} &= \int_{A \cap \mathcal{U}_n} R_{n-1} + \int_{A \cap \mathcal{V}_n} R_{T_n} \\ (\mathcal{V}_n \in \mathcal{F}_{n-1}) & \\ &= \int_{A \cap \mathcal{U}_n} R_{n-1} + \int_{A \cap \mathcal{V}_n} \mathbb{E}[\mathbb{E}[R_{T_n} \mid \mathcal{F}_n] \mid \mathcal{F}_{n-1}] \\ (Y_n = \mathbb{E}[R_{T_n} \mid \mathcal{F}_n] \text{ by induction}) & \end{aligned}$$

$$\begin{aligned} &\int_{A \cap \mathcal{U}_n} R_{n-1} + \int_{A \cap \mathcal{V}_n} \mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]. \\ (\text{use (1.4.20a) and (1.4.20b)}) & \\ &= \int_A \max\{R_{n-1}, \mathbb{E}[Y_n \mid \mathcal{F}_{n-1}]\} = \int_A Y_{n-1}. \end{aligned}$$

□

Remark 1.4.15. The procedure for determining the optimal time T_1 outlined in the above example is a bit counterintuitive. The maximal expected reward is $\mathbb{E}[Y_1]$. By construction, the random variable Y_1 is $\mathcal{F}_1$ -measurable, by construction, and thus has the form $f(X_1)$ for some Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$. Thus we can determine Y_1 knowing only the initial input X_1 . On the other hand the definition of Y_1 by descending induction used the knowledge of the entire stream $X_1, \dots, X_N$, not just the initial input X_1 .

What it is true is that we can compute the maximal expected reward without running the stream. On the other hand, the moment we stop, and the actual reward when we stop are *random quantities*. It is conceivable that if we do not stop when T_1 tells us to stop we could get a higher reward later on. However, on average, we cannot beat the stopping strategy T_1 .

We will illustrate this process on the classical *secretary problem*. □

Example 1.4.16 (The secretary problem). Suppose we have a box with N prizes with values $v_1 < \dots < v_N$. Bob would like to pick the most valuable item but he does not know the actual values v_n . He is allowed to sample them successively without replacement. At the j -th draw he is told the value V_j of the j -th prize. He can either accept the j -th prize or he can decline it and ask to sample another one. A prize once declined cannot be accepted later on. We are interested in a strategy that maximizes the probability that Bob picks the most valuable prize.¹¹ My presentation is inspired from [36, Sec.3.3].

Consider the relative rankings

$$X_n := \#\{j \leq n; V_j \leq V_n\}. \quad (1.4.21)$$

Thus, X_n counts how many gifts unveiled up to the moment n are at most as valuable as the n -gift revealed. In particular, if $X_n = n$, then V_n is the largest of the observed values $V_1, \dots, V_n$.

We might be tempted to set the reward $R_n = \mathbf{I}_{\{V_n = v_N\}}$, but this is not $\mathcal{F}_n$ -measurable. We can fix this issue by setting

$$R_n := \mathbb{E}[\mathbf{I}_{\{V_n = v_N\}} \mid X_n].$$

Observe that for any stopping time T we have

$$\begin{aligned} \mathbb{E}[R_T] &= \sum_{n=1}^N \int_{T=n} R_n = \sum_{n=1}^N \int_{T=n} \mathbf{I}_{\{V_n = v_N\}} \\ &= \sum_{n=1}^N \mathbb{P}[V_n = v_N, T = n] = \mathbb{P}[V_T = v_N]. \end{aligned}$$

We want to find a stopping time T that maximizes $\mathbb{E}[R_T]$, i.e., the probability that Bob pick the biggest prize. Let us make a few remarks.

1. Observe that rankings $(X_n)_{n \in \mathbb{N}}$ defined in (1.4.21) are *independent* and

$$\mathbb{P}[X_n = j] = \frac{1}{n}, \quad \forall 1 \leq j \leq n \leq N. \quad (1.4.22)$$

Indeed, the random vector $(V_1, \dots, V_N)$ can be identified with a random permutation $\varphi \in \mathfrak{S}_N$ of $\mathbb{I}_N$

$$(V_1, \dots, V_N) = (v_{\varphi(1)}, \dots, v_{\varphi(N)}).$$

¹¹Think of N secretaries interviewing for a single job and the values $v_1, \dots, v_N$ rank their job suitability, the higher the value the more suitable. The interviewer learns the value v_k only at the time of the interview.

The rank X_n is then a function of φ

$$X_n(\varphi) := \#\{j \leq n; \varphi(j) \leq \varphi(n)\}.$$

To reach the desired conclusion observe that the map

$$\vec{X} : \mathfrak{S}_N \rightarrow \mathbb{I}_1 \times \mathbb{I}_2 \times \cdots \times \mathbb{I}_N, \quad \varphi \mapsto (X_1(\varphi), \dots, X_N(\varphi)) \quad (1.4.23)$$

is a bijection. Let us justify this claim.

Consider the following apparently more general situation. For any finite set of real numbers A and any $a \in A$ we define its rank

$$r(a) = r_A(a) := \#\{a' \in A; a' \leq a\}.$$

Given a set of real numbers A_N of cardinality N , a bijection $\varphi : A_N \rightarrow \mathbb{I}_N$, and $a \in A_N$ we define

$$X_a(\varphi) := \#\{a'; a' \leq a, \varphi(a') \leq \varphi(a)\} \in \mathbb{I}_{r(a)}.$$

Denote by $\text{Bij}(A, B)$ the set of bijections between two sets.

For any $\varphi_0 \in \text{Bij}(A_N, \mathbb{I}_N)$ we have

$$\text{argmax } \varphi_0 = \varphi_0^{-1}(N) = \max \{a : X_a(\varphi_0) = r_{A_N}(a)\}.$$

In other words, $\varphi_0^{-1}(N)$ is uniquely determined by its rank $(X_a(\varphi_0))_{a \in A_N}$.

Define

$$A_{N-1} := A_N \setminus \{\varphi_0^{-1}(N)\}.$$

The set A_{N-1} is uniquely determined by its ranks $(X_a(\varphi_0))_{a \in A_N}$. We obtain a new bijection $\varphi_1 : A_{N-1} \rightarrow \mathbb{I}_{N-1}$, $\varphi_1 = \varphi_0|_{A_{N-1}}$ such that

$$X_a(\varphi_1) = X_a(\varphi_0), \quad a \in A_{N-1}.$$

Then,

$$\text{argmax } \varphi_1 = \varphi_0^{-1}(N-1) = \max \{a \in A_{N-1} : X_a(\varphi_1) = r_{A_{N-1}}(a)\}.$$

Hence $\varphi_0^{-1}(N-1)$ is also uniquely determined by its ranks $(X_a(\varphi_0))_{a \in A_N}$.

We deduce iteratively that a bijection $\varphi_0 \in \text{Bij}(A_N, \mathbb{I}_N)$ is uniquely determined by its ranks $X_a(\varphi_0)$, $a \in A_N$. In other words the map

$$\vec{X} : \mathfrak{S}_N \rightarrow \mathbb{I}_1 \times \mathbb{I}_2 \times \cdots \times \mathbb{I}_N$$

defined in (1.4.23) is injective. It is also surjective since $\mathfrak{S}_N$ and $\prod_{n=1}^N \mathbb{I}_n$ have the same cardinality. This proves the claim (1.4.23).

2. We have

$$R_n = \frac{n}{N} \mathbf{I}_{\{X_n=n\}}. \quad (1.4.24)$$

Indeed, the conditional expectation $R_n = \mathbb{E}[\mathbf{I}_{\{V_n=v_N\}} | X_n]$ is a function of $x_n \in \mathbb{I}_n$ and we have

$$R_n(x_n) = \mathbb{E}[\mathbf{I}_{\{V_n=v_N\}} | X_n = x_n] = \mathbb{P}[V_n = v_N | X_n = x_n].$$

This probability is zero if $x_n < n$. Now observe that

$$\begin{aligned} \mathbb{P}[V_n = v_N | X_n = n] &= \frac{\mathbb{P}[V_n = v_N, X_n = n]}{\mathbb{P}[X_n = n]} \\ &= \frac{\mathbb{P}[V_n = v_N]}{\mathbb{P}[X_n = n]} = \frac{\frac{1}{N}}{\frac{1}{n}} = \frac{n}{N}. \end{aligned} \quad (1.4.25)$$

This shows that the reward R_n depends only on the random variable X_n . Denote by $\mathcal{F}_n$ the sigma-algebra generated by the independent random variables $X_1, \dots, X_n$.

Following (1.4.17a) and (1.4.18a) we set $y_n = \mathbb{E}[Y_n]$. The quantity y_n is the probability of Bob obtaining the largest prize among the strategies that reject the first $(n-1)$ shown prizes. We have

$$Y_N = R_N = \mathbf{I}_{\{V_N=v_N\}}, \quad y_N = \frac{1}{N}.$$

Since $\{X_N = N\}$ is independent of $\mathcal{F}_{N-1}$ deduce

$$\begin{aligned} Y_N &= \mathbb{E}[\mathbf{I}_{\{X_N=N\}} \mid \mathcal{F}_{N-1}] = \mathbb{E}[\mathbf{I}_{\{X_N=N\}}] = \frac{1}{N} = y_N, \\ Y_{N-1} &= \max \left\{ R_{N-1}, y_N \right\} = \max \left\{ \frac{N-1}{N} \mathbf{I}_{\{X_{N-1}=N-1\}}, y_N \right\} \\ &= \frac{N-1}{N} \mathbf{I}_{\{X_{N-1}=N-1\}} + \mathbf{I}_{\{X_{N-1}<N-1\}} y_N, \\ y_{N-1} &= \frac{1}{N} + \frac{N-2}{N-1} y_N. \end{aligned}$$

Similarly

$$\mathbb{E}[Y_{N-1} \mid \mathcal{F}_{N-2}] = y_{N-1}$$

and

$$\begin{aligned} Y_{N-2} &= \max \{ R_{N-2}, Y_{N-1} \} = \frac{N-2}{N-1} \\ &= \max \{ (N-2)/N, y_{N-1} \} \mathbf{I}_{\{X_{N-2}=N-2\}} + y_{N-1} \mathbf{I}_{\{X_{N-2}<N-2\}}, \\ y_{N-2} &\stackrel{(1.4.22)}{=} \max \{ (N-2)/N, y_{N-1} \} \frac{1}{N-2} + \frac{N-3}{N-2} y_{N-1} \end{aligned}$$

Iterating we deduce

$$\begin{aligned} Y_n &= \max \{ R_n, y_{n+1} \} = \max \{ n/N, y_{n+1} \} \mathbf{I}_{\{X_n=1\}} + y_{n+1} \mathbf{I}_{\{X_n>1\}}, \\ y_n &= \max \{ n/N, y_{n+1} \} \frac{1}{n} + \frac{n-1}{n} y_{n+1}. \end{aligned}$$

While it is difficult to find an explicit formula for y_n , the above equalities can be easily implemented on a computer. The optimal probability is $p_N = y_1$.

As long as $y_{n+1} \leq \frac{n}{N}$ we have

$$y_n = \frac{1}{N} + \frac{n-1}{n} y_{n+1} > y_n.$$

If $y_{n+1} > \frac{n}{N}$, then $y_n = y_{n+1}$. Set

$$N_* := \min \left\{ n; \quad y_{n+1} \leq \frac{n}{N} \right\}.$$

The above discussion shows that

$$y_N < y_{N-1} < \dots < y_{N_*} = y_{N_*-1} = \dots = y_1.$$

Let us determine N_* . We set

$$q_n = \frac{y_n}{n-1}.$$

We deduce

$$q_n - q_{n+1} = \frac{1}{N(n-1)}.$$

Observing that $q_N = \frac{1}{N(N-1)}$ we deduce that

$$q_{N-j} = \frac{1}{N} \left(\frac{1}{N-1-j} + \cdots + \frac{1}{N-1} \right),$$

This is valid as long as $q_{N-j+1} \leq \frac{1}{N}$, that is

$$\frac{1}{N-j} + \cdots + \frac{1}{N-1} \leq 1.$$

We set $n = N - j$. Then

$$\frac{1}{N-j} + \cdots + \frac{1}{N-1} = \frac{1}{n} + \cdots + \frac{1}{N-1} =: g_N(n).$$

As long as $g_N(n) \leq 1$ we have

$$y_n = \frac{n-1}{N} \left(\frac{1}{n-1} + \cdots + \frac{1}{N-1} \right) = \frac{n-1}{N} g_N(n-1).$$

Thus

$$N^* = \min \{ n; g_N(n) \leq 1 \}$$

and

$$y_1 = \cdots = y_{N^*} = \frac{N^* - 1}{N} g_N(N^* - 1).$$

Thus, the maximal probability is

$$p_N = \frac{N^* - 1}{N} g_N(N^* - 1) \tag{1.4.26}$$

Note that

$$g_N(N^* - 1) = \frac{1}{N^* - 1} + g_N(N^*)$$

so

$$p_N = \frac{1}{N} + \frac{N^* - 1}{N} g_N(N^*).$$

Let us analyze the behavior of p_N as $N \rightarrow \infty$.

Using Riemann sums we deduce that for $j > 1$ we have

$$\log \frac{N}{n} = \int_j^N \frac{1}{x} dx < g_N(n) < \int_{n-1}^N \frac{1}{x} dx = \log \frac{N}{j-1}.$$

Note that

$$\log \frac{N}{n-1} - \log \frac{N}{n} = \log \frac{n}{n-1} = \log \left(1 + \frac{1}{n-1} \right).$$

Note that

$$g_N(N^*) \leq 1, \quad g_N(N^* - 1) = g_N(N^*) + \frac{1}{N^* - 1} > 1,$$

that is

$$1 - \frac{1}{N^* - 1} \leq g_N(N^*) \leq 1.$$

Hence

$$\log \frac{N}{N^*} \leq g_N(N^*) \leq 1, \quad 1 - \frac{1}{N^* - 1} \leq g_N(N^*) \leq \log \frac{N}{N^* - 1},$$

so that

$$\log \frac{N}{N^*} \leq 1 \leq \log \frac{N}{N^* - 1} + \frac{1}{N^* - 1} = \log \frac{N}{N^*} + \log \frac{N^*}{N^* - 1} + \frac{1}{N^* - 1}.$$

In other words

$$1 - \frac{2}{N^* - 1} \leq 1 - \log \frac{N^*}{N^* - 1} - \frac{1}{N^* - 1} < \log \frac{N}{N^*} < 1.$$

We deduce that

$$\frac{N}{e} < N^* < \frac{N}{e} \cdot e^{\frac{2}{N^* - 1}},$$

and so $N^* \rightarrow \infty$ as $N \rightarrow \infty$ and,

$$N^* \sim \frac{N}{e}, \text{ as } N \rightarrow \infty.$$

Hence $p_N \sim \frac{1}{e}$ as $N \rightarrow \infty$.

Here is a less than optimal¹² but simple R code that computes the vector $(y_1, \dots, y_N)$, given N .

```
optim<-function(N) {
  p<-1/N
  m<-N-1
  y<-c(p)
  for (i in 1:m) {
    p<-max((N-i), N^*p) / (N*(N-i)) + ((N-i-1) / (N-i)) * p
    y<-c(y, p)
  }
  y=y[N:1]
  y
}
```

The optimal probability p_N of choosing the largest among N prizes is obtained using the command

```
optim(N)[1]
```

For example, for $N = 10$ we have

n	1	2	3	4	5	6	7	8	9	10
y_n	0.3986	0.3986	0.3986	0.3986	0.3982	0.372	0.32	0.26	0.18	0.1

(1.4.27)

In this case $N_* = 4$ and the optimal strategy corresponds to the stopping time T_5 : reject the first 3 gifts and then accept the first gift more valuable than any of the previously chosen. In this case the probability of choosing the most valuable gifts is $p_{10} \approx 0.3986$.

Here are some results. Below, p_N denotes the

N	3	4	5	6	8	100	200
p_N	0.5	0.458	0.433	0.4277	0.4098	0.3710	0.3694

The optimal strategy is given by the stopping time T_{N_*} : reject the first $N_* - 1$ selected gifts and then pick the first gift that is more valuable than any of the preceding ones. One can compute N_* as $\text{stops}(N)$, where

¹²A word of warning! This script is numerically unstable for large N due to the errors introduced by the conversion of fractions into decimals. It is exact up to 4 decimals at least.

```

stops=function(N) {
  M=N-1
  x=c(M:1)
  x= 1/x
  x= cumsum(x)
  j=1
  while (x[j]<1) {
    j=j+1
  }
  N-j+1
}

```

Here are some results.

N	3	4	8	10	50	100	1000
N_*	2	2	4	4	19	38	369

This table is in perfect agreement with the earlier computations in [88, Table 2, p.40].

As explained in [88], a (nearly) optimal strategy is as follows. Denote by m the largest integer satisfying

$$\frac{N-1/2}{e} + \frac{1}{2} \leq m \leq \frac{N-1/2}{e} + \frac{3}{2}.$$

Reject the first m prizes and accept the next prize more valuable than any of the preceding ones. $\square$

1.4.3. Conditional independence. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space.

Definition 1.4.17. Fix a sigma-subalgebra $\mathcal{G}$ of $\mathcal{S}$. The family $(\mathcal{F}_i)_{i \in I}$ of sigma-subalgebras of $\mathcal{S}$ is said to be *conditionally independent given $\mathcal{G}$* if, for any finite subset $J \subset I$ and any events $F_j \in \mathcal{F}_j$, $j \in J$, we have

$$\mathbb{E} \left[\prod_{j \in J} \mathbf{1}_{F_j} \mid \mathcal{G} \right] = \prod_{j \in J} \mathbb{E} \left[\mathbf{1}_{F_j} \mid \mathcal{G} \right] \quad \text{a.s..}$$

Given sigma algebras $\mathcal{F}, \mathcal{G}, \mathcal{H} \subset \mathcal{S}$ we use the notation $\mathcal{F} \perp_{\mathcal{G}} \mathcal{H}$ to indicate that $\mathcal{F}$ is independent of $\mathcal{H}$ given $\mathcal{G}$. $\square$

The next proposition generalizes the result in Exercise 1.10.

Proposition 1.4.18 (Doob-Markov). *Given sigma algebras $\mathcal{F}_{\pm}, \mathcal{F}_0 \subset \mathcal{S}$ the following are equivalent.*

- (i) $\mathbb{E}[X_+ \mid \mathcal{F}_- \vee \mathcal{F}_0] = \mathbb{E}[X_+ \mid \mathcal{F}_0] \quad \text{a.s., } \forall X_+ \in L^1(\Omega, \mathcal{F}_+, \mathbb{P}).$
- (ii) $\mathcal{F}_+ \perp_{\mathcal{F}_0} \mathcal{F}_-.$

Proof. The condition (i) is equivalent to

$$\mathbb{E}[X X_+] = \mathbb{E} \left[X \mathbb{E}[X_+ \mid \mathcal{F}_0] \right], \quad \forall X \in L^\infty(\Omega, \mathcal{F}_0 \vee \mathcal{F}_-, \mathbb{P}). \quad (1.4.28)$$

The condition (ii) equivalent to

$$\mathbb{E}[X_+ X_- \mid \mathcal{F}_0] = \mathbb{E}[X_+ \mid \mathcal{F}_0] \mathbb{E}[X_- \mid \mathcal{F}_0], \quad \forall X_{\pm} \in L^\infty(\Omega, \mathcal{F}_{\pm}, \mathbb{P}).$$

Note that since $\mathbb{E}[X_+ \mid \mathcal{F}_0]$ is an $\mathcal{F}_0$ -measurable random variable we have

$$\mathbb{E}[X_+ \mid \mathcal{F}_0] \mathbb{E}[X_- \mid \mathcal{F}_0] = \mathbb{E} \left[X_- \left(\mathbb{E}[X_+ \mid \mathcal{F}_0] \right) \mid \mathcal{F}_0 \right].$$

Thus, (ii) is equivalent to

$$\mathbb{E}[X_+X_- \mid \mathcal{F}_0] = \mathbb{E}\left[X_- \left(\mathbb{E}[X_+ \mid \mathcal{F}_0] \right) \mid \mathcal{F}_0\right],$$

i.e., for any nonnegative, bounded, $\mathcal{F}_0$ -measurable random variable X_0 we have

$$\mathbb{E}[X_0X_-X_+] = \mathbb{E}\left[X_0X_- \mathbb{E}[X_+ \mid \mathcal{F}_0]\right].$$

Since $\mathcal{F}_0 \vee \mathcal{F}_-$ coincides with the sigma-algebra generated collection of random variables X_0X_- , $X_0 \in L^\infty(\Omega, \mathcal{F}_0, \mathbb{P})$, $X_- \in L^\infty(\Omega, \mathcal{F}_-, \mathbb{P})$ we deduce that the last equality is equivalent to (1.4.28), i.e., (i) is equivalent to (ii). $\square$

Remark 1.4.19. You should think of a system evolving in time. Then $\mathcal{F}_0$ collects the present information about the system, $\mathcal{F}_-$ collects the past information and $\mathcal{F}_+$ collects the future information. Roughly speaking, the above proposition shows that the information about a future event given the present and the past coincides with the information given the present if and only if the future is independent of the past given the present. $\square$

1.4.4. Kernels and regular conditional distributions. Suppose that $(\Omega_0, \mathcal{F}_0)$ and $(\Omega_1, \mathcal{S}_1)$ are two measurable spaces.¹³ A *kernel* from $(\Omega_0, \mathcal{F}_0)$ to $(\Omega_1, \mathcal{S}_1)$ is a function

$$K : \Omega_0 \times \mathcal{S}_1 \rightarrow [0, \infty], \quad (\omega_0, S_1) \mapsto K_{\omega_0}[S_1]$$

with the following properties.

(K₁) For each $\omega_0 \in \Omega_0$, the map

$$S_1 \ni S_1 \mapsto K_{\omega_0}[S_1] \in [0, \infty]$$

is a measure. We will denote this measure by $K_{\omega_0}[d\omega_1]$.

(K₂) For each $S_1 \in \mathcal{S}_1$ the function

$$\Omega_0 \ni \omega_0 \mapsto K_{\omega_0}[S_1] \in [0, \infty]$$

is $\mathcal{F}_0$ -measurable. We will denote this random variable by $K_{\square}[S_1]$

The kernel K is called a *probability kernel* or a *Markovian kernel* if $K_{\omega_0}[-]$ is a probability measure on $(\Omega_1, \mathcal{S}_1)$, for any $\omega_0 \in \Omega_0$.

We will use the notation $K : (\Omega_0, \mathcal{F}_0) \rightsquigarrow (\Omega_1, \mathcal{S}_1)$ to indicate that K is a kernel from $(\Omega_0, \mathcal{F}_0)$ to $(\Omega_1, \mathcal{S}_1)$

The condition (K₁) above shows that a kernel is a family $(K_{\omega_0}[-])_{\omega_0 \in \Omega_0}$ of measures on $(\Omega_1, \mathcal{S}_1)$ parametrized by Ω_0 . Condition (K₂) is a measurability condition on this family. For this reason kernels are also known as *random measures*.

Example 1.4.20. Consider the Bernoulli measure

$$\beta_p := q\delta_0 + p\delta_1 \in \text{Prob}(\mathbb{R}), \quad p \in [0, 1], \quad q = 1 - p.$$

To obtain a random measure we let p be a random quantity. More precisely, if $f : (\Omega, \mathcal{S}) \rightarrow [0, 1]$ is a measurable function, then

$$\beta_{f(\omega)} = (1 - f(\omega))\delta_0 + f(\omega)\delta_1$$

¹³In the story of kernels, the sigma-algebras $\mathcal{F}_0, \mathcal{S}_1$ play rather different roles and, for this reason, we chose to indicate them using visually distinctive notation.

defines a Markov kernel $K : (\Omega, \mathcal{S}) \rightsquigarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$,

$$K_{\omega}[B] = (1 - f(\omega))\delta_0[B] + f(\omega)\delta_1[B]. \quad \square$$

Given a measure μ on the measurable space $(\Omega, \mathcal{F})$ and a nonnegative measurable function $f \in \mathcal{L}_+^0(\Omega, \mathcal{F})$ we set

$$\langle \mu, f \rangle := \mu[f] = \int_{\Omega} f(\omega) \mu[d\omega] \in [0, \infty].$$

Theorem 1.4.21. *Suppose that $K : (\Omega_0, \mathcal{F}_0) \rightsquigarrow (\Omega_1, \mathcal{S}_1)$ is a kernel.*

(i) *For any $f \in \mathcal{L}_+^0(\Omega_1, \mathcal{S}_1)$ we define its pullback by K to be the function*

$$K^*f : \Omega_0 \rightarrow [0, \infty], \quad K^*f(\omega_0) = \int_{\Omega_1} f(\omega_1) K_{\omega_0}[d\omega_1]$$

Then $K^*f \in \mathcal{L}_+^0(\Omega_0, \mathcal{F}_0)$.

(ii) *For any measure $\mu : \mathcal{F}_0 \rightarrow [0, \infty]$ we define its push-forward by K to be the function $K_*\mu : \mathcal{S}_1 \rightarrow [0, \infty]$ defined by*

$$K_*\mu[F_1] := \int_{\Omega_0} K_{\omega_0}[S_1] \mu[d\omega_0] \in [0, \infty], \quad S_1 \in \mathcal{S}_1. \quad (1.4.29)$$

Then $K_*\mu$ is a measure on $(\Omega_1, \mathcal{S}_1)$.

(iii) *The pullback and push-forward by K are adjoints of each other. More precisely, for any measure μ on $(\Omega_0, \mathcal{F}_0)$ and any measurable function $f \in \mathcal{L}_+^0(\Omega_1, \mathcal{S}_1)$ we have*

$$\langle \mu, K^*f \rangle = \langle K_*\mu, f \rangle. \quad (1.4.30)$$

Proof. (i) For any $S \in \mathcal{S}_1$ we have $K^*\mathbf{I}_S(\omega_0) = K_{\omega_0}[S]$ so $K^*\mathbf{I}_S \in \mathcal{L}^0(\Omega_0, \mathcal{F}_0)$. Clearly the correspondence $f \mapsto K^*f$ is monotone and the conclusion follows from the fact that a nonnegative function is measurable iff it is the limit of an increasing sequence of simple functions.

The statement (ii) follows from the Monotone Convergence theorem and $(\mathbf{K}_1)$. For part (iii), fix the measure μ . Observe that for $S \in \mathcal{S}_1$ we have

$$\begin{aligned} \langle \mu, K^*\mathbf{I}_S \rangle &= \int_{\Omega_0} K^*\mathbf{I}_S(\omega_0) \mu[d\omega_0] \\ &= \int_{\Omega_0} \left(\int_{\Omega_1} \mathbf{I}_S(\omega_1) K_{\omega_0}[d\omega_1] \right) \mu[d\omega_0] \\ &= \int_{\Omega_0} K_{\omega_0}[S] \mu[d\omega_0] = K_*\mu[S] = \langle K_*\mu, \mathbf{I}_S \rangle. \end{aligned}$$

Thus (1.4.30) holds for $f = \mathbf{I}_S$, $S \in \mathcal{S}_1$. The general case follows by invoking the Monotone Class Theorem. $\square$

When K is a Markovian kernel and μ is a probability measure, then the pushforward $K_*\mu$ is also a probability measure. For any $S_1 \in \mathcal{S}_1$ the scalar $K_*\mu[S_1]$ is the expectation of the nonnegative random variable

$$\Omega_0 \ni \omega_0 \mapsto K_{\omega_0}[S_1] \in [0, \infty]$$

with respect to μ . The measure $K_*\mu$ is said to be a *mixture* of the random measure $\omega_0 \mapsto K_{\omega_0}[-]$ driven by μ .

Example 1.4.22. (a) Suppose that $(\Omega_0, \mathcal{F}_0), (\Omega_1, \mathcal{F}_1)$ are two measurable spaces and

$$T : (\Omega_0, \mathcal{F}_0) \rightarrow (\Omega_1, \mathcal{F}_1)$$

is a measurable map. Then T defines a kernel $K^T : (\Omega_0, \mathcal{F}_0) \rightsquigarrow (\Omega_1, \mathcal{F}_1)$

$$K_{\omega_0}^T [F_1] = \delta_{T(\omega_0)} [F_1],$$

where δ_{ω_1} denotes the Dirac measure on $(\Omega_1, \mathcal{F}_1)$ concentrated at ω_1 ; see Example 1.2.6(a).

Observe that for any measure μ on $\mathcal{F}_0$ and any $f \in \mathcal{L}_+^0(\Omega_1, \mathcal{F}_1)$ we have

$$K_*^T \mu = T_{\#} \mu, \quad (K^T)^* f = T^* f := f \circ T.$$

Thus, (1.4.30) contains as a special case the change in variables formula (1.2.25).

(b) Any measurable function $f : (\Omega, \mathcal{S}) \rightarrow [0, 1]$ defines as in Example 1.4.20 the random Bernoulli measure

$$K_{\omega} [-] = (1 - f(\omega)) \delta_0 + f(\omega) \delta_1.$$

Given a probability measure μ on $(\Omega, \mathcal{S})$ we have

$$K_* \mu = \text{Ber}(\bar{f}) = (1 - \bar{f}) \delta_0 + \bar{f} \delta_1, \quad \bar{f} := \mathbb{E}_{\mu} [f].$$

(c) Suppose that $\mathfrak{X}$ is a finite or countable set. A kernel $(\mathfrak{X}, \mathfrak{P}) \rightsquigarrow (\mathfrak{X}, \mathfrak{P})$ is defined by a function (matrix) $K : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, \infty]$, via the equality

$$K_x [S] = \sum_{s \in S} K(x, s), \quad \forall x \in \mathfrak{X}, \quad S \subset \mathfrak{X}.$$

The kernel is Markovian if

$$\sum_{x' \in \mathfrak{X}} K(x, x') = 1, \quad \forall x \in \mathfrak{X}.$$

(d) Suppose that $f : \mathbb{R}^2 \rightarrow [0, \infty)$ is an integrable function such that

$$\int_{\mathbb{R}} f(x, y) dy = 1, \quad \forall x \in \mathbb{R}.$$

It defines a Markovian kernel $K : (\mathbb{R}, \mathcal{B}_{\mathbb{R}}) \rightsquigarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$

$$K_x [B] = \int_B f(x, y) dy,$$

$\forall x \in \mathbb{R}$ and any Borel subset $B \subset \mathbb{R}$. The measurability of the map $x \mapsto K_x [B]$ follows from Fubini's theorem. We can rewrite this as $K_x [dy] = f(x, y) dy$.

(e) Suppose that ν is a finite Borel measure on $\mathbb{R}$. It defines a kernel

$$K_{\nu} : (\mathbb{R}, \mathcal{B}_{\mathbb{R}}) \rightsquigarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}}), \quad K_{\nu, y} [B] = \nu [B - y].$$

In Exercise 1.63 we ask the reader to prove that the map $y \mapsto K_{\nu, y} [B]$ is measurable for any Borel set $B \subset \mathbb{R}$. Then, for any finite Borel measure μ on $\mathbb{R}$ we have $(K_{\nu})_* \mu = \mu * \nu$. $\square$

Example 1.4.23 (Skhorokhod). Given real numbers $u \leq 0 < v$ we define the *dipole concentrated at u, v* to be the probability measure

$$D_{u,v} = \frac{v}{v-u} \delta_u - \frac{u}{v-u} \delta_v = \frac{v}{v+|u|} \delta_{-|u|} + \frac{|u|}{v+|u|} \delta_v \in \text{Prob}(\mathbb{R}).$$

Note that the center of mass of a dipole is the origin, i.e.,

$$\int_{\mathbb{R}} x \mathbf{D}_{u,v}[dx] = 0.$$

If we denote by Q the quadrant $(-\infty, 0] \times (0, \infty) \subset \mathbb{R}^2$, then the correspondence

$$Q \ni (u, v) \mapsto \mathbf{D}_{u,v} \in \text{Prob}(\mathbb{R})$$

is a Markov kernel. For any $\nu \in \text{Prob}(Q)$ we get a mixture of dipoles

$$\mathbf{D}_* \nu \in \text{Prob}(\mathbb{R}), \quad \mathbf{D}_* \nu[B] = \int_Q (\mathbf{D}_{u,v}[B]) \nu[du dv], \quad \forall B \in \mathcal{B}_{\mathbb{R}}.$$

Suppose that X is a random variable such that

$$\mathbb{E}[X] = 0, \quad \text{Var}[X] = \mathbb{E}[X^2] = 1.$$

Denote by μ the distribution of X . We want to show that μ can be represented as a mixture of dipoles.

Set

$$K := \frac{1}{2} \mathbb{E}[|X|] = \int_{(0,\infty)} v \mu[dv] = - \int_{(-\infty,0]} u \mu[du].$$

Set $\mu_+ := \mathbf{I}_{(0,\infty)} \cdot \mu$, $\mu_- = \mathbf{I}_{(-\infty,0]} \cdot \mu$. Consider the Borel measure on $\mathbb{R}^2$

$$\mathbf{m}_{\mu}[du dv] := \frac{v-u}{K} \mu_- \otimes \mu_+[du dv]. \quad (1.4.31)$$

A direct computation shows that $\mathbf{m}_{\mu}$ is a probability measure. Note that

$$\int_{\mathbb{R}^2} |uv| \mathbf{m}_{\mu}[du dv] = 1. \quad (1.4.32)$$

Indeed

$$\begin{aligned} \int_{\mathbb{R}^2} |uv| \mathbf{m}_{\mu}[du dv] &= \frac{1}{K} \int_{(-\infty,0] \times (0,\infty)} |uv|(u+|v|) \mu \otimes \mu[du dv] \\ &= \frac{1}{K} \int_{(-\infty,0] \times (0,\infty)} |u|v^2 \mu \otimes \mu[du dv] + \frac{1}{K} \int_{(-\infty,0] \times (0,\infty)} |u|^2 v \mu \otimes \mu[du dv] \\ &= \frac{1}{K} \left(\int_{(-\infty,0]} |u| \mu[du] \right) \left(\int_{(0,\infty)} v^2 \mu[dv] \right) \\ &\quad + \frac{1}{K} \left(\int_{(-\infty,0]} |u|^2 \mu[du] \right) \left(\int_{(0,\infty)} v \mu[dv] \right) < \infty \end{aligned}$$

(use the definition of K)

$$= \int_{(-\infty,0]} u^2 \mu[du] + \int_{(0,\infty)} v^2 \mu[dv] = \text{Var}[X] = 1.$$

We claim that

$$\mathbf{D}_* \mathbf{m}_{\mu} = \mu. \quad (1.4.33)$$

More explicitly, this means that μ can be expressed as a mixture of dipoles driven by $\mathbf{m}_{\mu}$

$$\mu = \int_{\mathbb{R}^2} \mathbf{D}_{u,v} \mathbf{m}_{\mu}[du dv]. \quad (1.4.34)$$

To verify (1.4.33) it suffices to check that for any $c \in \mathbb{R}$ we have

$$\mathbf{D}_* \mathbf{m}_{\mu}[(-\infty, c]] = \mu[(-\infty, c]].$$

We have

$$\begin{aligned}
D_* m_\mu [(-\infty, c)] &= \int_{(-\infty, 0] \times (0, \infty)} D_{u,v} [(-\infty, c)] \frac{v-u}{K} \mu^{\otimes 2} [dudv] \\
&= \frac{1}{K} \int_{(-\infty, 0] \times (0, \infty)} \left(v \delta_u [(-\infty, c)] - u \delta_v [(-\infty, c)] \right) \mu^{\otimes 2} [dudv] \\
&= \frac{1}{K} \int_{(-\infty, 0] \times (0, \infty)} \left(v \mathbf{I}_{(-\infty, c]}(u) - u \mathbf{I}_{(-\infty, c]}(v) \right) \mu^{\otimes 2} [dudv] \\
&= \frac{1}{K} \mu [(-\infty, c \wedge 0)] \underbrace{\int_{(0, \infty)} v \mu [dv]}_{=K} + \frac{1}{K} \mu [(0, c \vee 0)] \underbrace{\int_{(-\infty, 0]} |u| \mu [du]}_{=K} \\
&= \mu [(-\infty, c \wedge 0)] + \mu [(0, c \vee 0)] = \mu [(-\infty, c)].
\end{aligned}$$

□

Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $\mathcal{F} \subset \mathcal{S}$ is a sigma subalgebra. For every event $S \in \mathcal{S}$ the *random variable*

$$\mathbb{P}[S \parallel \mathcal{F}] := \mathbb{E}[\mathbf{I}_S \parallel \mathcal{F}]$$

is called the *conditional probability* of S given $\mathcal{F}$. The random variable $\mathbb{P}[S \parallel \mathcal{F}]$ is unique up to equality off a negligible set.

Note that for any increasing family $(S_n)_{n \geq 1} \subset \mathcal{S}$ there exists a negligible set $\mathcal{N} \subset \Omega$ such that

$$\lim_n \mathbb{P}[S_n \parallel \mathcal{F}](\omega) = \mathbb{P}[\lim_n S_n \parallel \mathcal{F}](\omega), \quad \forall \omega \in \Omega \setminus \mathcal{N}.$$

A priori, the negligible set $\mathcal{N}$ depends on the family $(S_n)_{n \geq 1}$, and there might not exist one negligible set that works for all such increasing families. When such a thing is possible we say that the conditional probability $\mathbb{P}[- \parallel \mathcal{F}]$ admits a *regular version*. Here is the precise definition.

Definition 1.4.24. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and $\mathcal{F} \subset \mathcal{S}$ a sigma-subalgebra. A *regular version* of $\mathbb{P}[- \parallel \mathcal{F}]$ is a kernel $Q : (\Omega, \mathcal{F}) \rightsquigarrow (\Omega, \mathcal{S})$ such that, for any $S \in \mathcal{S}$, the random variable $\Omega \ni \omega \mapsto Q_\omega[S]$ is a version of $\mathbb{P}[S \parallel \mathcal{F}]$. In other words,

- the map $\omega \mapsto Q_\omega[S]$ is $\mathcal{F}$ -measurable and
- for any $S \in \mathcal{S}$, $F \in \mathcal{F}$ we have

$$\mathbb{P}[S \cap F] = \int_F Q_\omega[S] \mathbb{P}[d\omega].$$

□

Proposition 1.4.25. If $Q : (\Omega, \mathcal{F}) \rightsquigarrow (\Omega, \mathcal{S})$ is a regular version of $\mathbb{P}[- \parallel \mathcal{F}]$, then $\forall X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$,

$$\mathbb{E}[X \parallel \mathcal{F}] = Q^* X,$$

i.e.,

$$\mathbb{E}[X \parallel \mathcal{F}]_\omega = \int_\Omega X(\eta) Q_\omega[d\eta] = Q^* X(\omega) \quad \text{a.s..} \quad (1.4.35)$$

Proof. Note that (1.4.35) holds in the special case $X = \mathbf{I}_S$ because

$$Q^* \mathbf{I}_S(\omega) = Q_\omega[S] = \mathbb{P}[S \parallel \mathcal{F}](\omega) = \mathbb{E}[\mathbf{I}_S \parallel \mathcal{F}](\omega).$$

The general case follows from the Monotone Class theorem. □

The equality (1.4.35) can be written in the less precise, but more intuitive way

$$\mathbb{E}[X \mid \mathcal{F}] = \int_{\Omega} X(\eta) \mathbb{P}[d\eta \mid \mathcal{F}]. \quad (1.4.36)$$

More generally, consider a measurable map $T : (\tilde{\Omega}, \tilde{\mathcal{S}}) \rightarrow (\Omega, \mathcal{S})$. Let $\tilde{\mathbb{P}}$ be a probability measure on $(\tilde{\Omega}, \tilde{\mathcal{S}})$ and suppose that $\tilde{\mathcal{F}} \subset \tilde{\mathcal{S}}$ is a sigma subalgebra. For every $S \in \mathcal{S}$ we set

$$\mathbb{P}_T[S \mid \tilde{\mathcal{F}}] := \tilde{\mathbb{P}}[T \in S \mid \tilde{\mathcal{F}}] = \mathbb{E}_{\tilde{\mathbb{P}}}[T^* \mathbf{I}_S \mid \tilde{\mathcal{F}}] = \mathbb{E}_{\tilde{\mathbb{P}}}[\mathbf{I}_{T^{-1}(S)} \mid \tilde{\mathcal{F}}]. \quad (1.4.37)$$

We will refer to $\mathbb{P}_T[- \mid \tilde{\mathcal{F}}]$ as the *conditional distribution* of T given $\tilde{\mathcal{F}}$. Observe that when

$$(\tilde{\Omega}, \tilde{\mathcal{S}}) = (\Omega, \mathcal{S}), \quad \tilde{\mathbb{P}} = \mathbb{P} \quad \text{and} \quad T = \mathbb{1}_{\Omega},$$

then

$$\mathbb{P}_{\mathbb{1}_{\Omega}}[- \mid \tilde{\mathcal{F}}] = \mathbb{P}[- \mid \tilde{\mathcal{F}}].$$

Note that for any increasing family $(S_n)_{n \geq 1} \subset \mathcal{S}$ we have

$$\lim_{n \rightarrow \infty} \mathbb{P}_T[S_n \mid \tilde{\mathcal{F}}] = \mathbb{P}_T[\lim_n S_n \mid \tilde{\mathcal{F}}] \text{ a.s..}$$

We say that $\mathbb{P}_T[- \mid \tilde{\mathcal{F}}]$ admits a regular version if we can choose representatives for each $\mathbb{P}_T[F \mid \tilde{\mathcal{F}}]$, $F \in \mathcal{F}$ so that the above equality holds for any increasing sequence (S_n) . Here is a more precise definition.

Definition 1.4.26. Let $(\tilde{\Omega}, \tilde{\mathcal{S}}, \tilde{\mathbb{P}})$ be a probability space and $T : (\tilde{\Omega}, \tilde{\mathcal{S}}) \rightarrow (\Omega, \mathcal{S})$ be a measurable map. Fix a sigma-subalgebra $\tilde{\mathcal{F}} \subset \tilde{\mathcal{S}}$. A *regular version* of the conditional probability distribution $\mathbb{P}_T[- \mid \tilde{\mathcal{F}}]$ of the map T conditioned on $\tilde{\mathcal{F}}$ is a kernel $Q : (\tilde{\Omega}, \tilde{\mathcal{F}}) \rightsquigarrow (\Omega, \mathcal{S})$ such that, for any $S \in \mathcal{S}$, the random variable $Q_{\square}[S]$ is a version of $\mathbb{P}_T[S \mid \tilde{\mathcal{F}}]$. In other words,

- the random variable $Q_{\square}[S]$ (on $\tilde{\Omega}$) is $\tilde{\mathcal{F}}$ -measurable and
- for any $\tilde{F} \in \tilde{\mathcal{F}}$, $S \in \mathcal{S}$ we have

$$\tilde{\mathbb{P}}[\tilde{F} \cap T^{-1}(S)] = \int_{\tilde{F}} Q_{\tilde{\omega}}[S] \tilde{\mathbb{P}}[d\tilde{\omega}]. \quad (1.4.38)$$

□

A conditional probability distribution need not admit a regular version. For that to happen we have to impose conditions on $\mathcal{S}$, the sigma-algebra in the target space. This requires a brief topological digression.

Definition 1.4.27. A *Lusin space* is a topological space homeomorphic to a Borel subset of a compact metric space. □

Remark 1.4.28. (a) The above is not the usual definition of a Lusin space but it has the advantage that emphasizes the compactness feature we need in the proof of Kolmogorov's existence theorem.

There are plenty of Lusin spaces. In fact, a topological space that is not Lusin is rather unusual. We refer to [21, 45, 50] for a more in depth presentation of these spaces and their applications in measure theory and probability. To give the reader a taste of the fauna of Lusin spaces we list a few examples.

- The Euclidean spaces $\mathbb{R}^n$ are Lusin spaces.

- A Borel subset of a Lusin space is also Lusin space.
- The Cartesian product of two Lusin spaces is a Lusin space.
- A less obvious example is that of *Polish spaces*, i.e., complete separable metric spaces. More precisely every Polish space is homeomorphic to a countable intersection of open subsets of $[0, 1]^{\mathbb{N}}$; see [25], Chap. IX, Sec.6.1, Corollary 1.
- A Hausdorff space is Lusin iff it is the image of a continuous bijection from a Polish space.
- A Hausdorff space is Lusin if and only if it is homeomorphic to a Borel subset of a Polish space.

(b) From a measure theoretic point of view the Lusin spaces are indistinguishable from the Polish spaces. More precisely, for any Lusin space X , there exists a Polish space Y and a Borel measurable bijection $\Phi : X \rightarrow Y$ such that the inverse is also Borel measurable; see [45, Prop. 8.6.13].

The Polish spaces have another important property. More precisely, a Polish space equipped with the σ -algebra of Borel subsets is *isomorphic as a measurable space* to a Borel subset E of $[0, 1]$ equipped with the σ -algebra of Borel subsets. For a proof we refer to [166, Sec.I.2]. Moreover, any two Borel subsets of $\mathbb{R}$ are measurably isomorphic if and only if they have the same cardinality, [166, Ch.I, Thm.2.12].

On the other hand, it is known that the continuum hypothesis holds for the Borel subsets of a Polish space; see [50, Appendix III.80] or [126, XII.6]. In particular, any Borel subset of $\mathbb{R}$ is either finite, countable or has the continuum cardinality. We deduce from this a theorem of Kuratowski that a Lusin space is isomorphic as a measurable space with either a finite set, $\mathbb{N}$, or $[0, 1]$ equipped with their natural Borel sigma-algebra. Hence *any Lusin space is Borel isomorphic to a compact metric space!* $\square$

We have the following general existence result.

Theorem 1.4.29 (Existence of regular conditional probabilities). *Suppose that*

- $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space,
- $\mathcal{Y}$ is a Lusin space and
- $\mathcal{B}_{\mathcal{Y}}$ is the sigma-algebra of Borel subsets of $\mathcal{Y}$.

Then, for every measurable map $Y : (\Omega, \mathcal{S}) \rightarrow (\mathcal{Y}, \mathcal{B}_{\mathcal{Y}})$, and every σ -subalgebra $\mathcal{F} \subset \mathcal{S}$ there exists a regular version $Q : (\Omega, \mathcal{F}) \rightsquigarrow (\mathcal{Y}, \mathcal{B}_{\mathcal{Y}})$, $(\omega, B) \mapsto Q_{\omega}[B]$, of the conditional distribution $\mathbb{P}_Y[-|\mathcal{F}]$. This means that

$$Q_{\omega}[B] = \mathbb{P}[Y \in B | \mathcal{F}] \text{ a.s., } \forall B \in \mathcal{B}_{\mathcal{Y}}.$$

Moreover, for any measurable function $f : (\mathcal{Y}, \mathcal{B}_{\mathcal{Y}}) \rightarrow \mathbb{R}$, we have

$$\mathbb{E}[f \circ Y | \mathcal{F}](\omega) = \int_{\mathcal{Y}} f(y) Q_{\omega}[dy], \quad \forall \omega \in \Omega. \quad (1.4.39)$$

Idea of proof. For a complete proof we refer to [43, Th. IV2.10], [50, III.71], [51, IX.11] or [180, II.89].

We can assume that $\mathcal{Y}$ is a compact metric space. Fix a dense countable subset $\mathcal{U} \subset C(Y)$ such that $1 \in \mathcal{U}$ and $\mathcal{U}$ is a vector space over $\mathbb{Q}$. We can find representatives $\Phi(u)$ of $\mathbb{E}[u(Y) | \mathcal{F}]$ such that the map

$$\mathcal{U} \ni u \mapsto \Phi(u) \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$$

is $\mathbb{Q}$ -linear, $\Phi(1) = 1$ and $\Phi(u) \geq 0$ if $u \geq 0$. For every nonnegative $f \in C(U)$ we set

$$\Phi^*(f) := \sup \{ \Phi(u); u \in \mathcal{U}, 0 \leq u \leq f \}.$$

One can show that

$$\Phi^*(f) := \inf \{ \Phi(u); u \in \mathcal{U}, u \geq f \}.$$

For arbitrary $f \in C(\mathcal{Y})$ we set

$$\Phi^*(f) = \Phi^*(f^+) - \Phi^*(f^-).$$

One can show that the resulting map

$$C(\mathcal{Y}) \ni f \mapsto \Phi^*(f) \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$$

is $\mathbb{R}$ -linear, $\Phi^*(1) = 1$ and $\Phi^*(f) \geq 0$ if $f \geq 0$. The Riesz Representation Theorem 1.2.65 implies that for any $\omega \in \Omega$ there exists a probability measure $\mu_\omega : \mathcal{B}_Y \rightarrow [0, 1]$ such that

$$\Phi^*(f)(\omega) = \int_Y f(y) \mu_\omega[dy].$$

One then shows that for any $B \in \mathcal{B}_Y$ the map $\Omega \ni \omega \mapsto \mu_\omega[B] \in [0, 1]$ is $\mathcal{F}$ -measurable and thus it is a regular version of the conditional distribution of Y given $\mathcal{F}$. $\square$

In the special case when $\mathcal{F}$ is the σ -algebra generated by a measurable map $X : \Omega \rightarrow \mathfrak{X}$, where $\mathfrak{X}$ is some measurable space, we use the notation

$$\mathbb{P}_Y[dy \| X] := \mathbb{P}_Y[dy \| \sigma(X)]$$

to denote a regular version for the conditional distribution of Y given X . This is a random Borel measure on $\mathcal{Y}$.

Example 1.4.30. Consider the special case of Theorem 1.4.29 where $\mathcal{Y} = \mathbb{R}$ and $Y \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. For any sigma subalgebra $\mathcal{F} \subset \mathcal{S}$ there exists a kernel $Q : (\Omega, \mathcal{F}) \rightsquigarrow (\mathbb{R}, \mathcal{B}_\mathbb{R})$ such that

$$\mathbb{P}[Y \leq y \| \mathcal{F}] = Q_\square[(-\infty, y]].$$

Moreover

$$\mathbb{E}[Y \| \mathcal{F}] = \int_{\mathbb{R}} y Q_\square[dy], \quad \mathbb{P} - \text{a.s. on } \Omega.$$

$\square$

Example 1.4.31. Suppose that X_0, Y_0, X_1, Y_1 are random variables and $T : \mathbb{R}^2 \rightarrow \mathbb{R}^k$ is a Borel measurable map. Denote by $\mathbb{P}^0$ the joint probability distribution of (X_0, Y_0) . Suppose that the joint distribution of (X_1, Y_1) has the form

$$\mathbb{P}^1[dxdy] = g(T(x, y)) \mathbb{P}^0[dxdy]$$

for some nonnegative measurable function $g : \mathbb{R}^k \rightarrow [0, \infty)$.

We denote by $\mathbb{P}^i[- \| T]$ the regular conditional probability $\mathbb{P}^i[- \| \sigma(T)]$. In other words, for any bounded nonnegative measurable function $f : \mathbb{R}^k \rightarrow [0, \infty)$ and any Borel set $B \subset \mathbb{R}^2$ we have $\mathbb{P}^i[B \| T] \in \mathcal{L}_+^0(\mathbb{R}^2, \sigma(T))$ and

$$\int_{\mathbb{R}^2} \mathbf{I}_B f(T(x, y)) \mathbb{P}^i[dxdy] = \int_{\mathbb{R}^2} \mathbb{P}^i[B \| T] f(T(x, y)) \mathbb{P}^i[dxdy], \quad i = 0, 1.$$

Note that

$$\int_{\mathbb{R}^2} \mathbf{I}_B f(T(x, y)) \mathbb{P}^1[dxdy] = \int_{\mathbb{R}^2} \mathbf{I}_B f(T(x, y)) g(T(x, y)) \mathbb{P}^0[dxdy]$$

$$\begin{aligned}
&= \int_{\mathbb{R}^2} \mathbb{P}^0[B \parallel T] f(T(x, y)) g(T(x, y)) \mathbb{P}^0[dx dy] \\
&= \int_{\mathbb{R}^2} \mathbb{P}^0[B \parallel T] f(T(x, y)) \mathbb{P}^1[dx dy].
\end{aligned}$$

Hence

$$\mathbb{P}^1[B \parallel T] = \mathbb{P}^0[A \parallel T], \quad \forall B \in \mathcal{B}_{\mathbb{R}^2}.$$

Suppose that the distribution $\mathbb{P}^0$ is known and would like to get information about the distribution of (X_1, Y_1) by investigating $T(X_0, Y_0)$. The above equality shows that knowledge of T adds nothing to our understanding of the density $g(T(x, y))$ beyond what we know from (X_0, Y_0) . $\square$

1.4.5. Disintegration of measures. Suppose that $(\Omega_i, \mathcal{S}_i)$, $i = 0, 1$ are two measurable spaces and

$$K : (\Omega_0, \mathcal{S}_0) \rightsquigarrow (\Omega_1, \mathcal{S}_1)$$

is a kernel from $(\Omega_0, \mathcal{S}_0)$ to $(\Omega_1, \mathcal{S}_1)$. Then any measure μ_0 on $(\Omega_0, \mathcal{S}_0)$ defines a measure $\mu = \mu_{K, \mu_0}$ on $(\Omega, \mathcal{S}) := (\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$ via the equality

$$\mu[S] = \int_{\Omega_0} \left(\int_{\Omega_1} \mathbf{I}_S(\omega_0, \omega_1) K_{\omega_0}[d\omega_1] \right) \mu_0[d\omega_0]. \quad (1.4.40)$$

We say that a measure μ on $(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$ is *disintegrated* by μ_0 , or that μ_0 *disintegrates* μ if μ is of the form μ_{K, μ_0} defined above. In this case K is called a *disintegration kernel*, and we say that K *disintegrates* μ with respect to μ_0 . Often we will use the notation

$$\mu[d\omega_0 d\omega_1] = \mu_0[d\omega_0] K_{\omega_0}[d\omega_1] \quad (1.4.41)$$

Observe that if K is a Markovian kernel and μ_0 is a probability measure, then μ_{K, μ_0} is a probability measure. In this case, for emphasis, we use the notation $\mathbb{P}_{K, \mu_0}$

Example 1.4.32. For any probability measures μ_i on $(\Omega_i, \mathcal{S}_i)$, $i = 0, 1$, the product measure $\mu = \mu_0 \otimes \mu_1$ is disintegrated by μ_0 since

$$\mu = \mathbb{P}_{K, \mu_0}, \quad K_{\omega_0}[-] = \mu_1[-].$$

$\square$

Example 1.4.33. Consider a measure ν on $(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$, and a measure μ on $(\Omega_0, \mathcal{S}_0)$. Suppose that $f : (\Omega_0, \mathcal{S}_0) \rightarrow [0, \infty)$ is a nonnegative measurable function. Denote by μ_f the measure $\mu_f[d\omega_0] = f(\omega_0)\mu[d\omega_0]$.

If ν is disintegrated by μ_f then it is also disintegrated by μ . Indeed if K is the disintegration kernel of ν with respect to μ_f , $K = K_{\omega_0}[d\omega_1]$ so that

$$\nu[d\omega_0 d\omega_1] = \mu_f[d\omega_0] K_{\omega_0}[d\omega_1] = \mu[d\omega_0] f(\omega_0) K_{\omega_0}[d\omega_1].$$

Hence, the kernel K^f given by $K_{\omega_0}^f[d\omega_1] = f(\omega_0) K_{\omega_0}[d\omega_1]$ disintegrates ν with respect to μ . $\square$

Consider two measurable spaces $(\Omega_i, \mathcal{S}_i)$, $i = 0, 1$. We have natural projections

$$\pi_i : \Omega \rightarrow \Omega_i, \quad \pi_i(\omega_0, \omega_1) = \omega_i, \quad i = 0, 1,$$

and we set $\tilde{\mathcal{S}}_0 := \pi_0^{-1}(\mathcal{S}_0) \subset \mathcal{S} := \mathcal{S}_0 \otimes \mathcal{S}_1$.

Suppose that the probability measure μ on $(\Omega, \mathcal{S}) := (\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$ is disintegrated by $\mu_0 := (\pi_0)_\# \mu$, i.e., $\mu = \mu_{K, \mu_0}$. We can rewrite (1.4.41) as

$$\mu[d\omega_0 d\omega_1] = (\pi_0)_\# \mu[d\omega_0] K_{\omega_0}[d\omega_1]. \quad (1.4.42)$$

Note that if $\mu_1 := (\pi_1)_\# \mu$, then, for any $S_1 \in \mathcal{S}_1$, we have

$$\mu_1[S_1] = \mu[\Omega_0 \times S_1] = \int_{\Omega_0} K_{\omega_0}[S_1] \mu_0[d\omega_0].$$

In other words, $\mu_1 = K_* \mu_0$. Thus, μ_1 is a mixture of the measures $(K_{\omega_0}[-])_{\omega_0 \in \Omega_0}$ driven by μ_0 .

Observe next that for any $\tilde{S}_0 = S_0 \times \Omega_1 \in \tilde{\mathcal{S}}_0$, and any $S_1 \in \mathcal{S}_1$, we have

$$\mu[\pi_1^{-1}(S_1) \cap \tilde{S}_0] = \mu[S_0 \times S_1] \stackrel{(1.4.40)}{=} \int_{S_0} K_{\omega_0}[S_1] \mu_0[d\omega_0].$$

This shows that the the map

$$\tilde{K} : \Omega \times \mathcal{S}_1 \rightarrow [0, 1], \quad ((\omega_0, \omega_1), S_1) \rightarrow \tilde{K}_{(\omega_0, \omega_1)}[S_1] = K_{\omega_0}[S_1]$$

a regular version of the conditional distribution of the measurable map π_1 conditioned on $\tilde{\mathcal{S}}_0$; see (1.4.38).

Conversely, any regular version of the conditional distribution $\mathbb{P}_{\pi_1}[-|\tilde{\mathcal{S}}_0]$ of π_1 given $\tilde{\mathcal{S}}_0$ produces a disintegration kernel of the measure μ . Indeed, if $Q_{(\omega_0, \omega_1)}[-]$ is such a regular distribution, then its $\tilde{\mathcal{S}}_0$ -measurability implies that for any $S_1 \in \mathcal{S}_1$ the function

$$(\omega_0, \omega_1) \mapsto Q_{(\omega_0, \omega_1)}[S_1]$$

is independent¹⁴ of ω_1 . Then, writing $Q_{\omega_0} := Q_{(\omega_0, \omega_1)}$, we have

$$\begin{aligned} \mu[S_0 \times S_1] &= \mu[\pi_1^{-1}(S_1) \cap \tilde{S}_0] \stackrel{(1.4.38)}{=} \int_{S_0 \times \Omega_1} Q_{\omega_0}[S_1] \mu[d\omega_0 d\omega_1] \\ &= \int_{S_0} Q_{\omega_0}[S_1] \mu_0[d\omega_0], \quad \mu_0 := (\pi_0)_\# \mu. \end{aligned}$$

Thus μ is disintegrated by μ_0 and Q is the disintegration kernel. Theorem 1.4.29 implies the next result.

Corollary 1.4.34. *If $(\Omega_1, \mathcal{S}_1)$ is isomorphic, as a measurable space, with a Lusin space equipped with the Borel sigma-algebra then, for any measurable space $(\Omega_0, \mathcal{S}_0)$, any probability measure by $\mathbb{P}$ on $(\Omega_0 \times \Omega_1, \mathcal{S}_0 \otimes \mathcal{S}_1)$ is disintegrated by its marginal $\mathbb{P}_0 := (\pi_0)_\# \mathbb{P}$. $\square$*

Example 1.4.35. Consider a random 2-dimensional vector (X, Y) with joint distribution

$$\mathbb{P}_{X,Y} \in \text{Prob}(\mathbb{R}^2).$$

According to Corollary 1.4.34, the distribution $\mathbb{P}_X$ of X disintegrates the joint distribution $\mathbb{P}_{X,Y}$. Suppose that $K_x[dy]$ is a disintegration kernel of $\mathbb{P}_{X,Y}$, i.e.,

$$\mathbb{P}_{X,Y}[dxdy] = K_x[dy] \mathbb{P}_X[dx].$$

¹⁴For any $\tilde{S}_0 \in \tilde{\mathcal{S}}_0$, the indicator $\mathbf{I}_{\tilde{S}_0}(\omega_0, \omega_1)$ is independent of ω_1 and thus any $\tilde{\mathcal{S}}_0$ -elementary function is independent of ω_1 . Alternatively, invoke Dynkin's Theorem 1.1.24.

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function such that $f(Y) \in L^1$. Then $\mathbb{E}[f(Y) | X]$ is well defined and has the form $\mathbb{E}[f(Y) | X] = h(X)$, for some measurable function h . Traditionally $h(x)$ is denoted by $\mathbb{E}[f(Y) | X = x]$.

We can give a more explicit description of $\mathbb{E}[f(Y) | X = x]$ using the disintegration kernel. More precisely, we will show that

$$\mathbb{E}[f(Y) | X = x] = \int_{\mathbb{R}} f(y) K_x[dy] =: g(x). \quad (1.4.43)$$

A Monotone Class argument shows that the $g(x)$ is Borel measurable. For any $x_0 \in \mathbb{R}$ we have

$$\begin{aligned} \mathbb{E}[f(Y) \mathbf{I}_{\{X \leq x_0\}}] &= \int_{\mathbb{R}^2} f(y) \mathbf{I}_{(-\infty, x_0]}(x) \mathbb{P}_{X,Y}[dxdy] \\ &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(y) K_x[dy] \right) \mathbf{I}_{(-\infty, x_0]}(x) \mathbb{P}_X[dx] = \int_{\mathbb{R}} g(x) \mathbf{I}_{(-\infty, x_0]}(x) \mathbb{P}_X[dx] \\ &= \mathbb{E}[g(X) \mathbf{I}_{\{X \leq x_0\}}]. \end{aligned}$$

Since the sets $\{X \leq x_0\}$ form a π -system that generate $\sigma(X)$ we deduce that

$$\mathbb{E}[f(Y) \mathbf{I}_S] = \mathbb{E}[g(X) \mathbf{I}_S], \quad \forall S \in \sigma(X).$$

Thus

$$g(X) = \mathbb{E}[f(Y) | X].$$

We write this as

$$\mathbb{E}[f(Y) | X] = \int_{\mathbb{R}} f(y) K_X[dy].$$

Hence the conditional expectations $\mathbb{E}[f(Y) | X]$ are determined by the kernel K that disintegrates the joint probability distribution $\mathbb{P}_{X,Y}$.

In particular, if $B \subset \mathbb{R}$ is a Borel set, and $f = \mathbf{I}_B$ we have the *law of total probability*

$$\mathbb{P}[Y \in B] = \mathbb{E}[\mathbf{I}_B(Y)] = \int_{\mathbb{R}} \mathbb{E}[\mathbf{I}_B(Y) | X = x] \mathbb{P}_X[dx],$$

where

$$\mathbb{E}[\mathbf{I}_B(Y) | X = x] = \mathbb{P}[Y \in B | X = x] = \int_B K_x[dy].$$

This proves that *the disintegration kernel $K_x[dy]$ is a regular conditional distribution of Y given X , i.e.,*

$$\begin{aligned} K_x[dy] &= \mathbb{P}[Y \in [y, y + dy] | X \in [x, x + dx]] \\ &= \frac{\mathbb{P}[X \in [x, x + dx], Y \in [y, y + dy]]}{\mathbb{P}[X \in [x, x + dx]]}. \end{aligned}$$

For this reason $K_x[dy]$ is called the conditional distribution of Y given that $X = x$ and it is sometimes denoted by $\mathbb{P}_{Y|X=x}[dy]$. Hence we can rewrite (1.4.43) as

$$\mathbb{E}[f(Y) | X = x] = \int_{\mathbb{R}} f(y) \mathbb{P}_{Y|X=x}[dy]. \quad (1.4.44)$$

We often think of $\mathbb{P}_{Y|X=x} = K_x[dy]$ the distribution of a random variable $Y|X = x$,

Observe that if $\mathbb{P}_{X,Y}$ is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^2$ so that

$$\mathbb{P}_{X,Y}[dxdy] = p(x, y) dxdy,$$

then

$$\mathbb{P}_{Y|X=x}[dy] = \frac{p(x,y)}{p_0(x)} dy, \quad p_0(x) = \int_{\mathbb{R}} p(x,y) dy,$$

where we set $\frac{p(x,y)}{p_0(x)} = 0$ if $p_0(x) = 0$. Then

$$\mathbb{E}[f(Y)|X=x] = \int_{\mathbb{R}} f(y) \frac{p(x,y)}{p_0(x)} dy.$$

□

Example 1.4.36. Suppose that $X_1, \dots, X_n$ are independent and uniformly distributed in the interval $[0, L]$. Set

$$X_{(n)} := \max_{1 \leq k \leq n} X_k, \quad X_{(1)} := \min_{1 \leq k \leq n} X_k$$

Note that

$$\mathbb{P}[X_{(n)} \leq x] = \mathbb{P}[X_k \leq x, \forall k = 1, \dots, n] = \left(\frac{x}{L}\right)^n,$$

so that the probability distribution of $X_{(n)}$ is

$$\mathbb{P}_n[dx] = n \frac{x^{n-1}}{L^n} \mathbf{I}_{[0,L]}(x) dx.$$

Note that

$$\frac{n+1}{n} \mathbb{E}[X_{(n)}] = L.$$

Note also that

$$\mathbb{P}\left[\frac{n+1}{n} X_{(n)} < L - \varepsilon\right] = \left(\frac{n(L-\varepsilon)}{(n+1)L}\right)^n,$$

and, since

$$\sum_n \left(\frac{n(L-\varepsilon)}{(n+1)L}\right)^n < \infty$$

we deduce from the Borel-Cantelli lemma that $\frac{n+1}{n} X_{(n)} \rightarrow L$, a.e.. For these reasons, the statisticians refer to the sequence of random variables $X_{(n)}$ as an *unbiased estimator* of L .

Similarly,

$$\mathbb{P}[X_{(1)} > x] = \mathbb{P}[X_k > x, \forall k = 1, \dots, n] = \left(\frac{L-x}{L}\right)^n,$$

so the probability distribution of $X_{(1)}$ is

$$\mathbb{P}_1[dx] = n \underbrace{\frac{(L-x)^{n-1}}{L^n}}_{=: \rho_1(x)} \mathbf{I}_{[0,L]}(x) dx.$$

Let us compute the conditional distribution $\mathbb{P}_{X_{(n)}|X_{(1)}=x_1}[dx_n]$. We begin by computing the random variables.

$$\mathbb{P}[X_{(n)} \leq x_n \mid X_{(1)} = x_1], \quad 0 \leq x_n \leq L.$$

Observe first that $\forall 0 \leq x_1, x_n \leq L$,

$$\mathbb{E}[\mathbf{I}_{X_{(n)} \leq x_n} \mathbf{I}_{X_{(1)} \geq x_1}] = \mathbb{P}[x_1 \leq X_1, \dots, X_n \leq x_n] = \frac{(x_n - x_1)_+^n}{L^n}.$$

We need to find a function $f(x_1) = f_{x_n}(x_1)$ such that

$$\mathbb{E}[f(X_{(1)})\mathbf{I}_{X_{(1)} \geq x_1}] = \frac{(x_n - x_1)_+^n}{L^n}, \quad \forall x_1,$$

i.e.,

$$\int_{[x_1, L]} f(x) \rho_1(x) dx = \frac{(x_n - x_1)_+^n}{L^n}, \quad \forall x_1.$$

Derivating with respect to x_1 we deduce

$$f(x_1) \rho_1(x_1) = n \frac{(x_n - x_1)_+^{n-1}}{L^n}.$$

Hence

$$\mathbb{P}[X_{(n)} \leq y | X_{(1)} = x_1] = n \frac{(x_n - x_1)_+^{n-1}}{L^n \rho_1(x_1)} = \frac{(y - x_1)_+^{n-1}}{(L - x_1)^{n-1}}.$$

Thus, the conditional distribution of $X_{(n)}$ given that $X_{(1)} = x_1$ is

$$\mathbb{P}_{X_{(n)}|X_{(1)}=x_1}[dx_n] = \frac{(n-1)(x_n - x_1)_+^{n-2}}{(L - x_1)^{n-1}} dx_n.$$

We define the *empirical gap* or *sample range* to be the random variable $G = X_{(n)} - X_{(1)}$. To find the distribution of G we condition on $X_{(1)}$ and we have

$$\begin{aligned} \mathbb{P}[G \leq g] &= \int_{[0, L]} \mathbb{P}[X_{(n)} \leq x_1 + g | X_{(1)} = x_1] \mathbb{P}_{X_{(1)}}[dx_1] \\ &= \int_{[0, L]} \mathbb{P}[X_{(n)} \leq X_{(1)} + g | X_{(1)} = x_1] \rho_1(x_1) dx_1. \end{aligned}$$

Now observe that

$$\begin{aligned} \mathbb{P}[X_{(n)} \leq X_{(1)} + g | X_{(1)} = x_1] &= \int_{[0, \min(L, x_1 + g)]} \mathbb{P}_{X_{(n)}|X_{(1)}=x_1}[dx_n] \\ &= \int_{[0, \min(L, x_1 + g)]} \frac{(n-1)(x_n - x_1)_+^{n-2}}{(L - x_1)^{n-1}} dx_n \\ &= \frac{g^{n-1}}{(L - x_1)^{n-1}} \mathbf{I}_{[0, L-g]}(x_1) + \mathbf{I}_{[L-g, L]}(x_1). \end{aligned}$$

Thus

$$\mathbb{P}[G \leq g] = \frac{ng^{n-1}}{L^n} \int_0^{L-g} dx_1 + \int_{[L-g, L]} \rho_1(x_1) dx_1 = \frac{ng^{n-1}(L-g)}{L^n} + \frac{g^n}{L^n}.$$

We deduce

$$\frac{d}{dg} \mathbb{P}[G \leq g] = \frac{n(n-1)g^{n-2}}{L^{n-1}} + \frac{n^2 g^{n-1}}{L^n} - \frac{ng^{n-1}}{L^n} = \frac{n(n-1)g^{n-2}}{L^{n-1}} \left(1 - \frac{g}{L}\right).$$

Thus, the probability distribution of G is

$$P_G[dg] = \frac{n(n-1)g^{n-2}}{L^{n-1}} \left(1 - \frac{g}{L}\right) \mathbf{I}_{[0, L]}(g) dg.$$

If $L = 1$, then the above distribution is the Beta distribution $\text{Beta}(n-1, 2)$.

□

Example 1.4.37. Suppose that $f : [0, 1] \rightarrow \mathbb{R}$ is a C^1 -function whose graph has length L , i.e.,

$$L = \int_0^1 \sqrt{1 + |f'(x)|^2} dx.$$

Define a random measure $K : ([0, 1], \mathcal{B}) \rightsquigarrow (\mathbb{R}, \mathcal{B})$, $K_x = \delta_{f(x)}$.

Let

$$\mu_0[dx] = \frac{\sqrt{1 + |f'(x)|^2}}{L} \cdot \lambda[dx] \in \text{Prob}([0, 1]).$$

Then the Borel probability measure $\mathbb{P}_{K, \mu_0}$ on $[0, 1] \times \mathbb{R}$ corresponds to the integration with respect to the normalized arclength along the graph of f . $\square$

Example 1.4.38. Suppose that $X_1, \dots, X_n$ are independent random variables with common distribution $p(x)\lambda[dx]$. Denote by $\mathbf{X}$ the random vector $(X_1, \dots, X_n)$. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a Borel measurable function. Denote by $\mathbb{P}$ the distribution of the random vector $(\mathbf{X}, f(\mathbf{X}))$. This is disintegrated by the distribution $\mu_0 := \mathbb{P}_{\mathbf{X}}$ of the random vector $\mathbf{X}$. The disintegration kernel K is the conditional distribution of $f(\mathbf{X})$ given $\mathbf{X}$. We deduce that

$$K_{x_1, \dots, x_n}[-] = \delta_{f(x_1, \dots, x_n)}.$$

If B_0 is a Borel subset of $\mathbb{R}^n$ and B_1 is a Borel subset of $\mathbb{R}$, then

$$\mathbb{P}[B_0 \times B_1] = \int_{B_0} \mathbf{I}_{B_1}(f(x_1, \dots, x_n)) p(x_1) \cdots p(x_n) dx_1 \cdots dx_n.$$

Using a notation dear to theoretical physicists we can rewrite the above equality as

$$\mathbb{P}[dx_1 \cdots dx_n dy] = (\delta(y - f(x_1, \dots, x_n)) p(x_1) \cdots p(x_n)) dy dx_1 \cdots dx_n,$$

where $\delta(z)$ denotes the Dirac “function” on the real axis. $\square$

Remark 1.4.39. We refer to [33] for a very enlightening presentation of a more general concept of disintegration and some of its application to statistics. $\square$

1.5. What are stochastic processes?

We have already met stochastic processes though we have not called them so. This section has a rather restricted goal namely, to explain what they are, describe a few basic features and more importantly, show that stochastic processes with prescribed statistics do exist as mathematical objects.

1.5.1. Definition and examples. A *stochastic process* or *random process* is simply a family $(X_t)_{t \in T}$ of random variables parametrized by a set T . They are all defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. The variables could be real valued, vector valued or we can allow them to be valued in a measurable space $(\mathbb{X}, \mathcal{F})$, where $\mathcal{F}$ is a sigma-algebra of subsets of $\mathbb{X}$. Frequently $\mathbb{X} = \mathbb{R}^n$ for some n but, as we will see below, it is very easy to produce more complicated examples

Obviously stochastic processes exist, but once we impose some restriction on their behavior, the existence of such stochastic processes is less obvious. A classical situation, intensely investigated in probability, is that of families $(X_t)_{t \in \mathbb{T}}$ of real valued random variables that are *independent, identically distributed* (or i.i.d. for brevity). We denote by $\mathbb{P}_X$ their common distribution.

A basic question arises. Given a Borel probability measure μ on $\mathbb{R}$ and a set T , can we find a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and *independent* random variables

$$X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad t \in T,$$

such that $\mathbb{P}_{X_t} = \mu, \forall t \in T$?

When T is finite, say $T := \{1, 2, \dots, n\}$ the answer is positive. As probability space we can take

$$(\Omega, \mathcal{S}, \mathbb{P}) := (\mathbb{R}^n, \mathcal{B}_{\mathbb{R}^n}, \mu^{\otimes n}).$$

The random variables are then the coordinate functions

$$X_k : \mathbb{R}^n \rightarrow \mathbb{R}, \quad X_k(x_1, \dots, x_n) = x_k, \quad k = 1, \dots, n.$$

Using the notation $\mathbb{R}^T$ instead of $\mathbb{R}^n$ we see that we have defined a probability measure on the space of functions $T \rightarrow \mathbb{R}$.

If T is infinite, say $T = \mathbb{N}$, the question is then about the existence of a sequence $(X_n)_{n \in \mathbb{N}}$ of i.i.d. random variables with common probability distribution μ . A substantial portion of probability is devoted to such sequences and it would be embarrassing, to say the least, if it turned out they do not exist. We will see that this is not the case.

It is also very easy to stumble into situations in which the random variables are not independent, or take value in some infinite dimensional space. We have encountered a such a situation already.

Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $\mathcal{F} \subset \mathcal{S}$ is a sigma-subalgebra. For any $S \in \mathcal{S}$ choose a version $X_S \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ of the conditional probability $\mathbb{P}[S \mid \mathcal{F}]$. The collection $(X_S)_{S \in \mathcal{S}}$ is a stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$ parametrized by $\mathcal{S}$. We can view it as a map

$$X : \Omega \rightarrow [0, 1]^{\mathcal{S}} = \text{the space of functions } \mathcal{S} \rightarrow [0, 1].$$

Here is another such situation, of a different nature.

Example 1.5.1. Suppose that $A_0, A_1, \dots, A_n$ is a family of i.i.d. (real valued) random variables defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For every $t \in [0, 1]$ we set

$$X_t := A_0 + A_1 t + \dots + A_n t^n.$$

We now have on our hands a family of random variables $(X_t)_{t \in [0, 1]}$. These are dependent. To understand why, suppose for simplicity that the variables A_k have mean zero and variance 1. Then X_t has mean zero and for any $s, t \in [0, 1]$

$$\text{Cov}[X_s, X_t] = \mathbb{E}[X_s X_t] = 1 + (st) + \dots + (st)^n > 1.$$

Thus the random variables $(X_t)_{t \in [0, 1]}$ are dependent.

Let $\mathbb{X}$ denote the Banach space $C([0, 1])$ equipped with the sup norm. The family (X_t) defines a map

$$X : \Omega \rightarrow \mathbb{X}, \quad \Omega \ni \omega \mapsto X_t(\omega) = \sum_{k=0}^n A_k(\omega) t^k \in \mathbb{X}.$$

The space $C([0, 1])$ comes with a natural family of (linear) evaluation functionals

$$\mathbf{Ev}_t : C([0, 1]) \rightarrow \mathbb{R}, \quad t \in [0, 1], \quad \mathbf{Ev}_t(f) = f(t), \quad \forall f \in C([0, 1]).$$

Note that $X_t = \mathbf{Ev}_t \circ X$. The Borel sigma-algebra of $C([0, 1])$ coincides with the sigma-algebra generated by the collection of functions $\mathbf{Ev}_t, t \in [0, 1]$; see Exercise 1.4. This implies that the map $X : \Omega \rightarrow \mathbb{X}$ is measurable with respect to the Borel sigma-algebra of $\mathbb{X}$. The push-forward of $\mathbb{P}$ via the map X defines a Borel probability $\mathbb{P}_X$ measure on $\mathbb{X}$, so $(\mathbb{X}, \mathcal{B}_{\mathbb{X}}, \mathbb{P}_X)$ is a probability space. Thus

we can view $X_\bullet$ as a *random continuous function*. The measure $\mathbb{P}_X$ is called the *distribution* of this stochastic process. $\square$

Suppose now that $(X_t)_{t \in T}$ is a general family of random variables

$$X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{F}),$$

where $(\mathbb{X}, \mathcal{F})$ is a measurable space. This family defines a map

$$X : T \times \Omega \rightarrow \mathbb{X}, \quad T \times \Omega \ni (t, \omega) \mapsto X(t, \omega) := X_t(\omega) \in \mathbb{X},$$

such that X_t is measurable for any t .

Equivalently, we can view this as a map

$$X : \Omega \rightarrow \mathbb{X}^T = \text{the space of functions } f : T \rightarrow \mathbb{X}, \quad (1.5.1)$$

where to each $\omega \in \Omega$ we associate the function $X(\omega) : T \rightarrow \mathbb{X}, t \mapsto X_t(\omega)$.

It is convenient to regard $\mathbb{X}^T$ as a product of copies $\mathbb{X}_t$ of $\mathbb{X}$, $t \in T$,

$$\mathbb{X}^T = \prod_{t \in T} \mathbb{X}_t.$$

Each copy $\mathbb{X}_t$ is equipped with a copy $\mathcal{F}_t$ of the sigma-algebra $\mathcal{F}$.

The map (1.5.1) is measurable with respect to the sigma-algebra $\mathcal{F}^T$ in $\mathbb{X}^T$, the smallest sigma-algebra $\mathcal{S}$ in $\mathbb{X}^T$ such that all the evaluation maps

$$\mathbf{Ev}_t : (\mathbb{X}^T, \mathcal{S}) \rightarrow (\mathbb{X}, \mathcal{F}), \quad \mathbf{Ev}_t(f) := f(t),$$

are measurable. Equivalently,

$$\mathcal{F}^T = \bigvee_{t \in T} \mathbf{Ev}_t^{-1}(\mathcal{F}).$$

Any measurable map $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}^T, \mathcal{F}^T)$ defines a stochastic process

$$X_t(\omega) = \mathbf{Ev}_t(X(\omega)).$$

The pushforward of $\mathbb{P}$ via X is called the *distribution* the stochastic process and we denote it by $\mathbb{P}_X$. Thus we can view $(X_t)_{t \in T}$ as a random element of $\mathbb{X}^T$, i.e., a *random function*.

Note that any probability measure on $\mathcal{F}^T$ is the distribution of a stochastic process, namely the tautological process

$$\mathbb{1} : (\mathbb{X}^T, \mathcal{F}^T, \mathbb{P}) \rightarrow (\mathbb{X}^T, \mathcal{F}^T, \mathbb{P}), \quad \mathbb{1}_t(\omega) = \omega(t), \quad \forall \omega \in \mathbb{X}^T.$$

Suppose that $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}^T, \mathcal{F}^T)$ is a stochastic process. For any finite set

$$I = \{t_1, \dots, t_m\} \subset T$$

we have a sigma-algebra $\mathcal{F}^I$ in $\mathbb{X}^I$,

$$\mathcal{F}^I = \mathcal{F}_{t_1} \otimes \dots \otimes \mathcal{F}_{t_m},$$

and we obtain a random “vector”

$$X^I : (\Omega, \mathcal{S}) \rightarrow (\mathbb{X}^I, \mathcal{F}^I), \quad \omega \mapsto (X_{t_1}(\omega), \dots, X_{t_m}(\omega)) \in \mathbb{X}^I.$$

We denote by $\mathbb{P}_I \in \text{Prob}(\mathbb{X}^I, \mathcal{F}^I)$ its probability distribution

$$\mathbb{P}_I := (X^I)_\# \mathbb{P} : \mathcal{F}^I \rightarrow [0, 1].$$

Note that we have a tautological measurable projection $\Pi_I : \mathbb{X}^T \rightarrow \mathbb{X}^I$, and

$$\mathbb{P}_I = (\Pi_I)_\#(\mathbb{P}_X).$$

Suppose now that $J \subset T$ is another finite set containing I

$$J = \{t_1, \dots, t_m, t_{m+1}, \dots, t_n\}, \quad n > m.$$

We get in a similar fashion a probability measure on $\mathbb{X}^J$. We have a canonical projection

$$\mathcal{P}_{IJ} : \mathbb{X}^J \rightarrow \mathbb{X}^I, \quad (x_{t_1}, \dots, x_{t_m}, x_{t_{m+1}}, \dots, x_{t_n}) \mapsto (x_{t_1}, \dots, x_{t_m}).$$

and, since $\mathbb{X}^I = \mathcal{P}_{IJ}(\mathbb{X}^J)$, we have

$$(\mathcal{P}_{IJ})_\# \mathbb{P}_J = \mathbb{P}_I. \quad (1.5.2)$$

Observe that $\mathcal{F}^T$ is generated by the collection of subsets $\Pi_I^{-1}(F_I)$, $I \subset T$ finite, $F_I \in \mathcal{F}^I$. This collection is an algebra of subsets of $\mathbb{X}^T$. Proposition 1.2.4 shows that the distribution $\mathbb{P}_X$ of the stochastic process X is the unique probability measure $\bar{\mathbb{P}}$ on $\mathbb{X}^T$ such that for any finite subset $I \subset T$, and any $F_I \subset \mathcal{F}^I$ we have

$$\bar{\mathbb{P}}[\Pi_I^{-1}(F_I)] = \mathbb{P}_I[F_I].$$

Equivalently, this means

$$\mathbb{P}_I = (\Pi_I)_\#(\bar{\mathbb{P}}).$$

A family of measures $\mathbb{P}_I$ on $\mathbb{X}^I$, I finite subset of T , constrained by the compatibility condition (1.5.2) for any finite subsets $I \subset J \subset T$ is said to be a *projective* or *consistent family*.

We have thus shown that to any probability measure $\bar{\mathbb{P}}$ on $(\mathbb{X}^T, \mathcal{F}^T)$ we can naturally associate a projective the family of probability measures $\mathbb{P}_I := (\pi_I)_\#(\bar{\mathbb{P}})$. Moreover, $\bar{\mathbb{P}}$ is uniquely determined by this projective family.

There are other ways of constructing projective families.

Example 1.5.2. Suppose that we are given a sequence of measurable spaces $(\mathfrak{X}_n, \mathcal{F}_n)_{n \geq 0}$ is a measurable space. Recall that for any real numbers $a < b$ we have $\llbracket a, b \rrbracket := [a, b] \cap \mathbb{Z}$.

$$\mathfrak{X}^{\llbracket 0, n \rrbracket} := \prod_{k=0}^n \mathfrak{X}_k, \quad \mathcal{F}^{\llbracket 0, n \rrbracket} := \bigotimes_{k=0}^n \mathcal{F}_k.$$

Consider a family of Markovian kernels (see Subsection 1.4.4)

$$K_n : (\mathfrak{X}^{\llbracket 0, n \rrbracket}, \mathcal{F}^{\llbracket 0, n \rrbracket}) \rightsquigarrow (\mathfrak{X}_{n+1}, \mathcal{F}_{n+1}), \quad n \in \mathbb{N}_0 = \mathbb{Z}_{\geq 0}.$$

In other words, we have random probability measure

$$\mathfrak{X}^{\llbracket 0, n \rrbracket} \ni (x_0, \dots, x_n) \rightarrow K_{x_0, x_1, \dots, x_n}[dx_{n+1}]$$

on $(\mathfrak{X}_{n+1}, \mathcal{F}_{n+1})$. Then, starting with a probability measure μ_0 on $(\mathfrak{X}, \mathcal{F})$, we obtain a family of probability measures $\mathbb{P}_n$ on $\mathfrak{X}^{\llbracket 0, n \rrbracket}$ described inductively by the disintegration formula (1.4.40)

$$\mathbb{P}_0 = \mu_0, \quad \mathbb{P}_{n+1} = \mathbb{P}_{K_n, \mathbb{P}_n}. \quad (1.5.3)$$

This means that for any $S \in \mathcal{F}^{\llbracket 0, n+1 \rrbracket}$ we have

$$\mathbb{P}_{n+1}[S] = \int_{\mathfrak{X}^{\llbracket 0, n \rrbracket}} \left(\int_{\mathfrak{X}} K_{\vec{x}}[dx_{n+1}] \mathbf{I}_S(\vec{x}, x_{n+1}) \right) \mathbb{P}_n[d\vec{x}], \quad \vec{x} = (x_0, \dots, x_n).$$

Equivalently, $\mathbb{P}_n$ disintegrates $\mathbb{P}_{n+1}$ and K_n is the disintegration kernel.

Denote by $\mathcal{P}_{n,n+1}$ the natural projection $\mathfrak{X}^{\llbracket 0,n+1 \rrbracket} \rightarrow \mathfrak{X}^{\llbracket 0,n \rrbracket}$,

$$(x_0, x_1, \dots, x_n, x_{n+1}) \mapsto (x_0, x_1, \dots, x_n).$$

Since K_n is a *Markovian* kernel, i.e.,

$$\int_{\mathfrak{X}} K_{\vec{x}}[dx'] = 1, \quad \forall \vec{x} \in \mathfrak{X}^{\llbracket 0,n \rrbracket},$$

we deduce that $\mathbb{P}_n = (\mathcal{P}_{n,n+1})_{\#} \mathbb{P}_{n+1}$, $\forall n \in \mathbb{N}_0$. This shows that the collection $(\widehat{\mathbb{P}}_n)_{n \in \mathbb{N}_0}$ is a projective family of probability measures.

Note that if $K_{x_0, \dots, x_n}[-]$ is independent of $x_0, \dots, x_n$, then we can think of K_n as a probability measure μ_n on $\mathfrak{X}$. In this case

$$\mathbb{P}_n = \mu_0 \otimes \dots \otimes \mu_n.$$

If $(\mathfrak{X}_n, \mathcal{F}_n) = (\mathfrak{X}, \mathcal{F})$ for all $n \geq 0$ can obtain kernels K_n as above starting from a single Markovian kernel $K = (\mathfrak{X}, \mathcal{F}) \rightsquigarrow (\mathfrak{X}, \mathcal{F})$

$$K : \mathfrak{X} \times \mathcal{F} \rightarrow [0, 1], \quad (x, F) \mapsto K_x[F].$$

More precisely, we set $K_{x_0, \dots, x_n}[dx] := K_{x_n}[dx]$.

In this case the measures $\mathbb{P}_n$ on $\mathcal{F}^{\llbracket 0,n \rrbracket}$ are defined by

$$\mathbb{P}_n[dx_0 dx_1 \dots dx_n] = \mu_0[dx_0] K_{x_0}[dx_1] \dots K_{x_{n-1}}[dx_n].$$

More precisely, for any $S \in \mathcal{F}^{\llbracket 0,n \rrbracket}$ we have

$$\mathbb{P}_n[S] = \int_{\mathfrak{X}^{\llbracket 0,n \rrbracket}} \left(\int_{\mathfrak{X}^{\llbracket 0,n-1 \rrbracket}} \mathbf{I}_S(x_0, \dots, x_{n-1}, x_n) \mathbb{P}_{n-1}[dx_0 \dots dx_{n-1}] \right) K_{x_{n-1}}[dx_n]. \quad (1.5.4)$$

Such a situation occurs in the context of Markov chains. □

1.5.2. Kolmogorov's existence theorem. Fix a topological space $\mathbb{X}$ and a parameter set T . Recall that $\mathbb{X}^T$ denotes the space of functions $T \rightarrow \mathbb{X}$.

We denote by $\mathcal{2}_0^T$ the collection of *finite* subsets of T . Denote by $\mathcal{B}$ the Borel sigma algebra on $\mathbb{X}$. For $I \in \mathcal{2}_0^T$ we denote by $\mathcal{B}_I$ the product σ -algebra in $\mathbb{X}^I$, i.e., the smallest sigma-algebra in $\mathbb{X}^I$ such that, for any $i \in I$, the natural projection

$$\mathbb{X}^I \ni (\underline{x} : I \rightarrow \mathbb{X}) \rightarrow x(i) \in \mathbb{X}$$

is $\mathcal{B}$ -measurable.

For any finite subsets $I \subset J \subset T$ we denote by $\mathcal{P}_{IJ}$ the natural projection $\mathbb{X}^J \rightarrow \mathbb{X}^I$. This associates to a function $J \rightarrow \mathbb{X}$ its restriction to I .

For $t \in T$ we denote by π_t the natural projection

$$\pi_t : \mathbb{X}^T \rightarrow \mathbb{X}, \quad \pi_t(\underline{x}) = x_t.$$

More generally, for any $I \in \mathcal{2}_0^T$ we define $\pi_I : \mathbb{X}^T \rightarrow \mathbb{X}^I$ by setting

$$\mathbb{X}^T \ni \underline{x} \mapsto \pi_I(\underline{x}) = \underline{x}|_I = (x_i)_{i \in I} \in \mathbb{X}^I.$$

Definition 1.5.3. The *natural* σ -algebra $\mathcal{E}_T$ in $\mathbb{X}^T$ is the smallest σ -algebra $\mathcal{E} \subset \mathcal{2}^{\mathbb{X}^T}$ such that all the maps π_t , $t \in T$, are $(\mathcal{E}, \mathcal{B}_{\mathbb{X}})$ -measurable, i.e., the σ -algebra generated by the family of σ -algebras $\pi_t^{-1}(\mathcal{B}_{\mathbb{X}})$. □

A *cylinder* is a subset of $\mathbb{X}^T$ of the form

$$\pi_I^{-1}(S) = S \times \mathbb{X}^{T \setminus I}, \quad I \in \mathcal{I}_0^T, \quad S \in \mathcal{B}_I.$$

We denote by $\mathcal{C}_T$ the collection of cylinders. Clearly $\mathcal{C}_T$ is an algebra of sets that generates the natural σ -algebra $\mathcal{E}_T$.

Definition 1.5.4. A *projective family* of probability measures on $\mathbb{X}^T$ is a family $\mathbb{P}_I$ of probability measures on $(\mathbb{X}^I, \mathcal{B}_I)$, $I \in \mathcal{I}_0^T$, such that for any $I \subset J$ in $\mathcal{I}_0^T$ we have

$$\mathbb{P}_I = (\mathcal{P}_{IJ})_{\#} \mathbb{P}_J. \quad (1.5.5)$$

□

As discussed in the previous subsection any Borel measure on $\mathbb{X}^T$ defines a canonical projective family. *Kolmogorov's existence (or consistency) theorem* states that, under mild topological constraints on $\mathbb{X}$, all the projective families are obtained in this fashion.

Theorem 1.5.5 (Kolmogorov existence theorem). *Suppose that $\mathbb{X}$ is a Lusin space, i.e., a Borel subset of a compact metric space; see Definition 1.4.27. For any projective family $(\mathbb{P}_I)_{I \in \mathcal{I}_0^T}$ of Borel probability measures on $\mathbb{X}^I$ there exists a probability measure $\hat{\mathbb{P}}$ on $\mathcal{E}_T$ uniquely determined by the requirement:*

$$\forall I \in \mathcal{I}_0^T, \quad \mathbb{P}_I = (\mathcal{P}_I)_{\#}(\hat{\mathbb{P}}).$$

This means that for any $B_I \in \mathcal{B}_I$,

$$\hat{\mathbb{P}}[\pi_I^{-1}(B_I)] = \mathbb{P}_I[B_I]. \quad (1.5.6)$$

Proof. The uniqueness follows from Proposition 1.2.4.

The existence is a rather deep result ultimately based on Tikhonov's compactness result. We follow the approach in [180, Sec. 30, 31].

Observe that C is a cylinder if and only if

$$\exists I \in \mathcal{I}_0^T \text{ and } B_I \in \mathcal{B}_I \text{ such that } C = \pi_I^{-1}(B_I).$$

For $I \in \mathcal{I}_0^T$ we set $\mathcal{C}_T^I := \pi^{-1}(\mathcal{B}_I) \subset \mathcal{E}_T$. Note that

$$C \in \mathcal{C}_T^I \iff C = B_I \times \mathbb{X}^{T \setminus I}, \quad B_I \in \mathcal{B}_I, \quad (1.5.7a)$$

$$C \in \mathcal{C}_T^I \cap \mathcal{C}_T^J \neq \emptyset \Rightarrow C \in \mathcal{C}_T^{I \cap J}. \quad (1.5.7b)$$

Define

$$\hat{\mathbb{P}}_I : \mathcal{C}_T^I \rightarrow [0, \infty), \quad \hat{\mathbb{P}}_I[C] = \mathbb{P}_I[\pi_I(C)].$$

Note that if $C \in \mathcal{C}_T^I \cap \mathcal{C}_T^J$, then, according to (1.5.7b), $C \in \mathcal{C}_T^K$ for some $K \subset I \cap J$. Then

$$\pi_I(C) = \mathcal{P}_{KI}^{-1}(\pi_K(C)), \quad \pi_J(C) = \mathcal{P}_{KJ}^{-1}(\pi_K(C)).$$

Thus

$$\mathbb{P}_I[\pi_I(C)] = \mathbb{P}_I[\mathcal{P}_{KI}^{-1}(\pi_K(C))] = (\mathcal{P}_{KI})_{\#} \mathbb{P}_I[\pi_K(C)] \stackrel{(1.5.5)}{=} \mathbb{P}_K[\pi_K(C)],$$

and, similarly,

$$\mathbb{P}_J[\pi_J(C)] = \mathbb{P}_J[\mathcal{P}_{KJ}^{-1}(\pi_K(C))] = (\mathcal{P}_{KJ})_{\#} \mathbb{P}_J[\pi_K(C)] \stackrel{(1.5.5)}{=} \mathbb{P}_K[\pi_K(C)],$$

Hence, if $C \in \mathcal{C}_T^I \cap \mathcal{C}_T^J$, then $\hat{\mathbb{P}}_I[C] = \hat{\mathbb{P}}_J[C]$.

We have thus defined a *finitely additive* measure $\hat{\mathbb{P}}$ on the algebra

$$\mathcal{C}_T = \bigcup_{I \in 2_0^T} \mathcal{C}_T^I.$$

To invoke Carathéodory's extension theorem (Theorem 1.2.17) it suffices to show that $\hat{\mathbb{P}}$ is countably additive on $\mathcal{C}_T$. We will achieve this step by relying on Alexandrov's Theorem 1.2.15, but to complete this step we need to make a brief foundational digression.

Digression 1.5.6 (Regularity of Borel measures). When dealing with measures on topological spaces there are several desirable compatibility conditions between the measure-theoretic objects and the topological ones.

Definition 1.5.7. Let X be a topological space and μ a Borel measure on X .

- (i) The measure μ is called *outer regular* if for any Borel set $B \in \mathcal{B}_X$ we have

$$\mu[B] = \inf_{\substack{U \supset B, \\ U \text{ open}}} \mu[U].$$

- (ii) The measure μ is called *inner regular* if for any Borel set $B \in \mathcal{B}_X$ we have

$$\mu[B] = \sup_{\substack{C \subset B, \\ C \text{ closed}}} \mu[C].$$

- (iii) The measure μ is called *regular* if it is both inner and outer regular.

- (iv) The measure μ is called *Radon* if it is outer regular, and for any Borel set $B \in \mathcal{B}_X$, we have

$$\mu[B] = \sup_{\substack{K \subset B, \\ K \text{ compact}}} \mu[K].$$

Note that a finite Borel measure is regular iff it is inner regular. From the above definition it is clear that

$$\mu \text{ is Radon} \Rightarrow \mu \text{ is regular.}$$

A deep result in measure theory states that any Borel probability measure on a Lusin space is Radon, [21, Thm. 7.4.3]. For our immediate needs we can get away by with a lot less. We have the following useful result, [166, Chap. II, Thm.1.2]. A proof is outlined in Exercise 1.66.

Theorem 1.5.8. *Any Borel probability measure on a metric space is regular.*

From Theorem 1.5.8 we deduce the following result.

Lemma 1.5.9. *Let Y be a compact metric space. Then any Borel probability measure on Y is Radon.* $\square$

This concludes our digression. $\square$

As mentioned in Remark 1.4.28(b), any Lusin space is Borel isomorphic to a compact metric space. Thus it suffices to prove Kolmogorov's theorem only in the special when $\mathbb{X}$ is a compact metric space. In this case Kolmogorov's theorem follows from Alexandrov's Theorem 1.2.15.

Note first that Tikhonov's compactness theorem implies that the space $\mathbb{X}^T$ is compact with respect to the product topology. Suppose that $C \in \mathcal{C}_T$ is a cylinder. Thus, there exists a *finite* subset $I \subset T$ and a Borel subset B_I of $\mathbb{X}^I$ such that $C = \pi_I^{-1}(B_I)$. The regularity Theorem 1.5.8 implies that for any $\varepsilon > 0$ there exists a closed subset $K_\varepsilon \in \mathbb{X}^I$ such that

$$\mathbb{P}_I[B_I \setminus K_\varepsilon] < \varepsilon.$$

Note that the set $F_\varepsilon := \pi_I^{-1}(K_\varepsilon)$ is also a cylinder contained in C , it is closed as a subset of $\mathbb{X}^T$ and

$$\widehat{\mathbb{P}}[C \setminus F_\varepsilon] = \mathbb{P}_I[B_I \setminus K_\varepsilon] < \varepsilon.$$

Alexandrov's Theorem 1.2.15 implies that $\widehat{\mathbb{P}}$ is a premeasure and thus extends to a probability measure on $\mathcal{T}$. $\square$

The real axis $\mathbb{R}$ is a Lusin space. Given a Borel probability measure $\mathbb{P}$ on $\mathbb{R}$ we can construct trivially a projective family $\mathbb{P}_I$, $I \in 2_0^\mathbb{N}$. More precisely $\mathbb{P}_I = \mathbb{P}^{\otimes |I|}$ on $\mathbb{R}^I$. We deduce that we have a natural Borel probability measure $\mathbb{R}^\mathbb{N}$. We have natural random variables on this probability space

$$X_n : \mathbb{R}^\mathbb{N} \rightarrow \mathbb{R}, \quad X_n(\underline{x}) = x_n, \quad \forall \underline{x} = (x_1, x_2, \dots) \in \mathbb{R}^\mathbb{N}.$$

Note that $\mathbb{P}_{X_n} = \mathbb{P}$, $\forall n$ and the joint distribution of $X_1, \dots, X_n$ is $\mathbb{P}^{\otimes n}$. Thus, the random variables (X_n) are independent and have identical distributions. We have thus proved the following fact.

Corollary 1.5.10. *For any probability measure $\mathbb{P} \in \text{Prob}(\mathbb{R}, \mathcal{B}_\mathbb{R})$, there exists a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and a sequence of independent identically distributed (or i.i.d. for brevity) random variables $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, with common distribution $\mathbb{P}$.* $\square$

Remark 1.5.11. (a) An earlier version of the Existence Theorem 1.5.5 was proved by P. J. Daniell [48]. We refer to [3] for an interesting historical perspective on this theorem. The existence theorem can be substantially generalized; see e.g. [21, Sec. 7.7].

(b) The proof of Theorem 1.5.5 uses in an essential fashion the topological nature of the projective family of measures $(\mathbb{P}_I)_{I \in 2_0^T}$. I want to emphasize that in this theorem the set of parameters T is arbitrary.

If the set of parameters T is countable, say $T = \mathbb{N}_0$, then one can avoid the topological assumptions.

Consider for example the projective family of measures $\mathbb{P}_n$ constructed in Example 1.5.2. Recall briefly its construction that we are given a sequence of measurable spaces $(\mathfrak{X}_n, \mathcal{F}_n)_{n \geq 0}$ and measures $\mathbb{P}_n$ on

$$(\mathfrak{X}_0 \times \dots \times \mathfrak{X}_n, \mathcal{F}_0 \otimes \dots \otimes \mathcal{F}_n)$$

such that $\mathbb{P}_n$ disintegrates $\mathbb{P}_{n+1}$, $\forall n \geq 0$. (Observe that this condition is automatically satisfied if each $\mathfrak{X}_n$ is a Lusin space.) Set

$$\mathfrak{X}^\infty := \prod_{n=0}^{\infty} \mathfrak{X}_n,$$

denote by π_n the natural projection $\mathfrak{X}^\infty \rightarrow \mathfrak{X}_n$ and by $\mathcal{F}^{\otimes \infty}$ the sigma-algebra

$$\mathcal{F}^{\otimes \infty} := \bigvee_{n \geq 0} \pi_n^{-1}(\mathcal{F}_n).$$

A theorem of C. Ionescu-Tulcea states that there exists a unique probability measure $\mathbb{P}_\infty$ on $\mathcal{F}^{\otimes\infty}$ such that

$$(\mathcal{P}_n)_\# \mathbb{P}_\infty = \mathbb{P}_n, \quad \forall n \geq 0,$$

where $\mathcal{P}_n$ denotes the natural projection $\mathfrak{X}^\infty \rightarrow \mathfrak{X}_0 \times \cdots \times \mathfrak{X}_n$. For a particularly readable proof of this result I refer to [156, Prop.V.1.1].

As a special case of this result let us mention an infinite-dimensional version of Theorem 1.3.41 (i): given measures μ_n on $\mathcal{F}_n$, there exists a unique measure μ_∞ on $\mathcal{F}^{\otimes\infty}$ such that

$$(\mathcal{P}_n)_\# \mu_\infty = \bigotimes_{k=0}^n \mu_k.$$

For this reason we will denote the measure μ_∞ by $\bigotimes_{n=0}^\infty \mu_n$. The resulting measurable maps $\pi_n : \mathfrak{X}^\infty \rightarrow \mathfrak{X}_n$ are independent and have the (prescribed) distribution μ_n . $\square$

1.6. Exercises

Exercise 1.1. Let $\mathcal{S}_0, \mathcal{S}_1$ be two sigma-algebras of a set Ω . Prove that the following are equivalent.

- (i) The union $\mathcal{S}_0 \cup \mathcal{S}_1$ is a sigma-algebra.
- (ii) Either $\mathcal{S}_0 \subset \mathcal{S}_1$ or $\mathcal{S}_1 \subset \mathcal{S}_0$.

□

Exercise 1.2. Construct a bijection $F : [-1, 1] \rightarrow \mathbb{R}$ such that both F and F^{-1} are Borel measurable.

□

Exercise 1.3. Fix a set Ω . Denote by $\mathbf{B}(\Omega)$ the space of bounded functions $\Omega \rightarrow \mathbb{R}$. Let $\mathfrak{F} \subset \mathbf{B}(\Omega)$ be a vector subspace with the following property: if (f_n) is a nondecreasing sequence of nonnegative functions in $\mathfrak{F}$ converging pointwisely to a function $f_\infty \in \mathbf{B}(\Omega)$, then $f_\infty \in \mathfrak{F}$. Suppose that $\mathcal{M} \subset \mathfrak{F}$ is a collection closed under multiplication. Then $\mathfrak{F}$ contains every bounded $\sigma(\mathcal{M})$ -measurable function.

□

Exercise 1.4. Let Ω denote the space $C([0, 1])$ of continuous functions $\omega : [0, 1] \rightarrow \mathbb{R}$ equipped with the topology defined by the sup-norm

$$\|\omega\| := \sup_{t \in \mathbb{T}} |\omega(t)|.$$

Denote by $\mathcal{B}$ the resulting Borel sigma-algebra. For each $t \in [0, 1]$ we have an *evaluation map*

$$E_t : \Omega \rightarrow \mathbb{R}, \quad E_t(\omega) = \omega_t.$$

Denote by $\mathcal{E}$ the sigma-algebra generated by the evaluation maps

$$\mathcal{E} := \bigvee_{t \in [0, 1]} E_t^{-1}(\mathcal{B}_{\mathbb{R}}).$$

Prove that $\mathcal{B} = \mathcal{E}$.

Hint. Prove first that

$$\|\omega\| = \sup_{t \in [0, 1] \cap \mathbb{Q}} |E_t(\omega)|, \quad \forall \omega \in C([0, 1]).$$

Use next the fact that the Banach space $C([0, 1])$ is separable.

□

Exercise 1.5. Suppose that (X, d) is a complete, separable metric space. Denote by $\mathcal{B}_X$ the Borel sigma-algebra generated by the open subsets of X , and by $\mathcal{B}_{X \times X}$ the Borel sigma-algebra generated by the product topology on $X \times X$. Prove that

$$\mathcal{B}_{X \times X} = \mathcal{B}_X \otimes \mathcal{B}_X.$$

□

Exercise 1.6. Fix a set Ω of finite cardinality m and a probability measure $\mathbb{P}$ on Ω . Assume that $\mathbb{P}[\{\omega\}] \neq 0, \forall \omega \in \Omega$. Set $\Omega^\infty := \Omega^{\mathbb{N}}$ so the elements of Ω^∞ are functions $\underline{\omega} : \mathbb{N} \rightarrow \Omega, n \mapsto \omega_n := \underline{\omega}(n)$. For every $n \in \mathbb{N}$ define

$$\pi_n : \Omega^\infty \rightarrow \Omega^n, \quad \pi_n(\underline{\omega}) = (\omega_1, \dots, \omega_n),$$

and denote by $\mathcal{C}_n$ the collection of sets of the form

$$C = \pi_n^{-1}(S), \quad S \subset \Omega^n, \quad n \in \mathbb{N}.$$

Note that $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \dots$. Set

$$\mathcal{C} := \bigcup_{n \in \mathbb{N}} \mathcal{C}_n.$$

The sets in $\mathcal{C}$ are called *cylinders*.

- (i) Show that $\mathcal{C}_n$ is a σ -algebra of subsets of Ω^∞ , $\forall n \in \mathbb{N}$.
- (ii) For any $n \in \mathbb{N}$ define $\beta_n = \beta_{n,\pi} : \mathcal{C}_n \rightarrow [0, 1]$,

$$\beta_n[\pi_n^{-1}(S)] := \pi^{\otimes n}[S] = \sum_{(\omega_1, \dots, \omega_n) \in S} \prod_{j=1}^n \mathbb{P}[\{\omega_j\}].$$

Show that β_n is a well defined measure on $\mathcal{C}_n$ and

$$\beta_{n+1}|_{\mathcal{C}_n} = \beta_n.$$

- (iii) Equip Ω^∞ with the metric

$$d(\underline{\omega}, \underline{\eta}) = \sum_{n \in \mathbb{N}} \frac{1}{2^n} h(\omega_n, \eta_n), \quad h(\omega, \eta) = \begin{cases} 0, & \omega = \eta, \\ 1, & \omega \neq \eta. \end{cases}$$

Prove that (Ω^∞, d) is a compact metric space. **Hint.** Use the diagonal procedure to show that any sequence if Ω admits a convergent subsequence.

- (iv) Define $\beta = \beta_{\mathbb{P}} : \mathcal{C} \rightarrow [0, 1]$,

$$\beta|_{\mathcal{C}_n} = \beta_n.$$

Show that β is a well defined premeasure on $\mathcal{C}$. **Hint.** Use Theorem 1.2.15. Prove first that any cylinder is simultaneously closed and open.

- (v) Prove that $\sigma(\mathcal{C})$ coincides with the Borel sigma-algebra of the metric space (Ω^∞, d) .
- (vi) Denote by $\bar{\beta} = \bar{\beta}_{\mathbb{P}}$ the extension of β as measure to the σ -algebra $\sigma(\mathcal{C})$. (Its existence is guaranteed by the Caratheodory extension theorem.) For $\omega_0 \in \Omega$ we set

$$\Delta_{\omega_0} := \{ \underline{\omega} \in \Omega : \exists m \in \mathbb{N} \text{ such that } \omega_n = \omega_0, \forall n > m \}.$$

Show that $\Delta_{\omega_0} \in \sigma(\mathcal{C})$ and $\bar{\beta}[\Delta_{\omega_0}] = 0$.

- (vii) Define $X_n : \Omega^\infty \rightarrow \Omega$, $X_n(\underline{\omega}) = \omega_n$. Show that the collection of random variables $(X_n)_{n \in \mathbb{N}}$ is independent and have the same distribution $\mathbb{P}$.
- (viii) Let $\Omega = \{0, 1\}$, $\mathbb{P}$ = the uniform measure on $\{0, 1\}$, and consider $\Omega^\infty = \{0, 1\}^{\mathbb{N}}$ equipped with the measure $\bar{\beta} = \bar{\beta}_{\mathbb{P}}$ constructed as above. Show that the map

$$B : (\Omega^\infty, \sigma(\mathcal{C})) \rightarrow ([0, 1], \mathcal{B}_{[0,1]}), \quad B = \sum_{n \in \mathbb{N}} \frac{1}{2^n} X_n$$

is measurable and find $B_{\#} \bar{\beta}$. □

Exercise 1.7. Suppose that $(\Omega, \mathcal{F}, \mu)$ is a measured space and (S, d) a metric space. Consider a function

$$F : S \times \Omega \rightarrow \mathbb{R}, \quad (s, \omega) \mapsto F_s(\omega)$$

satisfying the following properties.

- (i) For any $s \in S$ the function $\Omega \ni \omega \mapsto F_s(\omega) \in \mathbb{R}$ is measurable.
- (ii) For any $\omega \in \Omega$ the function $S \ni s \mapsto F_s(\omega) \in \mathbb{R}$ is continuous.

(iii) There exists $h \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ such that $|F_s(\omega)| \leq h(\omega)$, $\forall (s, \omega) \in S \times \Omega$.

Prove that $F_s \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$, $\forall s \in S$, and the resulting function

$$S \ni s \mapsto \int_{\Omega} F_s(\omega) \mu[d\omega] \in \mathbb{R}$$

is continuous. **Hint.** Use the Dominated Convergence Theorem. $\square$

Exercise 1.8. Suppose that $(\Omega, \mathcal{F}, \mu)$ is a measured space and $I \subset \mathbb{R}$ is an open interval. Consider a function

$$F : I \times \Omega \rightarrow \mathbb{R}, \quad (t, \omega) \mapsto F(t, \omega)$$

satisfying the following properties.

(i) For any $t \in I$ the function $F(t, -) : \Omega \rightarrow \mathbb{R}$ is integrable,

$$\int_{\Omega} |F(t, \omega)| \mu[d\omega] < \infty.$$

(ii) For any $\omega \in \Omega$ the function $I \ni t \mapsto F(t, \omega) \in \mathbb{R}$ is differentiable at $t_0 \in I$. We denote by $F'(t_0, \omega)$ its derivative.

(iii) There exists $h \in \mathcal{L}^1(\Omega, \mathcal{S}, \mu)$ and $c > 0$ such that

$$|F(t, \omega) - F(t_0, \omega)| \leq h(\omega)|t - t_0|, \quad \forall (t, \omega) \in I \times \Omega.$$

Prove that the function

$$I \ni t \mapsto \int_{\Omega} F(t, \omega) \mu[d\omega] \in \mathbb{R}$$

is differentiable at t_0 and

$$\left. \frac{d}{dt} \right|_{t=t_0} \left(\int_{\Omega} F(t, \omega) \mu[d\omega] \right) = \int_{\Omega} F'(t_0, \omega) \mu[d\omega]. \quad \square$$

Exercise 1.9. Suppose that $(\Omega, \mathcal{S}, \mu)$ is a *finite*¹⁵ measured space and $\mathcal{A} \subset \mathcal{S}$ a countable π -system that generates $\mathcal{S}$, $\sigma(\mathcal{A}) = \mathcal{S}$. Assume $\Omega \in \mathcal{A}$. Denote by $\mathbb{R}[\mathcal{A}]$ the vector space spanned by $\mathbf{1}_A$, $A \in \mathcal{A}$. Fix $p \in [1, \infty)$ and denote by $\mathcal{M}_p$ the intersection of $L^\infty(\Omega, \mathcal{S}, \mu)$ with the L^p -closure of $\mathbb{R}[\mathcal{A}]$.

(i) Prove $\mathcal{M}_p = L^\infty(\Omega, \mathcal{S}, \mu)$.

(ii) Prove that $\mathbb{R}[\mathcal{A}]$ is dense in $L^p(\Omega, \mathcal{S}, \mu)$.

(iii) Prove that $L^p(\Omega, \mathcal{S}, \mu)$ is separable. $\square$

Exercise 1.10 (Markov). Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a sample space and A_-, A_0, A_+ , $\mathbb{P}[A_0 \cap A_-] \neq 0$. We say that A_+ is *independent of* A_- given A_0 if

$$\mathbb{P}[A_+ \cap A_- | A_0] = \mathbb{P}[A_+ | A_0] \mathbb{P}[A_- | A_0].$$

Show that A_+ is independent of A_- given A_0 if and only if

$$\mathbb{P}[A_+ | A_0 \cap A_-] = \mathbb{P}[A_+ | A_0]. \quad \square$$

Exercise 1.11. Prove the equality (??). $\square$

¹⁵The sigma-finite situation follows from the finite situation in a standard fashion.

Exercise 1.12 (M. Gardner). A family has two children. Find the conditional probability that both children are boys in each of the following situations.

- (i) One of the children is a boy.
- (ii) One of the children is a boy born on a Thursday. □

Exercise 1.13. A random experiment is performed repeatedly and the outcome of an experiment is independent of the outcomes of the previous experiments. While performing these experiments we keep track of the occurrence of the mutually exclusive events A and B , i.e., $A \cap B = \emptyset$. We assume that A and B have positive probabilities.¹⁶ What is the probability that A occurs before B ? **Hint.** Consider the event $C = (A \cup B)^c$ = neither A , nor B . Condition on the result of the first experiment which can be A , B or C . □

Exercise 1.14. Prove that the random variables $N_1, \dots, N_m$ that appear in Example 1.3.28 on the coupon collector problem can be realized as measurable functions defined on the same probability space. **Hint.** Use Exercise 1.6. □

Exercise 1.15. Construct a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and random variables

$$X, Y : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (0, \infty)$$

such that

$$\mathbb{E}[X] < \mathbb{E}[Y] < \infty \text{ and } \mathbb{P}[X > Y] < \mathbb{P}[X < Y]. \quad \square$$

Exercise 1.16 (Your neighbor has more neighbors).¹⁷ Consider a connected finite unoriented graph. Denote by $\mathcal{V}$ its set of vertices and by $\mathcal{E}$ the set of edges. For each $v \in V$ we denote by $\deg(v)$ the number of neighbors of v . Pix a vertex $A \in \mathcal{V}$ uniformly randomly, and then choose a neighbor B of A , equally likely among the $\deg(A)$ neighbors of A . Prove that

$$\mathbb{E}[\deg(A)] \leq \mathbb{E}[\deg(B)]. \quad \square$$

Exercise 1.17. Suppose that $X, Y : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ are two random variables whose ranges $\mathfrak{X}$ and $\mathfrak{Y}$ are countable subsets of $\mathbb{R}$. Assume additionally that $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$. We set

$$\mathbb{E}[X \parallel Y] = \sum_{y \in \mathfrak{Y}} \mathbb{E}[X \mid Y = y] \mathbf{I}_{\{Y=y\}} \in \mathcal{L}^0(\Omega, \sigma(Y), \mathbb{P}),$$

where

$$\mathbb{E}[X \mid Y = y] := \sum_{x \in \mathfrak{X}} x \mathbb{P}[X = x \mid Y = y] = \frac{1}{\mathbb{P}[\{Y = y\}]} \int_{\{Y=y\}} X(\omega) \mathbb{P}[d\omega].$$

The random variable $\mathbb{E}[X \parallel Y]$ is called the *conditional expectation of X given Y* . Prove that

$$\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X \parallel Y]]. \quad \square$$

¹⁶For example if we roll a pair of dice, A could be the event “the sum is 4” and B could be the event “the sum is 7”. In this case

$$\mathbb{P}[A] = \frac{3}{36} = \frac{1}{12}, \quad \mathbb{P}[B] = \frac{6}{36} = \frac{1}{6}.$$

¹⁷This fact was first observed empirically in 1991 by Scott L. Feld, a sociologist.

Exercise 1.18 (Polya's urn). An urn U contains r_0 red balls and g_0 green balls. At each stage a ball is selected at random from the urn, we observe its color, we return it to the urn and then we add another ball of the same color. We denote by R_n the number of red balls and by G_n the number of green balls at stage n . Finally, we denote by C_n the “concentration” of red balls at stage n ,

$$C_n = \frac{R_n}{R_n + G_n}.$$

- (i) Show that $\mathbb{E}[C_{n+1} \mid R_n] = C_n$, where the conditional expectation $\mathbb{E}[C_{n+1} \mid R_n]$ is defined in Exercise 1.17.
- (ii) Show that $\mathbb{E}[C_n] = \frac{r_0}{r_0 + g_0}, \forall n \in \mathbb{N}$.

□

Exercise 1.19. Prove the claim about the events S_k at the end of Example 1.3.26.

□

Exercise 1.20 (Banach's matchbox problem). An eminent mathematician fuels a smoking habit by keeping matches in both trouser pockets. When impelled by need, he reaches a hand into a randomly selected pocket and grabs about for a match. Suppose he starts with n matches in each pocket. What is the probability that when he first discovers a pocket to be empty of matches the other pocket contains exactly m matches?

□

Exercise 1.21. Suppose that $X_n \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$, $n \in \mathbb{N}$, is a sequence of independent and identically distributed (i.i.d.) random variables and $T \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ is a random variable with range contained in $\mathbb{N}$ and independent of the variables X_n . Define $S_T : \Omega \rightarrow \mathbb{R}$

$$S_T(\omega) = \sum_{n=1}^{T(\omega)} X_n(\omega).$$

Prove Wald's formula

$$\mathbb{E}[S_T] = \mathbb{E}[T] \mathbb{E}[X_1]. \quad (1.6.1)$$

□

Exercise 1.22. A box contains n identical balls labelled $1, \dots, n$. Draw one ball, uniformly random, and record its label N . Next flip a fair coin N times. What is the expected number of heads you roll?

Hint. Use Wald's formula.

□

Exercise 1.23. Suppose that $X \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ is a *nonnegative* random variable. Prove that if the range of $\mathfrak{X}$ is contained in $\mathbb{N}_0$, then

$$\mathbb{E}[X] - 1 \leq \sum_{n \geq 0} \mathbb{P}[X > n] \leq \mathbb{E}[X].$$

In particular, conclude that

$$X \in L^1(\Omega, \mathcal{S}, \mathbb{P}) \iff \sum_{n \geq 0} \mathbb{P}[X > n] < \infty.$$

Hint. Use (1.3.50)

□

Exercise 1.24. Let X be a random variable with range contained in $\{0, 1, \dots, n\}$.

- (i) Prove that for any $k \in \{0, \dots, n\}$

$$\mathbb{P}[X = k] = \sum_{j=0}^{n-k} (-1)^j \binom{k+j}{j} \mathbb{E}[B_{k+j}(X)],$$

where,

$$B_m(x) = \binom{x}{m} := \frac{1}{m!} x(x-1) \cdots (x-m+1) \in \mathbb{R}[x].$$

Hint. Set $p_k = \mathbb{P}[X = k]$. Then $\mathbb{E}[B_m(X)] = \sum_{k=m}^n \binom{k}{m} p_k$. Conclude using the binomial inversion trick, Remark 1.3.33.

- (ii) Let $A_1, \dots, A_n$ be a collection of measurable subsets of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and set

$$X := \sum_{k=1}^n \mathbf{I}_{A_k}.$$

Prove that

$$\mathbb{E}[B_k(X)] = s_k^n$$

for any $k \in \{0, 1, \dots, n\}$, where s_k^n is defined in (1.3.27). **Hint.** Use binomial inversion.

□

Exercise 1.25. Consider the standard random walk on $\mathbb{Z}$ started at 0. More precisely, are given a sequence of i.i.d random variables $(X_n)_{n \in \mathbb{N}}$ such that $\mathbb{P}[X_n = 1] = \mathbb{P}[X_n = -1] = \frac{1}{2}$, $\forall n$ and we set

$$S_n := X_1 + \cdots + X_n.$$

Let T denote the time of the first return to 0,

$$T := \min\{n \in \mathbb{N}; S_n = 0\},$$

where $\min \emptyset := \infty$. Set $f_n = \mathbb{P}[T = n]$, $u_n := \mathbb{P}[S_n = 0]$.

- (i) Prove that $u_{2n} = \mathbb{P}[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$. Deduce that $f_{2n} = u_{2n-2} - u_{2n}$.
Hint. Use André's reflection trick in Example 1.2.38.
- (ii) Prove that $\mathbb{P}[T < \infty] = 1$, but $\mathbb{E}[T] = \infty$. **Hint.** Use (i) and (1.3.50).
- (iii) Visualize the random walk as a zig-zag of the kind depicted in Figure 1.2. For such a zigzag we denote by $L_n(z)$ the number of its first n segments that are above the x axis. Equivalently,

$$L_n(z) := \#\{k; 1 \leq k \leq n; \max(S_{k-1}, S_k) > 0\}.$$

For example, for the zig-zag z in Figure 1.2 we have

$$L_8(z) = L_9(z) = L_{10}(z) = 8.$$

Show that

$$\mathbb{P}[L_{2n} = m] = \begin{cases} u_{2k} u_{2n-2k}, & m = 2k \leq 2n, \\ 0, & m \equiv 1 \pmod{2}. \end{cases}$$

- (iv) Prove that $\mathbb{P}[L_{2n} = 2k | S_{2n} = 0] = \frac{1}{n+1}$.

□

Exercise 1.26. Consider the group $\mathfrak{S}_n$ of permutation of n ordered objects. We equip it with the uniform probability measure. Let Z_n denote the number of inversions of a random permutation, i.e.,

$$Z_n(\varphi) := \#\{ (i, j); \ 1 \leq i < j \leq n, \ \varphi(i) > \varphi(j) \}.$$

Compute the mean and variance of Z_n . **Hint.** For $\varphi \in \mathfrak{S}_n$ and $1 \leq k \leq n-1$ set

$$Z_{n,k}(\varphi) = \#\{ j; \ n \geq j > k \text{ and } \varphi(j) < \varphi(k) \}.$$

Prove that $Z_n = \sum_{k=1}^{n-1} Z_{n,k}$ and that the random variables $Z_{n,1}, \dots, Z_{n,n-1}$ are independent. You will also need to use the classical identities

$$1 + 2 + \dots + m = \frac{m(m+1)}{2}, \quad 1^2 + 2^2 + \dots + m^2 = \frac{m(m+1)(2m+1)}{6}.$$

□

Exercise 1.27. There are n unstable molecules $m_1, \dots, m_n$ in a row. One of the $n-1$ pairs of neighbors, chosen uniformly at random, combine to form a stable dimer. This process continues until there remain U_n isolated molecules, no two of which are adjacent.

(i) Show that the probability p_n that m_1 remains uncombined satisfies

$$(n-1)p_n = p_1 + p_2 + \dots + p_{n-2}.$$

Deduce that

$$p_n = \sum_{k=0}^{n-1} \frac{(-1)^k}{k!} \rightarrow e^{-1} \text{ as } n \rightarrow \infty.$$

Hint. Condition on the first pair of molecules (m_r, m_{r+1}) that gets combined.

(ii) Show that the probability $q_{r,n}$ that the molecule m_r remains uncombined is $p_r p_{n-r+1}$.

(iii) Show that

$$\mathbb{E}[U_n] = \sum_{r=1}^n q_{r,n}.$$

(iv) Show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[U_n] = e^{-2}.$$

Hint. At some point you may need to take for granted the result in Exercise 2.6.

□

Exercise 1.28. Let $N = N_m$ be the random variable defined in the coupon collector problem described in Example 1.3.28. Show that

$$\text{Var}[N_m] = m \sum_{k=1}^m \frac{m-k}{k^2}.$$

□

Exercise 1.29 (The Birthday Problem). Let $N \in \mathbb{N}$. Consider a sequence $(X_n)_{n \in \mathbb{N}}$ of independent random variables uniformly distributed on the finite set $\{1, \dots, N\}$. Define B_N to be the *birthday random variable*¹⁸

$$B_N(\omega) = \min \{ j \in \mathbb{N} : \exists 1 \leq i < j \text{ such that } X_j(\omega) = X_i(\omega) \}.$$

¹⁸You should think of B_N as follows. Suppose that you have an urn with N balls labelled $1, \dots, N$. Suppose we perform the following experiment: draw a ball at random, record its label, put it back in the box, and then repeat until you notice that the label you've drawn has appeared before. The random variable B_N is the first moment when you've noticed a label that was drawn before. Note that $B_N \leq N+1$. The classical birthday problem is the special case $N = 365$.

Compute the probabilities

$$\mathbb{P}[B_N \leq k], \quad k = 1, \dots, N. \quad \square$$

Exercise 1.30 (Buffon's Problem). A needle of length ℓ is thrown at random on a plane ruled by parallel lines distance d apart. Denote by N_ℓ the number of lines that intersect the needle.

- (i) Compute $\mathbb{P}[N_\ell = 1]$ when $\ell \leq d$.
- (ii) Prove that $\mathbb{E}[N_{\ell_0 + \ell_1}] = \mathbb{E}[N_{\ell_0}] + \mathbb{E}[N_{\ell_1}]$, $\forall \ell_0, \ell_1 > 0$.
- (iii) Compute $\mathbb{E}[N_\ell]$, $\ell > 0$.

□

Exercise 1.31. Suppose that I is an interval of the real axis and $f : I \rightarrow \mathbb{R}$ is a continuous function. Prove that the following are equivalent.

- (i) For any $x, y \in I$, and any $t \in (0, 1)$ we have $f((1-t)x + ty) \leq (1-t)f(x) + tf(y)$.
- (ii) For any $x_0 \in I$ there exists a linear function $\ell : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\ell(x_0) = f(x_0), \quad \ell(x) \leq f(x), \quad \forall x \in I.$$

□

Exercise 1.32. Suppose that $X \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ is an integrable random variable and consider the function

$$\varphi : \mathbb{R} \rightarrow [0, \infty), \quad \varphi(x) = \mathbb{E}[|X - x|].$$

Prove that if m is a median of X (see Definition 1.2.23) then

$$\varphi(m) \leq \varphi(x), \quad \forall x \in \mathbb{R}.$$

□

Exercise 1.33 (Hermite polynomials). Suppose that $X \sim N(0, 1)$ so

$$\mathbb{P}_X[dx] = \Gamma_1[dx] := \gamma_1(x)\lambda[dx], \quad \gamma_1(x) = \frac{1}{\sqrt{2\pi}}e^{-\frac{x^2}{2}}.$$

For $k \in \mathbb{N}_0$ we denote by $\mathbb{R}[x]$ the space of polynomial with real coefficients. Define the linear operators

$$\begin{aligned} P, Q : \mathbb{R}[x] &\rightarrow \mathbb{R}[x], \\ (Pf)(x) &= f'(x), \quad (Qf)(x) = -f'(x) + xf(x). \end{aligned} \quad (1.6.2)$$

The operator P is called the *annihilation* operator and the operator Q is called the *creation* operator.

- (i) Prove that for any $f \in \mathbb{R}[x]$ we have

$$(PQ - QP)f = f.$$

- (ii) Denote by $H_0 \in \mathbb{R}[x]$ the constant polynomial identically equal to 1. Show that for any $n \in \mathbb{N}$ the function

$$H_n := Q^n H_0$$

is a degree n polynomial satisfying

$$PH_n = nH_{n-1}, \quad QPH_n = nH_n, \quad \forall n \in \mathbb{N},$$

and

$$H_n = xH(x)_{n-1} - (n-1)H_{n-2}(x), \quad \forall n \geq 2.$$

Deduce that

$$H_n(x) = n! \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^r}{2^r r! (n-2r)!} x^{n-2r}. \quad (1.6.3)$$

The polynomials $H_n(x)$ are called the *Hermite polynomials*. The operator $-QP$ is called the *Ornstein-Uhlenbeck operator*.

(iii) Show that for any $f, g \in \mathbb{R}[x]$

$$\int_{\mathbb{R}} Pf(x)g(x) \Gamma_1[dx] = \int_{\mathbb{R}} f(x)Qg(x) \Gamma_1[dx]. \quad (1.6.4)$$

(iv) Show that

$$H_n(x) = (-1)^n e^{\frac{x^2}{2}} P^n(e^{-\frac{x^2}{2}}) = (-1)^n e^{\frac{x^2}{2}} \frac{d^n}{dx^n} (e^{-\frac{x^2}{2}}), \quad \forall n \in \mathbb{N}. \quad (1.6.5)$$

(v) Show that for any $m, n \in \mathbb{N}_0$ we have

$$\int_{\mathbb{R}} H_n(x)H_m(x) \Gamma_1[dx] = n! \delta_{mn}.$$

(vi) Show that

$$\sum_{n \geq 0} H_n(x) \frac{\lambda^n}{n!} = e^{\lambda x - \lambda^2/2}. \quad (1.6.6)$$

(vii) Suppose that $f \in \mathbb{R}[x]$, $\deg f \leq n$. Recall that $X \sim N(0, 1)$. Prove that

$$f(x) = \sum_{k=1}^n \frac{1}{k!} \mathbb{E}[f^{(k)}(X)] H_k(x). \quad (1.6.7)$$

□

Exercise 1.34. Suppose that $X \sim N(0, 1)$, i.e.,

$$\mathbb{P}_X[dx] = \gamma_1(x)dx, \quad \gamma_1(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}.$$

Set $\bar{\Phi}(x) := \mathbb{P}[X > x]$. Prove the *Mills ratio inequalities* (1.3.44), i.e.,

$$\frac{x}{x^2 + 1} \gamma_1(x) \leq \bar{\Phi}(x) \leq \frac{1}{x} \gamma_1(x), \quad \forall x > 0.$$

Hint. For the upper bound observe that

$$-Q\bar{\Phi} = \int_x^\infty \bar{\Phi}(t)dt > 0,$$

where Q is the operator defined in (1.6.2). Next express

$$\int_x^\infty Q\bar{\Phi}(t)dt \leq 0$$

in terms of $\bar{\Phi}$ and γ_1 .

□

Exercise 1.35. We denote by $\text{Dens}(\mathbb{R})$ the space of probability densities on $\mathbb{R}$, i.e., functions $p \in L^1(\mathbb{R}, \lambda)$ such that

$$\int_{\mathbb{R}} p(x) dx = 1 \quad \text{and} \quad p(x) \geq 0 \quad \text{almost everywhere.}$$

For $p \in \text{Dens}(\mathbb{R})$ we set

$$\mathbb{E}[p] := \int_{\mathbb{R}} xp(x) dx, \quad \text{Var}[p] := \int_{\mathbb{R}} x^2 p(x) dx - \mathbb{E}[p]^2.$$

The *entropy*¹⁹ of $p \in \text{Dens}(\mathbb{R})$ is the quantity

$$\text{Ent}[p] := - \int_{\mathbb{R}} p(x) \log p(x) dx \in [0, \infty],$$

where we set $0 \cdot \log 0 = 0$.

(i) Show that if

$$\gamma_1(x) := \frac{1}{\sqrt{2\pi}} e^{-x^2/2},$$

then

$$\text{Ent}[\gamma_1] = \frac{1 + \log 2\pi}{2}.$$

(ii) Show that if $p, q \in \text{Dens}(\mathbb{R})$ and $q(x) > 0, \forall x \in \mathbb{R}$, then

$$\text{Ent}[p] \leq - \int_{\mathbb{R}} p(x) \log q(x) dx$$

if the integral on the right hand side is finite. Moreover equality holds iff $p = q$.

Hint. Show that $p(x) - p(x) \log p(x) \leq q(x) - p(x) \log q(x), \forall x \in \mathbb{R}$.

(iii) Show that if $p \in \text{Dens}(\mathbb{R})$ satisfies

$$\mathbb{E}[p] = 0 = \mathbb{E}[\gamma_1], \quad \text{Var}[p] = 1 = \text{Var}[\gamma_1],$$

then $\text{Ent}[p] \leq \text{Ent}[\gamma_1]$ with equality iff $p = \gamma_1$.

□

Exercise 1.36. Let $X : (\Omega, \mathcal{S}, \mathbb{P} \rightarrow \mathbb{N}_0)$ be a random variable and $\lambda > 0$. Prove that the following are equivalent.

(i) $X \sim \text{Poi}(\lambda)$.

(ii) $\mathbb{E}[\lambda f(X+1) - X f(X)] = 0$, for any bounded function $f : \mathbb{N}_0 \rightarrow \mathbb{R}$.

□

Exercise 1.37. Prove Proposition 1.3.20.

□

¹⁹The entropy is a measure of disorder or randomness of the probability density: the higher the entropy the less predictable is the associated random variable.

Exercise 1.38. Show that

$$\mathbb{M}_N(t) = \frac{pe^t}{1 - qe^t} \quad \text{if } N \sim \text{Geom}(p),$$

$$\mathbb{M}_N(t) = e^{\lambda(e^t - 1)} \quad \text{if } N \sim \text{Poi}(\lambda),$$

and

$$\mathbb{M}_X(t) = \frac{\lambda}{\lambda - t} \quad \text{if } X \sim \text{Exp}(\lambda).$$

□

Exercise 1.39. Let $Y \sim N(0, 1)$ be a standard normal random variable and set $X := \exp(Y)$.

(i) Show that

$$\mathbb{E}[X^n] = e^{n^2/2}, \quad \forall n \in \mathbb{N}.$$

(ii) Prove that the probability distribution $\mathbb{P}_X$ of X is given by the *log-normal* law

$$\mathbb{P}_X[dx] = p(x)dx, \quad p(x) = \begin{cases} \frac{1}{x\sqrt{2\pi}} e^{-\frac{1}{2}(\log x)^2}, & x > 0, \\ 0, & x \leq 0, \end{cases}$$

where $\log$ denotes the natural logarithm.

(iii) For $\alpha \in [-1, 1]$ we set

$$p_\alpha(x) = \begin{cases} p(x)(1 + \alpha \sin(2\pi \log x)), & x > 0, \\ 0, & x \leq 0. \end{cases}$$

Prove that for any $\alpha \in [-1, 1]$ and any $n \in \mathbb{N}_0$ we have

$$\int_{\mathbb{R}} x^n p_\alpha(x) dx = e^{n^2/2}.$$

Thus, for any $\alpha \in [-1, 1]$, the function $p_\alpha(x)dx$ is a probability density on $\mathbb{R}$ and the probability measure $p_\alpha(x)$ has the same moments as X ,

□

Exercise 1.40. Let $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ be a random variable with range contained in

$$\mathbb{N}_0 = \{0, 1, 2, \dots\}.$$

Its *probability generating function* (or *pgf* for brevity) is the formal power series

$$PG_X(s) = \sum_{n \geq 0} \mathbb{P}[X = n] s^n.$$

(i) Show that the power series defining PG_X is convergent for any $|s| < 1$. Moreover, $\forall t \leq 0$ we have

$$\mathbb{M}_X(t) = PG_X(e^t).$$

(ii) Compute PG_X when $X \sim \text{Bin}(n, p)$, $X \sim \text{Geom}(p)$, $X \sim \text{Poi}(\lambda)$.

□

Exercise 1.41. Show that

$$\text{Gamma}(\nu_0, \lambda) * \text{Gamma}(\nu_1, \lambda) = \text{Gamma}(\nu_0 + \nu_1, \lambda), \quad \forall \nu_0, \nu_1 > 0, \quad (1.6.8a)$$

$$N(0, v_0) * N(0, v_1) = N(0, v_0 + v_1), \quad \forall v_0, v_1 > 0, \quad (1.6.8b)$$

$$\text{Poi}(\lambda_0) * \text{Poi}(\lambda_1) = \text{Poi}(\lambda_0 + \lambda_1), \quad \forall \lambda_0, \lambda_1 > 0. \quad (1.6.8c)$$

Hint. Use Theorem 1.3.23, Corollary 1.3.21 and Corollary 1.3.51. $\square$

Exercise 1.42. Let $\mu_0, \mu_1 \in \text{Prob}([0, 1])$ be two Borel probability measures. Prove that the following statements are equivalent.

(i)

$$\int_0^1 x^n \mu_0[dx] = \int_0^1 x^n \mu_1[dx], \quad \forall n \in \mathbb{N}.$$

(ii) For any Borel subset $B \subset [0, 1]$, $\mu_0[B] = \mu_1[B]$.

$\square$

Exercise 1.43. Suppose that μ is a Borel probability measure on $\mathbb{R}_{\geq 0}$. We define its *Laplace transform* to be the function

$$\mathcal{L}_\mu : [0, \infty) \rightarrow \mathbb{R}, \quad \mathcal{L}_\mu(\lambda) = \int_{\mathbb{R}_{\geq 0}} e^{-\lambda x} \mu[dx].$$

Consider the measurable map $F : [0, \infty) \rightarrow [0, 1]$, $F(x) = e^{-x}$. We set $\tilde{\mu} := F_\# \mu$.

(i) Prove that

$$\mathcal{L}_\mu(\lambda) = \int_{[0, 1]} y^\lambda \tilde{\mu}[dy].$$

(ii) Prove that the measure μ is uniquely determined by its Laplace transform $\mathcal{L}_\mu$.

$\square$

Exercise 1.44. Denote by $\text{Prob} = \text{Prob}(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ the space of probability measures on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$. Show that $(\text{Prob}, *)$ is a commutative semigroup with unit δ_0 , the Dirac measure concentrated at 0. $\square$

Exercise 1.45. Consider the interval $[-\pi/2, \pi/2]$ equipped with the probability measure

$$\mathbb{P}[dx] = \frac{1}{\pi} \lambda[dx],$$

λ = the Lebesgue measure. We regard the function

$$X : [-\pi/2, \pi/2] \rightarrow \mathbb{R}, \quad X(t) = \sin^2 t$$

as a random variable on this probability spaces. Prove that $X \sim \text{Beta}(1/2, 1/2)$. $\square$

Exercise 1.46. For any $a, b > 0$ we define the *incomplete Beta function*

$$B_{a,b} : (0, 1) \rightarrow \mathbb{R}, \quad B_{a,b}(x) = \frac{1}{B(a, b)} \int_0^x t^{a-1} (1-t)^{b-1} dt,$$

where $B(a, b)$ is the Beta function (A.1.2).

(i) Prove that

$$\frac{x^a(1-x)^b}{aB(a,b)} = B_{a,b}(x) - B_{a+1,b}(x). \quad (1.6.9a)$$

$$\frac{x^a(1-x)^b}{bB(a,b)} = B_{a,b+1}(x) - B_{a,b}(x). \quad (1.6.9b)$$

(ii) Show that if $k, n \in \mathbb{N}$, $k < n$ we have

$$B_{k,n+1-k}(x) = \sum_{a=k}^n \binom{n}{a} x^a (1-x)^{n-a}. \quad (1.6.10)$$

□

Exercise 1.47. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent Bernoulli variables, $X_n \sim \text{Ber}(p_n)$, $0 \leq p_n < 1$. As in Example 1.3.58, this sequence defines a random subset of $\mathbb{N}$,

$$S := \{n \in \mathbb{N}; X_n = 1\}.$$

Suppose that there exists a function $f : (C, \infty) \rightarrow (0, \infty)$, $C > 0$, such that $\log f$ is convex

$$\lim_{n \rightarrow \infty} \frac{nf(n)^2}{\log n} = \infty,$$

and

$$p_n > f(n), \quad \forall n \in \mathbb{N} \cap (C, \infty).$$

Denote by T_n the event

$$T_n := \{n \notin S + S\}, \quad S + S = \{j + k; j, k \in S\}.$$

Prove that

$$\mathbb{P}[T_n \text{ i.o.}] = 0.$$

so that

$$\mathbb{P}[\exists N \in \mathbb{N}; \forall n \geq N : n \in S + S] = 1.$$

Hint. Prove that for any $k > 1$, $\mathbb{P}[T_n] = O(n^{-k})$ as $n \rightarrow \infty$.

□

Exercise 1.48. Suppose that $X_1, \dots, X_n : (\Omega, \mathcal{S}, \mu) \rightarrow \mathbb{R}$ are random variables with joint probability distribution

$$\begin{aligned} \mathbb{P}_{X_1, \dots, X_n} [dx_1 \cdots dx_n] &= p(x_1, \dots, x_n) dx_1 \cdots dx_n, \\ p &\geq 0, \quad \int_{\mathbb{R}^n} p(x_1, \dots, x_n) dx_1 \cdots dx_n = 1. \end{aligned}$$

Consider the new random variables

$$Y_i = \sum_{k=1}^n a_{ik} X_k, \quad a_{ik} \in \mathbb{R}$$

where the matrix $A = (a_{ij})_{1 \leq i, j \leq n}$ is invertible with inverse $A^{-1} = (a^{ij})_{1 \leq i, j \leq n}$. Prove that the joint distribution of $Y_1, \dots, Y_n$ is given by the density

$$q(y_1, \dots, y_n) = \frac{1}{|\det A|} p(a^{11}y_1 + \cdots + a^{1n}y_n, \dots, a^{n1}y_1 + \cdots + a^{nn}y_n). \quad \square$$

Exercise 1.49. Suppose that $X_1, \dots, X_N$ are independent standard normal random variables. For $n = 1, \dots, N$, we denote by R_n^2 the random variable $X_1^2 + \cdots + X_n^2$.

(i) Prove that

$$R_n^2 \sim \chi^2(n) := \text{Gamma}(\nu, \lambda), \quad \nu = \frac{n}{2}, \quad \lambda = \frac{1}{2}.$$

(ii) Prove that

$$\frac{R_n^2}{R_N^2} \sim \text{Beta}(a, b), \quad \text{where } a = \frac{n}{2}, \quad b = \frac{N-n}{2}.$$

□

Exercise 1.50. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Show that $L^0(\Omega, \mathcal{S}, \mathbb{P})$ equipped with the metric dist defined in (1.3.60) is a complete metric space. More precisely, show that if a sequence of random variables $X_n \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ is Cauchy in probability, i.e.,

$$\lim_{m, n \rightarrow \infty} \mathbb{P}[|X_m - X_n| > r] = 0, \quad \forall r > 0,$$

then there exists a random variable $X \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ such that $X_n \rightarrow X$ in probability. □

Exercise 1.51. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Prove that if a sequence of random variables $X_n \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ converges a.s. to a random variable $X \in L^0(\Omega, \mathcal{S}, \mathbb{P})$ iff it satisfies

$$\lim_{m, n \rightarrow \infty} \mathbb{P}\left[\sup_{m < k \leq n} |X_k - X_m| > r\right] = 0, \quad \forall r > 0.$$

□

Exercise 1.52. Prove the claim in Remark 1.4.4. □

Exercise 1.53. Suppose that X, Y are independent random variables with distributions $\mathbb{P}_X$ and respectively $\mathbb{P}_Y$. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a Borel measurable function such that $f(X, Y)$ is integrable. Show that

$$\mathbb{E}[f(X, Y) | X] = h(X),$$

where

$$h(x) = \int_{\mathbb{R}} f(x, y) \mathbb{P}_Y[dy].$$

□

Exercise 1.54. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space, $\mathcal{F} \subset \mathcal{S}$ a sigma-subalgebra and $X \in \mathcal{L}^0(\Omega, \mathcal{S})$, $Y \in \mathcal{L}^0(\Omega, \mathcal{F})$. Prove that the following are equivalent.

(i) $X = Y$ a.s..

(ii) For any bounded Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$, $\mathbb{E}[f(X) | \mathcal{F}] = f(Y)$ a.s.

□

Exercise 1.55. Suppose that the sequence of independent random variables $(X_n)_{n \in \mathbb{N}}$ converges in probability to a random variable X . Prove that X is a.s. constant. **Hint.** Use Kolmogorov's zero-one theorem. □

Exercise 1.56 (Strong memoryless property). Suppose that T is an exponential random variable and $T_0, S \geq 0$ are nonnegative random variables so that T, T_0, S are pairwise independent. Then

$$\mathbb{P}[T > T_0 + S | T > S] = \mathbb{P}[T > T_0].$$

Note that when T_0, S are deterministic we recover the memoryless property (1.3.51). □

Exercise 1.57. For $n \in \mathbb{N}$ we denote by C_n the cone in $\mathbb{R}^n$ defined by

$$C_n := \{ (x_1, \dots, x_n) \in \mathbb{R}^n : x_1 \leq x_2 \leq \dots \leq x_n \}.$$

Define $\text{ord} : \mathbb{R}^n \rightarrow C_n$

$$(x_1, \dots, x_n) \mapsto \text{ord}(x_1, \dots, x_n) = (x_{(1)}, x_{(2)}, \dots, x_{(n)}),$$

where

$$x_{(1)} = \min\{x_1, \dots, x_n\}, \quad x_{(2)} = \min\left(\{x_1, \dots, x_n\} \setminus \{x_{(1)}\}\right), \dots$$

In other words, $x_{(1)}, \dots, x_{(n)}$ are the numbers $x_1, \dots, x_n$ rearranged in increasing order.

Suppose $X_1, \dots, X_n$ are n i.i.d. random variables with common cdf

$$F(x) = \int_{-\infty}^x p(s) ds, \quad p \in L^1(\mathbb{R}, \lambda).$$

The *order statistics* of the random variables $X_1, \dots, X_n$ is the random vector

$$\text{ord}(\mathbf{X}) := (X_{(1)}, \dots, X_{(n)}),$$

where $\mathbf{X} = (X_1, \dots, X_n)$.

(i) Show that the distribution of $\text{ord}(\mathbf{X})$ is

$$\mathbb{P}_{\text{ord}(\mathbf{X})}[dx_1 \cdots dx_n] = n! p(x_1) \cdots p(x_n) \mathbf{I}_{C_n}(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

(ii) Denote by $F_{(j)}$ the cdf of the component $X_{(j)}$, $F_{(j)}(x) = \mathbb{P}[X_{(j)} \leq x]$. Prove that

$$F_{(j)}(x) = \sum_{k=j}^n \binom{n}{k} F(x)^k (1 - F(x))^{n-k}.$$

(iii) Suppose that $X_1, \dots, X_n \sim \text{Unif}(0, 1)$. Show that

$$X_{(j)} \sim \text{Beta}(j, n+1-j), \quad \mathbb{E}[X_{(j)}] = \frac{j}{n+1}.$$

(iv) Suppose that $X_1, \dots, X_n \sim \text{Unif}(0, 1)$ and consider the random vector

$$Y = (X_{(2)}, \dots, X_{(n)}).$$

Compute the conditional distribution of Y given $X_{(1)}$

$$\mathbb{P}_Y[dy_2 \cdots dy_n \mid X_{(1)} = x].$$

(v) Suppose that $X_1, \dots, X_n \sim \text{Exp}(\lambda)$. Show that²⁰

$$X_{(1)} \sim \text{Exp}(n\lambda), \quad \mathbb{E}[X_{(1)}] = \frac{1}{n\lambda}.$$

(vi) Suppose that $X_1, \dots, X_n \sim \text{Exp}(\lambda)$. Show that

$$nX_{(1)}, (n-1)(X_{(2)} - X_{(1)}), \dots, 2(X_{(n-1)} - X_{(n-2)}), X_{(n)} - X_{(n-1)}$$

are independent $\text{Exp}(\lambda)$ random variables. **Hint.** Use (i) and Exercise 1.48 to prove first that the spacings

$$S_1 = X_{(1)}, \quad S_2 = X_{(2)} - X_{(1)}, \dots, S_n = X_{(n)} - X_{(n-1)}$$

are independent exponential random variables.

²⁰To appreciate how surprising then conclusion (v) think that an institution buys a large number n of computers, all of the same brand, and $X_1, \dots, X_n$ denote the lifetimes of these machines. Each is expected to last $1/\lambda$ years. The random variable $X_{(1)}$ is the lifetime of the first computer that breaks down. The result in (v) show that we should expect the first break down pretty soon, in $\frac{1}{n\lambda}$ years!

□

Exercise 1.58. Suppose that $X_1, \dots, X_{n-1}$ are independent and uniformly distributed in $[0, 1]$. Consider their order statistics

$$X_{(1)} \leq \dots \leq X_{(n-1)}$$

and the corresponding spacings²¹

$$S_1 = X_{(1)}, S_2 = X_{(2)} - X_{(1)}, \dots, S_n = 1 - X_{(n-1)}.$$

Denote by L_n the largest spacing, $L_n = \max(S_1, \dots, S_n)$.

(i) Prove that $(S_1, \dots, S_n)$ is uniformly distributed in the simplex

$$\Delta_n := \left\{ (s_1, \dots, s_n) \in [0, 1]^n; \sum_{k=1}^n s_k = 1 \right\}.$$

Deduce that $\mathbb{E}[S_k] = \frac{1}{n}, \forall k = 1, \dots, n$.

(ii) Show that

$$\mathbb{E}[L_n] = \frac{1}{n} \sum_{k=1}^n (-1)^{k+1} \frac{1}{k} \binom{n}{k}.$$

Hint. Let For $x \in [0, 1]$ denote by $E_k = E_k(x)$ the event $\{S_k > x\}$. Then $\{L_n > x\} = \bigcap_{k=1}^n E_k(x)$. Conclude using inclusion-exclusion, (i) and (1.3.49).

(iii) Let $Y_1, \dots, Y_n$ be independent $\text{Exp}(1)$ random variables. Set $T_n = Y_1 + \dots + Y_n$. Find the joint distribution of $(Y_1, \dots, Y_n, T_n)$ and show that the random variables

$$\frac{Y_1}{T_n}, \dots, \frac{Y_n}{T_n}$$

has the same joint distribution as the spacings $S_1, \dots, S_n$. Deduce that L_n has the same distribution as

$$\frac{\max_{1 \leq k \leq n} Y_k}{T_n} = \frac{Y_{(n)}}{T_n}.$$

(iv) Prove that L_n and

$$\frac{1}{T_n} \sum_{k=1}^n \frac{Y_k}{k}$$

have the same distribution. **Hint.** Use (iii) and Exercise 1.57(vi). Deduce that²²

$$\mathbb{E}[L_n] := \frac{1}{n} \sum_{k=1}^n \frac{1}{k}.$$

□

Remark 1.6.1. Observe that the above exercise produces a strange identity,

$$\sum_{k=1}^n \frac{1}{k} = \sum_{k=1}^n (-1)^{k+1} \frac{1}{k} \binom{n}{k}.$$

□

²¹The $n-1$ points $X_1, \dots, X_{n-1}$ divide the interval $[0, 1]$ into n subintervals and the spacings are the lengths of these subintervals.

²²This equality shows that $\mathbb{E}[L_n] \sim \frac{\log n}{n}$, which is substantially higher than the mean of each individual spacing, $\mathbb{E}[S_k] = \frac{1}{n}, \forall k$.

Exercise 1.59. Consider the Poisson process $(N(t))_{t \geq 0}$ with intensity λ described in Example 1.3.7.

- (i) Find the distribution of $W_t = N(t) + 1 - t$.
- (ii) Show that $N(t+h) - N(t) \sim \text{Poi } \lambda h, t \geq 0, h > 0$.

□

Exercise 1.60. Consider the Poisson process $(N(t))_{t \geq 0}$ with intensity λ described in Example 1.3.7. Let S be a nonnegative random variable independent of the arrival times $(T_n)_{n \geq 0}$ of the Poisson process. For any arrival time T_n we denote by $Z_{T_n, S}$ the number of arrival times located in the interval $(T_n, T_n + S]$

$$Z_{T_n, S} := \#\{k > n; T_n < T_k \leq T_n + S\}.$$

Prove that

$$\mathbb{P}[Z_{T_n, S} = k] = \int_0^\infty e^{-k\lambda s} \frac{(\lambda s)^k}{k!} \mathbb{P}_S[ds].$$

□

Exercise 1.61. Suppose that $N(t)$ is a Poisson process (see Example 1.3.7) with intensity λ and arrival times

$$T_1 \leq T_2 \leq \dots$$

Fix $t > 0$ and let $(X_n)_{n \geq 1}$ be i.i.d. random variables uniformly distributed in $[0, t]$. Prove that, conditional on $N(t) = n$, the random vectors

$$(T_1, \dots, T_n) \text{ and } (X_{(1)}, \dots, X_{(n)})$$

have the same distribution.

□

Exercise 1.62. Suppose that the 20 contestants at a quiz show are each given the same question, and that each answers it correctly, independently of the others, with probability P . However, the probability of success P itself is a random variable.²³ Suppose, for the sake of illustration, that P is uniformly distributed over the interval $(0, 1]$.

- (i) What is the probability that exactly two of the contestants answer the question correctly?
- (ii) What is the expected number of contestants that answer the question correctly?

□

Exercise 1.63. Let ν be a Borel probability measure on $\mathbb{R}$. Prove that for any Borel subset $B \subset \mathbb{R}$ the map $\Psi_B : \mathbb{R} \rightarrow \mathbb{R}, \Psi_B(y) = \nu[B - y]$ is measurable.

□

Exercise 1.64. Given sigma algebras $\mathcal{F}_\pm, \mathcal{F}_0 \subset \mathcal{S}$, prove that the following are equivalent.

- (i) $\mathcal{F}_+ \perp_{\mathcal{F}_0} \mathcal{F}_-$.
- (ii) $\mathcal{F}_+ \perp_{\mathcal{F}_0} \mathcal{F}_0 \vee \mathcal{F}_-$.

□

Exercise 1.65. Given sigma algebras $\mathcal{F}_\pm, \mathcal{F}_0 \subset \mathcal{S}$, prove that the following are equivalent.

- (i) $\mathcal{F}_+ \perp \mathcal{F}_0 \vee \mathcal{F}_-$

²³Think of P as a random Bernoulli measure of the kind discussed in Example 1.4.20. The source of randomness could be due to the fact that the difficulty of the questions could change randomly from one show to another.

(ii) $\mathcal{F}_+ \perp \mathcal{F}_0$ and $\mathcal{F}_+ \perp_{\mathcal{F}_0} \mathcal{F}_-$.

□

Exercise 1.66. Suppose that μ is a Borel probability measure on the metric space (X, d) . Denote by $\mathcal{C}$ the collection of Borel subsets S of X satisfying the regularity property: for any ε_0 there exists a closed subset $C_\varepsilon \subset S$ and an open subset $\mathcal{O}_\varepsilon \supset S$ such that

$$\mu[\mathcal{O}_\varepsilon \setminus C_\varepsilon] < \varepsilon.$$

- (i) Show that $S \in \mathcal{C} \Rightarrow S^c := X \setminus S \in \mathcal{C}$.
- (ii) Show that any closed set belongs to $\mathcal{C}$.²⁴
- (iii) Show that $\mathcal{C}$ is a π -system.
- (iv) Show that $\mathcal{C}$ is a λ -system.
- (v) Show that $\mathcal{C}$ coincides with the family of Borel subsets.

□

Exercise 1.67. Suppose that (X, d) is a compact metric space and μ is a finite Borel measure on X . Prove that for any $p \in [1, \infty)$ the space $C(X)$ of continuous functions on X is dense in $L^p(X, \mu)$.

Hint. Use Exercise 1.66 to show that for any Borel subset $B \subset X$ the indicator function $\mathbf{I}_B$ can be approximated in L^p by continuous functions.

□

²⁴This is where the fact is a X metric space plays an important role.

Limit theorems

The limit theorems have preoccupied mathematicians from the dawns of probability. The first law of large numbers goes back to Jacob Bernoulli at the end of the seventeenth century. The Golden Theorem in his *Ars Conjectandi* is what we call today a weak law of large numbers. Bernoulli considers an urn that contains a large number of black and white balls. If $p \in (0, 1)$ is the proportion of white balls in the urn and we draw with replacement a large number n of balls, then the proportion p_n of white balls among the extracted ones is with high confidence within a given open interval containing p .

His result lacked foundations since the concept of probability lacked a proper definition. The situation improved at the beginning of the twentieth century when E. Borel proved a strong form of Bernoulli's law. Borel too lacked a good definition of a probability space, but he worked rigorously. In modern terms, he used the interval $[0, 1]$ with the Lebesgue measure as probability space. He then proceeded to construct explicitly a sequence of functions $X_n : [0, 1] \rightarrow \mathbb{R}$ which, viewed as random variables are i.i.d. with common distribution $\text{Bin}(1/2)$.

It took the efforts of Hinchin and Kolmogorov to settle the general case. The strong law of large numbers states that if $(X_n)_{n \in \mathbb{N}}$ are i.i.d. random variables with finite mean μ , then the empirical mean

$$M_n = \frac{1}{n} \sum_{k=1}^n X_k$$

converges a.s. to the theoretical mean μ .

This chapter is devoted to these limit theorems. In the first section we investigate the SLLN= Strong Law of Large Numbers. The approach we use is due to Kolmogorov. It reduces this law to the convergence of random series of independent random variables.

The second section is devoted to the Central Limit Theorem stating that the distribution of M_n is very close to the distribution of a Gaussian random variable with the same mean and variance as M_n . The third section, is more modern, and it is devoted to concentration inequalities. These state in a quantitative fashion that the probability that M_n deviates from the mean μ by a certain amount is extremely small under certain conditions. The fourth section is devoted to uniform limit theorem of the Glivenko-Cantelli type. We have included this section due to its applications in machine learning.

In particular, we show how such results coupled with the concentration inequalities lead to **Probably Approximately Correct**, or PAC, learning.

The last section of this chapter is devoted to a brief introduction to the Brownian motion. This is such a fundamental object that we thought that any student of probability ought to make its acquaintance as soon as possible. As always, along the way we present many, we hope, interesting examples.

2.1. The Law of Large Numbers

This section is devoted to the (Strong) Law of Large numbers. We follow Kolmogorov's approach based on random series, a subject of independent interest.

2.1.1. Random series. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and consider a sequence of *independent* random variables

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad n \in \mathbb{N}.$$

The independence of the random variables (X_n) allows us to invoke Kolmogorov's 0-1 theorem and conclude that the random series

$$\sum_{n \in \mathbb{N}} X_n \tag{2.1.1}$$

either converges almost surely, or diverges almost surely. We want to describe by describing one simple sufficient condition for convergence.

Theorem 2.1.1 (Kolmogorov's one series). *Suppose that*

$$\mathbb{E}[X_n] = 0, \quad \forall n \in \mathbb{N}, \tag{2.1.2a}$$

$$\sum_{n \geq 1} \text{Var}[X_n] < \infty. \tag{2.1.2b}$$

Then the series (2.1.1) converges almost surely and in L^2 .

Proof. For $n \in \mathbb{N}$ we denote by S_n the n -th partial sum of the series (2.1.1),

$$S_n := \sum_{k=1}^n X_k.$$

The L^2 -convergence follows immediately from (2.1.2b) which, coupled with the independence of the random variables (X_n) implies that the sequence (S_n) is Cauchy in L^2 since

$$\|S_{n+k} - S_n\|_{L^2}^2 = \sum_{j=1}^k \text{Var}[X_{n+j}], \quad \forall k, n \in \mathbb{N}.$$

The proof of the a.s. convergence is more difficult. It relies on a fundamental inequality which we will further generalize in the next chapter. The independence of the random variables (X_n) is used crucially in its proof.

Lemma 2.1.2 (Kolmogorov's maximal inequality). *Set*

$$M_n := \max_{1 \leq k \leq n} |S_k|.$$

Then, for all $a > 0$, we have

$$\mathbb{P}[M_n > a] \leq \frac{1}{a^2} \text{Var}[S_n] = \frac{1}{a^2} \sum_{k=1}^n \text{Var}[X_k]. \quad (2.1.3)$$

Additionally if $\exists c > 0$ such that $|X_n| \leq c$, $\forall n$, then

$$1 - \frac{(a+c)^2}{\text{Var}[S_n]} \leq \mathbb{P}[M_n > a]. \quad (2.1.4)$$

Proof of Kolmogorov's maximal inequality. Define

$$N : \Omega \rightarrow \mathbb{N} \cup \{\infty\}, \quad N(\omega) := \inf \{ n \geq 1; |S_n(\omega)| > a \}.$$

Notice that $N(\omega)$ is the first $n \in \mathbb{N} \cup \{\infty\}$ such that $S_n(\omega) > a$, i.e.,

$$N(\omega) = k \iff S_1(\omega), \dots, S_{k-1}(\omega) \leq a \text{ and } S_k(\omega) > a.$$

This shows that the event $A_k = \{N = k\}$ is in the σ -algebra generated by $X_1, \dots, X_k$. Since $S_n - S_k = X_{k+1} + \dots + X_n$ we deduce that $\mathbf{I}_{A_k}, \mathbf{I}_{A_k} S_k$ are independent of $S_n - S_k$. We have

$$\begin{aligned} \text{Var}[S_n] &= \mathbb{E}[S_n^2] \geq \mathbb{E}[S_n^2 \mathbf{I}_{\{M_n \geq a\}}] \\ &= \sum_{k=1}^n \mathbb{E}[\mathbf{I}_{A_k} S_n^2] = \sum_{k=1}^n \mathbb{E}[\mathbf{I}_{A_k} (S_k^2 + 2S_k(S_n - S_k) + (S_n - S_k)^2)] \\ &\quad (\mathbf{I}_{A_k}, \mathbf{I}_{A_k} S_k \perp\!\!\!\perp S_n - S_k) \\ &= \sum_{k=1}^n \left(\mathbb{E}[\mathbf{I}_{A_k} S_k^2] + 2\mathbb{E}[\mathbf{I}_{A_k} S_k] \underbrace{\mathbb{E}[S_n - S_k]}_{=0} + \underbrace{\mathbb{E}[\mathbf{I}_{A_k}] \mathbb{E}[(S_n - S_k)^2]}_{\geq 0} \right) \\ &\geq \sum_{k=1}^n \underbrace{\mathbb{E}[\mathbf{I}_{A_k} S_k^2]}_{S_k^2 \geq a^2 \text{ on } A_k} \geq a^2 \sum_{k=1}^n \mathbb{P}[A_k] = a^2 \mathbb{P}[M_n \geq a]. \end{aligned}$$

This proves (2.1.3).

To prove (2.1.4) we argue as in [137, Sec.17.2] and we set

$$B_0 := \Omega, \quad B_k := \{N > k\} = \{M_k \leq a\}.$$

Observe that, $\forall k = 1, \dots, n$, $B_{k-1} \supset B_k$ and

$$A_k = \{N = k\} = \{N > k-1\} \setminus \{N > k\} = B_{k-1} \setminus B_k,$$

$$S_{k-1} \mathbf{I}_{B_{k-1}} + X_k \mathbf{I}_{B_{k-1}} = S_k \mathbf{I}_{B_{k-1}} = S_k \mathbf{I}_{B_k} + S_k \mathbf{I}_{A_k}.$$

Since $B_{k-1} \in \sigma(X_1, \dots, X_{k-1})$ we have $X_k \perp\!\!\!\perp S_{k-1} \mathbf{I}_{B_{k-1}}$. Hence

$$\mathbb{E}[(S_{k-1} \mathbf{I}_{B_{k-1}}) \cdot (X_k \mathbf{I}_{B_{k-1}})] = \mathbb{E}[(S_{k-1} \mathbf{I}_{B_{k-1}}) \cdot X_k] = 0.$$

We deduce that

$$\begin{aligned} \mathbb{E}[(S_{k-1} \mathbf{I}_{B_{k-1}} + X_k \mathbf{I}_{B_{k-1}})^2] &= \mathbb{E}[(S_{k-1} \mathbf{I}_{B_{k-1}})^2] + \mathbb{E}[(X_k \mathbf{I}_{B_{k-1}})^2] \\ &\quad (X_k \perp\!\!\!\perp \mathbf{I}_{B_{k-1}}) \\ &= \mathbb{E}[(S_{k-1} \mathbf{I}_{B_{k-1}})^2] + \text{Var}[X_k^2] \mathbb{P}[B_{k-1}]. \end{aligned}$$

On the other hand, $\mathbf{I}_{B_k} \mathbf{I}_{A_k} = 0$ so

$$\mathbb{E}[(S_k \mathbf{I}_{B_k} + S_k \mathbf{I}_{A_k})^2] = \mathbb{E}[(S_k \mathbf{I}_{B_k})^2] + \mathbb{E}[(S_k \mathbf{I}_{A_k})^2].$$

Hence

$$\mathbb{E}[(S_{k-1}\mathbf{I}_{B_{k-1}})^2] + \text{Var}[X_k^2]\mathbb{P}[B_{k-1}] = \mathbb{E}[(S_k\mathbf{I}_{B_k})^2] + \mathbb{E}[(S_k\mathbf{I}_{A_k})^2].$$

Since $|X_k| \leq c$ and $|S_{k-1}| \leq a$ on A_k we deduce

$$|S_k\mathbf{I}_{A_k}| \leq |S_{k-1}|\mathbf{I}_{A_k} + |X_k|\mathbf{I}_{A_k} \leq |S_{k-1}|\mathbf{I}_{A_k} + c\mathbf{I}_{A_k} \leq (a+c)\mathbf{I}_{A_k}$$

Observe that $\mathbb{P}[B_{k-1}] \geq \mathbb{P}[B_n]$. We deduce

$$\leq \mathbb{E}[(S_{k-1}\mathbf{I}_{B_{k-1}})^2] + \text{Var}[X_k^2]\mathbb{P}[B_n] \leq \mathbb{E}[(S_k\mathbf{I}_{B_k})^2] + (a+c)^2\mathbb{P}[A_k].$$

Hence

$$\text{Var}[X_k^2]\mathbb{P}[B_n] \leq \mathbb{E}[(S_k\mathbf{I}_{B_k})^2] - \mathbb{E}[(S_{k-1}\mathbf{I}_{B_{k-1}})^2] + (a+c)^2\mathbb{P}[A_k]$$

and

$$\begin{aligned} \sum_{k=1}^n \text{Var}[X_k^2]\mathbb{P}[B_n] &\leq \mathbb{E}[(S_n\mathbf{I}_{B_n})^2] + (a+c)^2 \sum_{k=1}^n \mathbb{P}[A_k] \\ &\leq a^2\mathbb{P}[B_n] + (a+c)^2\mathbb{P}[B_n^c] \leq (a+c)^2 \end{aligned}$$

In other words,

$$\text{Var}[S_n](1 - \mathbb{P}[M_n > a]) \leq (a+c)^2$$

This proves (2.1.4). $\square$

We can now complete the proof of Theorem 2.1.1. Using Kolmogorov's maximal inequality for the sequence $(X_{m+n})_{n \in \mathbb{N}}$ we deduce that for any $n \in \mathbb{N}$ we have

$$\begin{aligned} \mathbb{P}\left[\max_{1 \leq k \leq n} |S_{m+k} - S_m| > \varepsilon\right] &\leq \frac{1}{\varepsilon^2} \text{Var}[S_{m+n} - S_m] = \frac{1}{\varepsilon^2} \sum_{k=1}^n \text{Var}[X_{m+k}] \\ &\leq \frac{1}{\varepsilon^2} \underbrace{\sum_{k=1}^n \text{Var}[X_{m+k}]}_{=: r_m}. \end{aligned}$$

Thus

$$\mathbb{P}\left[\sup_{n \geq 1} |S_{m+n} - S_m| > \varepsilon\right] \leq \frac{r_m}{\varepsilon^2}. \quad (2.1.5)$$

We set

$$Y_m := \sup_{i, j \geq m} |S_i - S_j|, \quad Z_m := \sup_{n \geq 1} |S_{m+n} - S_m|.$$

Now observe that S_m converges a.s. iff $Y_m \rightarrow 0$ a.s. The sequence Y_m is nonincreasing and thus it converges a.s. to a random variable $Y \geq 0$. We will show that $Y = 0$ a.s..

Note that, for $i, j > m$ we have

$$|S_i - S_j| \leq |S_i - S_m| + |S_j - S_m| \leq 2Z_m,$$

so $Y_m \leq 2Z_m, \forall m$ so

$$Y_m > 2\varepsilon \Rightarrow Z_m > \varepsilon \Rightarrow \mathbb{P}[Y_m > 2\varepsilon] \leq \mathbb{P}[Z_m > \varepsilon] \stackrel{(2.1.5)}{\leq} \frac{r_m}{\varepsilon^2} \quad \forall m \geq 1, \forall \varepsilon > 0.$$

Hence

$$\lim_{m \rightarrow \infty} \mathbb{P}[Y_m > \varepsilon] = \lim_{m \rightarrow \infty} \mathbb{P}[Z_m > \varepsilon] = 0.$$

In other words, the sequence (Y_m) converges in probability to 0. Since it also converges a.s. to Y we deduce that $Y = 0$ a.s.. $\square$

Kolmogorov also established necessary and sufficient conditions for convergence in his *three series theorem*. Before we state it let us introduce a convenient notation. For any random variable X and any positive constant C we denote by X^C the truncation

$$X^C := X \mathbf{I}_{\{|X| \leq C\}} = \begin{cases} X, & |X| \leq C, \\ 0, & |X| > C. \end{cases} \quad (2.1.6)$$

Theorem 2.1.3 (Kolmogorov's three series theorem). *Consider a sequence of independent random variables $X_n \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$. The following statements are equivalent.*

(i) *The series*

$$\sum_{n \geq 1} X_n \quad (2.1.7)$$

converges almost surely.

(ii) *There exists $C > 0$ such that the following three series are convergent.*

$$\sum_{n \geq 1} \mathbb{P}[|X_n| > C] = \sum_{n \geq 1} \mathbb{P}[X_n \neq X_n^C], \quad (2.1.8a)$$

$$\sum_{n \geq 1} \mathbb{E}[X_n^C], \quad \sum_{n \geq 1} \text{Var}[X_n^C]. \quad (2.1.8b)$$

Proof. (ii) $\Rightarrow$ (i) Note that that condition (2.1.8a) coupled with the first Borel-Cantelli lemma (Theorem 1.3.55(i)) implies that $\mathbb{P}[X_n \neq X_n^C \text{ i.o.}] = 0$. Hence the series $\sum_{n > 0} X_n$ converges a.s. iff the series $\sum_{n > 0} X_n^C$ converges. The convergence of the latter follows from (2.1.8b) and Kolmogorov's one series theorem.

(i) $\Rightarrow$ (ii) Since the series $\sum_{n > 0} X_n$ is a.s. convergent we deduce that $X_n \rightarrow 0$ a.s.. Thus, for any $c > 0$

$$\mathbb{P}[|X_n| > c \text{ i.o.}] = 0$$

and the second Borel-Cantelli Lemma (Theorem 1.3.55(ii)) implies

$$\sum_{n > 0} \mathbb{P}[|X_n| > C \text{ i.o.}] < \infty.$$

This proves (2.1.8a). Hence $\mathbb{P}[X_n \neq X_n^C \text{ i.o.}] = 0$. Since the series $\sum_{n > 0} X_n$ converges a.s. we deduce that $\sum_{n > 0} X_n^C$ converges a.s.. The conditions (2.1.8b) are now a consequence of the following result of independent interest.

Lemma 2.1.4. *Suppose that $(Y_n)_{n \in \mathbb{N}}$ is a sequence of independent random variables that are uniformly bounded and such the series $\sum_n Y_n$ converges a.s.. Then the numerical series*

$$\sum_n \mathbb{E}[Y_n] \quad \text{and} \quad \sum_n \text{Var}[Y_n]$$

converge.

Proof of Lemma 2.1.4. We follow the approach in [137, Sec.17.3]. The proof uses a clever symmetrization trick.

Choose a sequence of independent random variables (Y'_n) , that are independent of (Y_n) and such that Y_n and Y'_n have the same distribution for any n . We set $Y_n^* := Y_n - Y'_n$. The random variables Y_n^* are symmetric in the sense that for any Borel subset $B \subset \mathbb{R}$ we have $\mathbb{P}[Y_n^* \in B] = \mathbb{P}[-Y_n^* \in B]$.

Fix $C > 0$ such that $|Y_n| \leq c, \forall n$. Then

$$|Y_n^*| \leq |Y_n| + |Y'_n| \leq 2C, \quad \mathbb{E}[Y_n^*] = 0, \quad \text{Var}[Y_n^*] = 2 \text{Var}[Y_n].$$

Since $\sum_n Y_n$ converges a.s. so does $\sum_n Y_n^*$. We set

$$S_n^* = \sum_{k=1}^n Y_k^*.$$

For $m \leq n$

$$M_{m,n} = \max_{1 \leq k} |S_{m+k}^* - S_m^*|.$$

Using (2.1.4) we deduce that for any $\varepsilon > 0$ and any $0 \leq m < n$ we have

$$1 - \frac{(\varepsilon + c)^2}{\text{Var}[S_n^* - S_m^*]} \leq \mathbb{P}[M_{m,n} > \varepsilon].$$

Since S_n^* converges a.s. we deduce that for any $\varepsilon > 0$

$$\lim_{m,n \rightarrow \infty} \mathbb{P}[M_{m,n} > \varepsilon] = 0.$$

Choose $m_0 > 0$ such that $\mathbb{P}[M_{m_0,n} > \varepsilon] < \frac{1}{2}, \forall n > m_0$. Hence, $\forall n > m_0$

$$2 \sum_{m_0 < k \leq n} \text{Var}[Y_k] = \sum_{m_0 < k \leq n} \text{Var}[Y_k^*] = \text{Var}[S_n^* - S_{m_0}^*] \leq 2(\varepsilon + c)^2.$$

This proves that the series

$$\sum_{n>0} \text{Var}[Y_n]$$

is convergent. Kolmogorov's one series theorem implies that the series

$$\sum_{n>0} (Y_n - \mathbb{E}[Y_n])$$

converges a.s.. We deduce that $\sum_{n>0} \mathbb{E}[Y_n]$ is convergent since $\sum_{n>0} Y_n$ converges a.s.. $\square$

This completes the proof of Theorem 2.1.3. $\square$

Example 2.1.5. Consider a sequence of i.i.d. Bernoulli random variables $(B_n)_{n \in \mathbb{N}}$ with success probability $\frac{1}{2}$. The resulting random variables $R_n = (-1)^{B_n}$ are called *Rademacher random variables* and take only the values ± 1 with equal probabilities.

Suppose that

$$\sum_{n \geq 1} a_n$$

is a deterministic series with bounded positive terms. We obtain the random series

$$\sum_{n \geq 1} R_n a_n. \tag{2.1.9}$$

We have $\mathbb{E}[R_n a_n] = 0$ and Kolmogorov's one series theorem shows that if

$$\sum_{n \geq 1} a_n^2 < \infty$$

the random series (2.1.9) is a.s. convergent. Conversely, if the random series (2.1.9) is a.s. convergent, then Lemma 2.1.4 implies that $\sum_{n \geq 1} a_n^2 < \infty$.

As a special case, suppose that $a_n = \frac{1}{n}$, $\forall n \geq 1$. Consider the harmonic series with random signs

$$\sum_{n \geq 1} \frac{R_n}{n} = \pm 1 \pm \frac{1}{2} \pm \frac{1}{3} \pm \cdots, \quad (2.1.10)$$

We know that if all the terms are positive, a probability zero event, then we obtain the harmonic series which is divergent. On the other hand,

$$\sum_{k \geq 1} \frac{1}{k^2} < \infty,$$

and we deduce from Kolmogorov's one series theorem that the series (2.1.10) is a.s. convergent. Thus, if we flip a fair coin with two sides, a + side and a - side and we assign the signs in (2.1.10) according to the coin flips, the resulting series is convergent with probability 1! $\square$

Remark 2.1.6. The so called *Lévy's equivalence theorem*, [61, §III.2, Cor. 2], [64, §9.7], [134, §43], [137, Sec.18.2] or [216, Thm.3.9] states that a series with independent terms converges a.s. iff converges in probability, iff converges in distribution; see Definition 2.2.3(iii). The proof that the convergence in probability implies convergence a.s. is outlined in Exercises 2.1 and 2.2. In Example 3.2.8 we present a martingale-based proof of this theorem. $\square$

2.1.2. The Law of Large Numbers. The frequentist interpretation of probability asserts that the probability of an event is roughly the frequency of the occurrence of that event in a very large number of independent trials. The Law of Large Numbers formalizes this intuition. The surprising thing, at least to this author, is that reality respects the theory so closely: the Law of Large Numbers adds a surprising level of predictability to uncertainty!

Throughout this section $(X_n)_{n \geq 1}$ is a sequence of iid random variables $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$\mu := \mathbb{E}[X_n], \quad S_n := X_1 + \cdots + X_n.$$

The various versions of the Law of Large Numbers state that the empirical means S_n/n converge in an appropriate sense to the theoretical mean μ . The convergence in probability is usually referred to as the *Weak Law of Large Numbers* (or WLLN) while the a.s. convergence is known as the *Strong Law of Large Numbers* (or SLLN). We begin by presenting a few special, but historically important, cases.

Theorem 2.1.7 (Markov). *If $X_n \in L^2(\Omega, \mathcal{S}, \mathbb{P})$, then $\frac{1}{n}S_n \rightarrow \mu$ in L^2 and thus also in probability.*

Proof. Denote by σ^2 the common variance of the random variables X_n . Since they are independent we have $\text{Var}[S_n] = n\sigma^2$, so

$$\text{Var}[S_n/n] = \frac{1}{n^2} \text{Var}[S_n] = \frac{1}{n\sigma^2}.$$

Let $\varepsilon > 0$. Note that $\mathbb{E}[S_n/n] = \mu$ so

$$\|S_n/n - \mu\|_{L^2}^2 = \text{Var}[S_n/n] = \frac{1}{n\sigma^2}.$$

Hence $\frac{S_n}{n}$ converges to μ in L^2 . Proposition 1.3.65 implies that $S_n/n \rightarrow \mu$ in probability. $\square$

Theorem 2.1.8 (Cantelli). *If $X_n \in L^4(\Omega, \mathcal{S}, \mathbb{P})$, then $\frac{1}{n}S_n \rightarrow \mu$ almost surely.*

Proof. By replacing X_n with $Y_n := X_n - \mu$ we can assume $\mu = 0$. We set

$$\sigma^2 := \mu_2[X_k], \quad r^4 := \mu_4[X_k], \quad M_n := S_n/n.$$

Note that

$$\mathbb{P}[|M_n| > \varepsilon] = \mathbb{P}[|M_n|^4 > \varepsilon^4] \leq \frac{1}{\varepsilon^4} \mathbb{E}[M_n^4] = \frac{1}{n^4 \varepsilon^4} \mathbb{E}[S_n^4].$$

Observe that

$$\mathbb{E}[S_n^4] = \sum_{i,j,k,\ell=1}^n \mathbb{E}[X_i X_j X_k X_\ell]. \quad (2.1.11)$$

Let $i \neq j$. Due to the independence of the random variables $(X_n)_{n \in \mathbb{N}}$ we have

$$\mathbb{E}[X_i^2 X_j^2] = \mathbb{E}[X_i^2] \mathbb{E}[X_j^2] = \sigma^4, \quad \mathbb{E}[X_i X_j^3] = \mathbb{E}[X_i] \mathbb{E}[X_j^3] = 0$$

Similarly, for distinct i, j, k, ℓ we have

$$\mathbb{E}[X_i X_j X_k X_\ell] = 0.$$

Thus

$$\mathbb{E}[S_n] = n^4 r^4 + 2 \binom{4}{2} \sum_{j < k} \sigma^4 = n r^4 + 6 \binom{n}{2} \sigma^4 = O(n^2) \text{ as } n \rightarrow \infty.$$

Hence $\mathbb{E}[M_n^4] = O\left(\frac{1}{n^2}\right)$, so

$$\mathbb{P}[|M_n| > \varepsilon] = O\left(\frac{1}{n^2 \varepsilon^4}\right) \text{ as } n \rightarrow \infty.$$

Since the series $\sum_{n \geq 1} \frac{1}{n^2}$ is convergent we deduce that, for any $\varepsilon > 0$,

$$\sum_{n \geq 1} \mathbb{P}[|M_n| > \varepsilon] < \infty.$$

Corollary 1.3.57 implies that $M_n \rightarrow 0$ a.s.. □

Remark 2.1.9. The above Strong Law of Large Numbers is not the most general, but its proof makes the role of independence much more visible. More precisely the independence, or the small correlations force the fourth moment of S_n to be “unnaturally” small and thus the large fluctuations around the mean are highly unlikely, i.e. the $\mathbb{P}[|M_n| > \varepsilon]$ is very small for large n . □

The next result, due to Kolmogorov, generalizes both results above.

Theorem 2.1.10 (The Strong Law of Large Numbers). *Suppose that $(X_n)_{n \geq 1}$ is a sequence of iid random variables $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} S_n = \mu \text{ a.s..}$$

Proof. We accomplish this in several steps.

Step 1. Truncate. Set

$$Y_n := X_n \mathbf{I}_{\{|X_n| < n\}}, \quad T_n := Y_1 + \cdots + Y_n.$$

Obviously $|Y_n| \leq n, \forall n$. We claim that

$$\mathbb{P}[X_n \neq Y_n \text{ i.o.}] = 0. \quad (2.1.12)$$

Indeed, since the random variables (X_n) are identically distributed we have

$$\begin{aligned} \sum_{k \geq 1} \mathbb{P}[|X_k| > k] &= \sum_{k \geq 1} \mathbb{P}[|X_1| > k] \\ &\leq \int_0^\infty \mathbb{P}[|X_1| > t] dt \stackrel{(1.3.48)}{=} \mathbb{E}[|X_1|] < \infty. \end{aligned}$$

and Borel-Cantelli's Lemma implies implies that

$$\mathbb{P}[|X_k| > k \text{ i.o.}] = 0.$$

This is equivalent to (2.1.12). We deduce from (2.1.12) that

$$\lim_{n \rightarrow \infty} \frac{1}{n} |S_n - T_n| = 0 \text{ a.s..}$$

Thus, it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} T_n = \mu \text{ a.s..} \quad (2.1.13)$$

Step 2. Centering. The sequence $(\mathbb{E}[Y_k])_{k \geq 1}$ converges to $\mu = \mathbb{E}[X]$ as $k \rightarrow \infty$. Indeed, since the random variables are identically distributed we have

$$\mathbb{E}[Y_k] = \mathbb{E}[X_k \mathbf{I}_{\{|X_k| \leq k\}}] = \mathbb{E}[X_1 \mathbf{I}_{\{|X_1| \leq k\}}] \rightarrow \mathbb{E}[X_1],$$

where at the last step we used the Dominated Convergence theorem. It follows that the sequence $\mathbb{E}[Y_n]$ is also *Cèsaro convergent*¹ to the same limit, i.e.,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[T_n] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \mathbb{E}[Y_k] = \mu.$$

Thus, it suffices to prove that

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\frac{1}{n} \sum_{k=1}^n Y_k - \frac{1}{n} \sum_{k=1}^n \mathbb{E}[Y_k] \right) &= 0, \text{ a.s.} \\ Z_n &:= Y_n - \mathbb{E}[Y_n]. \end{aligned}$$

The random variables Z_n are bounded, centered, independent but not identically distributed. We have to prove that the *Cèsaro means* of Z_n converge to 0 a.s., i.e.,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n Z_k = 0 \text{ a.s..} \quad (2.1.14)$$

Step 3. Conclusion. We will rely on the following elementary result.

Lemma 2.1.11 (Kronecker's Lemma). *Suppose that $(a_n)_{n \in \mathbb{N}}$ and $(x_n)_{n \in \mathbb{N}}$ are sequences of real numbers satisfying the following conditions.*

- (i) *The sequence (a_n) is increasing, positive and unbounded.*
- (ii) *The series $\sum_{n \geq 1} \frac{x_n}{a_n}$ is convergent.*

Then

$$\lim_{n \rightarrow \infty} \frac{1}{a_n} \sum_{k=1}^n x_k = 0.$$

¹Use Exercise 2.6 with $p_{k,n} = 1/n$.

Assume temporarily the validity of Kronecker's lemma. Thus, to prove (2.1.14) it suffices to show that the random series

$$\sum_{n \geq 1} \frac{Z_n}{n}$$

is a.s. convergent. The independence assumption will finally play a role because we will invoke the one-series theorem. Clearly the random variables $\frac{Z_n}{n}$ are independent. We claim that

$$\sum_{k \geq 1} \frac{\text{Var}[Z_k]}{k^2} < \infty. \quad (2.1.15)$$

We have

$$\begin{aligned} \text{Var}[Z_k] &= \text{Var}[Y_k] = \mathbb{E}[Y_k^2] - \mathbb{E}[Y_k]^2 \leq \mathbb{E}[Y_k^2] \\ &\stackrel{(1.3.48)}{=} \int_0^\infty 2y \mathbb{P}[|Y_k| > y] dy = \int_0^\infty 2y \mathbb{P}[k \geq |X_k| > y] \mathbf{I}_{\{y < k\}} dy \\ &\leq \int_0^\infty 2y \mathbb{P}[|X_k| > y] \mathbf{I}_{\{y < k\}} dy. \end{aligned}$$

Thus

$$\begin{aligned} \sum_{k \geq 1} \frac{\text{Var}[Z_k]}{k^2} &\leq \sum_{k \geq 1} \frac{1}{k^2} \int_0^\infty 2y \mathbb{P}[|X_k| > y] \mathbf{I}_{\{y < k\}} dy \\ &= \int_0^\infty \left(\sum_{k \geq 1} \frac{1}{k^2} \mathbf{I}_{\{y \leq k\}} \right) 2y \mathbb{P}[|X_1| > y] dy \\ &= \int_0^\infty \underbrace{\left(\sum_{k \geq y} \frac{1}{k^2} \right)}_{=: w(y)} 2y \mathbb{P}[|X_1| > y] dy. \end{aligned}$$

We claim that

$$w(y) < 6, \quad \forall y \geq 0. \quad (2.1.16)$$

Indeed, for $y \leq 1$ we have

$$w(y) = 2y \sum_{k \geq 1} \frac{1}{k^2} \leq 4y < 4.$$

For $y \in (1, 2]$ we have

$$w(y) = 2y \sum_{k \geq 2} \frac{1}{k^2} < 2y \leq 4.$$

For $y > 2$ we have

$$\sum_{k \geq y} \frac{1}{k^2} \leq \int_{[y]-1}^\infty \frac{1}{t^2} dt = \frac{1}{[y]-1}$$

so

$$w(y) \leq \frac{2y}{[y]-1} \leq \frac{2[y]+2}{[y]-1} = 2 + \frac{4}{[y]-1} < 6.$$

Using (2.1.16) we deduce

$$\sum_{k \geq 1} \frac{\text{Var}[Z_k]}{k^2} < 6 \int_0^\infty \mathbb{P}[|X_1| > y] dy = 6\mathbb{E}[|X_1|] < \infty.$$

This proves (2.1.15) and completes the proof of the SLLN, assuming Lemma 2.1.11. $\square$

Proof of Lemma 2.1.11. Set

$$y_k := \frac{x_k}{a_k}, \quad s_0 = a_0 := 0, \quad s_k := \sum_{j=1}^k y_j, \quad k \geq 1,$$

so that the sequence $(s_n)_{n \geq 1}$ is convergent. We have to show that

$$\lim_{n \rightarrow \infty} \frac{1}{a_n} \sum_{k=1}^n a_k y_k = 0.$$

We have Abel's trick²

$$\begin{aligned} \sum_{k=1}^n a_k y_k &= \sum_{k=1}^n a_k (s_k - s_{k-1}) \\ &= a_1 (\boxed{s_1} - s_0) + a_2 (s_2 - \boxed{s_1}) + \cdots + a_n (s_n - \boxed{s_{n-1}}) \\ &= a_n s_n - \sum_{k=1}^n s_{k-1} (a_k - a_{k-1}). \end{aligned}$$

Now set

$$w_k := a_k - a_{k-1}, \quad p_{n,k} := \frac{w_k}{a_n}.$$

Since $(a_n)_{n \in \mathbb{N}}$ is increasing, positive and unbounded we deduce

$$\sum_{k=1}^n p_{n,k} = 1, \quad \forall n \geq 1, \quad \lim_{n \rightarrow \infty} p_{n,k} = 0, \quad \forall k. \quad (2.1.17)$$

Observe that

$$\frac{1}{a_n} \sum_{k=1}^n a_k y_k = s_n - \sum_{k=1}^n p_{n,k} s_{k-1}.$$

The conditions (2.1.17) imply that (see Exercise 2.6)

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n p_{n,k} s_{k-1} = \lim_{n \rightarrow \infty} s_n.$$

$\square$

Since a.s. convergence implies convergence in probability we deduce from the SLLN the *Weak Law of Large Numbers* (or WLLN)

Corollary 2.1.12. Suppose that $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$, $n \in \mathbb{N}$, is a sequence of i.i.d. random variables with common mean μ . We set

$$S_n = \sum_{k=1}^n X_k.$$

Then the empirical mean $\frac{1}{n} S_n$ converges in probability to μ . $\square$

²This is a discrete version of the integration by parts formula.

Remark 2.1.13. (a) With a bit more effort one can show that in Strong Law of Large Number the empirical mean $\frac{1}{n}S_n$ not just a.s. but also in L^1 ; see Corollary 3.2.65.

(b) Let us observe that in the Weak Law of Large Numbers Theorem 2.1.7 the random variables X_n need not be independent or identically distributed. Assuming all have mean 0, all we need for that for the Weak Law of Large numbers to hold is that the random variables are pairwise uncorrelated,

$$\mathbb{E}[X_m X_n] = \mathbb{E}[X_m] \mathbb{E}[X_n], \quad \forall m \neq n, \quad (2.1.18)$$

and the only constraint on their distribution is

$$\sup_n \mathbb{E}[X_n^2] < \infty.$$

In Exercise 2.9 we ask the reader to show that the WLLN holds even the random variables are not identically or dependent. It suffices to we assume something weaker than (2.1.18) namely that if $|m - n| \gg 1$, the random variables X_m and X_n are weakly correlated, i.e.,

$$\lim_{k \rightarrow \infty} \sup_{m \in \mathbb{N}} |\mathbb{E}[X_m X_{m+k}]| = 0. \quad (2.1.19)$$

If the random variables are independent, but not necessarily identically distributed, there are known necessary and sufficient conditions for the WLLN to hold. We refer to [76, IX], [89, §22.], or [169, Chap. 4] for details.

Similarly, in the Strong Law of Large Numbers the variables need not be independent. The theorem continues to hold if the variables are identically distributed, integrable and only pairwise independent. For a proof we refer to [68, Sec. 2.4]. In Chapters 4 and 5 we describe examples of identically distributed integrable random variables satisfying SLLN yet they are not even pairwise independent. Another principle is at work in these cases, namely ergodicity. In fact Theorem 2.1.10 is a special case of Birkhoff's Ergodic Theorem 5.1.22.

The arguments in the proof Theorem 2.1.8 show that the SLLN holds even when the variables X_n are neither independent, nor identically distributed. Assuming that all the variables have mean zero, the SLLN holds if any four of them are independent, and the only assumptions about their distributions is

$$\sup_n \mathbb{E}[X_n^4] < \infty.$$

This can be substantially strengthened. For example, this happens if $\exists C > 0$ such that

$$|\mathbb{E}[X_n X_m]| \leq \frac{C}{1 + |m - n|}, \quad \forall m, n. \quad (2.1.20)$$

We refer to [142] for more details. Note that (2.1.20) is a stronger version of (2.1.19).

A natural philosophical question arises. What makes the Law of Large Numbers possible? The above discussion suggests that it is a consequence of a mysterious form of “asynchronicity”: their fluctuations around the mean are not in resonance and they cancel each other out. These features can be observed in the other Laws of Large Numbers we will discuss in this text. $\square$

Remark 2.1.14. Suppose that $(X_n)_{n \geq 1}$ is a sequence of i.i.d. variables. The Strong Law of Large Numbers shows that if they have finite mean μ , then the empirical means

$$M_n = \frac{1}{n} (X_1 + \cdots + X_n)$$

converge a.s. to μ . If $\mu = \infty$ and M_n converge a.s. to a random variable M_∞ , then M_∞ is a.s. constant. Exercise 2.12 outlines a proof of this fact. $\square$

Example 2.1.15. Suppose we roll a fair die a large number n of times and we denote by S_n the number of times we roll a 1. Intuition tells us that if the die is fair, then for large n , the fraction of times we get a 1 should be close to $\frac{1}{6}$, i.e.,

$$\frac{S_n}{n} \approx \frac{1}{6} \text{ for } n \gg 0.$$

This follows from the SLLN. Indeed, the above experiment is encoded by a sequence $(X_n)_{n \in \mathbb{N}}$ of i.i.d. Bernoulli random variables with success probability $p = \frac{1}{6}$. Then

$$S_n = \sum_{k=1}^n X_k,$$

and the SLLN

$$\frac{S_n}{n} \rightarrow \mathbb{E}[X_1] = \frac{1}{6} \text{ a.s. as } n \rightarrow \infty.$$

It helps to visualize a computer simulation of such an experiment. Suppose we roll a die a large number N of times. For $i = 1, \dots, N$ we denote by f_i the frequency of 1-s during the first i trials, i.e.,

$$f_i = \frac{S_i}{i}.$$

The resulting vector $(f_i)_{1 \leq i \leq N} \in \mathbb{R}^N$ is called relative or cumulative frequency.

The R-code below simulates one such experiment when we roll the die 12,000 times.

```
N<-12000
x<-sample(1:6, N, replace=TRUE)
rolls<-x==1
rel_freq<-cumsum(rolls)/(1:N)

plot(1:N,rel_freq,type="l", xlab="Number of rolls",
     ylab="The frequency of occurrence of 1",
     main="Average number 1s during random rolls of a die")
abline(h=1/6,col="red")
```

The output is a plot of the collection of points (i, f_i) depicted in Figure 2.1. □




Figure 2.1. The frequencies f_i fluctuates wildly initially and then stabilizes around the horizontal line $y = 1/6$ in perfect agreement with SLLN.

Example 2.1.16 (The Monte-Carlo method). Consider a box (parallelepiped)

$$B_k := I_1 \times \cdots \times I_k \subset \mathbb{R}^k$$

where $I_1, \dots, I_k \subset \mathbb{R}$ are nontrivial bounded intervals. Consider independent random variables $X_1, \dots, X_k$, where X_j is uniformly distributed on I_j . The probability distribution of the random vector $\mathbf{X} = (X_1, \dots, X_k)$ is

$$\frac{1}{\lambda_k[B_k]} \mathbf{I}_{B_k} \lambda_k,$$

where we recall that λ_k denotes the Lebesgue measure on $\mathbb{R}^k$. If $f : B_k \rightarrow \mathbb{R}$ is integrable, then

$$\frac{1}{\lambda_k[B_k]} \int_{B_k} f(\mathbf{x}) \lambda_k(d\mathbf{x}) = \mathbb{E}[f(\mathbf{X})].$$

Suppose that $\mathbf{X}_n = (X_{n,1}, \dots, X_{n,k})$, $n \in \mathbb{N}$, is a sequence of i.i.d. random vectors uniformly distributed in B_k , then the sequence of random variables $(f(\mathbf{X}_n))_{n \in \mathbb{N}}$ is i.i.d., with the same distribution as $f(\mathbf{X})$. The SLLN implies that the sequence random variables

$$Z_n = \frac{1}{n} (f(\mathbf{X}_1) + \cdots + f(\mathbf{X}_n))$$

converges a.s. to

$$\frac{1}{\lambda_k[B_k]} \int_{B_k} f(\mathbf{x}) \lambda_k(d\mathbf{x}).$$

This fact can be used to produce approximations to integrals using probabilistic methods. When the dimension k is large these methods are, to this day, the only viable methods for approximating integrals of functions of many variables.

In Example A.3.19 we describe a computer implementation of this strategy using the programming language R. □

2.1.3. Entropy and compression. Let us describe a surprising application of the law of large numbers. Suppose that we are given a finite set $\mathfrak{X}$ equipped with a probability measure $\mathbb{P}$ defined by the function $p : \mathfrak{X} \rightarrow [0, 1]$

$$p(x) := \mathbb{P}[\{x\}].$$

We will refer to the pair $(\mathfrak{X}, p)$ as *alphabet*.

Example 2.1.17. A good example to have in mind is the “alphabet” of the English language. In this alphabet we throw in not just the letters, but also the punctuation signs and the blank space. The elements x_i are letters/symbols of the alphabet. The probabilities $p(x_i)$ can be viewed as the frequency of the symbol x_i in the written texts. One way to estimate these frequencies³ is to count the number of their occurrences in a large text, say Moby Dick.

Another good example is the alphabet $\{0, 1\}$ used in computer languages. The frequencies $p(0) = p(1) = \frac{1}{2}$. □

For a letter x_i of the alphabet we define the “surprise” or “information” contained in the letter x_i to be the quantity

$$S(x_i) := -\log_2 p(x_i).$$

³As a curiosity, the letter “e” is the most frequent letter of the English language; it appears 13% of the time in large texts. It is for this reason that it has the simplest Morse code, a dot.

The base 2 of the logarithm is the convention used in information theory and we will stick with it. The unit of measure of surprise/information is the *bit*. Note that $S(x_i) \in [0, \infty]$. Observe that the less likely the letter x_i , the bigger the surprise. The *Shanon entropy* or the *information entropy* of the alphabet is the quantity

$$\text{Ent}_2 [p] := \mathbb{E}_p [S] := - \sum_{x \in \mathfrak{X}} p(x) \log_2 p(x), \quad (2.1.21)$$

where we adhere to the convention $0 \cdot \log 0 = 0$. Thus the entropy is the expected “surprise” of the alphabet. For example, if an urn contains 99 black balls and only one white ball we would be extremely surprised if we randomly draw a ball from the urn it turns out to be the white one. The average amount of surprise in this case is

$$-0.99 \log_2(0.99) - 0.01 \log_2(0.01) \approx 0.08.$$

If p_0 is the uniform probability measure on $\mathfrak{X}$, then

$$\text{Ent}_2 [p_0] = \log_2 |\mathfrak{X}|.$$

Let $m := |\mathfrak{X}|$. Note that $\text{Prob}(\mathfrak{X})$ can be identified with the $(m-1)$ -dimensional simplex

$$\Delta_m = \{ p = (p_1, \dots, p_m) \in [0, \infty)^m; p_1 + \dots + p_m = 1 \}.$$

We can view the entropy as a function $\text{Ent}_2 : \Delta_{m-1} \rightarrow [0, \infty)$. One can check that it is concave since the function $[0, \infty) \ni x \mapsto f(x) = -x \log_2 x$ is strictly concave. We have

$$\text{Ent}_2 [p] = \sum_{i=1}^m f(p_i).$$

Jensen’s inequality shows that

$$\frac{1}{m} \sum_{i=1}^m f(p_i) \leq f\left(\frac{1}{m} \sum_{i=1}^m p_i\right) = f(1/m) = \frac{\log_2 m}{m},$$

with equality if and only if $p_1 = \dots = p_m = \frac{1}{m}$. We deduce

$$\text{Ent}_2 [p] \leq \log_2 |\mathfrak{X}|, \quad \forall p \in \text{Prob}(\mathfrak{X}), \quad (2.1.22)$$

with equality if and only if p is the uniform probability measure. We will see later that the above is a special case of the Gibbs’ inequality (2.3.9). Intuitively, this inequality says that among all the probability measures on a finite set, the uniform one is the the most “chaotic”, the least “predictable”.

We will refer to the elements of $\mathfrak{X}^n$ as *words* of length n . The term “word” is a bit misleading. For example, when $\mathfrak{X}$ is the English alphabet as above, an element of $\mathfrak{X}^n$ with large n can be thought of as the sequence of symbols appearing in a large text. On the other hand, we can think of $\mathfrak{X}^n$ itself as a new alphabet with frequencies

$$p_n(x_1, \dots, x_n) = p(x_1) \cdots p(x_n).$$

The amount of “surprise” of a word $(x_1, \dots, x_n)$ is

$$S(x_1, \dots, x_n) = \sum_{k=1}^n S(x_k).$$

The entropy of $(\mathfrak{X}^n, p_n)$ is

$$\text{Ent}_2 [p_n] = n \text{Ent}_2 [p].$$

We denote by $\mathfrak{X}^*$ the disjoint union of the sets $\mathfrak{X}^n$,

$$\mathfrak{X}^* = \bigsqcup_{n \in \mathbb{N}} \mathfrak{X}^n,$$

and we will refer to it as the *vocabulary* of the alphabet $\mathfrak{X}$

Fix an alphabet $(\mathfrak{X}, p)$. We want to describe an efficient way of encoding the words in $\mathfrak{X}^n$ by words in the vocabulary of the binary alphabet $\mathcal{B} := \{0, 1\}$. Thus, we want to construct a code map $\mathcal{C} : \mathfrak{X}^n \rightarrow \mathcal{B}^*$ such that the words $x \in \mathfrak{X}^n$ with high frequency are encoded by words in $\mathcal{B}^*$ of short length. Normally we would require that $\mathcal{C}$ be injective but we are willing to sacrifice precision a bit for the sake of efficiency. We would be happy if the probability that two different words have the same code is very small, i.e., the event

$$x, x' \in \mathfrak{X}^n, \quad x \neq x' \quad \text{and} \quad \mathcal{C}(x) = \mathcal{C}(x')$$

has a very small probability.

Definition 2.1.18. Let $\varepsilon > 0$. The ε -typical set $A_\varepsilon^{(n)}$ with respect to $p(x)$ is the set $A_\varepsilon^{(n)} \subset \mathfrak{X}^n$ consisting of words $(x_1, x_2, \dots, x_n)$ with the property

$$2^{-n(\text{Ent}_2[p] + \varepsilon)} \leq p(x_1, x_2, \dots, x_n) \leq 2^{-n(\text{Ent}_2[p] - \varepsilon)}. \quad (2.1.23)$$

□

Theorem 2.1.19 (Asymptotic Equipartition Property). *For any $\varepsilon > 0$ there exists $N = N(\varepsilon)$ such that for any $n > N(\varepsilon)$, the following hold.*

- (i) $p_n[A_\varepsilon^{(n)}] > 1 - \varepsilon$.
- (ii) $|A_\varepsilon^{(n)}| \leq 2^{n(\text{Ent}_2[p] + \varepsilon)}$.
- (iii) $|A_\varepsilon^{(n)}| \geq (1 - \varepsilon)2^{n(\text{Ent}_2[p] - \varepsilon)}$.

Proof. We sample $(\mathfrak{X}, p)$ according to the frequencies $p(x_k)$ and we obtain a sequence $(X_n)_{n \in \mathbb{N}}$ of i.i.d. $\mathfrak{X}$ -valued random variables distributed according to p . We obtain random words $(X_1, \dots, X_n)$, $n \in \mathbb{N}$. The average amount of surprise per letter in this word is

$$\frac{1}{n} S(X_1, \dots, X_n) = \frac{1}{n} \sum_{k=1}^n S(X_k).$$

The law of large numbers shows that the random variables $\frac{1}{n} S(X_1, \dots, X_n)$ converge in probability to $\text{Ent}_2[p]$. Now observe that

$$(X_1, \dots, X_n) \in A_\varepsilon^{(n)} \iff \text{Ent}_2[p] - \varepsilon \leq \frac{1}{n} \sum_{k=1}^n S(X_k) \leq \text{Ent}_2[p] + \varepsilon$$

so

$$\mathbb{P}_n[A_\varepsilon^{(n)}] = \mathbb{P}\left[\text{Ent}_2[p] - \varepsilon \leq \frac{1}{n} \sum_{k=1}^n S(X_k) \leq \text{Ent}_2[p] + \varepsilon\right] \rightarrow 1$$

as $n \rightarrow \infty$. Fix $N = N(\varepsilon)$ such that

$$\mathbb{P}_n[A_\varepsilon^{(n)}] > 1 - \varepsilon, \quad \forall n > N(\varepsilon).$$

Note that for $n > N(\varepsilon)$

$$1 = \sum_{x \in \mathcal{X}^n} p(x) \geq \sum_{x \in A_\varepsilon^{(n)}} p_n(x) \geq 2^{-n(\text{Ent}_2[p] + \varepsilon)} |A_\varepsilon^{(n)}|,$$

and thus we have

$$|A_\varepsilon^{(n)}| \leq 2^{n(\text{Ent}_2[p] + \varepsilon)}.$$

Finally, for $n > N(\varepsilon)$ we have

$$1 - \varepsilon < \mathbb{P}_n[A_\varepsilon^{(n)}] \leq \sum_{x \in A_\varepsilon^{(n)}} 2^{-n(\text{Ent}_2[p] - \varepsilon)} = 2^{-n(\text{Ent}_2[p] - \varepsilon)} |A_\varepsilon^{(n)}|,$$

and conclude that $|A_\varepsilon^{(n)}| \geq (1 - \varepsilon)2^{n(\text{Ent}_2[p] - \varepsilon)}$. $\square$

The *Asymptotic Equipartition Property* (or AEP) shows that a typical set has probability nearly 1, all its elements are nearly equiprobable, and its cardinality is nearly $2^{n \text{Ent}_2[p]}$. The inequality (2.1.22) shows that if p is not the uniform probability measure on $\mathcal{X}$, then

$$2^{\text{Ent}_2[p]} \ll |\mathcal{X}|.$$

Hence, if $\varepsilon > 0$ is sufficiently small, then

$$\frac{|A_\varepsilon^{(n)}|}{|\mathcal{X}^n|} \rightarrow 0$$

exponentially fast as $n \rightarrow \infty$. That is, the typical sets have high probability and are “extremely small” if the entropy is small.

This suggests the following coding procedure. Fix $\varepsilon > 0$ so that $1 - \varepsilon$ will be our confidence level. For $n > N(\varepsilon)$ the set $A_\varepsilon^{(n)}$ has about 2^L elements where $L = \lceil n \text{Ent}_2[p] \rceil$ elements and thus we can find an injection

$$\mathcal{J} : A_\varepsilon^{(n)} \rightarrow \mathcal{B}^L.$$

For $x \in A_\varepsilon^{(n)}$ we attach the symbol 1 at the beginning of the word $\mathcal{J}(x) \in \mathcal{B}^L$ and the resulting word in $\mathcal{B}^{L+1}$ will encode x . It uses $L + 1$ bits. The first bit is 1 and indicates that the word x is typical.

We are less careful with the atypical words. Choose *any* map

$$\mathcal{J} : \mathcal{X}^n \setminus A_\varepsilon^{(n)} \rightarrow \mathcal{B}^L$$

and we encode an atypical word x using the binary word $\mathcal{J}(x)$ with a prefix 0 attached to indicate that it is atypical. The resulting map $\mathcal{C} : \mathcal{X}^n \rightarrow \mathcal{B}^{L+1}$ is not injective, but if two words have the same code, they must be atypical and thus occur with very small frequency. This is an example of *compression*.

Take for example the English language. There are various estimates for its entropy, starting with the pioneering work of Claude Shannon. Most recent ones⁴ vary from 1 to 1.5 bits. How do we encode efficiently texts consisting of $n = 10^6$ symbols say? For example, the book “*Moby Dick*” has 206,052 words and the average length of an English word is 5 letters so “*Moby Dick*” consists of about 1.03 million symbols.

Forgetting capitalization and punctuation there are 26^n such texts and a brute encoding would require 26^n codewords to cover all the possibilities. The above result however says that roughly

⁴A Google search with the keywords “entropy of the English language” will provide many more details on this subject.

$2^{1.5n}$ texts suffice to capture nearly surely almost everything. The term compression is fully justified since this is a much smaller fraction of the total number of possible texts. Also we only need codewords of lengths 1.5 million. Thus we need is roughly 1.5 gigabits to encode such a text. If the letters of the alphabet were uniformly distributed in human texts⁵ then the entropy would be $\log_2(26) \approx 4.70 > 3 \times 1.5$ and we would need more than three times amount of memory to store it.

Remark 2.1.20. The story does not end here and much more precise results are available. To describe some of them note first that for any alphabet $\mathfrak{X}$ there is an obvious operation of concatenation

$$* : \mathfrak{X}^m \times \mathfrak{X}^n \rightarrow \mathfrak{X}^{m+n}, \quad (x, x') \mapsto x * x'$$

where the word $x * x'$ is obtained by writing in succession the word x followed by x' . Note that this code uses on average $\frac{L+1}{n} \approx \text{Ent}_2[p]$ bits per symbol in a word. This is an example of compression.

A binary code for the alphabet $(\mathfrak{X}, p)$ is an injection

$$C : \mathfrak{X} \rightarrow \mathcal{B}^*$$

For each $x \in \mathfrak{X}$ we denote by $L_C(x)$ the length of the code word $C(x)$. The expected length of a codeword is

$$\ell_C := \mathbb{E}[L_C] = \sum_{x \in \mathfrak{X}} L_C(x)p(x).$$

Note that C extends to a map

$$C^* : \mathfrak{X}^* \rightarrow \mathcal{B}^*, \quad C^*(x_1, \dots, x_n) = C(x_1) * \dots * C(x_n).$$

The code C is called *uniquely decodable* if its extension $C^* : \mathfrak{X}^* \rightarrow \mathcal{B}^*$ is also injective.

An important subclass of uniquely decodable codes are *instantaneous codes*. A code C is called *instantaneous* if no codeword is a prefix of some other code word. E.g., if one of the codewords is 10, then no other codeword can begin with 10.

Here is a very revealing example. Consider an alphabet $\mathcal{A}$ consisting of four letters $\mathcal{A} := \{a, b, c, d\}$ with frequencies

$$p_a = 1/2, \quad p_b = 1/3, \quad p_c = p_d = 1/12. \quad (2.1.24)$$

Consider the following instantaneous code

$$a \rightarrow 1, \quad b \rightarrow 01, \quad c \rightarrow 001 \quad d \rightarrow 000.$$

The expected code length is

$$\frac{1}{2} + \frac{2}{3} + \frac{3}{12} + \frac{3}{12} = \frac{5}{3} \approx 1.666$$

The entropy of alphabet is

$$\text{Ent}_2[\mathcal{A}] = \frac{\log 2}{2} + \frac{\log 3}{3} + \frac{\log 12}{6} \approx 1.625$$

Kraft's inequality shows that for any uniquely decodable code C we have

$$\ell_C \geq \text{Ent}_2[\mathcal{A}].$$

Moreover, there exist optimal codes C such that

$$\ell_C \leq \text{Ent}_2[\mathcal{A}] + 1.$$

⁵The famous monkey on a typewriter produces texts where the letters are uniformly distributed, but we can safely call the resulting texts highly atypical of the English texts humans are used to.

Such codes are called *Shannon codes*. The above code is a Shannon code. In fact it is a special example of the famous *Huffman code*, [47].

Let us discuss a particularly suggestive experiment that highlights a defining feature of Huffman codes and reveals one interpretation of the entropy of an alphabet.

Suppose we have an urn containing the letters a, b, c, d , in proportions (2.1.24). A person randomly draws a letter from the urn and you are supposed to guess what it is by asking YES/NO question. Think YES = 1, NO = 0. The above code describes an optimal guessing strategy. Here it is.

- (1) Ask first if the letter is $a \rightarrow 1$. If the answer is YES (= 1), the game is over. The game has length 1 with probability $1/2$.
- (01) If the answer is NO (= 0) the letter can only be b, c or d . Ask if the letter is $b \rightarrow 01$. If the answer is YES (= 1) the game is over. The game has length 2 with probability $1/3$
- (001) If the answer is NO (= 0) ask if the letter is $c \rightarrow 001$. The game has length 3 with probability $1/6$.

The average number of questions is then

$$\frac{1}{2} + \frac{2}{3} + \frac{3}{6} \approx 1.66.$$

This is not too far off $\text{Ent}_2[\mathcal{A}]$.

For more details about information theory and its application we refer to [47, 144]. For a more informal introduction to information theory we refer to [77]. The eminently readable [90] contains historical perspective on the evolution of information theory. Kolmogorov's brief but very rich in intuition survey [120] is a good place to start learning about the mathematical theory of information. $\square$

2.2. The Central Limit Theorem

The goal of this section is to prove a striking classical result that adds additional information to the Law of Large numbers.




Figure 2.2. Visualizing the Central Limit Theorem. The appropriate Gauss bell hugs closely the histogram of $\text{Bin}(n, p)$ for n large.

Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables with mean μ and *finite* variance σ^2 . Note that the sum $S_n := X_1 + \cdots + X_n$ has mean $n\mu$ and variance $n\sigma^2$. Loosely speaking, the central limit theorem states that for large n the probability distribution of S_n “resembles” very much a Gaussian with the same mean and variance.

For example, if the X_n -s are Bernoulli random variables with success probability p , then $\mu = p$, $\sigma^2 = pq$ and $S_n \sim \text{Bin}(n, p)$. In Figure 2.2 considered the case $p = 0.3$ and $n = 63$ we have illustrated the histogram of $\text{Bin}(n, p)$ and the Gauss bell centered at np with variance npq .

The vertical lines depict the probability mass function of the binomial distribution while the curve wrapping them is the Gaussian with the same mean and variance. They obviously do “resemble”. However, we need to define precisely what we mean by “resemble”.

2.2.1. Weak and vague convergence. Let (X, d) be a metric space. We will use the following notation.

- $\mathcal{B}_X$ denotes the Borel sigma-algebra of X ,
- $\text{Meas}(X)$ denotes the set of *finite* Borel measures on X ,
- $\text{Prob}(X) \subset \text{Meas}(X)$ denotes the space of Borel probability measures on X , and
- $\text{Prob}_s(X)$ denotes the space of subprobability measures⁶ on X , i.e., Borel measures μ on X such that $\mu[X] \leq 1$.

We denote by $C_{\text{cpt}}(X)$ the space of continuous functions $X \rightarrow \mathbb{R}$ with compact support and by $C_b(X)$ the space of bounded continuous functions $X \rightarrow \mathbb{R}$. This is a Banach space with respect to the sup-norm

$$\|f\|_\infty := \sup_{x \in X} |f(x)|.$$

For any $f \in C_b(X)$ and $\mu \in \text{Meas}(X)$ we set

$$\mu[f] := \int_X f(x) \mu[dx] < \infty.$$

Definition 2.2.1. Consider a sequence $(\mu_n)_{n \in \mathbb{N}}$ of finite Borel measures on X .

- (i) We say that the sequence (μ_n) *converges vaguely* to $\mu \in \text{Meas}(X)$, and we write this $\mu_n \dashrightarrow \mu$ if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} f(x) \mu_n[dx] = \int_{\mathbb{R}} f(x) \mu[dx], \quad \forall f \in C_{\text{cpt}}(X). \quad (2.2.1)$$

- (ii) We say that the sequence (μ_n) *converges weakly* to $\mu \in \text{Meas}(X)$, and we write this $\mu_n \Rightarrow \mu$ if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} f(x) \mu_n[dx] = \int_{\mathbb{R}} f(x) \mu[dx], \quad \forall f \in C_b(X). \quad (2.2.2)$$

- (iii) A sequence of measurable maps $\Phi_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (X, \mathcal{B}_X)$ is said to *converge in law* or *in distribution* to the random map $\Phi : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (X, \mathcal{B}_X)$ if

$$\mathbb{P}_{\Phi_n} \Rightarrow \mathbb{P}_\Phi \text{ in } \text{Prob}(X),$$

i.e.,

$$\lim_{n \rightarrow \infty} \mathbb{E}[f(\Phi_n)] = \mathbb{E}[f(\Phi)], \quad \forall f \in C_b(X). \quad (2.2.3)$$

⁶Some authors refer to subprobability measures as *defective distributions*.

We will use the notation $\Phi_n \xrightarrow{d} \Phi$ or $\Phi_n \Rightarrow \Phi$ to indicate that Φ_n converges to Φ in distribution. $\square$

Remark 2.2.2. The weak convergence of Borel probability measures on Polish space $(\mathbf{X}, d)$ admits an surprising characterization due to I.I. Skorokhod. More precisely, *Skorokhod's representation theorem* a sequence of Borel probability measures $\mu_n \in \text{Prob}(\mathbf{X})$, $n \in \mathbb{N}$, converges weakly to the Borel probability measure $\mu_\infty \in \text{Prob}(\mathbf{X})$ if and only if there exists a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and Borel measurable maps $X_n : \Omega \rightarrow \mathbf{X}$, $n \in \mathbb{N} \cup \{\infty\}$ such that $\mathbb{P}_{X_n} = \mu_n$, $\forall n \in \mathbb{N} \cup \{\infty\}$ and $d(X_n, X_\infty) \rightarrow 0$, $\mathbb{P}$ -a.s..

For a proof and more details we refer to [18, Thm. 6.7] or [64, Thm. 11.7.2]. Exercise 2.24 asks you to prove a refined version of this theorem in the special case $\mathbf{X} = \mathbb{R}$. $\square$

Theorem 2.2.3 (Mapping theorem). *Suppose that $(\mathbf{X}, d_X)$ and $(\mathbf{Y}, d_Y)$ are two metric spaces and $F : \mathbf{X} \rightarrow \mathbf{Y}$ is a continuous map.*

Let $\Phi_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbf{X}, \mathcal{B}_\mathbf{X})$, $n \in \mathbb{N}$ be a sequence of random map converging in distribution to the random map Φ . Then the sequence of random maps $\Psi_n = F(\Phi_n)$ converges in distribution to $\Psi = F(\Phi)$.

Proof. Let $f \in C_b(\mathbf{Y})$. Then $f \circ F \in C_b(\mathbf{X})$ and

$$\mathbb{E}[f(\Psi_n)] = \mathbb{E}[f \circ F(\Phi_n)] \rightarrow \mathbb{E}[f \circ F(\Phi)] = \mathbb{E}[f(\Psi)].$$

$\square$

Definition 2.2.4. A collection $\mathcal{F} \subset C_b(X)$ is called *separating* if given $\mu_0, \mu_1 \in \text{Meas}(X)$ such that $\mu_0[f] = \mu_1[f]$, $\forall f \in \mathcal{F}$, then $\mu_0 = \mu_1$. $\square$

As shown in Proposition 1.2.63, the collection $C_b(X)$ is separating so the above definition is not vacuous for any metric space.

Lemma 2.2.5. *Let $k \in \mathbb{N}$. The collection $C_{\text{cpt}}(\mathbb{R}^k)$ is separating.*

Proof. Let $\beta_n : [0, \infty) \rightarrow [0, 1]$ be the unique continuous piecewise linear function such that

$$\beta_n(r) = \begin{cases} 1, & r \leq n, \\ 0, & r \geq n+1. \end{cases}$$

Consider the functions $\varphi_n : \mathbb{R}^k \rightarrow \mathbb{R}$, $\varphi_n(\mathbf{x}) = \beta_n(\|\mathbf{x}\|)$. Then $\varphi_n \in C_{\text{cpt}}^0(\mathbb{R}^k)$, $\varphi_n \geq 0$ and $\varphi_n \nearrow 1$. The Monotone Convergence Theorem implies

$$\mu_0[\mathbb{R}^k] = \mu_0[1] = \lim_{n \rightarrow \infty} \mu_0[\varphi_n] = \lim_{n \rightarrow \infty} \mu_1[\varphi_n] = \mu_1[1] = \mu_1[\mathbb{R}^k].$$

According to Proposition 1.2.4 it suffices so show that for any compact subset $K \subset \mathbb{R}^n$ we have $\mu_0[K] = \mu_1[K]$. Set

$$S_n := \{x \in \mathbb{R}^k; \text{dist}(x, K) \geq 1/n\}.$$

For $n \in \mathbb{N}$ define $f_n : \mathbb{R} \rightarrow [0, 1]$

$$f_n(x) = \frac{\text{dist}(x, S_n)}{\text{dist}(x, K) + \text{dist}(x, S_n)}.$$

Observe that f_n is continuous, and $f_n|_{S_n=0}$, so f_n has compact support. Moreover and

$$\lim_{n \rightarrow \infty} f_n(x) = \mathbf{I}_K \text{ a.s..}$$

The Dominated Convergence Theorem implies that

$$\int_{\mathbb{R}} \mathbf{I}_K(x) \mu_0[dx] = \lim_{n \rightarrow \infty} \int_{\mathbb{R}} f_n(x) \mu_0[dx] = \lim_{n \rightarrow \infty} \int_{\mathbb{R}} f_n(x) \mu_1[dx] = \int_{\mathbb{R}} \mathbf{I}_K \mu_1[dx].$$

□

Lemma 2.2.5 shows a sequence of Borel probability measures on $\mathbb{R}^k$ has at most one vague limit, i.e., if $\mu_n \dashrightarrow$ and $\mu_n \dashrightarrow \mu'$, then $\mu = \mu'$.

Proposition 2.2.6. *Let $\mu_n \in \text{Prob}(\mathbb{R}^k)$, $n \in \mathbb{N}$ be a sequence of probability measures.*

- (i) *If μ_n converge vaguely to a measure $\mu \in \text{Meas}(\mathbb{R}^k)$, then μ is a subprobability measure.*
- (ii) *If μ_n converge weakly to a measure $\mu \in \text{Meas}(\mathbb{R}^k)$, then μ is a probability measure.*

Proof. (i) For each $\varepsilon > 0$ fix a radius R_ε such that $\mu[\mathbb{R}^k \setminus \bar{B}_{R_\varepsilon}(0)] \leq \varepsilon$.

Consider the continuous function $\varphi : \mathbb{R} \rightarrow [0, 1]$ uniquely determined by the requirements

$$\varphi(t) = \begin{cases} 1, & t \leq 0, \\ 0, & t \geq 1, \\ 1 - t, & t \in [0, 1]. \end{cases}$$

We set $\varphi_R(t) = \varphi(t - R)$ and define

$$\eta_R : \mathbb{R}^k \rightarrow [0, 1], \quad \eta_R(x) = \varphi_R(|x|) = \varphi(|x| - R).$$

Note that η_R is supported in $\bar{B}_{R+1}(0)$ and $\mathbf{I}_{B_R} \leq \eta_R \leq 1$. We have $\mu_n[\eta_{R_\varepsilon}] \leq 1, \forall n \in \mathbb{N}$. Letting $n \rightarrow \infty$ we deduce

$$\mu[\mathbb{R}^k] - \varepsilon \leq \mu[\bar{B}_{R_\varepsilon}] \leq \mu[\eta_{R_\varepsilon}] \leq 1, \quad \forall \varepsilon > 0.$$

This proves $\mu[\mathbb{R}^k] \leq 1$.

(ii) We have

$$\mu[\mathbb{R}^k] = \mu[1] = \lim_{n \rightarrow \infty} \mu_n[1] = 1.$$

□

Example 2.2.7. Let

$$\mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{k/n}.$$

Then

$$\mu_n \Rightarrow \mu = \mathbf{I}_{[0,1]}(x)dx \sim \text{Unif}(0, 1).$$

Indeed, if $f \in C_b(\mathbb{R})$, then

$$\int_{\mathbb{R}} f(x) \mu_n[dx] := \frac{1}{n} \sum_{k=1}^n f(k/n).$$

The sum in the right-hand-side of the above equality is a Riemann sum for f corresponding to the uniform partition

$$0 < \frac{1}{n} < \frac{2}{n} < \dots < \frac{n-1}{n} < 1.$$

Since f is Riemann integrable we deduce

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(k/n) = \int_0^1 f(x) dx = \int_{\mathbb{R}} f(x) \mu[dx].$$

□

Example 2.2.8. There exist vaguely convergent sequences of Borel probability measures on $\mathbb{R}$ that are not weakly convergent. Take for example $\mu_n = \delta_n$, $n \in \mathbb{N}$. Then $\mu_n \dashrightarrow 0$ yet μ_n does not converge weakly to 0 since $\mu_n[\mathbb{R}] = 1, \forall n$. □

Proposition 2.2.9. *If the random variables X_n converge in probability to X , then they also converge in law to X . In particular, if X_n converge in p -mean to X , then they also converge in law to X .*

Proof. We deduce from Corollary 1.3.62 that for any $f \in C_b(\mathbb{R})$ the random variables $f(X_n)$ converge in probability to $f(X)$. The Bounded Convergence Theorem implies

$$\lim_{n \rightarrow \infty} \mathbb{E}[f(X_n)] = \mathbb{E}[f(X)], \quad \forall f \in C_b(\mathbb{R}).$$

□

Example 2.2.10. Fix a standard normal random variable X . Then $\mathbb{P}_X = \mathbb{P}_{-X}$ so $-X$ is a standard normal random variable as well. Consider the constant sequence

$$X_n = X, \quad n \in \mathbb{N}.$$

Then $\mathbb{P}_{X_n} \Rightarrow \mathbb{P}_{-X}$, but X_n does not converge to $-X$ in probability. □

Theorem 2.2.11 (Portmanteau⁷ theorem). *Let $(\mathbf{X}, d)$ be a metric space and $\mu_n \in \text{Prob}(\mathbf{X})$, $n \in \mathbb{N}$, be a sequence of Borel probability measures on $\mathbf{X}$. The following statements are equivalent.*

- (i) *The sequence $(\mu_n)_{n \in \mathbb{N}}$ converges weakly to $\mu \in \text{Meas}(\mathbf{X})$.*
- (ii) *For any bounded and Lipschitz continuous function $f : \mathbf{X} \rightarrow \mathbb{R}$ we have*

$$\mu[f] = \lim_{n \rightarrow \infty} \mu_n[f].$$

- (iii) *For any closed set $C \subset \mathbf{X}$ we have*

$$\mu[C] \geq \limsup \mu_n[C].$$

- (iv) *For any open set $U \subset \mathbf{X}$ we have*

$$\mu[U] \leq \liminf \mu_n[U].$$

- (v) *For any Borel set $B \subset \mathbf{X}$ such that $\mu[\partial B] = 0$ we have*

$$\mu[B] = \lim_{n \rightarrow \infty} \mu_n[B].$$

Proof. This proof is inspired from [18, Sec.2, Thm.2.2]. Clearly (i) $\Rightarrow$ (ii).

- (ii) $\Rightarrow$ (iii) Let C be a closed set. For $\varepsilon > 0$ we set

$$C_\varepsilon = \{x \in \mathbf{X}; \text{dist}(x, C) \leq \varepsilon\}.$$

⁷portmanteau = large trunk or suitcase

The set C_ε is closed and contains $C = C_0$. Let

$$f_\varepsilon : \mathbf{X} \rightarrow \mathbb{R}, \quad f_\varepsilon(x) = \max(1 - \varepsilon^{-1} \text{dist}(x, C), 0).$$

The function f_ε is bounded and Lipschitz continuous. Moreover,

$$\mathbf{I}_C \leq f_\varepsilon \leq \mathbf{I}_{C_\varepsilon}.$$

Hence,

$$\mu_n[C] \leq \mu_n[f_\varepsilon].$$

Letting $n \rightarrow \infty$ we deduce that

$$\limsup \mu_n[C] \leq \mu[f_\varepsilon] \leq \mu[C_\varepsilon], \quad \forall \varepsilon > 0.$$

The desired conclusion follows from the fact that

$$\mu[C] = \lim_{\varepsilon \searrow 0} \mu[C_\varepsilon].$$

(iii) $\iff$ (iv) Follows from the following facts

- The set U is open iff U^c is closed
- For any Borel set $B \subset \mathbb{R}$, $\mu[B^c] = 1 - \mu[B]$.

(iii) + (iv) $\Rightarrow$ (v). Let $B \subset \mathbf{X}$ be a Borel set such that $\mu[\partial B] = 0$. Denote by U the interior of B and by C its closure so that $\partial B = C \setminus U$. We deduce

$$\mu[B] = \mu[C] = \mu[U].$$

Thus

$$\limsup \mu_n[C] \leq \mu[C] = \mu[B] = \mu[U] \leq \liminf \mu_n[U].$$

Since ∂B is closed we deduce

$$\limsup \mu_n[\partial B] \leq \mu[\partial B] = 0.$$

Hence

$$\mu_n[U] = \mu_n[C] + \mu_n[\partial B], \quad \lim_n \mu_n[\partial B] = 0,$$

so

$$\liminf \mu_n[U] = \liminf \mu_n[C].$$

Hence

$$\lim_n \mu_n[C] = \mu[C], \quad \lim_n \mu_n[B] = \lim_n \mu_n[C] + \lim_n \mu_n[\partial B] = \mu[B].$$

(iv) $\Rightarrow$ (i). Clearly it suffices to show that $\mu_n[f] \rightarrow \mu[f]$, for any *nonnegative*, bounded, continuous function f on $\mathbf{X}$.

Suppose that f be such a function. Set $K := \sup f$. For any $\nu \in \text{Prob}(\mathbf{X})$ we can regard f as a random variable $(\mathbf{X}, \mathcal{B}_\mathbf{X}, \nu) \rightarrow [0, K]$. The integral $\nu[f]$ is then the expectation of this random variable.

Using Proposition 1.3.43 with $p = 1$ we deduce that

$$\nu[f] = \mathbb{E}_\nu[f] = \int_0^K \nu[f > t] dt.$$

Note that

$$\nu[f = t] = 0 \Rightarrow \nu[\partial\{f > t\}] = 0.$$

Observe next that for any $n \in \mathbb{N}$ we have

$$\#\{t \in \mathbb{R}; \nu[f = t] \geq 1/n\} \leq n,$$

so, for any $\nu \in \text{Prob}(\mathbf{X})$ the set

$$\{t \in \mathbb{R}; \nu[f = t] > 0\}$$

is at most countable. We deduce from (iv) that

$$\lim_{n \rightarrow \infty} \mu_n[f > t] = \mu[f > t] \text{ for (Lebesgue) almost any } t.$$

The Dominated Convergence Theorem implies

$$\lim_{n \rightarrow \infty} \mu_n[f] = \lim_{n \rightarrow \infty} \int_0^K \mu_n[f > t] dt = \int_0^K \mu[f > t] dt = \mu[f].$$

□

Corollary 2.2.12. *Let X_n , $n \in \mathbb{N}$, be a sequence of random variables. Denote by $F_n(x)$ the cdf of X_n ,*

$$F_n(x) = \mathbb{P}[X_n \leq x], \quad x \in \mathbb{R}.$$

The following statements are equivalent.

- (i) *The random variables X_n converge in law to the random variable X .*
- (ii) *If $F(x)$ is the cdf of X , then*

$$\lim_{n \rightarrow \infty} F_n(x) = F(x),$$

for any point of continuity x of F .

Proof. Set $\mu_n := \mathbb{P}_{X_n}$, $\mu := \mathbb{P}_X$. The condition (ii) is a special case of condition (iv) of the Portmanteau Theorem so (i) $\Rightarrow$ (ii).

(ii) $\Rightarrow$ (i) Denote by $\mathfrak{X} \subset \mathbb{R}$ the set of points continuity of F . Note that its complement $\mathbb{R} \setminus \mathfrak{X}$ is at most countable so $\mathfrak{X}$ is dense. Note that for $a, b \in \mathfrak{X}$, $a < b$ we have

$$\mathbb{P}[a < X < b] = F(b) - F(a).$$

For any $a, b \in \mathbb{R}$, $a < b$ and any $\varepsilon > 0$ there exist $a_\varepsilon, b_\varepsilon \in \mathfrak{X}$, $a < a_\varepsilon < b_\varepsilon < b$ such that

$$F(b_\varepsilon) - F(a_\varepsilon) = \mathbb{P}[a_\varepsilon < X < b_\varepsilon] > \mathbb{P}[a < X < b] - \varepsilon.$$

Hence

$$\lim_{n \rightarrow \infty} (F_n(b_\varepsilon) - F_n(a_\varepsilon)) = F(b_\varepsilon) - F(a_\varepsilon) > \mathbb{P}[a < X < b] - \varepsilon.$$

On the other hand

$$\mathbb{P}[a < X_n < b] \geq \mathbb{P}[a_\varepsilon < X_n < b_\varepsilon], \quad \forall n,$$

so that

$$\liminf_{n \rightarrow \infty} \mathbb{P}[a < X_n < b] \geq \mathbb{P}[a < X < b] - \varepsilon, \quad \forall \varepsilon > 0,$$

i.e.,

$$\liminf_{n \rightarrow \infty} \mathbb{P}[a < X_n < b] \geq \mathbb{P}[a < X < b], \quad \forall a < b \in \mathbb{R}.$$

Thus, the sequence μ_n satisfies the condition (ii) in the Portmanteau Theorem 2.2.11, where U is any open interval of the real axis. Since any open set of the real axis is a disjoint union of countably many open intervals, we deduce that condition (ii) in the Portmanteau Theorem is satisfied for *all* the open sets $U \subset \mathbb{R}$.

Indeed, suppose that

$$U = \bigcup_{k \geq 1} I_k$$

where I_k are pairwise disjoint open intervals. For $K \in \mathbb{N}$ we set

$$U_K := \bigcup_{1 \leq k \leq K} I_k.$$

Then

$$\mathbb{P}[X \in U_K] \leq \liminf_{n \rightarrow \infty} \mathbb{P}[X_n \in U_K] \leq \liminf_{n \rightarrow \infty} \mathbb{P}[X_n \in U], \quad \forall K \in \mathbb{N}.$$

Letting $K \rightarrow \infty$ we deduce the desired conclusion. $\square$

Theorem 2.2.13 (Slutsky). *Let $\mathbf{F}$ be a Banach space with norm $\|\cdot\|$. Suppose that $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ are sequences of $\mathbf{F}$ -valued random vectors such that (X_n) converges in distribution to X and Y_n converges in probability to $c \in \mathbf{F}$. Then the sum $X_n + Y_n$ converges in distribution to $X + c$.*

Proof. Without loss of generality we can assume $c = 0$. We follow the argument in [18, Chap. 1, Sec. 3]. Fix a closed subset $C \subset \mathbf{F}$. For $\varepsilon > 0$ set

$$C_\varepsilon := \{x \in \mathbf{F}; \text{dist}(x, C) \leq \varepsilon\}.$$

The set C_ε is closed and we have

$$\{X_n + Y_n \in C\} \subset \{\|Y_n\| > \varepsilon\} \cup \{X_n \in C_\varepsilon\}.$$

and thus

$$\mathbb{P}[X_n + Y_n \in C] \leq \mathbb{P}[\|Y_n\| > \varepsilon] + \mathbb{P}[X_n \in C_\varepsilon].$$

Letting $n \rightarrow \infty$ we deduce from the assumptions and the Portmanteau Theorem that

$$\limsup_{n \rightarrow \infty} \mathbb{P}[X_n + Y_n \in C] \leq \limsup_{n \rightarrow \infty} \mathbb{P}[X_n \in C_\varepsilon] \leq \mathbb{P}[X \in C_\varepsilon].$$

Now let $\varepsilon \searrow 0$ observing that $C_\varepsilon \searrow C$. $\square$

We can now formulate and prove the main convergence criterion of this subsection.

Theorem 2.2.14. *Suppose that $(\mu_n)_{n \in \mathbb{N}}$ is a sequence of nonzero finite Borel measures on $\mathbb{R}^k$ and $\mu \in \text{Meas}(\mathbb{R}^k)$. The following statements are equivalent.*

- (i) *The sequence (μ_n) converges weakly to $\mu \in \text{Meas}(\mathbb{R}^k)$.*
- (ii) *The sequence (μ_n) converges vaguely to μ and*

$$\lim_{n \rightarrow \infty} \mu_n[\mathbb{R}^k] = \mu[\mathbb{R}^k].$$

- (iii) *There exists a collection $\mathcal{F} \subset C_b(\mathbb{R}^k)$ whose closure in $C_b(\mathbb{R}^k)$ contains $C_{\text{cpt}}(\mathbb{R}^k)$ and such that*

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^k} f(x) \mu_n[dx] = \int_{\mathbb{R}^k} f(x) \mu[dx], \quad \forall f \in \mathcal{F}, \quad (2.2.4a)$$

$$\lim_{n \rightarrow \infty} \mu_n[\mathbb{R}^k] = \mu[\mathbb{R}^k]. \quad (2.2.4b)$$

Proof. In each of the statements (i)-(iii) we have

$$0 < C := \sup_n \mu_n[\mathbb{R}^k] < \infty.$$

Replacing the measures μ_n by $\frac{1}{C}\mu_n$ we can assume that all the measures μ_n are subprobability measures.

Obviously (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii). It suffices to prove that (ii) $\Rightarrow$ (i) and (iii) $\Rightarrow$ (ii). We will need the following result.

Lemma 2.2.15. *Any finite Borel measure $\mu \in \text{Meas}(\mathbb{R}^k)$ is on $\mathbb{R}^k$ is Radon, i.e., for any Borel set $B \subset \mathbb{R}^k$ and any $\varepsilon > 0$, there exists a compact set $K \subset B$ such that $\mu[B \setminus K] < \varepsilon$.*

Proof. Let $B \subset \mathbb{R}^k$ be a Borel set and $\varepsilon > 0$. According to Theorem 1.5.8, the measure μ is regular. Hence, there exists a closed set $C \subset B$ such that

$$\mu[B \setminus C] < \frac{\varepsilon}{2}.$$

On the other hand, we can find $R > 0$ sufficiently large such that

$$\mu[\bar{B}_R(0)] > \mu[\mathbb{R}^k] - \frac{\varepsilon}{2}.$$

We set $K := \bar{B}_R \cap C$. The set K is clearly compact, contained in C , and

$$\mu[C \setminus K] \leq \mu[\mathbb{R}^k \setminus \bar{B}_R(0)] < \frac{\varepsilon}{2}.$$

Thus $\mu[B \setminus K] = \mu[B \setminus C] + \mu[C \setminus K] < \varepsilon$. □

(ii) $\Rightarrow$ (i) We will show that the sequence (μ_n) satisfies the condition (ii) in the Portmanteau Theorem. Now let $U \subset \mathbb{R}^k$ be an open set and $\varepsilon > 0$. Lemma 2.2.15 shows that for any $\varepsilon > 0$ there exists a compact set $K \subset U$ such that $\mu[K] > \mu[U] - \varepsilon$.

Choose $r < \frac{1}{2} \text{dist}(K, U^c)$ and set

$$C_r := \{x \in \mathbb{R}^k; \text{dist}(x, K) \geq r\}.$$

The set C_r is closed and its complement

$$V_r := \{x \in \mathbb{R}^k; \text{dist}(x, K) < r\} \subset U$$

is precompact. Consider the continuous function

$$\varphi : \mathbb{R}^k \rightarrow [0, 1], \quad \varphi(x) = \frac{\text{dist}(x, C_r)}{\text{dist}(x, K) + \text{dist}(x, C_r)}.$$

Observe that it vanishes on C_r and thus it has compact support contained in U . Moreover, $\varphi = 1$ on K . Thus $I_K \leq \varphi \leq I_U$ so that

$$\mu_n[K] \leq \mu_n[\varphi] \leq \mu_n[U].$$

Letting $n \rightarrow \infty$ we deduce

$$\mu[U] - \varepsilon < \mu[K] \leq \mu[\varphi] = \lim_n \mu_n[\varphi] \leq \liminf_n \mu_n[U], \quad \forall \varepsilon > 0.$$

This establishes condition (ii) of the Portmanteau Theorem.

(iii) $\Rightarrow$ (ii) Let $\varphi \in C_{\text{cpt}}(\mathbb{R}^k)$. For any $\varepsilon > 0$ choose $f_\varepsilon \in \mathcal{F}$ such that $\|\varphi - f_\varepsilon\|_\infty < \frac{\varepsilon}{2}$. Then

$$|\nu[f_\varepsilon] - \nu[\varphi]| < \frac{\varepsilon}{2}, \quad \forall \nu \in \text{Prob}(\mathbb{R}^k).$$

We deduce

$$\limsup_{n \rightarrow \infty} |\mu[f_\varepsilon] - \mu_n[\varphi]| = \limsup_{n \rightarrow \infty} |\mu_n[f_\varepsilon] - \mu_n[\varphi]| \leq \frac{\varepsilon}{2}$$

On the other hand

$$|\mu[\varphi] - \mu[f_\varepsilon]| < \frac{\varepsilon}{2}.$$

so that, $\forall \varepsilon > 0$,

$$\limsup_{n \rightarrow \infty} |\mu[\varphi] - \mu_n[\varphi]| \leq |\mu[\varphi] - \mu[f_\varepsilon]| + \limsup_{n \rightarrow \infty} |\mu[f_\varepsilon] - \mu_n[\varphi]| < \varepsilon.$$

Hence

$$\lim_{n \rightarrow \infty} |\mu_n[\varphi] - \mu[\varphi]| = 0.$$

□

Corollary 2.2.16. *Consider a sequence $\mu_n \in \text{Prob}(\mathbb{R}^k)$ and $\mu \in \text{Meas}(\mathbb{R}^k)$. Then the following are equivalent.*

- (i) *The sequence (μ_n) converges weakly to μ .*
- (ii) *For any bounded Lipschitz function $f : \mathbb{R}^k \rightarrow \mathbb{R}$ we have*

$$\mu_n[f] = \mu[f].$$

Proof. The implication (i) $\Rightarrow$ (ii) is obvious. To prove that (ii) $\Rightarrow$ (i) observe first that any compactly supported continuous function can be uniformly approximated by compactly supported smooth functions⁸ so the closure in $C_b(\mathbb{R}^k)$ of the set of bounded Lipschitz functions contains $C_{\text{cpt}}(\mathbb{R}^k)$. The measure μ is a probability measure since the constant function $\mathbf{I}_{\mathbb{R}^k}$ is bounded and Lipschitz and thus

$$\mu[\mathbf{I}_{\mathbb{R}^k}] = \lim_{n \rightarrow \infty} \mu_n[\mathbf{I}_{\mathbb{R}^k}] = 1.$$

The conclusion now follows from Theorem 2.2.14. □

Corollary 2.2.17. *If a sequence $\mu_n \in \text{Prob}(\mathbb{R}^k)$ converges vaguely to a probability measure, then it also converges weakly.* □

Corollary 2.2.18. *Suppose that $(X_n)_{n \in \mathbb{N}}$ and X are random variables with ranges contained in $\mathbb{Z}$. Then $X_n \Rightarrow X$ if and only if*

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_n = k] = \mathbb{P}[X = k], \quad \forall k \in \mathbb{Z}. \quad (2.2.5)$$

Proof. The condition (2.2.5) is clearly satisfied if $X_n \Rightarrow X$ since

$$\begin{aligned} \mathbb{P}[X = k] &= \mathbb{P}[k - 1/2 < X \leq k + 1/2] \\ &= \lim_{n \rightarrow \infty} \mathbb{P}[k - 1/2 < X_n \leq k + 1/2] = \lim_{n \rightarrow \infty} \mathbb{P}[X_n = k]. \end{aligned}$$

Conversely, if (2.2.5) is satisfied, then $\forall \varphi \in C_{\text{cpt}}(\mathbb{R})$ the set $\mathbb{Z} \cap \text{supp } \varphi$ is finite and thus

$$\mathbb{E}[\varphi(X_n)] = \sum_{k \in \mathbb{Z} \cap \text{supp } \varphi} \varphi(k) \mathbb{P}[X_n = k] \rightarrow \sum_{k \in \mathbb{Z} \cap \text{supp } \varphi} \varphi(k) \mathbb{P}[X = k]$$

⁸On simple way to see this is to use the convolution trick.

$$= \mathbb{E}[\varphi(X)].$$

The conclusion now follows from Theorem 2.2.14. $\square$

Corollary 2.2.19. *The topology of weak convergence on $\text{Prob}(\mathbb{R}^k)$ is metrizable, i.e., there exists a metric d on $\text{Prob}(\mathbb{R}^k)$ such that*

$$\mu_n \Rightarrow \nu \iff d(\mu_n, \nu) \rightarrow 0.$$

Proof. Let

$$\mathcal{F} = \{f_0 = 1, f_1, \dots\}$$

be a countable subset on $C_b(\mathbb{R})$ whose closure contains $C_{\text{cpt}}(\mathbb{R}^n)$. Define

$$d : \text{Prob}(\mathbb{R}) \times \text{Prob}(\mathbb{R}) \rightarrow [0, \infty), \quad d(\mu, \nu) = \sum_{\ell \geq 0} \frac{1}{2^\ell} \max(|\mu[f_\ell] - \nu[f_\ell]|, 1).$$

According to Theorem 2.2.14 we have

$$\mu_n \Rightarrow \nu \iff \mu_n[f_\ell] \rightarrow \nu[f_\ell], \quad \forall \ell \geq 0.$$

$\square$

Remark 2.2.20. For any metric space $\mathbf{X}$ there exists a metric d_{LP} on $\text{Prob}(\mathbf{X})$ called the *Lévy-Prokhorov metric* such that, the convergence with respect to this metric implies the weak convergence, i.e.,

$$d_{LP}(\mu_n, \mu) \rightarrow 0 \Rightarrow \mu_n \Rightarrow \mu.$$

If moreover the metric space $\mathbf{X}$ is *separable*, the convergence in the Lévy-Prokhorov metric is *equivalent* to the weak convergence. To describe this metric we need a bit of notation.

For any subset $S \subset \mathbf{X}$ and any $\varepsilon > 0$ we set

$$S^\varepsilon := \{x \in \mathbf{X}; \text{dist}(x, S) < \varepsilon\}.$$

The function $x \mapsto \text{dist}(x, S)$ is Lipschitz so S^ε is open for any $S \subset \mathbf{X}$. Given $\mu_0, \mu_1 \in \text{Meas}(\mathbf{X})$ we define

$$d_{LP}(\mu_0, \mu_1) := \inf \{ \varepsilon > 0; \mu_0[B] - \mu_1[B^\varepsilon], \mu_1[B] - \mu_0[B^\varepsilon] < \varepsilon, \forall B \in \mathcal{B}_X \}.$$

For more details and proofs we refer to [18, Sec.6] or [64, Sec.11.3]. $\square$

The next result generalizes Fatou's Lemma. However, our proof relies on Fatou's Lemma.

Proposition 2.2.21. *Suppose that the sequence of random variables $(X_n)_{n \in \mathbb{N}}$ converges in distribution to X . Then*

$$\mathbb{E}[|X|] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[|X_n|].$$

In particular, X is integrable if the sequence $(X_n)_{n \in \mathbb{N}}$ is bounded in L^1 , i.e.,

$$\sup_n \mathbb{E}[|X_n|] < \infty.$$

Proof. The Mapping Theorem 2.2.3 implies that the sequence $(|X_n|)_{n \in \mathbb{N}}$ converges in distribution to $|X|$. Thus

$$\lim_{n \in \mathbb{N}} \mathbb{P}[|X_n| > t] = \mathbb{P}[|X| > t],$$

for all t outside a countable subset of $[0, \infty)$. Using (1.3.48) we deduce

$$\mathbb{E}[|X|] = \int_0^\infty \mathbb{P}[|X| > t] dt, \quad \mathbb{E}[|X_n|] = \int_0^\infty \mathbb{P}[|X_n| > t] dt, \quad \forall n.$$

Fatou's Lemma implies

$$\int_0^\infty \mathbb{P}[|X| > t] dt \leq \liminf_{n \rightarrow \infty} \int_0^\infty \mathbb{P}[|X_n| > t] dt.$$

□

At this point it is profitable to look at the concept of weak convergence from a functional analytic viewpoint.

If (K, d) is a compact metric space, then Riesz's Representation Theorem 1.2.65 shows that $\text{Meas}(K)$ is a closed convex cone in $C(K)^*$, the topological dual of the Banach space $C(K)$ (with the sup-norm).

The weak convergence of finite measures corresponds to the convergence in the weak* topology on the dual space; see [29, Sec. 3.4]. Since $C(K)$ is separable, the weak* topology on $C(K)^*$ is defined by a countable family of seminorms and thus it is metrizable. The Banach-Alaoglu theorem [29, Thm. 3.16] implies that the unit ball in $C(K)^*$ is compact, so any abounded subsequence in $C(K)^*$ admits a convergent subsequence. In particular, this shows that any sequence $(\mu_n)_{n \in \mathbb{N}}$ such that

$$\sup_n \mu_n[K] < \infty$$

admits a subsequence that converges weakly to a finite Borel measure on K .

To see this principle at work consider the compactification $\bar{\mathbb{R}} = [-\infty, \infty]$ of $\mathbb{R}$. The map $\tan : (-\pi/2, \pi/2) \rightarrow \mathbb{R}$ induces a homeomorphism $[-\pi/2, \pi/2]$ and thus the compactification $[-\infty, \infty]$ is metrizable. The continuous functions on $\bar{\mathbb{R}}$ are the continuous functions on $\mathbb{R}$ that have finite limits at $\pm\infty$. In particular, $C(\bar{\mathbb{R}}) \subset C_b(\mathbb{R})$. A finite measure $\mu \in \text{Meas}(\mathbb{R})$ extends to a measure $\bar{\mu} \in \text{Meas}(\bar{\mathbb{R}})$, namely, $\bar{\mu}[\bar{B}] = \mu[\bar{B} \cap \mathbb{R}]$, for any Borel subset $\bar{B} \subset \bar{\mathbb{R}}$. We thus have an inclusion

$$\text{Meas}(\mathbb{R}) \subset \text{Meas}(\bar{\mathbb{R}}).$$

This inclusion is strict: the Dirac measures $\delta_{\pm\infty}$ do not belong to $\text{Meas}(\mathbb{R})$.

Suppose that $(\mu_n)_{n \geq 1}$ is a sequence in $\text{Prob}(\mathbb{R})$. The sequence $(\bar{\mu}_n)_{n \geq 1}$ in $\text{Prob}(\bar{\mathbb{R}})$ admits a subsequence $\bar{\mu}_{n_k}$ that converges weakly to a measure $\tilde{\mu}_\infty$. This defines a measure $\mu_\infty \in \text{Meas}(\mathbb{R})$ by setting

$$\mu_\infty[B] = \tilde{\mu}_\infty[B], \quad \forall B \in \mathcal{B}_\mathbb{R} \subset \mathcal{B}_{\bar{\mathbb{R}}}.$$

In particular, we deduce that for any compactly supported continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ we have

$$\lim_{k \rightarrow \infty} \mu_{n_k}[f] = \lim_{k \rightarrow \infty} \bar{\mu}_{n_k}[f] = \tilde{\mu}_\infty[f] = \mu_\infty[f].$$

Note that the limit μ_∞ need not be a probability measure since

$$\mu_\infty[\mathbb{R}] = \tilde{\mu}_\infty[\bar{\mathbb{R}} \setminus \{\pm\infty\}] = 1 - \tilde{\mu}_\infty[\{\pm\infty\}].$$

Theorem 2.2.22 (Helly's selection theorem). *Any sequence $(\nu_n)_{n \geq 1}$ of finite, nontrivial Borel probability measures on $\mathbb{R}$ such that*

$$\sup_n \nu_n[\mathbb{R}] < \infty$$

admits a vaguely convergent subsequence.

Proof. After extracting a subsequence we can assume that sequence $\nu_n[\mathbb{R}]$ converges to $\nu_\infty \geq 0$. Set

$$\mu_n := \frac{1}{\nu_n[\mathbb{R}]} \nu_n.$$

The above discussion shows that the sequence of probability measures μ_n admits a subsequence (μ_{n_k}) that converges vaguely to $\mu_\infty \in \text{Meas}(\mathbb{R})$. The subsequence $\nu_{n_k}[\]$ converges $\nu_\infty \cdot \mu_\infty[\]$ $\square$

Proposition 2.2.23. Suppose that (μ_n) is a sequence in $\text{Meas}(\mathbb{R})$ that converges vaguely to a measure $\mu_\infty \in \text{Meas}(\mathbb{R})$ and

$$\sup_n \mu_n[\mathbb{R}] < \infty.$$

Then following are equivalent

- (i) The sequence converges weakly to μ_∞ .
- (ii) $\mu_\infty[\mathbb{R}] = \lim_{n \rightarrow \infty} \mu_n[\mathbb{R}]$.
- (iii) The sequence (μ_n) is tight, i.e.,

$$\lim_{L \rightarrow \infty} \sup_{n \in \mathbb{N}} \mu_n[\mathbb{R} \setminus [-L, L]] = 0.$$

Proof. We proved the equivalence (i) $\Rightarrow$ (ii) In Theorem 2.2.14. Let us show that (iii) $\Rightarrow$ (ii). Fix $L > 0$ such that

$$\mu_\infty[\mathbb{R} \setminus [-L, L]] < \varepsilon.$$

Fix $L > 0$. As in the proof Theorem 2.2.14 choose $\varphi_L \in C_{\text{cpt}}(\mathbb{R})$ such that

$$\mathbf{I}_{[-L, L]} \leq \varphi_L \leq 1 = \mathbf{I}_{\mathbb{R}}$$

Hence

$$\mu_n[\mathbb{R}] - \varepsilon \leq \mu_n[\varphi_L] \leq \mu_n[\mathbb{R}], \quad \forall n \in \mathbb{N} \cup \{\infty\}$$

The tightness condition implies that for any $\varepsilon > 0$ we can choose $L = L(\varepsilon)$ such that

$$\forall n \in \mathbb{N} \cup \{\infty\}, \quad \mu_n[-L, L] \geq \mu_n[\mathbb{R}] - \varepsilon.$$

Letting $n \rightarrow \infty$ we deduce

$$\forall \varepsilon > 0, \quad \limsup_{n \rightarrow \infty} \mu_n[\mathbb{R}] - \varepsilon \leq \mu_\infty[\varphi_{L(\varepsilon)}] \leq \liminf_{n \rightarrow \infty} \mu_n[\mathbb{R}].$$

Since this holds for any $\varepsilon > 0$ we deduce that

$$\liminf_{n \rightarrow \infty} \mu_n[\mathbb{R}] = \lim_{n \rightarrow \infty} \mu_n[\mathbb{R}].$$

Thus, for any $\varepsilon > 0$

$$\left| \lim_{n \rightarrow \infty} \mu_n[\mathbb{R}] - \mu_\infty[\varphi_{L(\varepsilon)}] \right| < \varepsilon, \quad \left| \mu_\infty[\varphi_{L(\varepsilon)}] - \mu_\infty[\mathbb{R}] \right| < \varepsilon.$$

Hence

$$\lim_{n \rightarrow \infty} \mu_n[\mathbb{R}] = \mu_\infty[\mathbb{R}].$$

This proves (ii).

Finally, let us prove that (i) $\Rightarrow$ (iii). For each $L > 0$ choose as above $f_L \in C_{\text{cpt}}(\mathbb{R})$ such that $\mathbf{I}_{[-L, L]} \leq f_L \leq 1$. Next, for any $\varepsilon > 0$ choose $L = L(\varepsilon)$ such that

$$\mu_\infty[\mathbb{R}] \geq \mu_\infty[f_{L(\varepsilon)}] \geq \mu_\infty[[-L(\varepsilon), L(\varepsilon)]] > \mu_\infty[\mathbb{R}] - \frac{\varepsilon}{2}.$$

Let $g_\varepsilon = 1 - f_{L(\varepsilon)} \in C_b(\mathbb{R})$. Then $\mu_n[g_\varepsilon] \rightarrow \mu_\infty[g_\varepsilon]$ so that

$$\lim_{n \rightarrow \infty} (\mu_n[\mathbb{R}] - \mu_n[f_{L(\varepsilon)}]) = \mu_\infty[\mathbb{R}] - \mu_\infty[f_{L(\varepsilon)}] \leq \frac{\varepsilon}{2}.$$

Thus there exists $N = N(\varepsilon)$ such that, for any $n > N(\varepsilon)$

$$\mu_n[\mathbb{R}] - \mu_n[[-L(\varepsilon), L(\varepsilon)]] \leq \mu_n[\mathbb{R}] - \mu_n[f_{L(\varepsilon)}] < \varepsilon.$$

For each $k = 1, \dots, N(\varepsilon)$, choose $L_k(\varepsilon) > 0$ such that

$$\mu_k[\mathbb{R}] - \mu_k[[-L_k(\varepsilon), L_k(\varepsilon)]] < \varepsilon.$$

If we set

$$L_*(\varepsilon) = \max \{ L_1(\varepsilon), \dots, L_{N(\varepsilon)}(\varepsilon), L(\varepsilon) \},$$

then we deduce that for any $n \in \mathbb{N}$ and any $L > L_*(\varepsilon)$ we have

$$\mu_n[\mathbb{R}] - \mu_n[[-L, L]] < \varepsilon.$$

□

Proposition 2.2.24. *Suppose that $(X_n)_{n \in \mathbb{N} \cup \{\infty\}}$ is a family of random variables with the following properties*

(i) *For any $k \in \mathbb{N}$, and any $n \in \mathbb{N} \cup \{\infty\}$, $\mathbb{E}[|X_n|^k] < \infty$.*

(ii) *For any $k \in \mathbb{N}$*

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \mathbb{E}[X_\infty^k].$$

(iii) *The probability distribution of X_∞ is uniquely determined by its momenta. E.g., this happens if $\exists T > 0$ such that*

$$\mathbb{E}[e^{tX_\infty}] < \infty, \quad \forall |t| < T.$$

Then X_n converges in distribution to X_∞ .

Proof. Set $\mu_n := \mathbb{P}_{X_n}$. Observe that the family (μ_n) is tight. Indeed, if

$$M := \sup_{k \in \mathbb{N} \cup \{\infty\}} \mathbb{E}[X_n^2],$$

then we deduce from Chebyshev's inequality that for any $L > 0$ and any $n \in \mathbb{N}$ we have

$$\mu_n[\{|x| > L\}] \leq \frac{M}{L^2}.$$

We will first prove that the whole sequence μ_n converges vaguely to a finite measure μ_∞ .

Helly's Selection Theorem implies that any subsequence of μ_n has vaguely convergent subsequences. Thus it suffices to show that all the vaguely convergent subsequence of (μ_n) have the same limit.

Suppose that μ_∞ is the vague limit of a subsequence. To ease the presentation assume that the subsequence is (μ_n) . Since the sequence (μ_n) is tight the convergence is weak and μ_∞ is a probability measure. We will prove that μ_∞ has finite moments and, more precisely

$$\int_{\mathbb{R}} x^k \mu_\infty[dx] = \mathbb{E}[X_\infty^k], \quad \forall k \in \mathbb{N}.$$

Since the distribution of X_∞ is assumed to be uniquely determined by its moments we deduce that $\mu_\infty = \mathbb{P}_{X_\infty}$.

Fix $k \in \mathbb{N}$. Define the finite measures

$$\nu_n[dx] = \nu_{n,k}[dx] = (1 + x^{2k})\mu_n[dx].$$

We set $\nu_\infty[dx] := (1 + x^{2k})\mu_\infty[dx]$. Let us first show that ν_∞ is a *finite* measure.

To see this choose for any $L > 0$ a compactly supported functions such that

$$\mathbf{1}_{[-L,L]} \leq \varphi_L \leq 1.$$

Since $\varphi_L(x)(1 + x^{2k})$ is compactly supported we deduce that

$$\begin{aligned} \nu_\infty[-L, L] &\leq \nu_\infty[\varphi_L] = \int_{\mathbb{R}} \varphi_L(x)(1 + x^{2k})\mu_\infty[dx] \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}} \varphi_L(x)(1 + x^{2k})\mu_n[dx] \leq \lim_{n \rightarrow \infty} \int_{\mathbb{R}} (1 + x^{2k})\mu_n[dx] \\ &= (1 + \mathbb{E}[X_\infty^{2k}]). \end{aligned}$$

Hence

$$\nu_\infty[\mathbb{R}] = \lim_{L \rightarrow \infty} \nu_\infty[-L, L] \leq 1 + \mathbb{E}[X_\infty^{2k}] < \infty.$$

Since μ_n converges vaguely to μ_∞ and $\varphi(x)(1 + x^{2k})$ has compact support for any compactly supported φ we deduce that ν_n converges vaguely to the finite measure ν_∞ . We will show that in fact that the sequence (ν_n) is tight so that it converges weakly to ν_∞ .

For any $L > 0$ we have

$$\begin{aligned} L^{2k}\nu_n[\{|x| > L\}] &\leq \int_{|x|>L} x^{2k}\nu_n[dx] \\ &\leq \int_{\mathbb{R}} x^{2k}(1 + x^{2k})\mu_n[dx] \leq M_{2k} + M_{4k}, \end{aligned}$$

where

$$M_j := \sup_n \mathbb{E}[|X_n|^j], \quad \forall j \in \mathbb{N}.$$

Hence

$$\nu_n[\{|x| > L\}] \leq \frac{M_{2k} + M_{4k}}{L^{2k}}, \quad \forall n \in \mathbb{N}, \quad L > 0.$$

proving that the sequence (ν_n) is tight.

Consider now the bounded continuous function $f(x) = \frac{x^k}{1+x^{2k}}$. Then

$$\begin{aligned} \mathbb{E}[X_\infty^k] &= \lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \lim_{n \rightarrow \infty} \nu_n[f] \\ &= \nu_\infty[f] = \int_{\mathbb{R}} x^k \mu_\infty[dx]. \end{aligned}$$

Thus μ_∞ has finite moments of any order, all equal to the moments of $\mathbb{P}_{X_\infty}$. □

2.2.2. The characteristic function. Suppose that E is a finite dimensional real Euclidean space with inner product $(-, -)$ and associated norm $\| - \|$. Denote by E^* the dual of E , $E^* = \text{Hom}(E, \mathbb{R})$. For $\xi \in E^*$ and $x \in E$ we set

$$\langle \xi, x \rangle := \xi(x).$$

The inner product on E induces by duality an inner product and Euclidean norm on E^* denoted by the same corresponding symbols.

The key ingredient in the proof of the CLT is that of *Fourier transform* or *characteristic function* of a finite Borel measure $\mu \in \text{Meas}(E)$,

$$\hat{\mu} : E^* \rightarrow \mathbb{C}, \quad \hat{\mu}(\xi) = \int_E e^{i\langle \xi, x \rangle} \mu[dx].$$

Note that μ is a *probability* measure if and only if $\hat{\mu}(0) = 1$.

From the Dominated Convergence Theorem we deduce that $\hat{\mu}$ is a continuous function $E^* \rightarrow \mathbb{C}$. Thus, the Fourier transform is a map

$$\text{Prob}(E) \ni \mu \mapsto \hat{\mu} \in C_b(E^*, \mathbb{C}).$$

The characteristic function of an E -valued random vector X is the Fourier transform $\Phi_X(\xi)$ of its probability distribution $\mathbb{P}_X \in \text{Prob}(E)$,

$$\Phi_X(\xi) = \hat{\mathbb{P}}_X(\xi) = \mathbb{E}[e^{i\xi X}] = \int_E e^{i\langle \xi, x \rangle} \mathbb{P}_X[dx].$$

Note that

$$|\Phi_X(\xi)| \leq 1, \quad \forall \xi \in \mathbb{C}.$$

Moreover, $\Phi_X(0) = \mathbb{E}[1] = 1$.

Proposition 2.2.25. *Let X be an E valued random vector such that $\|X\| \in L^2(\Omega, \mathcal{S}, \mathbb{P})$. Suppose that $\dim E = m$. Fix orthonormal coordinates (x_k) on E with dual coordinates (ξ_k) on E^* so that $X = (X_1, \dots, X_m)$. Then $\Phi_X \in C^2(E^*)$ and, for any $j, k = 1, \dots, m$*

$$\partial_{\xi_k} \Phi_X(0) = i\mathbb{E}[X_k], \quad \partial_{\xi_k \xi_k}^2 \Phi_X(0) = -\mathbb{E}[X_j X_k].$$

Proof. Denote by $\mathbb{P}_X$ the probability distribution of X so $\mathbb{P}_X \in \text{Prob}(E)$. Then

$$\Phi_X(\xi) = \int_E e^{i\langle \xi, x \rangle} \mathbb{P}_X[dx].$$

Note that since $X \in L^2$ we have

$$\int_E \|x\| \mathbb{P}_X[dx], \quad \int_E \|x\|^2 \mathbb{P}_X[dx] < \infty.$$

Then, for any $k \in \llbracket 1, m \rrbracket$

$$\begin{aligned} \partial_{\xi_k} e^{i\langle \xi, x \rangle} &= i x_k e^{i\langle \xi, x \rangle} \in L^1(E, \mathbb{P}_X), \\ \partial_{\xi_j \xi_k}^2 e^{i\langle \xi, x \rangle} &= -x_j x_k e^{i\langle \xi, x \rangle} \in L^1(E, \mathbb{P}_X). \end{aligned}$$

This shows (see Exercise 1.8) that the integral

$$E^* \ni \xi \mapsto \int_E e^{i\langle \xi, x \rangle} \mathbb{P}_X[dx]$$

is twice differentiable with respect to the parameter ξ and we have

$$\partial_{\xi_k} \Phi_X(0) = -i \int_{\mathbb{R}} x_k \mathbb{P}_X[dx],$$

$$\partial_{\xi_k \xi_k}^2 \Phi_X(0) = - \int_{\mathbb{R}} x_j x_k \mathbb{P}_X[dx].$$

Using the Dominated Convergence Theorem we deduce that the function

$$\xi \mapsto - \int_E x_j x_k e^{i\langle \xi, x \rangle} \mathbb{P}_X[dx].$$

is continuous so $\Phi_X \in C^2(E)$. □

For $v > 0$ we denote $\Gamma_v \in \text{Meas}(E)$ the measure

$$\Gamma_v[dx] = \gamma_v(x) \lambda[dx], \quad \gamma_v(x) = \frac{1}{(2\pi v)^{\frac{\dim E}{2}}} e^{-\frac{\|x\|^2}{2v}}, \quad (2.2.6)$$

where λ is the Lebesgue measure on E .

Suppose that $m = \dim E$. Choose Euclidean coordinates $(x_1, \dots, x_m)$, $m = 1, \dots, m$. Then we observe Γ_v is the product of m Gaussian measures on $\mathbb{R}$ with mean 0 and variance v

$$\Gamma_v[dx] = \bigotimes_{k=1}^m \frac{1}{\sqrt{2\pi v}} e^{-\frac{x_k^2}{2v}} dx_k. \quad (2.2.7)$$

Thus Γ_v is a Borel probability measure on E .

Proposition 2.2.26. *Let $m := \dim E$. Then*

$$\int_E \|x\|^2 \Gamma_v[dx] = mv, \quad (2.2.8)$$

$$\widehat{\Gamma}_v(\xi) = e^{-\frac{v\|\xi\|^2}{2}} = \left(\frac{2\pi}{v}\right)^{\frac{m}{2}} \gamma_{1/v}(\xi), \quad \forall v > 0. \quad (2.2.9)$$

Proof. We choose an orthonormal frame of E with Euclidean coordinates $(x_1, \dots, x_m)$. Then

$$\int_E \|x\|^2 \Gamma_v[dx] = \sum_{k=1}^m \int_{\mathbb{R}^m} x_k^2 \Gamma_v[dx] \stackrel{(2.2.7)}{=} \sum_{k=1}^m \frac{1}{\sqrt{2\pi v}} \underbrace{\int_{\mathbb{R}} x_k^2 e^{-\frac{x_k^2}{2v}} dx_k}_{=v}.$$

$$\begin{aligned} \widehat{\Gamma}_v(\xi_1, \dots, \xi_m) &= \frac{1}{(2\pi v)^{\frac{m}{2}}} \int_{\mathbb{R}^m} \left(\prod_{k=1}^m e^{i\xi_k x_k - \frac{x_k^2}{2v}} \right) dx_1 \cdots dx_m \\ &= \prod_{k=1}^m \left(\frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} e^{i\xi_k x_k - \frac{x_k^2}{2v}} dx_k \right) = \left(\frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} e^{i\xi x - \frac{x^2}{2v}} dx \right)^m. \end{aligned}$$

Thus, it suffices to consider only the case $E = \mathbb{R}$. We have

$$\widehat{\Gamma}_v(\xi) = \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} e^{-\frac{x^2}{2v}} e^{i\xi x} dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\frac{y^2}{2}} e^{i\sqrt{v}\xi y} dy = \widehat{\Gamma}_1(\sqrt{v}\xi).$$

Hence only need to determine

$$f(\xi) = \widehat{\Gamma}_1(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\frac{x^2}{2}} e^{i\xi x} dx.$$

The imaginary part of the above integrand is an odd function (in x) so $f(\xi)$ is real, $\forall \xi$, i.e.,

$$f(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\frac{x^2}{2}} \cos(\xi x) dx.$$

The function

$$\frac{d}{d\xi} \left(e^{-\frac{x^2}{2}} \cos(\xi x) \right) = -x e^{-\frac{x^2}{2}} \sin(\xi x)$$

is integrable (in the x variable). This shows that $f(\xi)$ is differentiable (see Exercise 1.8) and

$$f'(\xi) = -\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} x e^{-\frac{x^2}{2}} \sin(\xi x) dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \frac{d}{dx} \left(e^{-\frac{x^2}{2}} \right) \sin(\xi x) dx$$

(integrate by parts)

$$= -\frac{\xi}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\frac{x^2}{2}} \cos(\xi x) dx = -\xi f(\xi).$$

Thus

$$f'(\xi) + \xi f(\xi) = 0$$

so that

$$\frac{d}{d\xi} \left(e^{\xi^2/2} f(\xi) \right) = 0 \iff f(\xi) = C e^{-\frac{\xi^2}{2}}.$$

Since $f(0) = 1$ we deduce $C = 1$ and thus $\widehat{\Gamma}_1(\xi) = e^{-\frac{\xi^2}{2}}$. $\square$

Theorem 2.2.27. A probability measure $\mu \in \text{Prob}(E)$ is uniquely determined by its characteristic function, i.e., the map

$$\text{Prob}(E) \ni \mu \mapsto \widehat{\mu} \in C_b(E^*, \mathbb{C}).$$

is injective.

Proof. For any $v > 0$ and $\mu \in \text{Prob}(E)$ we set $\mu_v := \Gamma_v * \mu$. We have

$$\mu_v[dx] := \rho_v(x) dx, \quad \rho_v(x) = \int_E \gamma_v(x-y) \mu[dy].$$

The theorem follows from the following two facts.

Fact 1. The family $(\mu_v)_{v>0}$ is completely determined by $\widehat{\mu}$.

Fact 2. The family $(\mu_v)_{v>0}$ converges weakly to μ as $v \searrow 0$, i.e.,

$$\lim_{v \searrow 0} \mu_v[f] = \mu[f], \quad \forall f \in C_b(\mathbb{R}).$$

Proof of Fact 1. The idea behind this fact is that the Fourier transform and the convolution interact in a nice way. More precisely we will show that

$$\rho_v(x) = \frac{1}{(2\pi v)^{\frac{m}{2}}} \int_{E^*} e^{ix\xi} \gamma_{1/v}(\xi) \widehat{\mu}(-\xi) \lambda[d\xi]. \quad (2.2.10)$$

Using (2.2.9) with the roles of x and ξ reversed (i.e., we think of E as the dual of E^*) we deduce

$$(2\pi v)^{\frac{m}{2}} \gamma_v(x) = e^{-\frac{x^2}{2v}} = \int_{E^*} e^{i\langle x, \xi \rangle} \gamma_{1/v}(\xi) \lambda[d\xi].$$

Hence

$$\rho_v(x) = \frac{1}{(2\pi v)^{\frac{m}{2}}} \int_E \left(\int_{E^*} e^{i(x-y)\xi} \gamma_{1/v}(\xi) \lambda[d\xi] \right) \mu[dy]$$

(use Fubini)

$$= \frac{1}{(2\pi v)^{\frac{m}{2}}} \int_{E^*} e^{ix\xi} \gamma_{1/v}(\xi) \left(\int_E e^{-iy\xi} \mu[dy] \right) \lambda[d\xi]$$

$$= \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} e^{ix\xi} \gamma_{1/v}(\xi) \widehat{\mu}(-\xi) \lambda[d\xi].$$

Proof of Fact 2. Let $f \in C_b(E)$. Using Fubini's theorem we deduce that

$$\begin{aligned} \int_E f(x) \mu_v[dx] &= \int_E f(x) \left(\int_E \gamma_v(x-y) \mu[dy] \right) \lambda[dx] \\ &= \int_E \underbrace{\left(\int_E \gamma_v(x-y) f(x) \lambda[dx] \right)}_{=: f_v(y)} \mu[dy]. \end{aligned}$$

The function $y \mapsto f_v(y)$ is obviously continuous. If $C := \sup_{x \in E} |f(x)|$, then, $\forall y \in E$ and any $v > 0$ we have

$$|f_v(y)| \leq M \int_E \gamma_v(x-y) \lambda[dx] \stackrel{x \rightarrow z+y}{=} C \int_E \gamma_v(z) \lambda[dz] = C.$$

On the other hand

$$\begin{aligned} f_v(y) &= \int_E \gamma_v(y-t) f(t) \lambda[dt] = \int_E \gamma_v(t-y) f(t) \lambda[dt] \\ &= \int_E \gamma_v(z) f(z+y) \lambda[dz] = \Gamma_v[T_y f], \end{aligned}$$

where $T_y f(z) := f(z+y)$. Fix $y, \varepsilon > 0$ and a $\delta = \delta(\varepsilon, y) > 0$ such that

$$\sup_{\|z\| < \delta} |f(z+y) - f(y)| < \frac{\varepsilon}{2}.$$

Then

$$\begin{aligned} |f_v(y) - f(y)| &= |\Gamma_v[T_y f] - f(y)| \\ &= \left| \int_{\mathbb{R}} (f(z+y) - f(y)) \Gamma_v[dz] \right| \\ &\leq \int_{\|z\| < \delta} |f(z+y) - f(y)| \Gamma_v[dz] + \int_{\|z\| \geq \delta} |f(z+y) - f(y)| \Gamma_v[dz] \\ &\leq \sup_{\|z\| < \delta} |f(z+y) - f(y)| + 2C \int_{\|z\| \geq \delta} \Gamma_v[dz] \\ &\leq \sup_{\|z\| < \delta} |f(z+y) - f(y)| + \frac{2C}{\delta^2} \int_{\|z\| \geq \delta} \|z\|^2 \Gamma_v[dz] \\ &\stackrel{(2.2.8)}{<} \frac{\varepsilon}{2} + \frac{2Cmv}{\delta^2}. \end{aligned}$$

Hence

$$\forall \varepsilon > 0, \limsup_{v \searrow 0} |f_v(y) - f(y)| \leq \frac{\varepsilon}{2}, \quad \forall \varepsilon > 0, \quad \forall y \in \mathbb{R},$$

so that

$$\lim_{v \searrow 0} f_v(y) = f(y), \quad \forall y \in \mathbb{R}.$$

The Dominated Convergence Theorem implies

$$\lim_{v \searrow 0} \mu_v[f] = \lim_{v \searrow 0} \mu[f_v] = \mu\left[\lim_{v \searrow 0} f_v\right] = \mu[f].$$

□

Remark 2.2.28. (a) In the above proof set

$$\text{osc}_f(y, \delta) := \sup_{\|z\| < \delta} |f(y+z) - f(y)|, \quad \text{osc}_f(\delta) = \sup_{y \in E} \text{osc}_f(y, \delta).$$

We proved that

$$|f_v(y) - f(y)| \leq \text{osc}_f(y, \delta) + \frac{2m\|f\|_\infty v}{\delta^2} \leq \text{osc}_f(\delta) + \frac{2m\|f\|_\infty v}{\delta^2}.$$

In particular, if f is *uniformly continuous*, i.e., $\lim_{\delta \rightarrow 0} \omega_f(\delta) = 0$, we deduce that f_v converges *uniformly* to f . More precisely, if we set $\delta = v^{1/4}$ we deduce

$$\|f_v - f\|_\infty \leq \text{osc}_f(v^{1/4}) + 2m\|f\|_\infty v^{1/2}. \quad (2.2.11)$$

(b) The above theorem implies that the collection of trigonometric functions

$$\{ \mathbb{R} \ni x \mapsto \cos(\xi x), \sin(\xi x); \quad \xi \in \mathbb{R} \}$$

is separating. However, the smaller family

$$\{ \mathbb{R} \ni x \mapsto \cos(\xi x), \sin(\xi x); \quad |\xi| < 1 \},$$

is *not separating*! More precisely, there exists two *distinct* probability measures $\mu_0, \mu_1 \in \text{Prob}(\mathbb{R})$ such that

$$\hat{\mu}_0(\xi) = \hat{\mu}_1(\xi), \quad \forall |\xi| < 1.$$

We refer to [137, Chap. IV, Sec. 15, p.231] for more details.

(b) The range of the Fourier transform

$$\text{Prob}(\mathbb{R}) \ni \mu \mapsto \hat{\mu} \in C_b(\mathbb{R})$$

can also be characterized. Note first that $\forall \mu \in \text{Prob}(\mathbb{R})$

$$\hat{\mu}(0) = \mu[\mathbb{R}] = 1, \quad \hat{\mu}(-\xi) = \overline{\hat{\mu}(\xi)}, \quad \forall \xi \in \mathbb{R}.$$

Additionally, the function $\hat{\mu}$ is *positive definite*. This means that, for any $n \in \mathbb{N}$ and any $\xi_1, \dots, \xi_n \in \mathbb{R}$, the hermitian matrix

$$(\hat{\mu}(\xi_i - \xi_j))_{1 \leq i, j \leq n}$$

is positive semidefinite, i.e., for any $z_1, \dots, z_n$ we have

$$\sum_{1 \leq i, j \leq n} \hat{\mu}(\xi_i - \xi_j) z_i \bar{z}_j \geq 0.$$

This follows by observing that

$$\sum_{1 \leq i, j \leq n} \hat{\mu}(\xi_i - \xi_j) z_i \bar{z}_j = \int_{\mathbb{R}} \left| \sum_{i=1}^n z_i e^{i\xi_i x} \right|^2 \mu[dx].$$

It turns out that these above necessary conditions characterize the range of the Fourier transform: it consists of continuous positive semidefinite functions $\sigma : \mathbb{R} \rightarrow \mathbb{C}$ such that $\sigma(0) = 1$. This is the content of the celebrated Bochner theorem. For various proofs we refer to [76, Sec. XIX.2], [86, §II.3], [180, I.24], [183, Sec. 1.4], [194, Thm. 9.17], or [220, Chap.6]. $\square$

Corollary 2.2.29. Suppose that $X_1, \dots, X_m$ are real random variables. Denote by $\mathbb{P}_{\vec{X}} \in \text{Prob}(\mathbb{R}^m)$ the distribution of the random vector $\vec{X} = (X_1, \dots, X_m)$. Then the following are equivalent

(i) The random variables $X_1, \dots, X_m$ are independent.

(ii) For any $\xi_1, \dots, \xi_m \in \mathbb{R}$

$$\widehat{\mathbb{P}}_{\vec{X}}(\xi_1, \dots, \xi_m) = \prod_{k=1}^m \Phi_{X_k}(\xi_k).$$

Proof. Note that

$$\widehat{\mathbb{P}}_{\vec{X}}(\xi_1, \dots, \xi_m) = \int_{\mathbb{R}^m} e^{i\langle \xi, x \rangle} \mathbb{P}_{\vec{X}}[dx].$$

We denote by $\mathbb{Q}_{\vec{X}}$ the product measure

$$\mathbb{Q}_{\vec{X}} = \bigotimes_{k=1}^m \mathbb{P}_{X_k}.$$

Note that

$$\widehat{\mathbb{Q}}_{\vec{X}}(\xi_1, \dots, \xi_m) = \int_{\mathbb{R}^m} e^{i\langle \xi, x \rangle} \mathbb{P}_{X_1} \otimes \dots \otimes \mathbb{P}_{X_m}[dx] = \prod_{k=1}^m \Phi_{X_k}(\xi_k).$$

The random variables $X_1, \dots, X_m$ are independent iff $\mathbb{P}_{\vec{X}} = \mathbb{Q}_{\vec{X}}$. The corollary now follows from Theorem 2.2.27. $\square$

Corollary 2.2.30. *Let X be a random variable on the sample space $(\Omega, \mathcal{S}, \mathbb{P})$ and $\mathcal{F} \subset \mathcal{S}$ a sigma-subalgebra. Then the following are equivalent.*

- (i) X is independent of $\mathcal{F}$.
- (ii) For any $\xi \in \mathbb{R}$

$$\mathbb{E}[e^{i\xi X} \mid \mathcal{F}] = \mathbb{E}[e^{i\xi X}].$$

Proof. The implication (i) $\Rightarrow$ (ii) follows from the definition of independence. Conversely, suppose that (ii) holds. Then for any $F \in \mathcal{F}$ we have

$$\mathbb{E}[e^{i\xi X} e^{i\xi \mathbf{I}_F}] = \mathbb{E}[e^{i\xi X}] \mathbb{E}[e^{i\xi \mathbf{I}_F}], \quad \forall \xi \in \mathbb{R}.$$

Corollary 2.2.29 implies that X is independent of $\mathbf{I}_F$, for any $F \in \mathcal{F}$, i.e., X is independent of F . $\square$

Theorem 2.2.31 (Lévy's Continuity Theorem). *Let $(\mu_n)_{n \in \mathbb{N}}$ be a sequence in $\text{Prob}(\mathbb{R})$ and $\mu \in \text{Prob}(\mathbb{R})$. The following statements are equivalent.*

- (i) The sequence $(\mu_n)_{n \in \mathbb{N}}$ converges weakly to μ .
- (ii) For any $\xi \in \mathbb{R}$

$$\lim_{n \rightarrow \infty} \widehat{\mu}_n(\xi) = \widehat{\mu}(\xi).$$

Proof. Our presentation is influenced by Le Gall's course notes [131].

(i) $\Rightarrow$ (ii) Since $\mu_n \Rightarrow \mu$ we deduce that for any $\xi \in \mathbb{R}$ we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}} \cos(\xi x) \mu_n[dx] &= \int_{\mathbb{R}} \cos(\xi x) \mu[dx], \\ \lim_{n \rightarrow \infty} \int_{\mathbb{R}} \sin(\xi x) \mu_n[dx] &= \int_{\mathbb{R}} \sin(\xi x) \mu[dx]. \end{aligned}$$

(ii) $\Rightarrow$ (i) For any $v > 0$ and any $f \in C_b(\mathbb{R})$ we define $f_v : \mathbb{R} \rightarrow \mathbb{R}$

$$f_v(x) = \int_{\mathbb{R}} f(x - y) \Gamma_v[dy].$$

It is easy to see that $f_v \in C_b(\mathbb{R})$. We set

$$\mathcal{F} := \{ f_v; f \in C_{\text{cpt}}(\mathbb{R}), v > 0 \}.$$

We will prove that the closure of $\mathcal{F}$ in $C_b(\mathbb{R})$ contains $C_{\text{cpt}}(\mathbb{R})$ and then

$$\lim_{n \rightarrow \infty} \mu_n[f_v] = \mu[f_v], \quad \forall v > 0, \quad \forall f \in C_{\text{cpt}}(\mathbb{R}). \quad (2.2.12)$$

Let $f \in C_{\text{cpt}}(\mathbb{R})$. Observe that

$$f_v(x) = \int_{\mathbb{R}} f(x-y) \gamma_v(y) dy = \int_{\mathbb{R}} f(z) \gamma_v(x-z) dz.$$

Since f has compact support f is uniformly continuous and according to Remark 2.2.28 (a), the function f_v converges uniformly to f . Thus the closure of $\mathcal{F}$ in $C_b(\mathbb{R})$ contains $C_{\text{cpt}}(\mathbb{R})$.

Let $\nu \in \text{Prob}(\mathbb{R})$. Then

$$\begin{aligned} \nu[f_v] &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(z) \gamma_v(z-x) dz \right) \nu[dx] = \int_{\mathbb{R}} f(z) \underbrace{\left(\int_{\mathbb{R}} \gamma_v(z-x) \nu[dx] \right)}_{\rho_v(z)} dz \\ &\stackrel{(2.2.10)}{=} \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{ix\xi} \gamma_{1/v}(\xi) \widehat{\nu}(-\xi) d\xi \right) f(x) dx \\ &= \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} \underbrace{\left(\int_{\mathbb{R}} e^{ix\xi} f(x) d\xi \right)}_{=: \widehat{f}(\xi)} \gamma_{1/v}(\xi) \widehat{\nu}(-\xi) d\xi = \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} \widehat{f}(\xi) \gamma_{1/v}(\xi) \widehat{\nu}(-\xi) d\xi. \end{aligned}$$

The function $\widehat{f}(\xi)$ is well defined since $f \in C_{\text{cpt}}(\mathbb{R})$. The Dominated Convergence theorem shows that $\widehat{f}$ is continuous. Moreover

$$|\widehat{f}(\xi)| \leq \int_{\mathbb{R}} |f(x)| dx.$$

We deduce that, $\forall n \in \mathbb{N}$,

$$\mu_n[f_v] = \frac{1}{\sqrt{2\pi v}} \int_{\mathbb{R}} \widehat{f}(\xi) \gamma_{1/v}(\xi) \widehat{\mu}_n(-\xi) d\xi.$$

Note that for any $\nu \in \text{Prob}(\mathbb{R})$

$$|\widehat{f}(\xi) \gamma_{1/v}(\xi) \widehat{\nu}(-\xi)| \leq \left(\sup_{x \in \mathbb{R}} |f(x)| \right) \cdot \gamma_{1/v}(\xi) \in L^1(\mathbb{R}, \lambda).$$

The Dominated Convergence theorem shows that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} \widehat{f}(\xi) \gamma_{1/v}(\xi) \widehat{\mu}_n(-\xi) d\xi = \int_{\mathbb{R}} \widehat{f}(\xi) \gamma_{1/v}(\xi) \widehat{\mu}(-\xi) d\xi = \mu[f_v].$$

As explained in Remark 2.2.28(a), if $f \in C_b(\mathbb{R})$ is uniformly continuous, then f_v converges to f uniformly as $v \searrow 0$. In particular, if f has compact support, then f_v converges uniformly to f as $v \rightarrow 0$. We deduce that the family

$$\mathcal{F} := \{ \varphi_v; v > 0, \varphi \in C_{\text{cpt}}(\mathbb{R}) \}$$

contains $C_{\text{cpt}}(\mathbb{R})$ in its closure and $\mu_n[f] \rightarrow \mu[f]$ for any $f \in \mathcal{F}$. The conclusion follows from Theorem 2.2.14. $\square$

Proposition 2.2.32. *If a sequence $\mu_n \in \text{Prob}(\mathbb{R})$ converges weakly to a probability measure μ , then $\widehat{\mu}_n(\xi)$ converges to $\widehat{\mu}(\xi)$ uniformly on compacts.*

Proof. The proof relies on the following technical result.

Lemma 2.2.33. *For any $\mu \in \text{Prob}(\mathbb{R})$ and any $\xi, \eta \in \mathbb{R}$ we have*

$$|\widehat{\mu}(\xi) - \widehat{\mu}(\eta)| \leq 2(1 - \mathbf{Re} \widehat{\mu}(\xi - \eta)). \quad (2.2.13)$$

□

Taking for granted this result we can now complete the proof of Proposition . For simplicity set

$$\Phi(\xi) := \widehat{\mu}(\xi) \quad \text{and} \quad \Phi_n(\xi) := \widehat{\mu_n}(\xi).$$

To prove that Φ_n converges to Φ uniformly on compacts it suffices to show that it is equicontinuous. Thus, we want to prove that for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\forall n \in \mathbb{N}, \quad \forall \xi \in \mathbb{R}, \quad |t| \leq \delta : \quad |\Phi_n(\xi + t) - \Phi_n(\xi)| < \varepsilon.$$

Since μ_n converges weakly to μ we deduce that for any $\varepsilon > 0$ there exists $R > 0$ such that

$$\mu_n[\{|x| > R\}] < \frac{\varepsilon}{2}, \quad \forall n.$$

Fix $\delta > 0$ such that

$$\forall (t, x) \in [-\delta, \delta] \times [-R, R]. \quad |1 - e^{itx}| < \frac{\varepsilon}{2}.$$

We deduce that for any $n \in \mathbb{N}$, any $\xi \in \mathbb{R}$, and any $|t| \leq \delta$ we

$$\begin{aligned} |\Phi_n(\xi + t) - \Phi_n(\xi)| &\stackrel{(2.2.13)}{\leq} 1 - \mathbf{Re} \Phi_n(t) \leq \int_{\mathbb{R}} |1 - e^{itx}| \mu_n[dx] \\ &= \int_{[-R, R]} \underbrace{|1 - e^{itx}|}_{< \varepsilon/2} \mu_n[dx] + \int_{\mathbb{R} \setminus [-R, R]} \underbrace{|1 - e^{itx}|}_{\leq 2} \mu_n[dx] < \varepsilon. \end{aligned}$$

We still have to prove (2.2.13). We have

$$\begin{aligned} |\Phi(\xi + t) - \Phi(\xi)|^2 &= \left| \int_{\mathbb{R}} (e^{i(\xi+t)x} - e^{i\xi x}) \mu[dx] \right|^2 \\ &= \left| \int_{\mathbb{R}} (e^{itx} - 1) e^{i\xi x} \mu[dx] \right|^2 \\ &\leq \left(\int_{\mathbb{R}} |e^{itx} - 1|^2 \mu[dx] \right) \underbrace{\left(\int_{\mathbb{R}} |e^{i\xi x}|^2 \mu[dx] \right)}_{=1} \\ &= \int_{\mathbb{R}} ((\cos tx - 1)^2 + (\sin tx)^2) \mu[dx] \\ &= 2 \int_{\mathbb{R}} (1 - \cos(tx)) \mu[dx] = 2(1 - \mathbf{Re} \Phi(t)). \end{aligned}$$

□

Remark 2.2.34. In Theorem 2.2.31 we assumed that the pointwise limit of the sequence of characteristic functions $(\widehat{\mu_n})_{n \in \mathbb{N}}$ is the characteristic function of a probability measure μ . This assumption is not necessary. A lot less suffices.

More precisely, the general version of Lévy's continuity theorem states the following.

If the characteristic functions of probability measures $\mu_n \in \text{Prob}(\mathbb{R})$ converge pointwisely to a function that is continuous at the origin, then the limit itself is the characteristic function of a probability measure $\mu \in \text{Prob}(\mathbb{R})$ and $\mu_m \Rightarrow \mu$ as $n \rightarrow \infty$.

This is not obvious and requires additional effort. In Exercise 2.44 we describe the main steps of a proof of this fact. In fact, as shown in [76, Sec. XIX.2] or [203, Thm. 1.1.10], one can use this stronger version of the continuity theorem to prove Bochner's theorem. $\square$

Remark 2.2.35. P. Lévy, [134, §17, p.47], introduced a metric d_L on $\text{Prob}(\mathbb{R})$. More precisely, given $\mu_0, \mu_i \in \text{Prob}(\mathbb{R})$ with cumulative distribution functions

$$F_i(x) = \mu_i[(-\infty, x]], \quad x \in \mathbb{R}, \quad i = 0, 1,$$

then the Lévy metric is the length of the largest segment cut-out by the graphs Γ_0, Γ_1 of F_0, F_1 along a line of the form $x + y = a$. The graphs are made continuous by adding vertical segments connecting $F_i(x - 0)$ to $F_i(x)$ at the points of discontinuity. Intuitively, the distance is the diagonal of the largest square with sides parallel to the axes that can be squeezed between the curves Γ_0 and Γ_1 .

More precisely

$$d_L(\mu_0, \mu_1) = \sup_{a \in \mathbb{R}} \text{dist}_{\mathbb{R}^2}(p_0(a), p_1(a)),$$

where $p_i(a)$ is the intersection of the graph Γ_i with the line $x + y = a$. Note that if we write $p_i(a) = (x_i, y_i)$,⁹ then $y_i = F_i(x_i)$, then

$$d_L(\mu_0, \mu_1) = \sup \{ \sqrt{2}|x_0 - x_1|; \quad x_0 + F_0(x_0) = x_1 + F_1(x_1) \}.$$

Lévy refers to the convergence with respect to the metric d_L as “convergence from the point of view of Bernoulli”. He shows (see [134, §17]) that a sequence of probability measures μ_n converges in the metric d_L to a probability measure μ if and only if the characteristic functions $\hat{\mu}_n$ converge to the characteristic function $\hat{\mu}$. Hence, the convergence in the metric d_L is the weak convergence so that d_L metrizes the weak convergence. $\square$

2.2.3. The Central Limit Theorem. We can now state and prove the main result of this section.

Theorem 2.2.36 (Central Limit Theorem). *Suppose that $(X_n \in L^2(\Omega, \mathcal{S}, \mathbb{P}))_{n \geq 1}$ is a sequence of i.i.d. random variables with common mean μ and common variance v . Set*

$$\bar{X}_n = X_n - \mu, \quad \bar{S}_n = \sum_{k=1}^n (X_k - \mu), \quad Z_n = \frac{1}{\sqrt{nv}} \bar{S}_n = \frac{1}{\sqrt{nv}} \left(\sum_{k=1}^n X_k - n\mu \right).$$

Then $Z_n \Rightarrow N(0, 1)$.

Proof. According to Lévy's continuity theorem it suffices to show that

$$\lim_{n \rightarrow \infty} \Phi_{Z_n}(\xi) = \Phi_{\Gamma_1}(\xi) = e^{-\frac{\xi^2}{2}}.$$

⁹At a point of discontinuity this reads $y_i \in (F_i(x_i - 0), F_i(x_i))$.

Observe that $\bar{X}_n$ are i.i.d. with mean 0 and variance v , while Z_n has mean 0 and variance 1. Denote by $\Phi(\xi)$ their common characteristic function, $\Phi(\xi) = \mathbb{E}[e^{i\xi\bar{X}_1}]$. We have

$$\Phi_{Z_n}(\xi) = \Phi_{\bar{S}_n/\sqrt{nv}}(\xi) = \Phi_{\bar{S}_n}(\xi/\sqrt{nv}) = \mathbb{E}\left[\prod_{k=1}^n \exp\left(i\frac{\xi}{\sqrt{nv}}\bar{X}_k\right)\right]$$

(the variables $\exp\left(i\frac{\xi}{\sqrt{nv}}\bar{X}_k\right)$, $1 \leq k \leq n$ are independent)

$$= \prod_{k=1}^n \mathbb{E}\left[\exp\left(i\frac{\xi}{\sqrt{nv}}\bar{X}_k\right)\right] = \Phi\left(\xi/\sqrt{nv}\right)^n.$$

Proposition 2.2.25 shows that the function $\Phi(\eta)$ is C^2 , so as $\eta \rightarrow 0$ we have

$$\Phi(\eta) = \Phi(0) + \Phi'(0)\eta + \frac{1}{2}\Phi''(0)\eta^2 + o(\eta^2) = 1 + i\mathbb{E}[\bar{X}_1]\eta - \frac{1}{2}\mathbb{E}[\bar{X}_1^2]\eta^2 + o(\eta^2)$$

$$(\mathbb{E}[\bar{X}_1] = 0, \mathbb{E}[\bar{X}_1^2] = \text{Var}[\bar{X}_1] = v)$$

$$= 1 - \frac{v}{2}\eta^2 + o(\eta^2).$$

Now let $\eta = \xi/\sqrt{nv}$, $n \gg 0$. We deduce

$$\Phi\left(\xi/\sqrt{nv}\right)^n = \left(1 - \frac{\xi^2}{2n} + o(1/n)\right)^n.$$

At this point we want to invoke the following result.

Lemma 2.2.37. Suppose that $(c_n)_{n \geq 1}$ is a convergent sequence of complex numbers and

$$c = \lim_{n \rightarrow \infty} c_n.$$

Then

$$\lim_{n \rightarrow \infty} \left(1 + \frac{c_n}{n}\right)^n = e^c.$$

Assuming Lemma 2.2.37 we deduce that, for any $\xi \in \mathbb{R}$ we have

$$\lim_{n \rightarrow \infty} \Phi_{Z_n}(\xi) = \lim_{n \rightarrow \infty} \left(1 - \frac{\xi^2}{2n} + o(1/n)\right)^n = e^{-\xi^2/2} = \Phi_{\Gamma_1}(\xi).$$

Proof of Lemma 2.2.37. Set $c = a + bi$, $c_n = a_n + b_n i$, so that $a_n \rightarrow a$, $b_n \rightarrow b$. We set

$$z_n = 1 + \frac{c_n}{n} = 1 + \frac{a_n}{n} + \frac{b_n}{n}i.$$

For large n $z_n = r_n e^{i\theta_n}$, where

$$r_n = \sqrt{(1 + a_n/n)^2 + b_n^2/n^2} = \left(1 + 2a/n + o(1/n)\right)^{1/2},$$

$$|\theta_n| < \frac{\pi}{2}, \quad \tan \theta_n = \frac{1}{n} \frac{b_n}{1 + a_n/n}.$$

Thus

$$\theta_n = \arctan\left(\frac{1}{n} \frac{b_n}{1 + a_n/n}\right) = \frac{b}{n} + o(1/n) \text{ as } n \rightarrow \infty.$$

We deduce that as $n \rightarrow \infty$ we have

$$z_n^n = \left(1 + 2a/n + o(1/n)\right)^{n/2} \cdot e^{i(b+o(1))} \rightarrow e^a \cdot e^{ib} = e^c.$$

□

2.2.4. Semigroup approach to CLT. I want to describe an alternate approach to the Central Limit Theorem that bypasses the usage of Fourier transform. The presentation is heavily inspired from [76, Chap. VIII].

Denote by $C_0(\mathbb{R})$ the space of continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\lim_{x \rightarrow \pm\infty} f(x) = 0.$$

This is a Banach space with respect to the sup-norm

$$\|f\| = \sup_{x \in \mathbb{R}} |f(x)|.$$

Denote by $\mathbf{B}$ the Banach space of bounded linear operators

$$T : C_0(\mathbb{R}) \rightarrow C_0(\mathbb{R}).$$

For any Borel probability measure $\mu \in \text{Prob}(\mathbb{R})$ and $f \in C_0(\mathbb{R})$ we denote by $\mathfrak{A}_\mu[f]$ the function $\mathbb{R} \rightarrow \mathbb{R}$ given by

$$\mathfrak{A}_\mu[f](x) = \int_{\mathbb{R}} f(x+y) \mu[dy].$$

The Dominated Convergence Theorem implies that $\mathfrak{A}_\mu[f] \in C_0(\mathbb{R})$. Note that

$$|\mathfrak{A}_\mu[f](x)| \leq \int_{\mathbb{R}} |f(x+y)| \mu[dy] \leq \|f\|, \quad \forall x \in \mathbb{R}$$

so $\mathfrak{A}_\mu$ is a bounded operator $C_0(\mathbb{R}) \rightarrow C_0(\mathbb{R})$ of norm ≤ 1 . We thus have a correspondence

$$\text{Prob}(\mathbb{R}) \ni \mu \mapsto \mathfrak{A}_\mu \in \mathbf{B}.$$

Clearly $\mathfrak{A}_{\delta_0} = \mathbb{1}$. Observe that if Y is a random variable with distribution μ then

$$\mathfrak{A}_\mu[f](x) = \mathbb{E}[f(x+Y)], \quad \forall x \in \mathbb{R}.$$

For this reason, for any random variable Y we set

$$\mathcal{A}_Y := \mathfrak{A}_{\mathbb{P}_Y}.$$

Proposition 2.2.38. *Suppose that $\mu_1, \mu_2 \in \text{Prob}(\mathbb{R})$. Then*

$$\mathfrak{A}_{\mu_1 * \mu_2} = \mathfrak{A}_{\mu_1} \mathfrak{A}_{\mu_2},$$

where $*$ denotes the convolution of probability measures on $\mathbb{R}$.

Proof. Let Y_1, Y_2 be independent random variables such that $\mathbb{P}_{Y_i} = \mu_i, i = 1, 2$. Fix $f \in C_0(\mathbb{R})$. For any $x \in \mathbb{R}$ we have

$$\begin{aligned} \mathfrak{A}_{\mu_1 * \mu_2}[f](x) &= \mathbb{E}[f(x + Y_1 + Y_2)] = \int_{\mathbb{R}^2} f(x + y_1 + y_2) \mu_1 \otimes \mu_2[dy_1 dy_2] \\ &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x + y_1 + y_2) \mu[dy_2] \right) \mu[dy_1] = \int_{\mathbb{R}} \mathfrak{A}_{\mu_2}[f](x + y_1) \mu[dy_1] \\ &= \mathfrak{A}_{\mu_1}[\mathfrak{A}_{\mu_2}[f]](x). \end{aligned}$$

□

Thus the map

$$\text{Prob}(\mathbb{R}) \ni \mu \mapsto \mathfrak{A}_\mu \in \mathbf{B}$$

is a morphism of semigroups.

Proposition 2.2.39. *Let $n \in \mathbb{N}$ and suppose that $\mu_i, \nu_j \in \text{Prob}(\mathbb{R})$, $1 \leq i, j \leq n$. Then, for any $f \in C_0(\mathbb{R})$ we have*

$$\|\mathfrak{A}_{\mu_1 * \dots * \mu_n} f - \mathfrak{A}_{\nu_1 * \dots * \nu_n} f\| \leq \sum_{i=1}^n \|\mathfrak{A}_{\mu_i} f - \mathfrak{A}_{\nu_i} f\|. \quad (2.2.14)$$

Proof. For $n = 2$ we have

$$\begin{aligned} \|\mathfrak{A}_{\mu_1} \mathfrak{A}_{\mu_2} f - \mathfrak{A}_{\nu_1} \mathfrak{A}_{\nu_2} f\| &\leq \|\mathfrak{A}_{\mu_1} (\mathfrak{A}_{\mu_2} - \mathfrak{A}_{\nu_2}) f\| + \|\mathfrak{A}_{\mu_1} \mathfrak{A}_{\nu_2} f - \mathfrak{A}_{\nu_1} \mathfrak{A}_{\nu_2} f\| \\ &= \|\mathfrak{A}_{\mu_1} (\mathfrak{A}_{\mu_2} - \mathfrak{A}_{\nu_2}) f\| + \|\mathfrak{A}_{\nu_2} (\mathfrak{A}_{\mu_1} - \mathfrak{A}_{\nu_1}) f\| \\ &\leq \|(\mathfrak{A}_{\mu_2} - \mathfrak{A}_{\nu_2}) f\| + \|(\mathfrak{A}_{\mu_1} - \mathfrak{A}_{\nu_1}) f\| \end{aligned}$$

since $\|\mathfrak{A}_{\mu_1}\|, \|\mathfrak{A}_{\nu_2}\| \leq 1$. The general case follows inductively using the inequality

$$\|\mathfrak{A}_{\mu_1 * \dots * \mu_n} f - \mathfrak{A}_{\nu_1 * \dots * \nu_n} f\| \leq \|\mathfrak{A}_{\mu_1} (\mathfrak{A}_{\mu_2 * \dots * \mu_n} f - \mathfrak{A}_{\nu_2 * \dots * \nu_n} f)\| + \|\mathfrak{A}_{\mu_2 * \dots * \mu_n} f - \mathfrak{A}_{\nu_2 * \dots * \nu_n} f\|.$$

□

Define inductively

$$C_0^k(\mathbb{R}) = \{ f \in C^1(\mathbb{R}) \cap C_0^{k-1}(\mathbb{R}); f' \in C_0^{k-1}(\mathbb{R}) \}.$$

Theorem 2.2.40. *Let $(\mu_n)_{n \in \mathbb{N}}$ be a sequence in $\text{Prob}(\mathbb{R})$ and $\mu \in \text{Prob}(\mathbb{R})$. The following statements are equivalent.*

- (i) *The sequence (μ_n) converges weakly to μ .*
- (ii) *For any $f \in C_0(\mathbb{R})$*

$$\lim_{n \rightarrow \infty} \|\mathfrak{A}_{\mu_n} f - \mathfrak{A}_{\mu} f\| = 0.$$

- (iii) *There exists $k \in \mathbb{N}_0$ such that, for any $f \in C_0^k(\mathbb{R})$*

$$\lim_{n \rightarrow \infty} \|\mathfrak{A}_{\mu_n} f - \mathfrak{A}_{\mu} f\| = 0.$$

Proof. Clearly (ii) $\Rightarrow$ (iii). To prove that (iii) $\Rightarrow$ (i) note that for any smooth compactly supported function $f \in C_0^k(\mathbb{R})$ we have

$$\mu_n[f] = \mathfrak{A}_{\mu_n}[f](0) \rightarrow \mathfrak{A}_{\mu}[f](0) = \mu[f].$$

Now conclude using Theorem 2.2.14.

(i) $\Rightarrow$ (ii) Let $f \in C_0(\mathbb{R})$. For each $x \in \mathbb{R}$ we define

$$f_x : \mathbb{R} \rightarrow \mathbb{R}, \quad f_x(y) = f(x + y), \quad \forall y \in \mathbb{R}.$$

Then

$$\mathfrak{A}_{\mu_n} f(x) = \mu_n[f_x].$$

Since f is uniformly continuous the map

$$\mathbb{R} \ni x \mapsto f_x \in C_0(\mathbb{R})$$

is also uniformly continuous with respect to the sup-norm.

Fix $\varepsilon > 0$. Since $\mu_n \Rightarrow \mu$ there exists $M > 0$ such that

$$\mu_n[\{|y| > M\}] = \mu_n[\{|y| > M\}] < \varepsilon, \quad \forall n \in \mathbb{N}$$

We can assume that

$$\mu_n[\{|Y| \leq M\}] \rightarrow \mu[\{|Y| \leq M\}].$$

We have

$$\begin{aligned} |\mu_n[f_x] - \mu[f]| &\leq \left| \int_{[-M,M]} f_x(y) \mu_n[dy] - \int_{[-M,M]} f_x(y) \mu[dy] \right| \\ &\quad + \int_{|y|>M} |f| \mu_n[dy] + \int_{|y|>M} |f| \mu[dy] \\ &\leq \left| \int_{[-M,M]} f_x(y) \mu_n[dy] - \int_{[-M,M]} f_x(y) \mu_n[dy] \right| + 2\varepsilon \|f\|. \end{aligned}$$

Hence

$$\sup_{x \in \mathbb{R}} |\mu_n[f_x] - \mu[f]| \leq \sup_{x \in \mathbb{R}} \left| \int_{[-M,M]} f_x(y) \mu_n[dy] - \int_{[-M,M]} f_x(y) \mu_n[dy] \right| + 2\varepsilon \|f\|.$$

Since $f \in C_0(\mathbb{R})$, $\forall \varepsilon > 0$ there exists $K > 0$ such that

$$\sup_{y \in [-M,M]} |f_x(y)| < \varepsilon, \quad \forall |x| > K.$$

Hence

$$\left| \int_{[-M,M]} f_x(y) \mu_n[dy] - \int_{[-M,M]} f_x(y) \mu_n[dy] \right| < 2\varepsilon, \quad \forall |x| > K, \quad \forall n \in \mathbb{N}. \quad (2.2.15)$$

We deduce from (2.2.15) that

$$\sup_{|x|>K} |\mu_n[f_x] - \mu[f]| \leq 2\varepsilon + 2\varepsilon \|f\|. \quad (2.2.16)$$

Consider now the continuous functions

$$g, g_n : [-K, K] \rightarrow \mathbb{R}, \quad g_n(x) = \int_{[-M,M]} f_x(y) \mu_n[dy], \quad g(x) = \int_{[-M,M]} f_x(y) \mu[dy].$$

We deduce

$$g_n(x) \rightarrow g(x), \quad \forall x \in [-K, K].$$

The sequence (g_n) is equicontinuous since $x \mapsto f_x$ is uniformly continuous with respect to the sup-norm. Hence g_n converges uniformly to g on $[-K, K]$, i.e.,

$$\lim_{n \rightarrow \infty} \sup_{|x| \leq K} |g_n(x) - g(x)| = 0.$$

We have

$$\sup_{|x| \leq K} |\mu_n[f_x] - \mu[f]| \leq \sup_{|x| \leq K} |g_n(x) - g(x)| + 2\varepsilon \|f\|.$$

Hence

$$\limsup_{n \rightarrow \infty} \sup_{|x| \leq K} |\mu_n[f_x] - \mu[f]| \leq 2\varepsilon \|f\|.$$

Using (2.2.16) we deduce that $\forall \varepsilon > 0$ we have

$$\limsup_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} |\mu_n[f_x] - \mu[f]| \leq 2\varepsilon + 2\varepsilon \|f\|.$$

This proves (ii). □

Proposition 2.2.41. Suppose that Y is a random variable such that

$$\mathbb{E}[Y^2] = \sigma^2 > 0, \quad \mathbb{E}[Y] = 0. \quad (2.2.17)$$

Then for any $f \in C_0^3(\mathbb{R})$, $t > 0$ and $r > 0$ we have

$$\left\| (\mathfrak{A}_{tY} f - f) - \frac{t^2 \sigma^2}{2} f'' \right\| \leq \left(\frac{1}{2} r (t\sigma)^2 + t^2 \int_{|y| > r/t} y^2 \mu[dy] \right) \|f\|_{C^3} \quad (2.2.18)$$

Proof. Set $\mu := \mathbb{P}_Y$. Let $f \in C_0^3(\mathbb{R})$. Using (2.2.17) we deduce that

$$\begin{aligned} (\mathfrak{A}_{tY} - \mathbb{1})f(x) &= \int_{\mathbb{R}} \left(f(x + ty) - f(x) - tyf'(x) \right) \mu[dy], \\ (\mathfrak{A}_{tY} - \mathbb{1})f(x) - \frac{t^2 \sigma^2}{2} f''(x) &= \int_{\mathbb{R}} \underbrace{\left(f(x + ty) - f(x) - f'(x)ty - \frac{1}{2} f''(x)ty^2 \right)}_{=U_t(x,y)} \mu[dy]. \end{aligned}$$

Taylor's formula with Lagrange remainder yields

$$f(x + ty) - f(x) - f'(x)ty = \frac{1}{2} f''(\xi)ty^2,$$

for some $\xi = \xi_{x,y} \in (x, x + ty)$. Hence

$$\begin{aligned} \left| f(x + ty) - f(x) - f'(x)ty - \frac{t^2}{2} f''(x)ty^2 \right| &= \left| \frac{1}{2} f''(\xi) - \frac{1}{2} f''(x) \right| \cdot ty^2 \\ &\leq \min \left(t^2 \|f\|_{C^2} |y|^2, \frac{1}{2} \|f\|_{C^3} t^3 |y|^3 \right) \\ &\leq t \|f\|_{C^3} \min \left(|y|^2, \frac{1}{2} t |y|^3 \right), \quad \forall t > 0, \quad x, y \in \mathbb{R}. \end{aligned}$$

Hence

$$0 \leq U_t(x, y) \leq t^2 \|f\|_{C^3} \min \left(|y|^2, \frac{1}{2} t |y|^3 \right), \quad \forall t > 0, \quad x, y \in \mathbb{R}.$$

For any $R > 0$ we have

$$\begin{aligned} \left\| (\mathfrak{A}_{tY} - \mathbb{1})f - \frac{t^2 \sigma^2}{2} f'' \right\| &\leq t^2 \|f\|_{C^3} \int_{\mathbb{R}} \min \left(|y|^2, \frac{1}{2} t |y|^3 \right) \mu[dy] \\ &\leq \left(\frac{1}{2} \int_{|y| < R} t^3 |y|^3 \mu[dy] + t^2 \int_{|y| > R} |y|^2 \mu[dy] \right) \|f\|_{C^3} \\ &\leq \left(\frac{1}{2} t^3 R \int_{\mathbb{R}} y^2 \mu[dy] + t^2 \int_{|y| > R} y^2 \mu[dy] \right) \|f\|_{C^3} \\ &= \left(\frac{1}{2} t R (t\sigma)^2 + t^2 \int_{|y| > R} y^2 \mu[dy] \right) \|f\|_{C^3}. \end{aligned}$$

Now set $R := r/t$.

□

Corollary 2.2.42. *Suppose that X is a random variable such that*

$$\mathbb{E}[X^2] = \sigma^2, \quad \mathbb{E}[X] = 0.$$

Then for any $f \in C_0^3(\mathbb{R})$ we have

$$\lim_{t \searrow 0} \left\| \frac{1}{t} (\mathfrak{A}_{t^{1/2}X} f - f) - \frac{\sigma^2}{2} f'' \right\| = 0.$$

Theorem 2.2.43 (Lindeberg). *Suppose that $(X_n)_{n \geq 1}$ be a sequence of independent random variables with mean zero and variances*

$$\mathbb{E}[X_n^2] = \sigma_n^2.$$

Set

$$S_n^2 = \sum_{k=1}^n \sigma_k^2$$

and assume that the variables (X_n) satisfy the Lindeberg condition

$$\forall \varepsilon > 0, \quad \lim_{n \rightarrow \infty} \frac{1}{S_n^2} \sum_{k=1}^n \int_{|x| > \varepsilon S_n} |x|^2 \mathbb{P}_{X_k} [dx] = 0. \quad (2.2.19)$$

Then the random variables

$$\bar{X}_n = \frac{1}{S_n} (X_1 + \cdots + X_n)$$

converge in distribution to a standard normal random variable.

Proof. We set

$$v_n := \sup_{1 \leq k \leq n} \frac{\sigma_k}{S_n}.$$

Lemma 2.2.44.

$$\lim_{n \rightarrow \infty} v_n = 0. \quad (2.2.20)$$

Proof. Observe that the Lindeberg condition can be rewritten as

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \int_{|z| > \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz] = 0$$

Observe that for any $\varepsilon > 0$ and any $1 \leq k \leq n$ we have

$$\begin{aligned} \frac{\sigma_k^2}{S_n^2} &= \text{Var} [S_n^{-1} X_k] = \int_{|z| \leq \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz] + \int_{|z| > \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz] \\ &\leq \varepsilon^2 + \int_{|z| > \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz] \leq \varepsilon^2 + \sum_{k=1}^n \int_{|z| > \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz]. \end{aligned}$$

Thus, for any $\varepsilon > 0$ and any $n \in \mathbb{N}$ we have

$$v_n^2 \leq \varepsilon^2 + \sum_{k=1}^n \int_{|z| > \varepsilon} z^2 \mathbb{P}_{S_n^{-1} X_k} [dz].$$

The equality (2.2.20) now follows from the Lindeberg condition. $\square$

Let $(Y_n)_{n \in \mathbb{N}}$ be independent *normal* variables with mean zero and variances

$$\text{Var} [Y_n^2] = \sigma_n^2.$$

Then

$$\bar{Y}_n = \frac{1}{S_n} (Y_1 + \cdots + Y_n)$$

is a standard normal random variable. It suffices to show that

$$\lim_{n \rightarrow \infty} \|\mathfrak{A}_{\bar{X}_n} - \mathfrak{A}_{\bar{Y}_n} f\| = 0, \quad \forall f \in C_0^3(\mathbb{R}). \quad (2.2.21)$$

Fix $f \in C_0^3(\mathbb{R})$ and $\varepsilon > 0$. Using (2.2.14) we deduce

$$\|\mathfrak{A}_{\bar{X}_n} - \mathfrak{A}_{\bar{Y}_n} f\| \leq \sum_{k=1}^n \|\mathfrak{A}_{S_n^{-1}X_k} - \mathfrak{A}_{S_n^{-1}Y_k} f\|$$

Using (2.2.18) with $r = \varepsilon$ and $t = S_n^{-1}$ we deduce that

$$\begin{aligned} \|\mathfrak{A}_{S_n^{-1}X_k} - \mathfrak{A}_{S_n^{-1}Y_k} f\| &\leq \left\| (\mathfrak{A}_{S_n^{-1}X_k} - \mathbb{1})f - \frac{\sigma_k^2}{2S_n^2} f'' \right\| + \left\| (\mathfrak{A}_{S_n^{-1}Y_k} - \mathbb{1})f - \frac{\sigma_k^2}{2S_n^2} f'' \right\| \\ &\leq \left(\frac{\varepsilon \sigma_k^2}{2S_n^2} + \frac{1}{S_n^2} \int_{|x| > \varepsilon S_n} x^2 \mathbb{P}_{X_k} [dx] \right) \|f\|_{C^3} \\ &\quad + \left(\frac{\varepsilon \sigma_k^2}{2S_n^2} + \frac{1}{S_n^2} \int_{|y| > \varepsilon S_n} y^2 \Gamma_{\sigma_k^2} [dy] \right) \|f\|_{C^3}, \end{aligned}$$

where $\Gamma_{\sigma_k^2}$ denotes the normal distribution with mean zero and variance σ_k^2 . Hence

$$\begin{aligned} \|\mathfrak{A}_{\bar{X}_n} - \mathfrak{A}_{\bar{Y}_n} f\| &\leq \underbrace{\left(\sum_{k=1}^n \frac{\varepsilon \sigma_k^2}{S_n^2} \right)}_{=:\varepsilon \|f\|_{C^3}} \|f\|_{C^3} \\ &\quad + \left(\sum_{k=1}^n \frac{1}{S_n^2} \int_{|x| > \varepsilon S_n} x^2 \mathbb{P}_{X_k} [dx] + \sum_{k=1}^n \frac{1}{S_n^2} \int_{|y| > \varepsilon S_n} y^2 \Gamma_{\sigma_k^2} [dy] \right) \|f\|_{C^3} \\ &= \varepsilon \|f\|_{C^3} + \underbrace{\left(\sum_{k=1}^n \frac{1}{S_n^2} \int_{|x| > \varepsilon S_n} x^2 \mathbb{P}_{X_k} [dx] \right)}_{=:A_n} \|f\|_{C^3} \\ &\quad + \underbrace{\left(\sum_{k=1}^n \frac{1}{S_n^2} \int_{|y| > \varepsilon S_n} y^2 \Gamma_{\sigma_k^2} [dy] \right)}_{=:B_n} \|f\|_{C^3}. \end{aligned}$$

The Lindeberg condition implies that $A_n \rightarrow 0$ as $n \rightarrow \infty$. To deal with B_n note that

$$\frac{1}{S_n^2} \int_{|y| > \varepsilon S_n} y^2 \Gamma_{\sigma_k^2} [dy] = \frac{\sigma_k^2}{S_n^2} \int_{|z| > \varepsilon S_n \sigma_k} y^2 \Gamma_1 [dz] \leq \frac{\sigma_k^2}{S_n^2} \int_{|z| > \varepsilon / v_n} y^2 \Gamma_1 [dz]$$

Hence

$$B_n \leq \int_{|z| > \varepsilon / v_n} y^2 \Gamma_1 [dz].$$

The equality (2.2.20) implies $B_n \rightarrow 0$. □

Remark 2.2.45. (a) The above argument is due to H. F. Trotter [210]. The correspondence

$$\text{Prob}(\mathbb{R}) \ni \mu \mapsto \mathfrak{A}_\mu \in \mathcal{B}$$

used in the above proof has a wider range of applications and we refer to [76] for more information.

(b) Note that if the random variables X_n are also identically distributed with common variances σ^2 , then $S_n^2 = n\sigma^2$. then

$$\frac{1}{S_n^2} \sum_{k=1}^n \mathbb{E}[\mathbf{I}_{\{|X_k| > tS_n\}} X_k^2] = \frac{1}{\sigma^2} \mathbb{E}[\mathbf{I}_{\{|X_1| > t\sigma\sqrt{n}\}} X_1^2] \rightarrow 0$$

as $n \rightarrow \infty$. Hence the Lindeberg's condition is satisfied when the random variables are i.i.d..

If $p > 2$, then Hölder's inequality implies

$$\begin{aligned} \mathbb{E}[\mathbf{I}_{\{|X_k| > tS_n\}} X_k^2] &\leq \mathbb{P}[\{|X_k| > tS_n\}]^{1-\frac{2}{p}} \mathbb{E}[|X_k|^p]^{\frac{2}{p}} \\ &\leq \left(\frac{\mathbb{E}[|X_p|^p]}{t^p S_n^p} \right)^{1-\frac{2}{p}} \mathbb{E}[|X_k|^p]^{\frac{2}{p}} = \frac{1}{t^{p-2} S_n^{p-2}} \mathbb{E}[|X_k|^p]. \end{aligned}$$

This shows that Lindeberg condition is also satisfied if the sequence satisfied *Lyapunov's condition* of order p

$$\lim_{n \rightarrow \infty} \frac{1}{S_n^p} \sum_{k=1}^n \mathbb{E}[|X_k|^p] = 0.$$

For even more general versions of the CLT we refer to [91, 169]. □

2.3. Concentration inequalities

Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables with mean 0. Let

$$S_n := X_1 + \cdots + X_n.$$

The Strong Law of Large Numbers shows that $\frac{1}{n}S_n \rightarrow 0$ a.s. A concentration inequality offers a quantitative information on the probability that $\frac{1}{n}S_n$ deviates from 0 by a given amount ε . More concretely, it gives an upper bound for the probability that $\frac{1}{n}|S_n| > \varepsilon$. If the random variables X_n have finite second moments, $\sigma^2 = \text{Var}[X_1]$, then we have seen that Chebyshev's inequality yields the estimate

$$\mathbb{P}[|S_n| > n\varepsilon] = \mathbb{P}[S_n^2 > n^2\varepsilon^2] < \frac{\text{Var}[S_n]}{n^2\varepsilon^2} = \frac{\sigma^2}{n\varepsilon^2}.$$

In the proof of Theorem 2.1.8 we have shown that if the variables X_n have a stronger integrability property namely $\mathbb{E}[X_n^4] < \infty$, then there exists a constant $C > 0$ such that for any $\varepsilon > 0$ and any $\varepsilon > 0$ we have

$$\mathbb{P}[|S_n| > n\varepsilon] \leq \frac{C}{n^2\varepsilon^4},$$

showing that $\frac{1}{n}S_n$ is even more concentrated around its mean. Loosely speaking, we expect higher concentration around if X_n have lighter tails, i.e., the probabilities

$$\mathbb{P}[|X_n| > x]$$

decay faster as $x \rightarrow \infty$.

In this section we want to describe some quantitative results stating that, under appropriate light-tail assumptions, for any $\varepsilon > 0$ the probability $\mathbb{P}[|S_n| > n\varepsilon]$ decays exponentially fast to 0 as

$n \rightarrow \infty$. The subject of concentration inequalities has witnessed an explosive growth in the last three decades so we will only be able to scratch the surface. For more on this subject I refer to [24].

2.3.1. The Chernoff bound. Many useful concentration inequalities are based on the *Chernoff method*. Let us describe its basics.

Suppose that X is a centered, i.e., mean zero, random variable such that

$$\mathbb{M}_X(\lambda) := \mathbb{E}[e^{\lambda X}] < \infty, \quad \forall \lambda \in J,$$

where J is an open interval containing the origin. We set

$$J_{\pm} := \{\lambda \in J; \pm \lambda > 0\}.$$

Note that this implies that X has moments of any order and thus it imposes severe restrictions on the tail of X . We define the *cumulant* of X to be the function,

$$\Psi_X : J \rightarrow \mathbb{R}, \quad \Psi_X(\lambda) := \log \mathbb{M}_X(\lambda).$$

The function $x \mapsto e^{tx}$ is convex and Jensen's inequality shows

$$\mathbb{E}[e^{\lambda X}] \geq e^{\lambda \mathbb{E}[X]} = 1$$

so $\Psi_X(\lambda) \geq 0$.

For $x > 0$ we have

$$\mathbb{P}[X > x] = \mathbb{P}[e^{\lambda X} > e^{\lambda x}] \leq \frac{1}{e^{\lambda x}} \mathbb{E}[e^{\lambda X}], \quad \forall \lambda \in J_+,$$

where at the last step we used Markov's inequality. The crucial idea of Chernoff's method is the fact that in the right-hand-side we have many λ 's to choose from so we will choose that λ that minimizes the right-hand-side. Here are the details.

We have

$$\mathbb{P}[X > x] \leq e^{-(x\lambda - \Psi_X(\lambda))}, \quad \forall \lambda \in J_+ = (0, r).$$

Set

$$I_+(x) := \sup_{\lambda \in J_+} (x\lambda - \Psi_X(\lambda)).$$

We obtain in this fashion the *Chernoff bound*

$$\mathbb{P}[X > x] \leq e^{-I_+(x)}, \quad I_+(x) := \sup_{\lambda \in (0, r)} (x\lambda - \Psi_X(\lambda)), \quad \forall x > 0. \quad (2.3.1)$$

Note that $I_+(x) \geq 0$ since $\Psi_X(\lambda) \geq 0$. Arguing in a similar fashion we deduce

$$\mathbb{P}[X < x] \leq e^{-I_-(x)}, \quad I_-(x) := \sup_{\lambda \in J_-} (x\lambda - \Psi_X(\lambda)), \quad \forall x < 0. \quad (2.3.2)$$

More generally, if X has a nonzero mean μ , then $\bar{X} = X - \mu$ is centered. If $\mathbb{E}[e^{\lambda X}]$ exists for $\lambda \in J$, then

$$\Psi_{\bar{X}}(\lambda) = \Psi_X(\lambda) - \lambda\mu,$$

and we deduce

$$\mathbb{P}[X > x + \mu] \leq e^{-I_+(x)}, \quad I_+(x) := \sup_{\lambda \in J_+} ((x + \mu)\lambda - \Psi_X(\lambda)), \quad \forall x > 0. \quad (2.3.3)$$

and

$$\mathbb{P}[X < x + \mu] \leq e^{-I_-(x)}, \quad I_-(x) := \sup_{\lambda \in J_-} ((x + \mu)\lambda - \Psi_X(\lambda)), \quad \forall x < 0. \quad (2.3.4)$$

Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables such that

$$\mathbb{M}(\lambda) = \mathbb{M}_{X_k}(\lambda) < \infty,$$

for any λ in an open interval J containing 0. Set

$$\mu := \mathbb{E}[X_k], \quad S_n := X_1 + \cdots + X_n.$$

Then

$$\mathbb{E}[S_n] = n\mu, \quad \mathbb{M}_{S_n}(\lambda) = \mathbb{M}(\lambda)^n, \quad \Psi_{S_n}(\lambda) = n\Psi(\lambda).$$

We deduce that

$$\sup_{\lambda \in J_+} ((nx + n\mu)\lambda - \Psi_{S_n}(\lambda)) = nI_+(x), \quad \forall x > 0,$$

and

$$\sup_{\lambda \in J_-} ((nx + n\mu)\lambda - \Psi_{S_n}(\lambda)) = nI_-(x), \quad \forall x < 0.$$

We deduce

$$\mathbb{P}\left[\frac{1}{n}S_n - \mu > x\right] = \mathbb{P}[S_n - n\mu > nx] \leq e^{-nI_+(x)}, \quad \forall x > 0, \quad (2.3.5a)$$

$$\mathbb{P}\left[\frac{1}{n}S_n - \mu < x\right] = \mathbb{P}[S_n - n\mu < nx] \leq e^{-nI_-(x)}, \quad \forall x < 0. \quad (2.3.5b)$$

In particular

$$\mathbb{P}\left[\left|\frac{1}{n}S_n - \mu\right| > x\right] \leq e^{-nI_+(x)} + e^{-nI_-(-x)}, \quad \forall x > 0. \quad (2.3.6)$$

We have reached a remarkable conclusion. The assumption $\mathbb{M}(\lambda) < \infty$ for λ in an open neighborhood of the origin implies that the probability that the empirical mean $\frac{1}{n}S_n$ deviates from the theoretical mean μ by a fixed amount x decays exponentially to 0 as $n \rightarrow \infty$. In other words, $\frac{1}{n}S_n$ is highly concentrated around its mean and the above inequalities quantify this fact.

To gain some more insight on the above estimates it is useful to list a few properties of the function $I_+(x)$

Proposition 2.3.1. *Suppose that the centered random variable X satisfies*

$$\mathbb{M}_X(\lambda) = \mathbb{E}[e^{\lambda X}] < \infty, \quad \forall \lambda \in J,$$

where $J \subset \mathbb{R}$ is an open interval containing 0. Set

$$J_{\pm} := \{\lambda \in J; \pm\lambda > 0\}, \quad \Psi_X(\lambda) := \log \mathbb{M}_X(\lambda).$$

Then the following hold.

- (i) $\mathbb{M}_X(0) = 1, \mathbb{M}'_X(0) = 0, \mathbb{M}''_X(0) = \text{Var}[X]$.
- (ii) The function $J \ni \lambda \mapsto \Psi_X(\lambda) \in \mathbb{R}$ is convex and nonnegative. Moreover $\Psi''_X(0) > 0$.
- (iii) The function

$$I : \mathbb{R} \rightarrow [0, \infty], \quad I(x) = \sup_{\lambda \in J} (\lambda x - \Psi_X(\lambda))$$

is convex. If

$$I_{\pm}(x) = I(x) = \sup_{\lambda \in J_{\pm}} (\lambda x - \Psi_X(\lambda)),$$

then $I_{\pm}(x) = I(x)$ for $\pm x > 0$.

- (iv) $I(x) > 0$ if $x \neq 0$.

Proof. (i) Proposition 1.3.20 implies that $\mathbb{M}_X^{(k)}(0) = \mathbb{E}[X^k]$, $\forall k = 0, 1, 2, \dots$

(ii) To prove that $\Psi_X(\lambda)$ is convex let $t_1, t_2 \in (0, 1)$ such that $t_1 + t_2 = 1$. Then, using Hölder's inequality with $p = \frac{1}{t_1}$ and $q = \frac{1}{t_2}$ we deduce that for any $\lambda_1, \lambda_2 \in \mathbb{R}$ we have

$$\mathbb{E}[e^{t_1 \lambda_1 X + t_2 \lambda_2 X}] \leq \mathbb{E}[(e^{t_1 \lambda_1 X})^{1/t_1}]^{t_1} \mathbb{E}[(e^{t_2 \lambda_2 X})^{1/t_2}]^{t_2} = \mathbb{E}[e^{\lambda_1 X}]^{t_1} \mathbb{E}[e^{\lambda_2 X}]^{t_2}.$$

Taking the logarithm of both sides of the above inequality we obtain the convexity of $\Psi_X(\lambda)$. Next observe that

$$\Psi'_X(0) = \frac{\mathbb{M}'_X(0)}{\mathbb{M}_X(0)} = 0.$$

Since $\Psi_X(\lambda)$ is convex its graph sits above the tangent at $\lambda = 0$ so $\Psi_X(\lambda) \geq 0$, $\forall \lambda \in J$.

(iii) For $t_1, t_2 \in (0, 1)$ such that $t_1 + t_2 = 1$ and for $x_1, x_2 > 0$ we have

$$\begin{aligned} I_+(t_1 x_1 + t_2 x_2) &= \sup_{\lambda \in (0, r)} ((t_1 x_1 + t_2 x_2) - \Psi_X(\lambda)) \\ &= \sup_{\lambda \in (0, r)} (t_1(x_1 - \Psi_X(\lambda)) - (t_2 x_2 - \Psi_X(\lambda))) \leq t_1 I_+(x_1) + t_2 I_+(x_2). \end{aligned}$$

Observe that for $x > 0$ we have

$$\lambda x - \Psi_X(\lambda) \leq 0, \quad \forall \lambda \leq 0$$

proving that

$$I(x) = \sup_{\lambda \in J} (\lambda x - \Psi_X(\lambda)) = \sup_{\lambda \in J_+} (\lambda x - \Psi_X(\lambda)).$$

(iv) Observe that

$$\Psi''_X(\lambda) = \frac{\mathbb{M}''_X(\lambda)\mathbb{M}_X(\lambda) - \mathbb{M}'_X(\lambda)^2}{\mathbb{M}_X(\lambda)^2} \quad (2.3.7)$$

so $\Psi''_X(0) = \mathbb{M}''_X(0) = \text{Var}[X] > 0$. This proves that $\lambda x - \Psi_X(\lambda) > 0$ for $|\lambda|$ small and $x \neq 0$ so $I(x) > 0$ if $x \neq 0$. $\square$

Remark 2.3.2. As explained in [179, §12], to any convex lower semicontinuous function $f : \mathbb{R}^n \rightarrow (0, \infty]$ we can associate a *conjugate*

$$f^* : \mathbb{R}^n \rightarrow (-\infty, \infty], \quad f^*(p) = \sup_{x \in \mathbb{R}^n} (\langle p, x \rangle - f(x)),$$

where $\langle -, - \rangle$ denotes the canonical inner product in $\mathbb{R}^n$. One can show that f^* is also convex and lower semicontinuous and $f = (f^*)^*$. The conjugate f^* is sometimes called the *Fenchel-Legendre conjugate* of f . Observe that $I(x)$ is the conjugate of the convex function $\Psi_X(\lambda)$. $\square$

Example 2.3.3. Suppose that $X \sim N(0, 1)$. Then, for any $\lambda \in \mathbb{R}$,

$$\mathbb{M}_X(\lambda) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{\lambda x} e^{-x^2/2} dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-(x^2 - 2\lambda x + \lambda^2)/2} e^{\lambda^2/2} dx = e^{\lambda^2/2}.$$

Note that $Y = \sigma X \sim N(0, \sigma^2)$ and

$$\mathbb{M}_Y(\lambda) = \mathbb{M}_X(\sigma \lambda) = e^{\sigma^2 \lambda^2/2}, \quad \Psi_Y(\lambda) = \frac{\sigma^2 \lambda^2}{2}.$$

The supremum

$$I(x) := \sup_{\lambda \in \mathbb{R}} \left(x\lambda - \frac{\sigma^2 \lambda^2}{2} \right)$$

is achieved for $\lambda = \lambda_x = \frac{x}{\sigma^2}$ and it is equal to

$$I(x) = \frac{x^2}{2\sigma^2}.$$

In other words, if $X \sim N(0, \sigma^2)$, then

$$\mathbb{P}[|X| > \varepsilon] \leq 2 \max(\mathbb{P}[X < -x], \mathbb{P}[X > x]) \leq 2e^{-\frac{x^2}{2\sigma^2}}.$$

□

Example 2.3.4. Suppose that $X \sim \text{Bin}(p)$. Then $\mathbb{E}[X] = p$, $\mathbb{M}_{X-p}(\lambda) = (q + pe^\lambda)e^{-p\lambda}$. For $x \in \mathbb{R}$ we set

$$f_x(\lambda) := x\lambda - \Psi_{X-p}(\lambda) = (x+p)\lambda - \log(q + pe^\lambda), \quad I(x) = \sup_{\lambda \in \mathbb{R}} f_x(\lambda).$$

We will show that

$$I(x) = \begin{cases} (x+p) \log \frac{x+p}{p} + (q-x) \log \frac{q-x}{q}, & x \in [-q, p], \\ \infty, & x \notin [-q, p]. \end{cases}$$

Observe that

$$f'_x(\lambda) := \frac{d}{d\lambda} f_x(\lambda) = x + p - \frac{pe^\lambda}{q + pe^\lambda}$$

and $f'_x(\lambda) = 0$ if

$$p(x+p-1)e^\lambda = -q(x+p), \quad \text{i.e., } pe^\lambda = q \frac{x+p}{q-x}.$$

This shows that if $\frac{x+p}{q-x} < 0$, i.e., $x \in \mathbb{R} \setminus (-p, q)$, then $f'_x(\lambda) > 0, \forall \lambda$ and $I(x) = \infty$.¹⁰ If $x \in (-p, q)$, then $f_x(\lambda) = 0$ iff

$$\lambda = \log q - \log p + \log(x+p) - \log(q-x) = \log \frac{x+p}{p} - \log \frac{q-x}{q}.$$

$$\begin{aligned} I(x) &= (x+p) \log \frac{x+p}{p} - (x+p) \log \frac{q-x}{q} + \log \frac{q-x}{q} \\ &= (x+p) \log \frac{x+p}{p} + (q-x) \log \frac{q-x}{q}, \quad x \in (-p, q). \end{aligned}$$

One can verify that $I(q) = -\log p$ and $I(-p) = -\log q$. □

Remark 2.3.5. Suppose that $\mathbb{P}, \mathbb{Q}$ are two Borel probability measures on a Lusin space $\mathfrak{X}$ that are mutually absolutely continuous,

$$\mathbb{P} \ll \mathbb{Q} \quad \text{and} \quad \mathbb{Q} \ll \mathbb{P}.$$

We denote by $\rho_{\mathbb{P}|\mathbb{Q}} := \frac{d\mathbb{P}}{d\mathbb{Q}}$ the density of $\mathbb{P}$ with respect to $\mathbb{Q}$. We define the *Kullback-Leibler divergence* (or KL-divergence for brevity)

$$\mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}] := \mathbb{E}_{\mathbb{P}} \left[\frac{d\mathbb{P}}{d\mathbb{Q}} \right] = \int_{\mathfrak{X}} \log \frac{d\mathbb{P}}{d\mathbb{Q}} d\mathbb{P}. \quad (2.3.8)$$

(a) Suppose that $\mathbb{P}$ is the probability distribution $\text{Bin}(p)$,

$$\mathbb{P} = q\delta_0 + p\delta_1.$$

¹⁰Can you think of a simple reason why this happens?

For $x \in (-p, q)$ consider the probability distribution

$$\mathbb{Q}_x = (q - x)\delta_0 + (p + x)\delta_1.$$

Then

$$\mathbb{D}_{KL}[\mathbb{Q}_x \parallel \mathbb{P}] = (x + p) \log \frac{x + p}{p} + (q - x) \log \frac{q - x}{q}.$$

This is the rate $I(x)$ we found in Example 2.3.4.

(b) Let X be a random variable with probability distribution $\mathbb{Q}$ and set $Z := \rho_{\mathbb{P}|\mathbb{Q}}(X)$. Then

$$\begin{aligned} \mathbb{E}[Z] &= \int_{\mathbb{R}} \frac{d\mathbb{P}}{d\mathbb{Q}} d\mathbb{Q} = \int_{\mathbb{R}} d\mathbb{P} = 1, \\ \mathbb{E}[Z \log Z] &= \int_{\mathbb{R}} \frac{d\mathbb{P}}{d\mathbb{Q}} \log \frac{d\mathbb{P}}{d\mathbb{Q}} d\mathbb{Q} = \int_{\mathbb{R}} \log \frac{d\mathbb{P}}{d\mathbb{Q}} d\mathbb{P} = \mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}]. \end{aligned}$$

Thus

$$\mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}] = \mathbb{E}[Z \log Z] - \mathbb{E}[Z] \log \mathbb{E}[Z],$$

showing that KL -divergence is a special case of φ -entropy (1.3.13). More precisely, the above equality shows that

$$\mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}] = \mathbb{H}_{\varphi}[Z], \quad \varphi(z) = z \log z, \quad z > 0.$$

In particular this yields *Gibbs' inequality* or *information inequality*

$$\mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}] \geq 0. \tag{2.3.9}$$

Note that this yields an alternate definition of the KL -divergence

$$\mathbb{D}_{KL}[\mathbb{P} \parallel \mathbb{Q}] := \mathbb{E}_{\mathbb{Q}} \left[\frac{d\mathbb{P}}{d\mathbb{Q}} \log \frac{d\mathbb{P}}{d\mathbb{Q}} \right] = \int_{\mathfrak{X}} \frac{d\mathbb{P}}{d\mathbb{Q}} \log \frac{d\mathbb{P}}{d\mathbb{Q}} d\mathbb{Q}. \tag{2.3.10}$$

Above, we could have used instead of the natural logarithm any logarithm in a base > 1 and reach the same conclusion. In particular, if we work with $\log_2$ and we set

$$\mathbb{D}_2[\mathbb{P} \parallel \mathbb{Q}] = \int_{\mathbb{R}} \log_2 \frac{d\mathbb{Q}}{d\mathbb{P}} d\mathbb{P}.$$

Then Gibbs' inequality continues to hold in this case as well

$$\mathbb{D}_2[\mathbb{P} \parallel \mathbb{Q}] \geq 0. \tag{2.3.11}$$

Let $\mathfrak{X}$ be a finite subset of $\mathbb{R}$. Assume that we are given a function $p : \mathfrak{X} \rightarrow (0, 1]$ such that

$$\sum_{x \in \mathfrak{X}} p(x) = 1$$

so p defines the probability measure

$$\mathbb{P}_p = \sum_{x \in \mathfrak{X}} p(x) \delta_x \in \text{Prob}(\mathbb{R}).$$

Recall that its *Shannon entropy* is (see (2.1.21)) is the quantity

$$\text{Ent}_2[p] = - \sum_{x \in \mathfrak{X}} p(x) \log_2 p(x).$$

The uniform probability measure on $\mathfrak{X}$ is

$$\mathbb{P}_0 = \sum_{x \in \mathfrak{X}} p_0(x) \delta_x = \frac{1}{|\mathfrak{X}|} \sum_{x \in \mathfrak{X}} \delta_x.$$

Note that $\mathbb{P}_p$ and $\mathbb{P}_0$ are mutually absolutely continuous. Gibbs' inequality shows that

$$\mathbb{D}_2[\mathbb{P} \parallel \mathbb{P}_0] \geq 0.$$

On the other hand

$$0 \leq \mathbb{D}_2[\mathbb{P} \parallel \mathbb{P}_0] = \sum_{x \in \mathfrak{X}} \log_2(|\mathfrak{X}| \cdot p(x)) p(x) = \log_2 |\mathfrak{X}| - \text{Ent}_2[p_0].$$

We have obtained again the inequality (2.1.22).

$$\text{Ent}_2[p] \leq \log_2 |\mathfrak{X}| = \text{Ent}_2[p_0]. \quad (2.3.12)$$

□

Example 2.3.6 (Differential entropy). Suppose that $\mathbb{P}$ is a Borel probability measure on $\mathbb{R}$ that is absolutely continuous with respect to the Lebesgue measure λ . Denote by $p(x)$ the corresponding probability density

$$\mathbb{P}[dx] = p(x)\lambda[dx]$$

The *differential entropy* of $\mathbb{P}$ is

$$\text{Ent}[\mathbb{P}] := -\mathbb{D}[\mathbb{P} \parallel \lambda] = - \int_{\mathbb{R}} p(x) \log p(x) dx.$$

Suppose that $\mathbb{P} \sim N(0, \sigma^2)$. Then

$$\log p(x) = -\frac{x^2}{2\sigma^2} - \frac{1}{2} \log(2\pi\sigma^2),$$

so that

$$\text{Ent}[\mathbb{P}] = - \int_{\mathbb{R}} p(x) \log p(x) dx = \frac{1}{2\sigma^2} \text{Var}[X] + \frac{\log(2\pi\sigma^2)}{2} = \frac{1}{2} \log(2\pi e\sigma^2).$$

□

2.3.2. Some applications. Often an explicit description of $\Psi_X(\lambda)$ may either not be possible, or it could be too complicated to be useful. That is why it is more practical to have simple ways of producing upper bounds for the moment generating function.

Definition 2.3.7. A random variable X with mean μ said to be *subgaussian* of type σ^2 , and we write this $X \in \mathbb{G}(\sigma^2)$, if $\mathbb{E}[e^{\lambda X}] < \infty$, $\forall \lambda \in \mathbb{R}$, and

$$\Psi_{X-\mu}(\lambda) \leq \Psi_{N(0, \sigma^2)} = \frac{\sigma^2 \lambda^2}{2}, \quad \forall \lambda \in \mathbb{R} \iff \mathbb{E}[e^{\lambda(X-\mu)}] \leq e^{\frac{\lambda^2 \sigma^2}{2}}, \quad \forall \lambda \in \mathbb{R}. \quad \square$$

Note that if $X \in \mathbb{G}(\sigma^2)$, and $\pm x > 0$, then

$$\sup_{\pm \lambda \geq 0} (x\lambda - \Psi_{X-\mu}(\lambda)) \geq \sup_{\pm \lambda \geq 0} (x\lambda - \frac{\lambda^2 \sigma^2}{2}) = \frac{x^2}{2\sigma^2},$$

and thus

$$\max(\mathbb{P}[X - \mu < -x], \mathbb{P}[X - \mu > x]) \leq e^{-\frac{x^2}{2\sigma^2}}, \quad \forall x > 0, \quad (2.3.13a)$$

$$\mathbb{P}[|X - \mu| > x] \leq 2e^{-\frac{x^2}{2\sigma^2}}, \quad \forall x > 0. \quad (2.3.13b)$$

Observe that if X_1, X_2 are independent random variables and $X_k \in \mathbb{G}(\sigma_k^2)$, $k = 1, 2$, then

$$a_1 X_1 + a_2 X_2 \in \mathbb{G}(a_1^2 \sigma_1^2 + a_2^2 \sigma_2^2), \quad \forall a_1, a_2 \in \mathbb{R}.$$

In particular, if $X_1, \dots, X_n$ are centered, independent random variables in $\mathbb{G}(\sigma^2)$, then we have

$$\frac{1}{n}(X_1 + \dots + X_n) \in \mathbb{G}(\sigma^2/n),$$

and thus we obtain *Hoeffding's inequality*

$$\mathbb{P}\left[\left|\frac{1}{n}(X_1 + \dots + X_n)\right| > x\right] \leq 2e^{-\frac{nx^2}{2\sigma^2}}, \quad \forall x > 0. \quad (2.3.14)$$

Example 2.3.8. Suppose that R is a Rademacher random variable, i.e., it takes only the values ± 1 with equal probabilities. Then

$$\mathbb{E}[e^{\lambda R}] = \cosh \lambda \leq e^{\lambda^2/2}, \quad (2.3.15)$$

where the last inequality is obtained by inspecting the Taylor series of the two terms and using the inequality $2^n n! \leq (2n)!$. Hence $R \in \mathbb{G}(1)$. Similarly, $cR \in \mathbb{G}(1)$, $\forall c \in [0, 1]$. $\square$

For these estimates to be useful we need to have some simple ways of recognizing subgaussian random variables.

Proposition 2.3.9. Suppose that X is a centered random variable, i.e., $\mathbb{E}[X] = 0$. If there exists $C > 0$ such that

$$\mathbb{E}[X^{2k}] \leq k!C^k, \quad \forall k \in \mathbb{N},$$

then $X \in \mathbb{G}(4C)$.

Proof. We rely on a very useful symmetrization trick. Choose a random variable X' independent of X but with the same distribution as X . Then the random variable $Y = X - X'$ is symmetric, i.e., Y and $-Y$ have the same probability distributions. Observe next that since $-X'$ is centered we have

$$\mathbb{E}[e^{-\lambda X'}] \geq e^{-\lambda \mathbb{E}[X']} = 1, \quad \forall \lambda \in \mathbb{R}.$$

We deduce

$$\mathbb{E}[e^{\lambda X}] \leq \mathbb{E}[e^{\lambda X}] \cdot \mathbb{E}[e^{-\lambda X'}] = \mathbb{E}[e^{\lambda(X-X')}] = \sum_{k=0}^{\infty} \frac{\lambda^{2k}}{(2k)!} \mathbb{E}[(X - X')^{2k}].$$

Since the function x^{2k} is convex we have

$$(x + y)^{2k} \leq 2^{2k-1}(x^{2k} + y^{2k}), \quad \forall x, y \in \mathbb{R}$$

so

$$\mathbb{E}[(X - X')^{2k}] \leq 2^{2k} \mathbb{E}[X^{2k}] \leq 2^{2k} k! C^k = \frac{(2k)!}{(2k-1)!!} (2C)^k \leq \frac{(2k)!}{k!} (2C)^k$$

Hence

$$\mathbb{E}[e^{\lambda X}] \leq \sum_{k=0}^{\infty} \frac{(2C\lambda^2)^k}{k!} = e^{2C\lambda^2}.$$

Hence $X \in \mathbb{G}(4C)$. $\square$

Example 2.3.10. Suppose that R is a Rademacher random variable. Clearly

$$\mathbb{E}[R^{2k}] = 1 \leq k!1^k, \quad \forall k \in \mathbb{N}$$

so that $R \in \mathbb{G}(4)$. We see that this estimate is not as good as the one in Example 2.3.8. $\square$

The next result offers a sharper estimate under certain conditions.

Proposition 2.3.11 (Hoeffding's lemma). *Suppose that X is a random variable such that $X \in [a, b]$ a.s.. Then $X \in \mathbb{G}(\sigma^2)$, where $\sigma = \frac{b-a}{2}$, i.e.,*

$$\mathbb{E}[e^{\lambda(X-\mu)}] \leq e^{\frac{\lambda^2(b-a)^2}{8}}, \quad \forall \lambda \in \mathbb{R}. \quad (2.3.16)$$

Proof. Let us first observe that any random variable Y such that $Y \in [a, b]$ a.s. satisfies

$$\text{Var}[Y] \leq \frac{(b-a)^2}{4}.$$

Indeed, if $\mu = \mathbb{E}[Y]$, then $Y - \mu \in [a - \mu, b - \mu]$. If

$$m = \frac{(a - \mu) + (b - \mu)}{2}$$

is the midpoint of $[a - \mu, b - \mu]$, then

$$|(Y - \mu) - m| \leq \frac{b - a}{2}$$

and

$$\text{Var}[Y] \leq \mathbb{E}[(Y - \mu)^2] + m^2 = \mathbb{E}[(Y - \mu - m)^2] \leq \frac{(b-a)^2}{4}.$$

Observe next that we can assume that X is centered. Indeed, if $\mu = \mathbb{E}[X]$, then the centered variable $X - \mu$ satisfies $X - \mu \in [a - \mu, b - \mu]$ and $(b - a) = (b - \mu) - (a - \mu)$.

Denote by $\mathbb{P}$ the probability distribution of X . For any $\lambda \in \mathbb{R}$ we denote by $\mathbb{P}_\lambda$ the probability measure on $\mathbb{R}$ given by

$$\mathbb{P}_\lambda[dx] = \frac{e^{\lambda x}}{\mathbb{E}[e^{\lambda X}]} \mathbb{P}[dx] \quad (2.3.17)$$

Note that $\mathbb{P}_\lambda$ is also supported on $[a, b]$. Since $\mathbb{E}[X] = 0$ we have $\Psi'_X(0) = 0$. We deduce from (2.3.7) that

$$\begin{aligned} \Psi''_X(\lambda) &\stackrel{(2.3.7)}{=} \frac{1}{\mathbb{E}[e^{\lambda X}]} \mathbb{E}[X^2 e^{\lambda X}] - \left(\frac{\mathbb{E}[X e^{\lambda X}]}{\mathbb{E}[e^{\lambda X}]} \right)^2 \\ &= \int_{\mathbb{R}} x^2 \mathbb{P}_\lambda[dx] - \left(\int_{\mathbb{R}} x \mathbb{P}_\lambda[dx] \right)^2. \end{aligned}$$

The last term is the variance of a random variable Z with probability distribution $\mathbb{P}_\lambda$. Since $\mathbb{P}_\lambda$ is supported in $[a, b]$ we have $Z \in [a, b]$ and we deduce

$$\Psi''_X(\lambda) = \text{Var}[Z] \leq \frac{(b-a)^2}{4}. \quad (2.3.18)$$

Using the Taylor approximation with Lagrange remainder we deduce that for some $\xi \in [0, \lambda]$ we have

$$\Psi_X(\lambda) = \underbrace{\Psi_X(0) + \lambda \Psi'_X(0)}_{=0} + \frac{1}{2} \Psi''_X(\xi) \lambda^2 \leq \frac{\lambda^2 (b-a)^2}{8}.$$

Hence $X \in \mathbb{G}((b-a)^2/4)$. □

Hoeffding's Lemma shows that if R is a Rademacher random variable, then $R \in \mathbb{G}(1)$ as in Example 2.3.8. which is an improvement over Proposition 2.3.9.

If $R_1, \dots, R_n$ are independent Rademacher random variables, then for any $c_1, \dots, c_n \in [-1, 1]$ we have $c_k R_k \in \mathbb{G}(1)$ and we deduce from Hoeffding's inequality that

$$\mathbb{P}\left[\left|\frac{1}{n}c_1 R_1 + \dots + c_n R_n\right| > r\right] \leq 2e^{-\frac{nr^2}{2}}. \quad (2.3.19)$$

Example 2.3.12 (The Poincaré phenomenon). Suppose that X is a standard normal random variable and $Y = X^2$

$$\mathbb{M}_Y(\lambda) = \mathbb{E}[e^{\lambda X^2}] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{\frac{(2\lambda-1)x^2}{2}} dt.$$

This integral converges only for $\lambda < \frac{1}{2}$ and in this case it is equal to

$$\mathbb{M}_Y(\lambda) = \frac{1}{\sqrt{1-2\lambda}}.$$

In particular, X^2 is not subgaussian since its moment generating function is not defined for all $\lambda \in \mathbb{R}$. Note that $\mathbb{E}[Y] = \mathbb{E}[X^2] = 1$. Hence

$$\mathbb{M}_{Y-1}(\lambda) = \frac{e^{-\lambda}}{\sqrt{1-2\lambda}}, \quad \Psi_{Y-1}(\lambda) = -\lambda - \frac{1}{2} \log(1-2\lambda).$$

Since $Y \geq 0$ we have $\mathbb{P}[Y-1 < y] = 0$ for $y \leq -1$. For $y \in (-1, \infty)$ the supremum

$$I(y) := \sup_{\lambda < 1/2} (\lambda y - \Psi_{Y-1}(\lambda))$$

is achieved when

$$\frac{d}{d\lambda} (\lambda y - \Psi_{Y-1}(\lambda)) = y + 1 - \frac{1}{1-2\lambda} = 0.$$

Solving this equation for λ we get

$$1-2\lambda = \frac{1}{y+1} \iff \lambda = \frac{y}{2(y+1)}.$$

and

$$I(y) = \frac{y^2}{2(y+1)} + \frac{y}{2(y+1)} - \frac{1}{2} \log(1+y) = \frac{y}{2} - \frac{1}{2} \log(y+1) \geq \frac{y^2}{4}, \quad \forall y > -1.$$

Hence

$$\mathbb{P}[|Y-1| > y] \leq 2e^{-\frac{y^2}{4}}, \quad \forall y \in (0, 1).$$

Suppose now that

$$\vec{X} = (X_1, \dots, X_n)$$

is a Gaussian random vector, where X_k are independent standard normal random variables. The square of its Euclidean norm is the chi-squared random variable

$$Z_n = \|\vec{X}\|^2 = \sum_{k=1}^n X_k^2.$$

We deduce that

$$\mathbb{P}\left[\left|\frac{1}{n}Z_n - 1\right| > y\right] < 2e^{-\frac{ny^2}{4}}, \quad \forall 0 \leq y < 1.$$

Thus, for large n the random vector $\frac{1}{\sqrt{n}}\vec{X}$ is highly concentrated around the unit sphere in $\mathbb{R}^n$. This is one facet of the so called *Poincaré phenomenon*. In Exercise 2.64 we describe another facet of this phenomenon. $\square$

We conclude this section with a remarkable application of the Poincaré phenomenon. Consider a Gaussian random vector in $\mathbb{R}^N$

$$\vec{X} = (X_1, \dots, X_N),$$

where the components X_k are independent standard normal random variables. Note that for any unit vector $\vec{u} = (u_1, \dots, u_N)$ the inner product

$$\langle \vec{u}, \vec{X} \rangle = u_1 X_1 + \dots + u_N X_N$$

is a mean zero Gaussian random variable. Moreover

$$\text{Var} [\langle \vec{u}, \vec{X} \rangle] = \mathbb{E} [|\langle \vec{u}, \vec{X} \rangle|^2] = 1 = \|\vec{u}\|^2.$$

Suppose that we are now given d such independent¹¹ random vectors

$$\vec{X}_j = (X_{1,j}, \dots, X_{N,j}), \quad 1 \leq j \leq d.$$

We obtain a random map

$$A : \mathbb{R}^N \rightarrow \mathbb{R}^d, \quad \mathbb{R}^N \ni \vec{u} \mapsto (V_1, \dots, V_d) := (\langle \vec{u}, \vec{X}_1 \rangle, \dots, \langle \vec{u}, \vec{X}_d \rangle). \quad (2.3.20)$$

If $\|\vec{u}\| = 1$ components of $A\vec{u}$ are independent standard normal random variables so that $\|A\vec{u}\|^2$ is a chi-squared random variable. We set $B := \frac{1}{\sqrt{d}}A$. We deduce from Example 2.3.12 that for any $\varepsilon \in (0, 1)$ and any unit vector $\vec{u}$ we have

$$\mathbb{P} \left[\left| \|B\vec{u}\|^2 - 1 \right| > \varepsilon \right] \leq 2e^{-\frac{d\varepsilon^2}{4}}.$$

Suppose now that we have a cloud of points in a large dimensional Euclidean space

$$C = \{x_1, \dots, x_m\} \subset \mathbb{R}^N, \quad N \gg 1.$$

For $1 \leq i < j \leq m$ we write $v_{ij} = x_j - x_i$. We deduce that

$$\mathbb{P} \left[1 - \varepsilon \leq \frac{\|Bv_{ij}\|}{\|v_{ij}\|} \leq 1 + \varepsilon, \quad \forall 1 \leq i < j \leq m \right] \leq 2 \binom{m}{2} e^{-\frac{d\varepsilon^2}{4}} \leq m^2 e^{-\frac{d\varepsilon^2}{2}}.$$

Now fix a confidence level $0 < p_0 < 1$ and observe that

$$m^2 e^{-\frac{d\varepsilon^2}{2}} < p_0 \iff d\varepsilon^2 > 4 \log \frac{m}{p_0} \iff d > \frac{4}{\varepsilon^2} \log \frac{m}{p_0}.$$

We have thus proved the following remarkable result.

Theorem 2.3.13 (Lindenstrauss-Johnson). *Fix $\varepsilon > 0$ and $p_0 \in (0, 1)$ and a cloud of C of m points in $\mathbb{R}^N$. If*

$$d = d(m, \varepsilon, p_0) := \left\lceil \frac{4}{\varepsilon^2} \log \frac{m}{p_0} \right\rceil, \quad (2.3.21)$$

then, with probability at least $1 - p_0$, the random Gaussian map $B = \frac{1}{\sqrt{d}}A$, where A is described by (2.3.20), distorts very little the relative distances between the points in C , i.e., with probability at least $1 - p_0$

$$(1 - \varepsilon)\|Bx - By\| \leq \|x - y\| \leq (1 + \varepsilon)\|Bx - By\|, \quad \forall x, y \in C.$$

¹¹Independence is meant in probabilistic sense, not linear independence.

□

Remark 2.3.14. Let us highlight some remarkable features of the above result. Note first that the dimension $d(m, \varepsilon, p_0)$ is *independent* of the dimension of the ambient space $\mathbb{R}^N$ where the cloud C resides. Moreover, $d(m, \varepsilon, p_0)$ is substantially smaller than the size N of the cloud.

For example, if we choose the confidence level $p_0 = 10^{-3}$, the distortion factor $\varepsilon = 10^{-1}$ and the size of the cloud $m = 10^{12}$, then

$$\frac{4}{\varepsilon^2} \log \frac{N}{p_0} = 60 \cdot 10^2 \log 10 < 14 \cdot 10^3 \ll 10^{12}.$$

The cloud C could even be chosen in an infinite dimensional Hilbert space and we can choose as ambient space the subspace $\text{span}(C)$ that has dimension $N \leq m$. In this case the vectors $Y_k := \frac{1}{\sqrt{N}} \vec{X}_k$, $k = 1, \dots, d$, have with high confidence norm 1.

$$\mathbb{P}[\|Y_k\| - 1 > \delta, \forall 1 \leq k \leq d] \leq 2de^{-\frac{m\delta^2}{4}}, \quad d \approx C \log m.$$

The vectors Y_k are also, with high confidence, mutually orthogonal. Indeed, Exercise 2.59 shows that for $|r| < \frac{1}{2}$

$$\mathbb{P}[|\langle Y_i, Y_j \rangle| > r, \forall i < j] \leq 2 \binom{d}{2} e^{-\frac{Nr^2}{12}}, \quad d \approx C \log N.$$

This shows that the operator $\frac{1}{\sqrt{d}}A$ is, with high confidence, very close to the orthogonal projection $P_{\vec{X}_1, \dots, \vec{X}_d}$ onto the random d -dimensional¹² subspace $\text{span}\{\vec{X}_1, \dots, \vec{X}_d\}$. This shows that, with high confidence, the operator

$$\sqrt{\frac{N}{d}} P_{\vec{X}_1, \dots, \vec{X}_d}$$

distorts very little the distances between the points in C . The projected cloud has identical size, similar geometry but lives in a subspace of much smaller dimension. □

2.4. Uniform laws of large numbers

Fix a Borel probability measure μ on $\mathbb{R}$. Suppose that

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad n \in \mathbb{N}$$

is a sequence of i.i.d. random variables with common probability distribution μ . For any Borel set $B \subset \mathbb{R}$ the random variables $\mathbf{I}_B(X_n)$ are i.i.d. and have finite means

$$m_B := \mathbb{P}[X_1 \in B] = \mu[B].$$

The Strong Law of Large Numbers shows that the empirical means

$$M_n[B] := \frac{1}{n} (\mathbf{I}_B(X_1) + \dots + \mathbf{I}_B(X_n)) = \frac{\#\{1 \leq k \leq n; X_k \in B\}}{n}$$

converge a.s. to $\mu[B]$. In particular, this provides an asymptotic confirmation of the “frequentist” interpretation of probability as the ratio of favorable cases to the number of possible cases.

If we choose B of the form $(-\infty, x]$, then we obtain the *empirical cdf*

$$F_n(x) = M_n[(-\infty, x]] = \frac{1}{n} \frac{\#\{1 \leq k \leq n; X_k \leq x\}}{n}.$$

¹²It is not hard to see that $\dim \text{span}\{\vec{X}_1, \dots, \vec{X}_d\} = d$ a.s.

This is a *random* quantity (variable), $F_n(x) = F_n(x, \omega)$, $\omega \in \Omega$. For each $n \in \mathbb{N}$, the collection $(F_n(x))_{x \in \mathbb{R}}$ is an example of *empirical process*.

For any $x \in \mathbb{R}$, the random variable $F_n(x)$ converges a.s. to $F(x)$, where F is the cdf of μ

$$F(x) = \mu[(-\infty, x]].$$

For $x \in \Omega$ the set $N_x \subset \Omega$ such that $F_n(x, \omega)$ does not converge to $F(x)$ is negligible but, since $\mathbb{R}$ is not countable, the union

$$N = \bigcup_{x \in \mathbb{R}} N_x$$

need not be negligible. In other words, the set of ω 's such that the *functions* $F_n(-, \omega)$ do not converge pointwisely to the function $F(-)$ need not be negligible. We will show that this is not the case.

2.4.1. The Glivenko-Cantelli theorem. Define

$$D_n = D_n^F : \Omega \rightarrow [0, \infty), \quad D_n(\omega) := \sup_{x \in \mathbb{R}} |F_n(x, \omega) - F(x)|. \quad (2.4.1)$$

For a fixed $\omega \in \Omega$ the sequence of functions $(F_n(-, \omega))_{n \in \mathbb{N}}$ converges uniformly to $F(-)$ if and only if $D_n(\omega) \rightarrow 0$. We will show that this is the case for almost all ω .

Denote by $U(y)$ the cdf of the uniform distribution on $[0, 1]$,

$$U(y) = \begin{cases} 0, & y < 0, \\ y, & y \in [0, 1], \\ 1, & y > 1, \end{cases}$$

and by Q the quantile of F defined in (1.2.5), $Q : [0, 1] \rightarrow \bar{\mathbb{R}}$

$$Q(\ell) := \inf \{x : \ell \leq F(x)\} = \inf F^{-1}([\ell, \infty)) = \inf F^{-1}([\ell, 1]).$$

Lemma 2.4.1. *The function D_n^F is measurable and $D_n^F \leq D_n^U$, with equality if F is continuous.*

Proof. Let us first show that D_n is indeed measurable. We will show that

$$D_n^F = \sup_{x \in \mathbb{Q}} |F_n(x) - F(x)| \quad (2.4.2)$$

According to Proposition 1.1.18(iii) the quantity in the right-hand-side is measurable.

Fix $\omega \in \Omega$. There exists then a sequence of real numbers $(x_k)_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} |F_n(x_k, \omega) - F(x_k)| = D_n(\omega).$$

Now observe that the functions $x \mapsto F_n(x, \omega)$, $F(x)$ are right-continuous so there exists a sequence of rational numbers $(q_k)_{k \in \mathbb{N}}$ such that $q_k > x_k$ and

$$\left| |F_n(x_k, \omega) - F(x_k)| - |F_n(q_k, \omega) - F(q_k)| \right| < \frac{1}{k}.$$

Hence

$$\lim_{k \rightarrow \infty} |F_n(q_k, \omega) - F(q_k)| = \lim_{k \rightarrow \infty} |F_n(x_k, \omega) - F(x_k)|$$

thus proving that the functions (2.4.2) are measurable.

Consider now a sequence of i.i.d. random variables $(Y_n)_{n \in \mathbb{N}}$ uniformly distributed on $[0, 1]$. Denote by U_n the associated empirical c.d.f.-s,

$$U_n(x) = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{(-\infty, x]}(Y_k).$$

Then $X_n = Q(Y_n)$ are i.i.d. with common cdf F . Note that

$$\begin{aligned} U_n(F(x)) - F(x) &= \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{\{Y_k \leq F(x)\}} - F(x) \\ &\stackrel{(1.2.6)}{=} \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{\{Q(Y_k) \leq x\}} - F(x) = F_n(x) - F(x). \end{aligned}$$

Thus

$$\begin{aligned} D_n^F &= \sup_{x \in \mathbb{R}} |F_n(x) - F(x)| = \sup_{x \in \mathbb{R}} |U_n(F(x)) - U(F(x))| \\ &\leq \sup_{y \in \mathbb{R}} |U_n(y) - U(y)| = D_n^U. \end{aligned}$$

Observe that if F is continuous, then $\forall y \in (0, 1)$, $\exists x \in \mathbb{R}$, such that $F(x) = y$ so

$$\sum_{x \in \mathbb{R}} |U(F_n(x)) - U(F(x))| = \sup_{y \in \mathbb{R}} |U_n(y) - U(y)|.$$

□

Theorem 2.4.2 (Glivenko-Cantelli). *Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables with common distribution μ and cdf F . Denote by $F_n(x)$ the empirical cdf-s*

$$F_n(x) = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{(-\infty, x]}(X_k).$$

Then, almost surely, $F_n(x)$ converges uniformly to $F(x)$, i.e.,

$$D_n^F \rightarrow 0 \text{ a.s. as } n \rightarrow \infty,$$

where D_n is defined by (2.4.1).

Proof. Lemma 2.4.1 shows that it suffices to prove the theorem only in the special case when that random variables are uniformly distributed. Thus we assume $F = U$. Note that $U_n(x) = U(x)$ for $x \in \mathbb{R} \setminus [0, 1]$. Thus it suffices to prove that $U_n(x) \rightarrow U(x)$ a.s. uniformly on $[0, 1]$. This is a manifestation of a more general phenomenon.

Lemma 2.4.3. *Suppose that $f_n : [0, 1] \rightarrow \mathbb{R}$ is a sequence nondecreasing functions that converges pointwisely to a function $f : [0, 1] \rightarrow \mathbb{R}$. If the limit function f is continuous, then f_n converges uniformly to f .*

Proof of Lemma 2.4.3. Set

$$D_n^f := \sup_{x \in [0, 1]} |f_n(x) - f(x)|.$$

we will show that $D_n^f \rightarrow 0$ as $n \rightarrow \infty$.

Fix a partition $\mathcal{P}$ of $[0, 1]$, $\mathcal{P} = \{0 = x_0 < x_1 < x_2 < \dots < x_m = 1\}$. Set

$$\|\mathcal{P}\| := \max_{1 \leq k \leq m} (x_k - x_{k-1}), \quad \|\mathcal{P}\|_f := \max_{1 \leq k \leq m} (f(x_k) - f(x_{k-1})).$$

For $x \in [x_{k-1}, x_k]$ and $n \in \mathbb{N}$ we have

$$\begin{aligned} f_n(x_{k-1}) &\leq f_n(x) \leq f_n(x_k), \\ f(x_k) - f_n(x_k) &\leq (f(x_k) - f(x)) + (f(x) - f_n(x)) \leq \|\mathcal{P}\|_f + (f(x) - f_n(x)), \\ f(x) - f_n(x) &\leq f(x) - f(x_{k-1}) + f(x_{k-1}) - f_n(x_{k-1}) \leq \|\mathcal{P}\|_f + f(x_{k-1}) - f_n(x_{k-1}), \end{aligned}$$

Hence

$$\begin{aligned} f(x_k) - f_n(x_k) - \|\mathcal{P}\|_f &\leq f(x) - f_n(x) \leq \|\mathcal{P}\|_f + f(x_{k-1}) - f_n(x_{k-1}), \\ f_n(x_{k-1}) - f(x_{k-1}) - \|\mathcal{P}\|_f &\leq f_n(x) - f(x) \leq \|\mathcal{P}\|_f + f_n(x_k) - f(x_k). \end{aligned}$$

If we set

$$\begin{aligned} D_n^+(\mathcal{P}) &:= \max_{0 \leq k \leq m} (f(x_k) - f_n(x_k)), \quad D_n^-(\mathcal{P}) := \max_{0 \leq k \leq m} (f_n(x_k) - f(x_k)), \\ D_n(\mathcal{P}) &:= \max(D_n^+(\mathcal{P}), D_n^-(\mathcal{P})) \end{aligned}$$

we deduce that for any partition $\mathcal{P}$ of $[0, 1]$ we have

$$D_n(\mathcal{P}) = \max_{0 \leq k \leq m} |f(x_k) - f_n(x_k)|,$$

and

$$0 \leq D_n^f \leq D_n(\mathcal{P}) + \|\mathcal{P}\|_f. \quad (2.4.3)$$

Since f is uniformly continuous, there exists a sequence $\mathcal{P}_k$ of partitions of $[0, 1]$ such that

$$\|\mathcal{P}_k\|_f < \frac{1}{k}, \quad \forall k \in \mathbb{N}.$$

Since f_n converges pointwisely to f we deduce

$$\forall k \in \mathbb{N}, \quad \lim_{n \rightarrow \infty} D_n(\mathcal{P}_k) = 0 \text{ a.s..}$$

Hence

$$0 \leq \liminf_{n \rightarrow \infty} D_n^f \leq \limsup_{n \rightarrow \infty} D_n^f \leq \|\mathcal{P}_k\|_f < \frac{1}{k}, \quad \forall k \in \mathbb{N}.$$

Letting $k \rightarrow \infty$ we deduce the desired conclusion. $\square$

The Strong Law of Large Numbers implies that, for any $x \in [0, 1]$,

$$U_n(x) \rightarrow U(x) \text{ a.s. as } n \rightarrow \infty.$$

Thus, for every partition $\mathcal{P} = \{0 = x_0 < \dots < x_m = 1\}$ of $[0, 1]$ there exists a negligible subset $\mathcal{N}_{\mathcal{P}} \subset \Omega$ such that, $\forall \omega \in \Omega \setminus \mathcal{N}_{\mathcal{P}}$ we have

$$D_n(\mathcal{P}, \omega) = \sup_{x \in \mathcal{P}} |U_n(x, \omega) - U(x)| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We deduce from (2.4.3) with $f(x) = U(x) = x$ and $f_n(x) = U_n(x, \omega)$

$$\forall \omega \in \Omega \setminus \mathcal{N}_{\mathcal{P}}, \quad 0 \leq \liminf_{n \rightarrow \infty} D_n^U(\omega) \leq \limsup_{n \rightarrow \infty} D_n^U(\omega) \leq \|\mathcal{P}\|_U = \|\mathcal{P}\|.$$

Now choose a sequence of partitions $\mathcal{P}_k$ such that $\|\mathcal{P}_k\| \rightarrow 0$ as $k \rightarrow \infty$. If we set

$$\mathcal{N} = \bigcup_k \mathcal{N}_{\mathcal{P}_k},$$

then we deduce that for any $\omega \in \Omega \setminus \mathcal{N}$ we have

$$\liminf_{n \rightarrow \infty} D_n^U(\omega) = \limsup_{n \rightarrow \infty} D_n^U(\omega) = 0.$$

□

Remark 2.4.4. (a) Lemma 2.4.3 resembles Dini's theorem and seems to be rather old. The earliest reference to this result that I could find is the 1908 paper [32] by H. E. Buchanan and T. H. Hildebrandt. For two different proofs of this lemma I refer to [176, Sec.0.1].

(b) Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables with common cdf $F(x)$. Form the empirical (cumulative) distribution function

$$F_n(x) = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{(-\infty, x]}(X_k),$$

and the corresponding deviation $D_n := \sup_{x \in \mathbb{R}} |F_n(x) - F(x)|$. The Glivenko-Cantelli theorem shows that $D_n \rightarrow 0$ a.s..

On the other hand, observe that for each $x \in \mathbb{R}$ the random variables $\mathbf{I}_{(-\infty, x]}(X_n)$ are i.i.d. random Bernoulli random variables with success probability $F(x)$. The central limit theorem shows

$$\sqrt{n}(F_n(x) - F(x)) \Rightarrow N(0, F(x)(1 - F(x))).$$

The *Kolmogorov-Smirnov theorem* states that

$$\sqrt{n}D_n \Rightarrow D_\infty, \quad \mathbb{P}[D_\infty > c] = 2 \sum_{m \geq 1} (-1)^{m-1} e^{-2c^2 m^2}.$$

For an “elementary” proof of this fact we refer to [74]. For a more sophisticated proof that reveals the significance of the strange series above we refer to [18] or [65]. □

2.4.2. More general uniform limit laws. We want to present a generalization of the Glivenko-Cantelli theorem based on ideas pioneered by V. N. Vapnik and A. Ja. Cervonenkis [214] that turned out to be very useful in machine learning. Our presentation follows [171, Chap. II]. For more recent developments we refer to [65, 89, 213, 219].

Fix a Borel probability measure μ on $\mathfrak{X} := \mathbb{R}^N$. Any sequence of i.i.d. random vectors

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X} = \mathbb{R}^N$$

with common distribution μ defines empirical probabilities

$$P_n := \frac{1}{n} \sum_{k=1}^n \delta_{X_k}.$$

The empirical probabilities are random measures on $(\mathfrak{X}, \mathcal{B}_{\mathfrak{X}})$. More precisely, for any Borel subset $B \subset \mathfrak{X}$, $P_n[B]$ is the random variable

$$P_n[B] = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_B(X_k).$$

Suppose we are given a family $\mathcal{F} := (B_t)_{t \in T}$ of Borel subsets of $\mathfrak{X} = \mathbb{R}^N$, $N \geq 1$, parametrized by a set T . We assume T is a Borel subset of another Euclidean space $\mathbb{R}^p$ and we denote by $\mathcal{B}_T$ its Borel algebra. For example, we can choose $\mathfrak{X} = \mathbb{R}$,

$$B_t = (-\infty, t], \quad t \in T = \mathbb{R}.$$

For each $n \in \mathbb{N}$ we obtain a stochastic process parametrized by T ,

$$P_n : T \times \Omega \rightarrow [0, 1], \quad P_n(t, \omega) = P_n[B_t](\omega) = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{B_t}(X_k(\omega)).$$

For each $n \in \mathbb{N}$ we obtain a random variable

$$P_n(t) : \Omega \rightarrow [0, 1], \quad \omega \mapsto P_n(t, \omega).$$

The collection of random variables $(P_n(-))_{t \in T}$ is an example of *empirical process*. Note that

$$\mathbb{E}[P_n(t)] = \mu[B_t], \quad \text{Var}[P_n(t)] = \frac{1}{n} \text{Var}[P_1(t)] = \frac{v_t}{n},$$

where

$$v_t := \mu[B_t](1 - \mu[B_t]) \leq \frac{1}{4}.$$

The Strong Law of Large Numbers implies that

$$Z_n(t) := P_n(t) - \mu[B_t] = \frac{1}{n} \sum_{k=1}^n (Y_k(t) - \mathbb{E}[Y_k(t)]) \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty.$$

Moreover, Chebyshev's inequality shows that

$$\mathbb{P}[|Z_n(t)| > \varepsilon] \leq \frac{v_t}{n\varepsilon^2} \leq \frac{1}{4n\varepsilon^2}. \quad (2.4.4)$$

Can we conclude that $Z_n(t) \rightarrow 0$ uniformly a.s. in the precise sense described in Glivenko-Cantelli's theorem?

To proceed further we will need to make some further assumptions on the family $(B_t)_{t \in T}$. Later we will have a few things to say about their feasibility. Set

$$D_n := \sup_{t \in T} |Z_n(t)| : \Omega \rightarrow [0, 1].$$

Here is our first measure theoretic assumption.

M₁. *The function D_n is measurable*

To prove that $D_n \rightarrow 0$ a.s. we will employ a different strategy than before. More precisely we intend to show that, under certain assumptions on the family $(B_t)_{t \in T}$, the probability $\mathbb{P}[D_n > \varepsilon]$ decays very fast as $n \rightarrow \infty$, for any $\varepsilon > 0$. This will guarantee that the series

$$\sum_{n \in \mathbb{N}} \mathbb{P}[D_n > \varepsilon]$$

is convergent for any $\varepsilon > 0$ and thus, according to Corollary 1.3.57, the sequence D_n converges a.s. to 0. To obtain these tail estimates we will rely on some clever symmetrization tricks.

To state the first symmetrization result choose an independent copy of $(X_n)_{n \in \mathbb{N}}$. This is another sequence of i.i.d. random variables

$$X'_n : \Omega \rightarrow \mathfrak{X}, \quad n \in \mathbb{N},$$

independent of $(X_n)_{n \in \mathbb{N}}$, but with the same distribution as $(X_m)_{m \in \mathbb{N}}$. Set

$$Y'_k(t) := \mathbf{I}_{B_t}(X'_k), \quad Z'_n(t) := \frac{1}{n} \sum_{k=1}^n (Y'_k(t) - \mu[Y'_k(t)]), \quad \forall n \in \mathbb{N}, \quad t \in T,$$

$$D_{n,n} := \sup_{t \in T} |Z'_n(t) - Z_n(t)|. \quad (2.4.5)$$

Equivalently,

$$D_{n,n} = \sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n (Y_{n+k}(t) - Y_k(t)) \right|.$$

Here are our next measure theoretic assumption.

M₁'. The function $D_{n,n}$ is measurable

M₂. For any $n > 0$ and any $\varepsilon > 0$ there exists a measurable map

$$\tau : (\Omega, \sigma(X_1, \dots, X_n)) \rightarrow (T, \mathcal{B}_T)$$

such that $|Z_n(\tau)| > \varepsilon$ on $\{D_n > \varepsilon\}$, i.e.,

$$D_n(\omega) > \varepsilon \Rightarrow |Z_n(\tau(\omega))| > \varepsilon. \quad (2.4.6)$$

One should think of τ as a measurable map $\tau : \mathfrak{X}^n \rightarrow T$ such that

$$D_n(X_1, \dots, X_n) > \varepsilon \Rightarrow Z_n(\tau(X_1, \dots, X_n)) > \varepsilon.$$

This happens if for example

$$Z_n(\tau(X_1, \dots, X_n)) \geq D_n(X_1, \dots, X_n).$$

Thus, one could think of τ of a measurable selection of one element

$$\tau(x_1, \dots, x_n) \in [D_n(x_1, \dots, x_n), \infty)$$

for each $(x_1, \dots, x_n) \in \mathfrak{X}^n$.

Lemma 2.4.5 (First symmetrization lemma).

$$\mathbb{P}[D_n > \varepsilon] \leq 2\mathbb{P}[D_{n,n} > \varepsilon/2], \quad \forall \varepsilon > 0, \quad \forall n > \frac{1}{2\varepsilon^2}. \quad (2.4.7)$$

Proof. Choose a measurable map $\tau : (\Omega, \sigma(X_1, \dots, X_n)) \rightarrow (T, \mathcal{B}_T)$ satisfying **M₂**. Then τ is independent of Z'_n and we deduce

$$\begin{aligned} \mathbb{E}[\mathbf{I}_{\{|Z'_n(\tau)| \leq \varepsilon/2\}} \mid \sigma(X_1, \dots, X_n)] &= \mathbb{E}[\mathbf{I}_{\{|Z'_n(\tau(x_1, \dots, x_n))| \leq \varepsilon/2\}}] \stackrel{(2.4.4)}{\geq} 1 - \frac{1}{n\varepsilon^2}, \\ \mathbb{P}[|Z'_n(\tau)| \leq \varepsilon/2 \mid D_n] &= \mathbb{E}[\mathbb{E}[\mathbf{I}_{\{|Z'_n(\tau)| \leq \varepsilon/2\}} \mid \sigma(X_1, \dots, X_n)] \mid D_n] \\ &\geq 1 - \frac{1}{n\varepsilon^2}. \end{aligned}$$

Integrating over $\{D_n > \varepsilon\}$ we deduce

$$\begin{aligned} \left(1 - \frac{1}{n\varepsilon^2}\right) \mathbb{P}[D_n > \varepsilon] &\leq \mathbb{P}[|Z'_n(\tau)| \leq \varepsilon/2, D_n > \varepsilon] \\ &\stackrel{(2.4.6)}{\leq} \mathbb{P}[|Z'_n(\tau)| \leq \varepsilon/2, |Z_n(\tau)| > \varepsilon] \leq \mathbb{P}[|Z'_n(\tau) - Z_n(\tau)| > \varepsilon/2] \\ &\leq \mathbb{P}\left[\sup_{t \in T} |Z'_n(t) - Z_n(t)| > \varepsilon/2\right]. \end{aligned}$$

The inequality (2.4.7) follows by observing that for $n > \frac{1}{2\varepsilon^2}$ we have $1 - \frac{1}{n\varepsilon^2} > \frac{1}{2}$. □

Note that the variables $(Y_n(t))_{n \in \mathbb{N}}$ are independent Bernoulli random variables with success probability $p_t = \mu[B_t]$. The random variables $(Y'_n(t))$ are also of the same kind and also independent of the Y -s. The key gain is that the random variables

$$\Xi_n = Y'_n(t) - Y_n(t)$$

are symmetric, i.e., Ξ_n and $-\Xi_n$ have the same distributions. They take only the values $-1, 0, 1$ with distributions

$$\mathbb{P}[\Xi_t = \pm 1] = p_t(1 - p_t), \quad \mathbb{P}[\Xi_t = 0] = 1 - 2p_t(1 - p_t).$$

The advantage of working with symmetric random variables will become apparent after describe our second symmetrization trick known as *Rademacher symmetrization*.

Recall that a Rademacher random variable is a random variable that takes the only the values ± 1 , with equal probabilities. Suppose that $(E_n)_{n \in \mathbb{N}}$ is sequence of independent Rademacher random variables that are also independent of the variables X_n and X'_n .

Observe that the random variables $\bar{Y}_n := E_n Y_n$ are also symmetric.

Lemma 2.4.6 (Rademacher symmetrization). *For any $n \in \mathbb{N}$ we have*

$$\mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n (Y'_k(t) - Y_k(t)) \right| > \frac{\varepsilon}{2}\right] \leq 2\mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n \bar{Y}_k(t) \right| > \frac{\varepsilon}{4}\right]. \quad (2.4.8)$$

Proof. The key observation is that, because $\Xi_k(t) = Y'_k(t) - Y_k(t)$ is symmetric, it has the same distribution as $E_k \Xi_k(t)$. Set

$$S_n(t) := \frac{1}{n} \sum_{k=1}^n E_k Y_k(t), \quad S'_n(t) := \frac{1}{n} \sum_{k=1}^n E_k Y'_k(t).$$

$$\begin{aligned} \mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n (Y'_k(t) - Y_k(t)) \right| > \frac{\varepsilon}{2}\right] &= \mathbb{P}\left[\sup_{t \in T} \frac{1}{n} |S_n(t) - S'_n(t)| > \frac{\varepsilon}{2}\right] \\ &\leq \mathbb{P}\left[\sup_{t \in T} \frac{1}{n} |S_n(t)| > \frac{\varepsilon}{4}\right] + \mathbb{P}\left[\sup_{t \in T} \frac{1}{n} |S'_n(t)| > \frac{\varepsilon}{4}\right] = 2\mathbb{P}\left[\sup_{t \in T} \frac{1}{n} |S_n(t)| > \frac{\varepsilon}{4}\right], \end{aligned}$$

where we used the fact that $E_k Y'_k(t)$ and $E_k Y_k(t)$ have the same distributions. $\square$

Putting together all of the above we deduce

$$\mathbb{P}[D_n > \varepsilon] \leq 4\mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n E_k Y_k \right| > \frac{\varepsilon}{4}\right], \quad \forall \varepsilon > 0, \quad n > \frac{1}{2\varepsilon^2}. \quad (2.4.9)$$

To make further progress we condition on the variables (X_n) and we deduce

$$\begin{aligned} &\mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n E_k Y_k(t) \right| > \frac{\varepsilon}{4}\right] \\ &= \int_{\mathfrak{X}^n} \mathbb{P}\left[\sup_{t \in T} \frac{1}{n} \left| \underbrace{\sum_{k=1}^n E_k y_k(t, \vec{x})}_{=: S_t(\vec{x})} \right| > \frac{\varepsilon}{4}\right] \mu^{\otimes n}[dx_1 \cdots dx_n], \end{aligned}$$

where $\vec{x} := (x_1, \dots, x_n) \in \mathfrak{X}^n$ and

$$y_k(t, \vec{x}) = \mathbf{I}_{B_t}(x_k) \in \{0, 1\}, \quad \forall k = 1, \dots, n, \quad t \in T.$$

Hence

$$\mathbb{P}[D_n > \varepsilon] \leq 4 \int_{\mathfrak{X}^n} \mathbb{P}\left[\sup_{t \in T} S_t(\vec{x}) > \frac{\varepsilon}{4}\right] \mu^{\otimes n}[dx_1 \cdots dx_n]. \quad (2.4.10)$$

For each $n \in \mathbb{N}$, $t \in T$ and $\vec{x} \in \mathfrak{X}^n$ we set $\mathbb{I}_n := \{1, \dots, n\}$,

$$C_t(\vec{x}) := \{k \in \mathbb{I}_n; y_k(t, \vec{x}) = 1\} = \{k \in \mathbb{I}_n; x_k \in B_t\}.$$

Roughly speaking, $C_t(\vec{x}) = B_t \cap \{x_1, \dots, x_n\}$.

$$\mathcal{C}_n(\vec{x}) := \{C \subset \mathbb{I}_n; \exists t \in T, C = C_t(\vec{x})\}.$$

For every $C \subset \mathbb{I}_n$ we set

$$S_C := \frac{1}{n} \sum_{k \in C} E_k,$$

so that $S_t(\vec{x}) = |S_{C_t(\vec{x})}|$. Hence

$$\mathbb{P}\left[\sup_{t \in T} S_t(\vec{x}) > \varepsilon/4\right] = \mathbb{P}\left[\sup_{C \in \mathcal{C}_n(\vec{x})} |S_C| > \varepsilon/4\right] \leq \sum_{C \in \mathcal{C}_n(\vec{x})} \mathbb{P}[|S_C| > \varepsilon/4].$$

We can now finally understand the role of the Rademacher symmetrization. The sums S_C are of the type appearing in Hoeffding's inequality (2.3.19). We deduce

$$\mathbb{P}[|S_C| > \varepsilon/4] \leq 2e^{-n^2\varepsilon^2/32}, \quad \forall C \subset \mathbb{I}_n.$$

We deduce

$$\mathbb{P}\left[\sup_{t \in T} S_t(\vec{x}) > \varepsilon/4\right] \leq 2|\mathcal{C}_n(\vec{x})|e^{-n\varepsilon^2/32}. \quad (2.4.11)$$

Using this in (2.4.10) we deduce

$$\mathbb{P}[D_n > \varepsilon] \leq 8e^{-n\varepsilon^2/32} \int_{\mathfrak{X}^n} |\mathcal{C}_n(\vec{x})| \mu^{\otimes n}[dx_1 \cdots dx_n]. \quad (2.4.12)$$

The rough bound $|\mathcal{C}_n(\vec{x})| \leq 2^n$ is not helpful. At this point we add our last and crucial assumption.

VC. The family $\mathcal{F} = (B_t)_{t \in T}$ satisfies VC-condition.¹³ This means that there exists $d \in \mathbb{N}$ such that

$$\sup_{\vec{x} \in \mathfrak{X}^n} |\mathcal{C}_n(\vec{x})| = O(n^d) \text{ as } n \rightarrow \infty.$$

With this assumption in place we deduce that there exists $K > 0$ such that

$$2|\mathcal{C}_n(\vec{x})| \leq K(n^d + 1), \quad \forall n \in \mathbb{N}, \quad \forall \vec{x} \in \mathfrak{X}^n$$

so that

$$\mathbb{P}[D_n > \varepsilon] \leq 8Ke^{-n\varepsilon^2/32}(n^d + 1). \quad (2.4.13)$$

In the above estimate the constant K is independent of the distribution μ . Since the series

$$\sum_{n \in \mathbb{N}} e^{-n\varepsilon^2/32}(n^d + 1) < \infty, \quad \forall \varepsilon > 0,$$

we deduce that $D_n \rightarrow 0$ a.s.. We have thus proved the following wide ranging generalization of the Glivenko-Cantelli theorem.

¹³ VC = Vapnik-Chervonenkis

Theorem 2.4.7 (Vapnik-Chervonenkis). *Suppose that $\mathcal{F} = (B_t)_{t \in T}$ is a family of Borel subsets of $\mathfrak{X} = \mathbb{R}^N$ parametrized by a Borel subset T of some Euclidean space, and μ is a Borel probability measure on $\mathfrak{X}$. Assume that $\mu, \mathcal{F}$ satisfy the conditions $\mathbf{M}_1, \mathbf{M}_1', \mathbf{M}_2$.*

Fix a sequence of independent random vectors $X_n : \Omega \rightarrow \mathfrak{X}$ with common distribution μ . Form the empirical measures

$$\mu_n : \Omega \times \mathcal{B}_X \rightarrow [0, \infty], \quad \mu_n^\omega[B] = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_B[X_k(\omega)].$$

If $\mathcal{F}$ satisfies the VC-condition, then, almost surely,

$$\mu_n[B] \rightarrow \mu[B] \quad \text{as } n \rightarrow \infty$$

uniformly in $B \in \mathcal{F}$, i.e.,

$$\lim_{n \rightarrow \infty} \sup_{B \in \mathcal{F}} |\mu_n[B] - \mu[B]| = 0 \quad \text{a.s..}$$

□

Remark 2.4.8. (a) The technical assumptions $\mathbf{M}_1, \mathbf{M}_1', \mathbf{M}_2$ are measure-theoretic in nature and are automatically satisfied if the space of parameters T is countable. There are quite general (and very technical) results that guarantee that these results hold in a rather broad range of situations, [171, Appendix C].

There are more sophisticated ways of bypassing $\mathbf{M}_1$ and $\mathbf{M}_1'$ and we refer to [65, 89, 213] for details.

In Subsection 3.1.7 I will explain how One can bypass the seemingly punishing assumption, The key is an upper estimate $\mathbf{M}_2$. Using a weaker form of the first symmetrization trick one can show

$$\mathbb{E}[D_n] \leq 2 \underbrace{\mathbb{E} \left[\sup_{t \in T} \frac{1}{n} \left| \sum_{k=1}^n \bar{Y}_k(t) \right| \right]}_{=: \mathcal{R}_n(T)}.$$

The sequence $\mathcal{R}_n(T)$ is called the *Rademacher complexity* of the family $(B_t)_{t \in T}$.

McDiarmid's inequality (3.1.22a), a refined concentration inequality, shows that D_n is highly concentrated around its mean. The VC condition can be used to show that the Rademacher complexity goes to 0 as $n \rightarrow \infty$. Thus the mean of D_n goes to 0 as $n \rightarrow \infty$. Combining these facts one can obtain an inequality very similar to (2.4.12).

(b) One can obtain bounds for the tails of D_n by a Chernoff-like technique, by obtaining bounds for $\mathbb{E}[\Phi(D_n)]$, where $\Phi : [0, \infty) \rightarrow \mathbb{R}$ is a convex increasing function; see Exercise 2.67. We refer to [172] or [213] for details. □

The key assumption is VC and we want to discuss it in some detail and describe several nontrivial examples of families of sets satisfying this condition.

Fix an ambient space $\mathfrak{X}$ and $\mathcal{F} \subset 2^{\mathfrak{X}}$ a family of subsets of $\mathfrak{X}$. The *shadow* of $\mathcal{F}$ on a subset A is the family

$$\mathcal{F}_A := \{F \cap A; F \in \mathcal{F}\} \subset 2^A.$$

Note that for a finite set A we have

$$|\mathcal{F}_A| \leq 2^{|A|}.$$

When we have equality above we say that A is *shattered* by $\mathcal{F}$. Thus, A is shattered by $\mathcal{F}$ if *any* subset of A is in the shadow of $\mathcal{F}$. We set

$$s_{\mathcal{F}}(n) := \max \{ |\mathcal{F}_A|; |A| = n \}.$$

Thus $s_{\mathcal{F}}(n)$ is the size of the largest shadow on a subset of $\mathfrak{X}$ of cardinality n . Note that $s_{\mathcal{F}}(n) \leq 2^n$.

For a nonempty family $\mathcal{F}$ we define its *VC-dimension* to be

$$\dim_{VC}(\mathcal{F}) := \max \{ n \in \mathbb{N}; s_{\mathcal{F}}(n) = 2^n \}.$$

If a subset A shattered by $\mathcal{F}$, then $\dim_{VC} \mathcal{F} \geq |A|$. If $k = \dim_{VC}(\mathcal{F})$, then for any $n \leq k$ we have

$$s_{\mathcal{F}}(n) = 2^n = \sum_{j=1}^n \binom{n}{j} = \sum_{j=1}^{\min(n,k)} \binom{n}{j}.$$

We have the following remarkable dichotomy. For proof we refer to [65, Thm. 4.1.2] or [89, Thm. 3.6.3].

Theorem 2.4.9 (Sauer Lemma). *If $\dim_{VC}(\mathcal{F}) = k < \infty$, then*

$$\forall n > k : s_{\mathcal{F}}(n) \leq P_k(n) := \sum_{j=0}^{\min(n,k)} \binom{n}{j}.$$

Note that $P_k(n)$ is a polynomial of degree k in n . □

Define the *density* of $\mathcal{F}$ to be

$$\text{dens}(\mathcal{F}) = \inf \{ r > 0; s_{\mathcal{F}}(n) = O(n^r), \text{ as } n \rightarrow \infty \}.$$

We see that the family $\mathcal{F}$ satisfies the condition **VC** if and only if $\text{dens}(\mathcal{F}) < \infty$. Sauer's lemma implies that $\text{dens}(\mathcal{F}) \leq \dim_{VC}(\mathcal{F})$.

We see that a family $\mathcal{F}$ satisfies the condition **VC** if and only if its VC-dimension is finite. A family with finite VC-dimension is called a *VC-family*.

Note that $\dim_{VC}(\mathcal{F}) < k$ if and only no set $A \subset \mathfrak{X}$ of cardinality k is shattered by $\mathcal{F}$. contains a subset A_0 with the property that any set in $\mathcal{F}$ that contains A_0 also contains an element in $A \setminus A_0$. Intuitively, the sets in $\mathcal{F}$ cannot separate A_0 from its complement in A . Let us give some examples of VC families.

- (i) Suppose that $\mathcal{F}$ consists of all the lower half-lines $(-\infty, t] \subset \mathbb{R}$, $t \in \mathbb{R}$. Note that if $A = \{a_1, a_2\}$, $a_1 < a_2$, then any half-line that contains a_2 must also contain a_1 so that $\dim_{VC}(\mathcal{F}) \leq 1$.
- (ii) Suppose that $\mathcal{F}$ consists of all the open-half spaces of the vector space $\mathbb{R}^n$. A classical theorem of Radon [147, Thm. 1.3.1] shows that any subset $A \subset \mathbb{R}^n$ of cardinality $n + 2$ contains a subset A_0 that cannot be separated from its complement $A \setminus A_0$ by a hyperplane. Thus $\dim_{VC}(\mathcal{F}) \leq n + 1$. With a bit more work one can show that in fact we have equality.
- (iii) The above example is a special case of the following general result, [65, Thm. 4.2.1].

Theorem 2.4.10. *Let $\mathfrak{X}$ be a set. Suppose that V is a finite dimensional vector space of functions $f : \mathfrak{X} \rightarrow \mathbb{R}$. The space V defines two families of subsets of $\mathfrak{X}$,*

$$\mathcal{F}_V^{>0} = \{ \{f > 0\}, f \in V \}, \quad \mathcal{F}_V^{\geq 0} = \{ \{f \geq 0\}, f \in V \}.$$

Then

$$\dim_{VC}(\mathcal{F}_V^{>0}) = \dim_{VC}(\mathcal{F}_V^{\geq 0}) = \dim V.$$

- (iv) If $\mathcal{F}_0, \mathcal{F}_1$ are two VC-families of subsets of a set $\mathfrak{X}$, then $\mathcal{F}_0 \cup \mathcal{F}_1$ is also a VC family. Moreover (see [65, Thm. 4.5.1])

$$\text{dens}(\mathcal{F}_0 \cup \mathcal{F}_1) = \max(\text{dens}(\mathcal{F}_0), \text{dens}(\mathcal{F}_1)),$$

and (see [65, Prop. 4.5.2])

$$\dim_{VC}(\mathcal{F}_0 \cup \mathcal{F}_1) \leq \dim \mathcal{F}_0 + \dim \mathcal{F}_1 + 1.$$

The above equality is optimal.

- (v) If $\mathcal{F}_0, \mathcal{F}_1$ are two VC-families of subsets of a set $\mathfrak{X}$ and we set

$$\mathcal{F}_0 \sqcap \mathcal{F}_1 := \{F_0 \cap F_1; F_k \in \mathcal{F}_k, k = 0, 1\},$$

then (see [65, Thm. 4.5.3])

$$\text{dens}(\mathcal{F}_0 \sqcap \mathcal{F}_1) \leq \text{dens}(\mathcal{F}_0) + \text{dens}(\mathcal{F}_1).$$

- (vi) If $\mathcal{F}_k$ is a VC family of subsets of $\mathfrak{X}_k$, $k = 0, 1$, and we define

$$\mathcal{F}_0 \otimes \mathcal{F}_1 := \{F_0 \times F_1; F_k \in \mathcal{F}_k, k = 0, 1\},$$

then $\mathcal{F}_0 \otimes \mathcal{F}_1$ is a VC family of $\mathfrak{X}_0 \times \mathfrak{X}_1$; see [65, Thm. 4.5.3]. Moreover

$$\text{dens}(\mathcal{F}_0 \otimes \mathcal{F}_1) \leq \text{dens}(\mathcal{F}_0) + \text{dens}(\mathcal{F}_1).$$

2.4.3. PAC learning. Let us explain why the above results are relevant in machine learning. Suppose that we are dealing with a 0-1 good/bad decision/classification problem.

More precisely we want to determine when a parameter $x \in \mathbb{R}^N$ is “good”, i.e., determine the set G of “good” parameters. For example, we know from other considerations that a parameter $x \in \mathbb{R}$ is good if and only if $x \leq t_0$, but we do not know the precise value of t_0 . However, we have some information about the “good” set: it is of the form $(-\infty, t]$, $t \in \mathbb{R}$.

More generally, for one reason or another we are lead to believe that the set G belongs to a family $(B_t)_{t \in T}$, where $T \subset \mathbb{R}^p$ and B_t is a Borel subset of $\mathbb{R}^N$. The family is $(B_t)_{t \in T}$ called a *hypothesis class*. Thus we seek $t_0 \in T$ such that $B_{t_0} = G$. One of the simplest hypothesis classes is that of *perceptrons*, i.e., the collection of open half-spaces in a given Euclidean space.

Consider a silly but suggestive example. Suppose that we want to decide when a banana is good. The goodness of a banana is decided by say, three parameters: Color, Flavor, Softness, or CFS. Hence the good bananas are characterized by some measurable subset G of $\mathbb{R}^3$, the CFS space. Suppose we have a collection $\mathcal{F}$ of categories of bananas, each category $F \subset \mathcal{F}$ being defined as a subset in the CFS space. We also hypothesize that the good category G belongs to the family $\mathcal{F}$.

We are allowed to ask an Oracle to pick banana at random and answer then following yes/no questions. Given that the banana belongs to a category $F \in \mathcal{F}$, is it good? However, the Oracle won't tell us which of the categories of bananas is the good category. Saying that a banana is good given that it belongs to a category $F \cap G$ only says that the banana belongs to $F \cap G$. We are suppose to learn the good category G by repeating the above experiment many, many times and recording the answers.

Technically, the Oracle puts at our disposal a sequence of i.i.d. $\mathbb{R}^N$ -valued random vectors ($\mathbb{R}^N$ plays the role of the CFS space)

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}^N, \quad n \in \mathbb{N},$$

and the values $Y_n = \mathbf{I}_G(X_n)$, $n \in \mathbb{N}$. However, we do not know the common probability distribution μ of these random vectors.

If we knew this probability distribution, then we could find $G = F_0$ as a minimizer of the *deterministic* functional $L_\mu : \mathcal{F} \rightarrow [0, 1]$

$$\begin{aligned} L_\mu[F] &= \frac{1}{n} \sum_{k=1}^n \mathbb{P}[\mathbf{I}_F(X_k) \neq Y_k] = \frac{1}{n} \sum_{k=1}^n \mathbb{P}[\mathbf{I}_F(X_k) \neq \mathbf{I}_G(X_k)] \\ &= \frac{1}{n} \sum_{k=1}^n \mathbb{P}[\mathbf{I}_{F \Delta G}(X_k) = 1] = \mu[F \Delta G]. \end{aligned}$$

In fact $L_\mu[G] = 0$. Note that

$$\mu[F \Delta G] = \mathbb{E}[\mathbf{I}_{F \Delta G}] = \mathbb{E}[\mathbf{I}_F + \mathbf{I}_G - 2\mathbf{I}_{F \cap G}].$$

The law of large numbers shows that $\mathbb{P}$ -a.s. we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n (\mathbf{I}_F(X_k) + \mathbf{I}_G(X_k) - 2\mathbf{I}_{F \cap G}(X_k)) \\ = \mathbb{E}[\mathbf{I}_F + \mathbf{I}_G - 2\mathbf{I}_{F \cap G}] = \mu[F \Delta G] = L_\mu[F]. \end{aligned}$$

Thus, even if we do not know μ we can estimate $L_\mu[F]$ using the *random* functionals

$$\begin{aligned} L_n[F] &= \frac{1}{n} \sum_{k=1}^n (\mathbf{I}_F(X_k) + \mathbf{I}_G(X_k) - 2\mathbf{I}_{F \cap G}(X_k)) \\ &= \frac{1}{n} \sum_{k=1}^n (\mathbf{I}_F(X_k) + Y_k - 2Y_k \mathbf{I}_F(X_k)). \end{aligned}$$

The random quantity $L_n[F]$ is known as *empirical risk* and $L_\mu[F]$ is known as *population risk*.

Note that

$$F_0 \in \arg \min_{F \in \mathcal{F}} L_\mu[F].$$

In other words, F_0 is a minimizer of the population risk. In fact $L_n[F_0] = 0$. Unfortunately, the population risk is defined in terms of the unknown distribution μ . How can we minimize something that we cannot compute?

Suppose that we can determine a minimizer F_n of the empirical risk. The difference

$$\mathcal{E}(F_n) := L_\mu[F_n] - L_\mu[F_0]$$

is called *excess risk*. Note that the excess risk is a random nonnegative quantity.

We have

$$\mathcal{E}(F_n) = \underbrace{(L_\mu[F_n] - L_n[F_n])}_A + \underbrace{(L_n[F_n] - L_n[F_0])}_B + \underbrace{(L_n[F_0] - L_\mu[F_0])}_C.$$

Note that $B \leq 0$ since F_n is a minimizer of L_n . On the other hand

$$A, C \leq \sup_{F \in \mathcal{F}} |L_n[F] - L_\mu[F]|$$

so

$$\mathcal{E}(F_n) \leq 2 \sup_{F \in \mathcal{F}} |L_n[F] - L_\mu[F]|.$$

If $\mathcal{F}$ is a VC-family, then so is the family $(F \cap G)_{F \in \mathcal{F}}$ and (2.4.12) shows that there exist constants $K, c > 0$, independent of the mysterious μ , such that

$$\mathbb{P} \left[\sup_{F \in \mathcal{F}} |L_n[F] - L_\mu[F]| > \varepsilon \right] \leq K e^{-c n \varepsilon^2}, \quad \forall n.$$

Hence

$$\begin{aligned} \mathbb{P}[\mathcal{E}(F_n) > \varepsilon] &\leq \mathbb{P}[|L_n[F_n] - L_\mu[F_n]| > \varepsilon/2] \\ &\leq \mathbb{P} \left[\sup_{F \in \mathcal{F}} |L_n[F] - L_\mu[F]| > \varepsilon/2 \right] \leq K e^{-c n \varepsilon^2/4}. \end{aligned}$$

Thus, for large n , $L_n[F_n]$ is, with high confidence, within ε of the absolute minimum $L_\mu[F_0] = 0$. Hopefully, this signifies that F_n is close to F_0 . In the language of machine learning, we say that the hypothesis class $\mathcal{F}$ is PAC learnable, where PAC stands for **P**robably **A**pproximately **C**orrect.

In concrete applications, we get a handle on L_n through a sample $(x, y) = \{(x_i, y_i)\}_{1 \leq i \leq n}$ of the i.i.d. sequence

$$(X_1, Y_1), \dots, (X_n, Y_n)$$

that produces a statistical estimate of L_n that takes the form of a deterministic functional

$$L_n[-, x, y] : \mathcal{F} \rightarrow [0, \infty), \quad L_n[F, x, y] = \frac{1}{n} \sum_{k=1}^n (\mathbf{I}_F(x_k) + y_k - 2y_k \mathbf{I}_F(x_k)).$$

For more details we refer to [189, 153, 215, 219].

Remark 2.4.11. The results in this section only scratch the surface of the vast subject concerned with the limits of empirical processes. We have limited our presentation to 0-1-functions. The theory is more general than that.

Suppose that $(\mathbb{U}, \mathcal{U})$ is a measurable space and

$$X_n : (\Omega, \mathcal{S}, \mu) \rightarrow (\mathbb{U}, \mathcal{U})$$

is a sequence of i.i.d. measurable maps with common distribution $\mathbb{P} = (X_n)_\# \mu$, $\forall n$. Fix a family $\mathcal{F}$ of measurable functions $\mathbb{U} \rightarrow \mathbb{R}$. We obtain a random measure

$$\mathbb{P}_n := \frac{1}{n} \sum_{k=1}^n \delta_{X_n}$$

We obtain a stochastic process parametrized by $f \in \mathcal{F}$

$$(\mathbb{P}_n - \mathbb{P})[f] := \frac{1}{n} \sum_{k=1}^n (f(X_n) - \mathbb{E}[f(X_n)]) \in L^\infty(\Omega, \mathcal{S}, \mu), \quad f \in \mathcal{F}.$$

When $\mathcal{F}$ consists of indicator functions of measurable sets we obtain the situation described in this section.

For each f the SLLN shows that

$$(\mathbb{P}_n - \mathbb{P})[f] \rightarrow 0 \quad \text{a.s.}$$

while the CLT shows that

$$\sqrt{n}(\mathbb{P}_n - \mathbb{P})[f] \Rightarrow N(0, v(f)), \quad v(f) := \text{Var}[f(X_n)], \quad \forall n.$$

What can be said about the limit of the *process* $\mathbb{P}_n - \mathbb{P}$?

Just like there are different flavors of convergence of random variables, there are many ways in which stochastic processes can converge. Various measurability issues make empirical processes trickier to handle. We refer to [65, 89, 171, 213, 219] for more details about this problem. $\square$

2.5. The Brownian motion

The Brownian motion bears the name of its discoverer, the botanist R. Brown who observed in 1827 the chaotic motion of a particle of pollen in a fluid. Its study took off at the beginning of the 20th century and has since witnessed dramatic growth. It popped up in many branches of sciences and has lead to the development of many new branches of mathematics. In the theory of stochastic processes it plays a role similar to the role of Gaussian random variables in classical probability. It is such a fundamental and rich object that I believe any student learning the basic principles of probability needs to have a minimal introduction to it.

I drew my inspiration from many sources and I want to mention a few that I used more extensively, [18, 68, 132, 135, 185, 199]. My approach is not the most “efficient” one since I wanted to use the discussion of the Brownian motion as an opportunity to introduce the reader to other several important concepts concerning stochastic processes.

2.5.1. Heuristics. To get a grasp on the Brownian motion on a line, we consider first a discretization. We assume that the pollen particle performs a random walk along the line starting at the origin. Every unit of time τ it moves to the right or to the left, with equal probabilities, a distance δ . We denote by $S_n^{\delta, \tau}$ its location after n steps, or equivalently, its location at time $n\tau$, assuming we start the clock when the motion begins.

When $\delta = \tau = 1$ we obtain the standard random walk on $\mathbb{Z}$

$$S_n^{1,1} = S_n := \sum_{k=1}^n X_k,$$

where $(X_n)_{n \geq 1}$ is a sequence of independent Rademacher variables, i.e., random variables taking the values ± 1 with equal probabilities.

We assume that during the $(n+1)$ -th jump the particle travels with constant speed 1 so we can assume that its location at time $t \in [n, n+1)$ is

$$W^1(t) = S_n + (t - n)X_{n+1} = S_{[t]} + (t - [t])X_{[t]+1}.$$

If we sample the random variables (X_n) , then $W^1(t)$ is a piecewise linear function with linear pieces of slopes ± 1 . Its graph is a zig-zag of the type depicted in Figure 2.3




Figure 2.3. The zig-zag with slopes ± 1 depicting a random walk.




Figure 2.4. Approximating the first second of Brownian motion by a more jagged zig-zag with ν nodes and slopes $\pm\sqrt{\nu}$, $\nu \gg 1$.

Suppose now that the pollen particle performs these random jumps at a much faster rate, say ν jumps per second, and the size (in absolute value) of the jump is δ meters. We choose δ to depend on the frequency ν and we intend to let $\nu \rightarrow \infty$. Assuming that during a jump its speed is constant we deduce that this speed is $\delta\nu$ meters per second and its location at time t will be

$$W^{\nu,\delta}(t) = \delta S_{\lfloor \nu t \rfloor} + \underbrace{\delta(\nu t - \lfloor \nu t \rfloor) X_{\lfloor \nu t \rfloor + 1}}_{=: R_{\nu,\delta}(t)}.$$

To understand this formula observe that in the time interval $[0, t]$ the particle performed $\lfloor \nu t \rfloor$ complete jumps of size δ . It completed the last one at time $\frac{\lfloor \nu t \rfloor}{\nu}$. From this moment to t it travels in the direction $X_{\lfloor \nu t \rfloor + 1}$ with speed $\delta\nu$ for a duration of time $t - \frac{\lfloor \nu t \rfloor}{\nu}$.

Assuming that in finite time the particle will stay within a bounded region it is reasonable to assume that

$$\forall t, \sup_{\nu} \mathbb{E}[W^{\nu,\delta}(t)^2] < \infty. \quad (2.5.1)$$

Now observe that $\delta S_{\lfloor \nu t \rfloor}$ and $R_{\nu,\delta}$ are mean zero independent random variables so that

$$\mathbb{E}[W^{\nu,\delta}(t)^2] = \delta^2 \mathbb{E}[S_{\lfloor \nu t \rfloor}^2] + \mathbb{E}[R_{\nu,\delta}(t)^2] = \delta^2 \lfloor \nu t \rfloor + \mathbb{E}[R_{\nu,\delta}(t)^2].$$

Clearly $\mathbb{E}[R_{\nu,\delta}(t)^2] \in [0, \delta]$ so for (2.5.1) to hold we need

$$\sup_{\nu} \delta^2 \nu < \infty.$$

We achieve this by setting $\delta = \nu^{-1/2}$ and we set

$$\begin{aligned} W^{\nu}(t) &:= W^{\nu,\nu^{-1/2}}(t) = \nu^{-1/2} S_{\lfloor \nu t \rfloor} + R_{\nu}(t), \\ R_{\nu}(t) &:= \nu^{-1/2}(\nu t - \lfloor \nu t \rfloor) X_{\lfloor \nu t \rfloor + 1}. \end{aligned} \quad (2.5.2)$$

In this case $W^{\nu}(t)$ looks like a zig-zag with steep slopes of sizes $\pm\sqrt{\nu}$; see Figure 2.4.

For each ν , the collection $(W^\nu(t))_{t \geq 0}$ is a real valued random process parametrized by $[0, \infty)$. Think of it as a random real valued function defined on $[0, \infty)$. It turns out that the random processes $(W^\nu(t))_{t \geq 0}$ have a sort of limit as $\nu \rightarrow \infty$. The next result states this in a more precise form.

Proposition 2.5.1. *Let $0 \leq s < t$. Then as $\nu \rightarrow \infty$ the random variable $W^\nu(t) - W^\nu(s)$ converges in distribution to a Gaussian random variable with mean zero and variance $t - s$. In particular, since $W^\nu(0) = 0$ we deduce that the limit*

$$W(t) = \lim_{\nu} W^\nu(t)$$

exists in distribution and it is a Gaussian random variable with mean zero and variance t . Moreover, if

$$0 \leq s_0 < t_0 \leq s_1 < t_1 \leq \cdots \leq s_k < t_k, \quad k \geq 1,$$

then the $(k+1)$ -dimensional random vectors

$$\mathbf{D}^\nu = (W^\nu(t_0) - W^\nu(s_0), W^\nu(t_1) - W^\nu(s_1), \dots, W^\nu(t_k) - W^\nu(s_k))$$

converge in probability to a random vector with independent components, $W(t_j) - W(s_j)$.

Proof. Fix $0 \leq s < t$. For ν sufficiently large we have $\lfloor \nu s \rfloor < \lfloor \nu t \rfloor$ and

$$W^\nu(t) - W^\nu(s) = \underbrace{\nu^{-1} (S_{\lfloor \nu t \rfloor} - S_{\lfloor \nu s \rfloor})}_{Y_\nu} + \underbrace{((R_\nu(t) - R_\nu(s)))}_{Z_\nu}.$$

Observe first that

$$\lim_{n \rightarrow \infty} \mathbb{E}[Z_\nu^2] = 0.$$

In particular, this shows that Z_ν converges in probability to 0. On the other hand

$$Y_\nu = \frac{\sqrt{\lfloor \nu t \rfloor - \lfloor \nu s \rfloor}}{\sqrt{\nu}} \cdot \underbrace{\frac{1}{\sqrt{\lfloor \nu t \rfloor - \lfloor \nu s \rfloor}} \sum_{k=\lfloor \nu s \rfloor + 1}^{\lfloor \nu t \rfloor} X_k}_{\bar{Y}_\nu}$$

The Central Limit Theorem shows that $\bar{Y}_\nu$ converges in distribution to a standard normal random variable. Since

$$\lim_{\nu \rightarrow \infty} \frac{\sqrt{\lfloor \nu t \rfloor - \lfloor \nu s \rfloor}}{\sqrt{\nu}} = \sqrt{t - s}$$

we deduce that Y_ν converges in distribution to a Gaussian random variable with mean zero and variance $t - s$. Invoking Slutsky's theorem (Theorem 2.2.13) we deduce that $Y_\nu + Z_\nu$ converges in distribution to a Gaussian random variable with mean zero and variance $t - s$.

Now let

$$0 \leq s_0 < t_0 \leq s_1 < t_1 \leq \cdots \leq s_k < t_k, \quad k \geq 1.$$

For any ν set

$$X_j^\nu := \nu^{-1/2} (S_{\lfloor \nu t_j \rfloor} - S_{\lfloor \nu s_j \rfloor}), \quad j = 0, 1, \dots, k$$

$$\mathbf{X}^\nu = (X_0^\nu, \dots, X_k^\nu).$$

The components of $\mathbf{X}^\nu$ are independent and the above argument shows that they converge in law to the Gaussians

$$X_j^\infty := W(t_j) - W(s_j), \quad j = 0, 1, \dots, k.$$

Corollary 2.2.12 implies that for any open intervals (a_j, b_j) , $0 \leq j \leq k$ we have

$$\begin{aligned} & \mathbb{P}[X_j^\nu \in (a_j, b_j), \forall 0 \leq j \leq k] \\ &= \prod_{j=0}^k \mathbb{P}[X_j^\nu \in (a_j, b_j)] \xrightarrow{\nu \rightarrow \infty} \prod_{j=0}^k \mathbb{P}[X_j^\infty \in (a_j, b_j)]. \end{aligned}$$

Thus X^ν converges in distribution to a random vector with independent components. Slutsky's theorem implies that the increment vectors D^ν converge in distribution to the same limit as X^ν . $\square$

Definition 2.5.2 (Pre-Brownian motion). A *pre-Brownian motion* on $[0, \infty)$ is a collection of real valued random variables $(W(t))_{t \geq 0}$ with the following properties.

(i) $W(0) = 0$.

(ii) For any

$$0 \leq s_0 < t_0 \leq s_1 < t_1 \leq \dots \leq s_k < t_k, \quad k \geq 1,$$

increments

$$W(t_0) - W(s_0), W(t_1) - W(s_1), \dots, W(t_k) - W(s_k)$$

are independent.

(iii) For any $0 \leq s < t$ the increment $W(t) - W(s)$ is a Gaussian random variable with mean zero and variance $t - s$.

$\square$

We have thus proved that a suitable rescaling of the standard random walk on $\mathbb{Z}$ converges to a pre-Brownian motion. In Figure 2.4 we have depicted the graph of a sample of $W^\nu(t)$ for $\nu = 100$. Its graph is also a piecewise linear curve, but its linear pieces are much steeper, of slopes $\pm \nu^{1/2}$.

2.5.2. Gaussian measures and processes. Suppose that $(W(t))_{t \geq 0}$ is a pre-Brownian motion on $[0, \infty)$. As explained in Subsection 1.5.1, this process defines a probability measure on $\mathbb{R}^{[0, \infty)}$ equipped with the product sigma-algebra $\mathcal{B}_{\mathbb{T}}^{[0, \infty)}$ called the distribution of the process. We want to show that any two pre-Brownian motions have the same distributions. This requires a small digression in the world of Gaussian measures and processes. In this subsection we survey some basic facts concerning these concepts. In Exercise 2.70 we ask the reader to fill in some of the details of this digression. We refer to [203] for a more in depth presentation of these topics.

Let V be an n -dimensional real vector space. We denote by V^* its dual, $V^* = \text{Hom}(V, \mathbb{R})$. We have a natural pairing

$$\langle -, - \rangle : V^* \times V \rightarrow \mathbb{R}, \quad \langle \xi, x \rangle := \xi(x), \quad \forall \xi \in V^*, \quad x \in V.$$

A Borel probability measure $\mu \in \text{Prob}(V)$ is called *Gaussian* if for every linear functional $\xi \in V^*$, the resulting random variable $\xi : (V, \mathcal{B}_V, \mu) \rightarrow \mathbb{R}$ is Gaussian with mean $m[\xi]$ and variance $v[\xi]$, i.e., (see Example 1.3.37)

$$\mathbb{P}_\xi[dx] = \Gamma_{m[\xi], v[\xi]}[dx] = \begin{cases} \frac{1}{(2\pi)^{1/2}} \cdot e^{-\frac{(x-m[\xi])^2}{2v[\xi]}} dx, & v[\xi] \neq 0, \\ \delta_{m[\xi]}, & v[\xi] = 0. \end{cases}$$

Equivalently, this means that the characteristic function of $\mathbb{P}_\xi$ is

$$\widehat{\mathbb{P}_\xi}(t) = \mathbb{E}[e^{it\xi}] = e^{-\frac{v[\xi]t^2}{2} + itm[\xi]}.$$

A random vector

$$\mathbf{X} : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow V$$

is called *Gaussian* if its probability distribution is a Gaussian measure on V . The random variables $X_1, \dots, X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ are called *jointly Gaussian* if the random vector

$$\vec{X} : \Omega \rightarrow \mathbb{R}^n, \quad \vec{X}(\omega) = (X_1(\omega), \dots, X_n(\omega)),$$

is Gaussian. This means that for any real constants $\xi_1, \dots, \xi_n$, the linear combination

$$\xi_1 X_1 + \dots + \xi_n X_n$$

is a Gaussian random variable.

For any Gaussian measure μ on the finite dimensional vector space V with mean $m_\mu[\xi]$ and variance $v_\mu[\xi]$ we define its *covariance form* to be

$$C = C_\mu : V^* \times V^* \rightarrow \mathbb{R},$$

$$\begin{aligned} C(\xi, \eta) &= \frac{1}{4} (v_\mu[\xi + \eta] - v_\mu[\xi - \eta]) \\ &= \mathbb{E}_\mu[(\xi - m_\mu[\xi])(\eta - m_\mu[\eta])]. \end{aligned}$$

Then (see Exercise 2.70(ii)+(iii)) the mean m_μ is a linear functional $m_\mu : V^* \rightarrow \mathbb{R}$ and the covariance C_μ is a symmetric and positive semidefinite bilinear form on V^* . Equivalently, we can view the covariance as an element in the tensor product $V \otimes V$.

Proposition 2.5.3. *A Gaussian measure on a vector space is uniquely determined by its mean and covariance form.* $\square$

The proof of the above result is based on the Fourier transform and its main steps are described in Exercise 2.70. In the sequel we will refer to the mean zero Gaussian measures as *centered*.

Example 2.5.4. (a) If $X_1, \dots, X_n$ are independent Gaussian random variables, then any linear combination

$$\xi_1 X_1 + \dots + \xi_n X_n$$

is also Gaussian, with mean $\sum_i \xi_i \mathbb{E}[X_i]$ and variance $\sum_i \xi_i^2 \text{Var}[X_i]$. In particular, if $X_1, \dots, X_n$ are independent standard normal random variables, then the random vector $\vec{X} = (X_1, \dots, X_n)$ is Gaussian and its distribution is the *standard Gaussian measure* on $\mathbb{R}^n$

$$\Gamma_1[dx] = \frac{1}{(2\pi)^{n/2}} e^{-\frac{1}{2}\|x\|^2} dx.$$

(b) If $\vec{X} = (X_1, \dots, X_n)$ is a Gaussian random vector, then the mean of its distribution is the vector

$$m[\vec{X}] := (\mathbb{E}[X_1], \dots, \mathbb{E}[X_n])$$

and variance described by the $n \times n$ matrix C with entries the covariances of the components, i.e.,

$$C_{ij} = \text{Cov}[X_i, X_j] = \mathbb{E}[(X_i - \mathbb{E}[X_i])(X_j - \mathbb{E}[X_j])], \quad 1 \leq i, j \leq n.$$

(c) If μ is Gaussian measure on a finite dimensional vector space and $A : U \rightarrow V$ is a linear map to another vector space then the pushforward $A_{\#}\mu$ is also a Gaussian measure on V . In particular, if

$$\vec{X} = (X_1, \dots, X_n)$$

is a Gaussian vector and A is an $m \times n$ matrix, then the vector $\vec{Y} = A\vec{X}$ is also Gaussian. Note that

$$\vec{Y} = (Y_1, \dots, Y_m), \quad Y_i = \sum_{j=1}^n a_{ij} X_j, \quad i = 1, \dots, m.$$

(d) Suppose $(-, -)$ is an inner product on the vector space V with associated norm $\| - \|$. We can then identify V^* with V and the symmetric bilinear forms on V^* with symmetric operators $V \rightarrow V$. If μ is a Gaussian measure on V , then its covariance form identifies with a symmetric operator

$$\text{Var}[\mu] : V \rightarrow V$$

called the *variance* of μ .

The centered Gaussian measure on V whose variance is the identity operator is

$$\Gamma_{\mathbb{I}}[dx] = \frac{1}{(2\pi)^{\dim V/2}} e^{-\frac{1}{2}\|x\|^2} dx.$$

If $A : V \rightarrow V$ is a symmetric linear operator, then pushforward $A_{\#}\Gamma_{\mathbb{I}}$ is the Gaussian measure with variance $C = A^2$. More precisely,

$$C(v_1, v_2) = (Av_1, Av_2) = (A^2v_1, v_2).$$

If, additionally A is invertible, then

$$A_{\#}\Gamma_{\mathbb{I}}[dx] = \frac{1}{\sqrt{\det(2\pi A^2)}} e^{-\frac{1}{2}\|A^{-2}x\|^2} dx.$$

We deduce that for any bilinear, symmetric positive semidefinite form

$$C : V^* \times V^* \rightarrow \mathbb{R}$$

there exists a centered Gaussian measure admitting C as covariance form.

Indeed, if we fix a metric on V then we can identify C with a symmetric, positive semidefinite operator $C \rightarrow V$. If $A = \sqrt{C}$, then the Gaussian measure $A_{\#}\Gamma_{\mathbb{I}}$ is centered and has covariance form C . $\square$

Definition 2.5.5 (Gaussian processes). A *Gaussian process* parametrized by a set T is a collection of random variables $(X(t))_{t \in T}$ defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that, for any finite subset $I = \{t_1, \dots, t_n\} \subset T$, the random vector $X_I := (X(t_1), \dots, X(t_n))$ is Gaussian. We denote by Γ_I its distribution.

The process is called *centered* if $\mathbb{E}[X(t)] = 0, \forall t \in T$. $\square$

Suppose that $(X(t))_{t \in T}$ is a Gaussian process. Its distribution is a probability measure on $\mathbb{R}^T$ uniquely determined by the Gaussian measures Γ_I, I finite subset of T . In turn, these probability measures are *uniquely determined* by the *mean function*

$$m : T \rightarrow \mathbb{R}, \quad m(t) = \mathbb{E}[X(t)]$$

and the *covariance kernel*

$$K : T \times T \rightarrow \mathbb{R}, \quad K(s, t) = \text{Cov}[X(s), X(t)]. \quad (2.5.3)$$

Example 2.5.6. Suppose that $(W(t))_{t \geq 0}$ is a pre-Brownian motion. For any

$$0 \leq t_1 < \dots < t_n,$$

the random vector

$$(X_1, \dots, X_n) = (W(t_1), W(t_2) - W(t_1), \dots, W(t_n) - W(t_{n-1}))$$

is Gaussian since its components are independent Gaussian random variables; see Example 2.5.4(a). Observing that

$$((W(t_1), \dots, W(t_n))) = (X_1, X_1 + X_2, \dots, X_1 + \dots + X_n)$$

we deduce from Example 2.5.4(c) that the vector $((W(t_1), \dots, W(t_n)))$ is also Gaussian as linear image of a Gaussian vector. Thus, any pre-Brownian motion is a Gaussian process. It is centered since all the random variables $W(t)$ have mean zero. Its distribution is a probability measure on then path space $\mathbb{R}^{[0, \infty)}$ uniquely determined by the covariance kernel

$$K : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}, \quad K(s, t) = \mathbb{E}[W(s)W(t)].$$

We claim that

$$K(s, t) = \min(s, t), \quad \forall s, t \geq 0. \quad (2.5.4)$$

Indeed, assume without any loss of generality that $s \leq t$. Then

$$\mathbb{E}[W(s)W(t)] = \mathbb{E}[W(s)^2] + \mathbb{E}[W(s)(W(t) - W(s))].$$

The first summand is equal to s according to property (iii) of a pre-Brownian motion. Property (ii) implies

$$\mathbb{E}[W(s)(W(t) - W(s))] = \mathbb{E}[W(s)] \cdot \mathbb{E}[W(t) - W(s)] = 0.$$

Hence

$$\mathbb{E}[W(s)W(t)] = s = \min(s, t).$$

We see that *all pre-Brownian motions have the same covariance form and thus they all have the same distribution.*

Conversely, suppose that $(X(t))_{t \geq 0}$ is a centered Gaussian process whose covariance form is given by (2.5.4). Then this process is a pre-Brownian motion. Indeed,

$$\mathbb{E}[X(0)^2] = K(0, 0) = 0$$

so $X(0) = 0$ a.s.. Next, observe that

$$\mathbb{E}[X(t)^2] = K(t, t) = t.$$

Each increment $X(t) - X(s)$, $s < t$, is Gaussian and

$$\text{Var}[X(t) - X(s)] = K(t, t) - 2K(s, t) + K(s, s) = t - s.$$

Finally suppose that $0 \leq s_1 < t_1 \leq \dots \leq s_n < t_n$. Then the n -dimensional random vector of increments

$$\vec{Y} := (X(t_1) - X(s_1), \dots, X(t_n) - X(s_n))$$

is centered Gaussian. The equality (2.5.4) implies that

$$\text{Cov}[Y_i, Y_j] = 0, \quad \forall i \neq j$$

and we deduce from Exercise 2.71 that the components of $\vec{Y}$ are independent. This proves that $(X(t))_{t \geq 0}$ is a pre-Brownian motion. $\square$

Remark 2.5.7 (Brownian events). Consider an *arbitrary* pre-Brownian motion

$$B_t : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow \mathbb{R}, \quad t \geq 0.$$

We define the σ -algebra of *Brownian events* to be the σ -subalgebra of $\mathcal{F}$ generated by the family of random variables $B_t, t \geq 0$. Concretely, any Brownian event E has the form

$$(B_{\tau(n)})_{n \in \mathbb{N}} \in S,$$

where $S \subset [0, \infty)^\mathbb{N}$ is a measurable subset and $\tau : \mathbb{N} \rightarrow [0, \infty)$ is an injection.

The restriction of $\mathbb{P}$ to the σ -algebra of Brownian events is *uniquely determined* by the distributions of the *Gaussian* random vectors

$$(B_{t_1}, \dots, B_{t_n}), \quad n \in \mathbb{N}, \quad t_1, \dots, t_n.$$

In turn, the distribution of such a vector is uniquely determined by the covariances

$$\mathbb{E}[B_s B_t] = \mathbb{E}[B_s(B_s + B_t - B_s)] = \mathbb{E}[B_s^2] = s = \min(s, t).$$

We see that these distributions are *independent* of the choice of pre-Brownian motion B . This shows that if

$$B^i : (\Omega^i, \mathcal{F}^i, \mathbb{P}^i) \rightarrow \mathbb{R}, \quad i = 1, 2,$$

are two pre-Brownian motions, then for any measurable set $S \subset [0, \infty)^\mathbb{N}$ and any injection $\tau : \mathbb{N} \rightarrow [0, \infty)$ we have

$$\mathbb{P}^1[(B_{\tau(n)}^1)_{n \in \mathbb{N}} \in S] = \mathbb{P}^2[(B_{\tau(n)}^2)_{n \in \mathbb{N}} \in S]. \quad \square$$

Example 2.5.8 (Gaussian random functions). Suppose that $f_n : T \rightarrow \mathbb{R}, n \in \mathbb{N}$, is a sequence of functions defined on a set T and $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent standard normal random variables defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For each $t \in T$ we have a series of random variables

$$F(t) = \sum_{n \in \mathbb{N}} X_n f_n(t).$$

We want to emphasize that $F(t)$ also depends on the random parameter $\omega \in \Omega$,

$$F(t) = F(t, \omega) = \sum_{n \in \mathbb{N}} X_n(\omega) f_n(t). \quad (2.5.5)$$

The above is a series of *real numbers*.

Observe that if the sequence of functions f_n satisfies the condition

$$\sum_{n \in \mathbb{N}} f_n(t)^2 < \infty, \quad \forall t \in T, \quad (2.5.6)$$

then the series defining $F(t)$ converges in $L^2(\Omega, \mathcal{S}, \mathbb{P})$, for any $t \in T$. To see this, consider the partial sums

$$F_n(t) = \sum_{k=1}^n X_k f_k(t).$$

Then, for $m < n$, we have

$$\mathbb{E}[(F_n(t) - F_m(t))^2] = \sum_{k=m+1}^n f_k(t)^2 \mathbb{E}[X_k^2] = \sum_{k=m+1}^n f_k(t)^2$$

This proves that the sequence $(F_n(t))_{n \in \mathbb{N}}$ is Cauchy in $L^2(\Omega, \mathcal{S}, \mathbb{P})$.

The family $F = (F(t))_{t \in T}$ is a centered Gaussian random process. It is convenient to think of F as a random function. Its value $F(t)$ at t is not a deterministic quantity, it is random.

The covariance kernel is

$$K(s, t) = K_F(s, t) = \mathbb{E}[F(s)F(t)] = \sum_{n \in \mathbb{N}} f_n(s)f(t).$$

The above series is absolutely convergent since

$$2|f_n(s)f(t)| \leq f_n(s)^2 + f_n(t)^2, \quad \forall n, s, t.$$

Note that since the random vector $(F(s), F(t))$ is Gaussian, the random variables are independent iff they are not correlated, i.e., $\mathbb{E}[F(s)F(t)] = 0$. Thus the covariance kernel can be viewed as a measure of dependency between the values of F at different points $s, t \geq 0$.

Using Kolmogorov's one series theorem we deduce from the L^2 convergence that for any $t \in T$ there exists a measurable subset $\mathcal{N}_t \subset \Omega$ such that $\mathbb{P}[\mathcal{N}_t] = 0$ and, for any $\omega \in \Omega \setminus \mathcal{N}_t$ the series $F(t, \omega)$ in (2.5.5) converges. We will denote by $F(t, \omega)$ its sum. Set

$$\mathcal{N} := \bigcup_{t \in T} \mathcal{N}_t.$$

For $\omega \in \Omega \setminus \mathcal{N}$ we obtain a genuine function

$$F_\omega : T \rightarrow \mathbb{R}, \quad F_\omega(t) = F(t, \omega).$$

The function F_ω is referred to as a *path* of the stochastic process. We encounter here one of the recurring headaches in the theory of stochastic processes. Namely, if T is not countable, the set $\mathcal{N}$ may not be negligible so the paths may not exist a.s..

If the parameter space T has additional structure, one could ask if the paths are compatible in some fashion with that structure. For example, if T is an interval of the real axis, we could ask if the paths are continuous functions of t . $\square$

Example 2.5.9. A *Gaussian white noise* is a triplet $(H, (\Omega, \mathcal{S}, \mathbb{P}), W)$, where

- H is a separable real Hilbert space,
- $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and,
- $W : H \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P}), h \mapsto W(h)$ is an isometry of H into $L^2(\Omega, \mathcal{S}, \mathbb{P})$ such that, for any $h \in H$, the random variable W_h is centered Gaussian.

Since X is an isometry we deduce that

$$\text{Var}[W(h)] = \mathbb{E}[W(h)^2] = \|h\|_H^2.$$

In particular, this also shows that the image $W(H)$ of X is a *closed* subspace of $L^2(\Omega, \mathcal{S}, \mathbb{P})$ consisting of centered Gaussian random variables. Such a subspace is called a *Gaussian Hilbert space*. Obviously there is a natural bijection between Gaussian white noises and Gaussian Hilbert spaces.

Here is how one can construct Gaussian white noises. Fix a separable Hilbert space H with inner product $(-, -)$ and associated norm $\|-\|$. Next, fix a Hilbert basis of $(e_n)_{n \in \mathbb{N}}$. Every element in H can then be decomposed along this basis

$$h = \sum_{n \in \mathbb{N}} a_n(h)e_n, \quad a_n(h) := (h, e_n).$$

Choose a sequence of independent standard normal random variables $(X_n)_{n \in \mathbb{N}}$ defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For $h \in H$ we set

$$W(h) = \sum_{n \in \mathbb{N}} a_n(h) X_n.$$

From Parseval's identity we deduce that

$$\sum_{n \in \mathbb{N}} a_n(h)^2 = \|h\|_H^2$$

proving that the series defining $W(h)$ converges in L^2 . Kolmogorov's one series theorem shows that it also converges a.s. to a normal random variable with mean zero and variance $\|h\|^2$.

The collection $(W(h))_{h \in H}$ is a Gaussian process. Indeed, given $h_1, \dots, h_n \in H$, the random vector $(W(h_1), \dots, W(h_n))$ is Gaussian because for any $\xi := (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$, the random variable

$$\sum_{k=1}^n \xi_k W(h_k)$$

is Gaussian since

$$\sum_{k=1}^n \xi_k W(h_k) = W(h_\xi), \quad h_\xi = \sum_{k=1}^n \xi_k h_k.$$

The covariance kernel of this process

$$K(h_0, h_1) = \mathbb{E}[W(h_0)W(h_1)] = \sum_{n \in \mathbb{N}} a_n(h_0)a_n(h_1) = (h_0, h_1).$$

In particular, this proves that the correspondence $h \mapsto W(h)$ is an isometry, and thus we have produced a Gaussian white noise.

As a special example, suppose that $H = L^2([0, \infty), \lambda)$. Fix a Hilbert basis $(f_n)_{n \in \mathbb{N}}$ and construct the Gaussian noise as above,

$$L^2([0, \infty), \lambda) \ni f \mapsto W(f) := \sum a_n(f) X_n, \quad a_n(f) := \int_0^\infty f(t) f_n(t) dt.$$

For each $t \in [0, \infty)$ we set

$$B(t) := W(\mathbf{I}_{[0,t]}) = \sum_{n \in \mathbb{N}} \left(\int_0^t f_n(s) ds \right) X_n. \quad (2.5.7)$$

Note that

$$\mathbb{E}[B(s)B(t)] = \int_0^\infty \mathbf{I}_{[0,s]}(x) \mathbf{I}_{[0,t]}(x) dx = \min(s, t).$$

This shows that $W(t)$ is a pre-Brownian motion.

Conversely, a choice of pre-Brownian motion

$$B : \Omega \times [0, \infty) \rightarrow \mathbb{R}, \quad (\omega, t) \mapsto B_\omega(t)$$

defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$ canonically induces an isometry

$$\mathcal{W} = \mathcal{W}_B : L^2([0, \infty), \lambda) \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P}),$$

uniquely defined by

$$\mathcal{W}[\mathbf{I}_{(s,t]}] := B(t) - B(s).$$

We refer to Exercise 2.78 for details.

This isometry is called the *Wiener integral* and, traditionally one uses the notation

$$\int_0^t f(s)dB(s) := \mathcal{W}[\mathbf{I}_{[0,t]}f]. \quad (2.5.8)$$

I want to say a few words about the intuition behind this notation.

Observe that if $s \neq t$ and $|u| < |t - s|/2$, then the random variables

$$\frac{1}{u}(B(s+u) - B(s)) \quad \text{and} \quad \frac{1}{u}(B(t+u) - B(t))$$

are independent.

Now we need to make a leap of faith and pretend we can derivate with respect to t . (We really cannot.) Letting $u \rightarrow 0$ we deduce that $B'(t)$ and $B'(s)$ are independent Gaussian random variables. Thus, the elusive $B'(t)$ is a random “function” of the kind described in Example 2.5.8 with one big difference: in this case the condition (2.5.6) is not satisfied. Observe that the “value” of B' at a point t is independent of its value at a point s . Thus, the “value” of B' at a point carries no information about its value at a different point so $B'(t)$ is a completely chaotic random “function” and it is what is commonly referred to as *white noise*.

Derivating with the same abandon the equality (2.5.7) we deduce

$$B'(t) = \sum_{n \in \mathbb{N}} f_n(t)X_n. \quad (2.5.9)$$

This random series does not converge in a classical sense. However, it can be shown to converge in the sense of distributions.

More precisely, for any continuous compactly supported function $u : [0, \infty) \rightarrow \mathbb{R}$ with support in some compact set $[0, T]$, the random series

$$\int_0^\infty u(t)B'(t)dt = \int_0^T u(t)B'(t)dt = \sum_{n \in \mathbb{N}} X_n \int_0^T f_n(t)u(t)dt.$$

converges L^2 and a.s. since

$$\sum_{n \in \mathbb{N}} \left(\int_0^T f_n(t)u(t)dt \right)^2 \leq \|u\|_{C^0}^2 \sum_n \left(\int_0^T f_n(t)u(t)dt \right)^2 \stackrel{(2.5.7)}{=} \|u\|_{C^0}^2 \|B(T)\|_{L^2}^2$$

Thus we could assign $dB = B'dt$ a meaning as a sort of random measure. It is a random distribution in the very precise sense described in [86]. However, its integral over a bounded Borel subset $S \subset [0, \infty)$ has a precise meaning, namely

$$\int_S dB = \mathcal{W}[\mathbf{I}_S]$$

In particular,

$$B(t) = \int_0^t dB(s).$$

Even though $B'(t)$ does not exist in any any meaningful way, the above intuition is nevertheless very important since it is what lead to the very important concepts of Ito integral and stochastic differential equations. $\square$

2.5.3. The Brownian motion. We have almost everything we need to define the concept of Brownian motion and prove its existence.

Definition 2.5.10. A stochastic process $(B(t))_{t \geq 0}$ defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ is called a *standard Brownian motion* or *Wiener process* if the following hold.

- (i) $B(t)$ is a pre-Brownian motion.
- (ii) For any $\omega \in \Omega$ the path

$$B_\omega : [0, \infty) \rightarrow \mathbb{R}, \quad t \mapsto B(t, \omega)$$

is continuous.

□

As detailed in Example 2.5.6, the fact that $B(t)$ is a pre-Brownian motion can be given two equivalent characterizations.

- The process $B(t)$ is Gaussian with mean zero and covariance

$$K(s, t) = \min(s, t),$$

or,

- $B(0) = 0$, the process $B(t)$ has independent increments and, for each $s < t$, the increment $B(t) - B(s)$ is Gaussian with mean zero and variance $t - s$.

To prove the existence of a standard Brownian motions we need a bit more terminology and another fundamental result of Kolmogorov.

Definition 2.5.11. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space, T a set, and $(\mathbb{X}, \mathcal{F})$ a measurable set. Consider stochastic processes

$$X, Y : T \times \Omega \rightarrow \mathbb{X}, \quad (t, \omega) \mapsto X_t(\omega), Y_t(\omega).$$

- (i) The process Y is said to be a *modification* or *version* X , and we denote this $X \sim Y$ if for any $t \in T$ there exists a negligible subset $\mathcal{N}_t \in \mathcal{S}$ such that

$$X_t(\omega) = Y_t(\omega), \quad \forall \omega \in \Omega \setminus \mathcal{N}_t.$$

- (ii) The processes X, Y are said to be *indistinguishable* and we denote this $X \approx Y$, if there exists a negligible subset $\mathcal{N}$ such that

$$X_t(\omega) = Y_t(\omega), \quad \forall t \in T, \quad \forall \omega \in \Omega \setminus \mathcal{N}.$$

- (iii) The processes X, Y are said to be *stochastically equivalent*, and we denote this $X \sim_s Y$, if for any finite subset $I \subset T$ the random vectors X^I and Y^I have the same distribution.

□

Note that $\approx, \sim, \sim_s$ are equivalence relations and

$$X \approx Y \implies X \sim Y \implies X \sim_s Y.$$

Clearly, any two pre-Brownian motions are stochastically equivalent. We want to prove something stronger namely, that any pre-Brownian motion admits a version whose paths are almost surely continuous maps $[0, \infty) \rightarrow \mathbb{R}$. We begin by proving a more general result.

Theorem 2.5.12 (Kolmogorov's Continuity Theorem). *Suppose that T is a compact interval of the real axis, $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space and*

$$X : T \times \Omega \rightarrow \mathbb{R}, \quad (t, \omega) \mapsto X_t(\omega)$$

is a stochastic process such that, there exist constant $q, r, K > 0$ with the property that

$$\mathbb{E}[|X_s - X_t|^q] \leq K|s - t|^{1+r}, \quad \forall s, t \in T. \quad (2.5.10)$$

Then, for any $\alpha \in (0, r/q)$, the process X admits a modification Y whose paths are almost surely Hölder continuous with exponent α . This means, that for any $\alpha \in (0, r/q)$ there exists a stochastic process $(Y_t)_{t \in T}$, a negligible subset $\mathcal{N}_\alpha \subset \Omega$ and a measurable function

$$C = C_\alpha : \Omega \rightarrow [0, \infty),$$

such that

- $\forall t \in T, X_t = Y_t$ a.s. and,
- for any $\omega \in \Omega \setminus \mathcal{N}_\alpha$, and any $s, t \in T$ we have

$$|Y_s(\omega) - Y_t(\omega)| \leq C(\omega)|s - t|^\alpha.$$

Proof. We follow the presentation in [185, Sec. 10.1]. In the sequel we will denote various evolving universal positive constants by the same symbol, K . Without loss of generality we can assume that $T = [0, 1]$. We denote by $\mathcal{D}$ the set of dyadic numbers in $[0, 1]$

$$\mathcal{D} = \bigcup_{n \geq 0} D_n, \quad D_n = \left\{ \frac{k}{2^n}; \quad 0 \leq k < 2^n \right\}.$$

(Note that we do not consider 1 a dyadic number.) For $r \in D_n$ we set

$$I_{r,n} = [r, r + 1/2^n)$$

Every $t \in [0, 1)$ admits a binary/dyadic decomposition

$$t = \sum_{k=1}^{\infty} \frac{\epsilon_k(t)}{2^k}, \quad \epsilon_k(t) \in \{0, 1\},$$

such that

$$\pi_n(t) := \sum_{k=1}^n \frac{\epsilon_k(t)}{2^k} \nearrow t \quad \text{as } n \rightarrow \infty.$$

More precisely, $\pi_n(t) \in D_n$ and $t \in I_{\pi_n(t), n}$, for any $t \in [0, 1)$, $\forall n \geq 1$. For $n \geq 1$ we set

$$C_n := \left\{ (u, v) \in D_n \times D_n; \quad |u - v| \leq 2^{-(n-1)} \right\}.$$

Note that there exists $K > 0$ such that, $\forall n$,

$$\#C_n \leq K2^n.$$

For example, we can take $K = 5$. Let

$$\text{osc}_n := \sup_{(u,v) \in C_n} |X_u - X_v|.$$

We deduce

$$\begin{aligned} \mathbb{E}[\text{osc}_n^q] &\leq \sum_{(u,v) \in C_n} \mathbb{E}[|X_u - X_v|^q] \\ &\stackrel{(2.5.10)}{\leq} (\#C_n) 2^{-(n-1)(1+r)} \leq K 2^{-nr}. \end{aligned} \quad (2.5.11)$$

Let $s, t \in [0, 1]$, $s \neq t$. We set

$$\mu(s, t) := \min \{k \in \mathbb{N}; \epsilon_k(t) \neq \epsilon_k(s)\} = \min \{k \in \mathbb{N}; \pi_k(t) \neq \pi_k(s)\}.$$

For $m \in \mathbb{N}$ define

$$P_m := \{(s, t) \in \mathcal{D}; \mu(s, t) = m\}.$$

Note that if

$$(s, t) \in P_m \iff 2^{-m} \leq |s - t| \leq 2^{-m+1}.$$

Let $(s, t) \in P_m$. Then

$$\begin{aligned} X_t &= X_{\pi_m(t)} + \sum_{n \geq m} (X_{\pi_{n+1}(t)} - X_{\pi_n(t)}), \\ X_s &= X_{\pi_m(s)} + \sum_{n \geq m} (X_{\pi_{n+1}(s)} - X_{\pi_n(s)}). \end{aligned}$$

Each of these sums involves only finitely many terms since any dyadic number $t \in \mathcal{D}$ has a finite binary expansion. We deduce that

$$\begin{aligned} X_t - X_s &= X_{\pi_m(t)} - X_{\pi_m(s)} \\ &+ \sum_{n \geq m} (X_{\pi_{n+1}(t)} - X_{\pi_n(t)}) - \sum_{n \geq m} (X_{\pi_{n+1}(s)} - X_{\pi_n(s)}). \end{aligned}$$

Since

$$(\pi_m(t), \pi_m(s)) \in C_m, \quad (\pi_n(t), \pi_{n+1}(t)), \quad (\pi_n(s), \pi_{n+1}(s)) \in C_n,$$

$\forall n \geq m$, we conclude that

$$\forall (s, t) \in P_m, \quad |X_t - X_s| \leq \text{osc}_m + 2 \sum_{n \geq m} \text{osc}_n \leq 3 \underbrace{\sum_{n \geq m} \text{osc}_n}_{=: T_m}.$$

We deduce from (2.5.11) that

$$\|T_m\|_{L^q} \leq K \sum_{n \geq m} 2^{-\frac{nr}{q}} \leq K 2^{-m \frac{r}{q}}. \quad (2.5.12)$$

We have

$$\sup_{(s,t) \in P_m} \frac{|X_s - X_t|}{|s - t|^\alpha} \leq 2^{\alpha m} \sup_{(s,t) \in P_m} |X_s - X_t| \leq 2^{\alpha m} T_m.$$

Invoking (2.5.12) we conclude that

$$\left\| \sup_{(s,t) \in P_m} \frac{|X_s - X_t|}{|s - t|^\alpha} \right\|_{L^q} \leq K 2^{m(\alpha - \frac{r}{q})}, \quad \forall m \in \mathbb{N}. \quad (2.5.13)$$

The set $\mathcal{D}$ is countable so, for any $\alpha > 0$, the function

$$f_\alpha := \sup_{\substack{s, t \in \mathcal{D}, \\ s \neq t}} \frac{|X_s - X_t|}{|s - t|^\alpha} = \sup_{m \geq 1} \sup_{(s,t) \in P_m} \frac{|X_s - X_t|}{|s - t|^\alpha}$$

is measurable. We will show that if $\alpha < \frac{r}{q}$, then f_α is a.s. finite.

Fix $\alpha \in (0, r/q)$. Then $2^{m(\alpha - \frac{r}{q})} < 1$, $\forall m \in \mathbb{N}$ and using Fatou's Lemma we deduce

$$\begin{aligned} \|f_\alpha\|_q &= \left\| \sup_{m \geq 1} \sup_{(s,t) \in P_m} \frac{|X_s - X_t|}{|s - t|^\alpha} \right\|_{L^q} \\ &\leq \sup_{m \geq 1} \left\| \sup_{(s,t) \in P_m} \frac{|X_s - X_t|}{|s - t|^\alpha} \right\|_{L^q} \stackrel{(2.5.13)}{\leq} K. \end{aligned}$$

This proves that the measurable function f_α is a.s. finite.

We conclude that there exists a measurable negligible subset $N \subset \Omega$ and a nonnegative function $C \in L^q(\Omega, \mathcal{S}, \mathbb{P})$ such that

$$\forall s, t \in \mathcal{D}, \quad \forall \omega \in \Omega \setminus N, \quad |X_t(\omega) - X_s(\omega)| \leq C(\omega)|t - s|^\alpha. \quad (2.5.14)$$

We can now produce the claimed modification.

For every $\omega \in \Omega \setminus N$ the α -Hölder continuous function

$$\mathcal{D} \ni t \mapsto X_t(\omega) \in \mathbb{R}$$

admits a unique α -Hölder continuous extension $T \ni t \mapsto Y_t(\omega) \in \mathbb{R}$. For $t_0 \in T$ and $\omega \in \Omega \setminus N$ we have

$$Y_{t_0}(\omega) = \lim_{\substack{t \rightarrow t_0 \\ t \in \mathcal{D}}} X_t(\omega).$$

Since

$$\lim_{\substack{t \rightarrow t_0 \\ t \in \mathcal{D}}} \mathbb{E}[|X_t - X_{t_0}|^q] = 0$$

we deduce that $X_{t_0} = Y_{t_0}$ a.s.. Hence the process $(Y_t)_{t \in T}$ is a modification of $(X_t)_{t \in T}$ whose paths are a.s. α -Hölder continuous. $\square$

Remark 2.5.13. (a) Using Exercise 2.77 one can modify the modification in Theorem 2.5.12 to be α -Hölder continuous for any $\alpha \in (0, r/q)$, not just for a fixed α in this range.

(b) The argument in the proof of Theorem 2.5.12 is an elementary incarnation of the *chaining technique*. For a wide ranging generalization of the continuity Theorem 2.5.12 and the chaining technique we refer to [130, Chap. 11] or [206]. $\square$

Corollary 2.5.14. Suppose that $(W_t)_{t \geq 0}$ is a pre-Brownian motion. Then for any $\alpha \in (0, 1/2)$ the process (W_t) admits a modification whose paths are a.s. α -Hölder continuous. In particular, Brownian motions exist.

Proof. Set $\delta := \frac{1}{2} - \alpha$. Note that since $W_t - W_s$ is Gaussian with mean 0 and variance $|t - s|$. Then $D := \frac{1}{\sqrt{|t-s|}}(W_t - W_s) \sim N(0, 1)$ so that, $\forall q \geq 1$, we have

$$\mathbb{E}[|W_t - W_s|^q] = |t - s|^{q/2} \mathbb{E}[|D|^q].$$

If we choose $q > \frac{1}{\delta}$, then we deduce that

$$\frac{q/2 - 1}{q} = \frac{1}{2} - \frac{1}{q} > \alpha$$

and Theorem 2.5.12 implies that (W_t) admits a modification $(\bar{W}_t)_{t \geq 0}$ whose paths are a.s. α -Hölder continuous.

Recall that this means that there exists a measurable negligible set $\mathcal{N} \subset \Omega$ such that $\forall \omega \in \Omega \setminus \mathcal{N}$ the path $t \mapsto \bar{W}_t(\omega)$ is continuous. Now define

$$B : [0, \infty) \times \Omega, (t, \omega) \mapsto B_t(\omega) = \begin{cases} \bar{W}_t(\omega), & \omega \in \Omega \setminus \mathcal{N}, \\ 0, & \omega \in \mathcal{N}. \end{cases}$$

Clearly $(B_t)_{t \geq 0}$ is a (standard) Brownian motion. $\square$

Remark 2.5.15. I want to say a few words about Paul Lévy's elegant construction of the Brownian motion, [135, Sec.1].

He produces the Brownian motion on $[0, 1]$ as a limit of random piecewise linear functions L_n with nodes on the dyadic sets

$$D_n := \left\{ \frac{k}{2^n}; 0 \leq k \leq 2^n \right\}, \quad n \geq 0.$$

They are successively better and approximations of the Brownian motion.

The 0-th order approximation is the random linear function $L_0(t)$ such that $L_0(0) = 0$ and $L_0(1)$ is a standard normal random variable.

The n -th order approximation L_n satisfies the following conditions.

- It is linear on each of the intervals $((k-1)/2^n, k/2^n)$, $L_n(0) = 0$.
- The increments

$$L_n(k/2^n) - L_n((k-1)/2^n), \quad k = 1, \dots, 2^n$$

are normal random variables with mean zero and variance $1/2^n$.

- $L_n(t) = L_{n-1}(t)$, $\forall t \in D_{n-1}$.

To explain how to produce $L_n(t)$ given $L_{n-1}(t)$ we only need to explain how to produce $L_n((2k-1)/2^n)$ given that

$$L_n(j/2^{n-1}) = L_{n-1}(j/2^{n-1}), \quad j = k-1, k.$$

To “guess” what $L_n((2k-1)/2^n)$ should be, we take our inspiration from the Brownian motion that we want to approximate.

Consider two moments of time $t_0 < t_1$ in $[0, 1]$. Then $B(t_0) \sim N(0, t_0)$, $B(t_1) \sim N(0, t_1)$ and $B(t_1) - B(t_0)$ is a normal random variable with mean 0, variance $t_1 - t_0$, independent of $B(t_0)$. Denote by t_* the midpoint of $[t_0, t_1]$, $t_* = (t_0 + t_1)/2$.

Consider the linear interpolation

$$Z = \frac{1}{2}(B(t_0) + B(t_1)).$$

The difference

$$\Delta := B(t_*) - Z = \frac{1}{2}(B(t_*) - B(t_0)) + \frac{1}{2}(B(t_*) - B(t_1))$$

is a sum of two independent normal random variables, that are also independent of $B(t_0)$. Thus Δ is a normal random variable with mean 0, variance $(t_1 - t_0)/4$, independent of $B(t_0)$. We write

$$B(t_*) = Z + \Delta = \frac{1}{2}(B(t_0) + B(t_1)) + \frac{\sqrt{t_1 - t_0}}{2}X, \quad (2.5.15)$$

where X is a standard normal random variable independent of $B(t_0)$. We can now describe Lévy's prescription. We set

$$\mathcal{D} := \bigcup_{n \geq 0} D_n,$$

and consider a family $(X_t)_{t \in \mathcal{D}}$ of independent standard normal random variables. Then

$$L_0(t) := tX_1,$$

The approximation L_{n+1} is obtained from L_n as follows. If $t_0 < t_1$ are two consecutive points in D_n and $t_* \in D_{n+1}$ is the midpoint of $[t_0, t_1]$, then $L_{n+1}(t_*)$ is obtained by mimicking (2.5.15), i.e.,

$$L_{n+1}(t_*) = \frac{1}{2}(L_n(t_0) + L_n(t_1)) + \frac{\sqrt{t_1 - t_0}}{2}X_{t_*} = L_n(t_*) + \frac{1}{2^{1+n/2}}X_{t_*}.$$

On each of the intervals $[t_0, t_*]$ and $[t_*, t_1]$ the function $t \mapsto L_{n+1}(t)$ is linear so it is uniquely determined by its values at endpoints.

To prove that the sequence $L_n(t)$ converges uniformly a.s. it suffices to show that the series of random variables

$$\sum_{n \geq 0} \underbrace{\sup_{t \in [0,1]} |L_{n+1}(t) - L_n(t)|}_{=: U_n}$$

converges a.s..

Denote by M_n the set of midpoints of the 2^n intervals determined by D_n , $M_n = D_{n+1} \setminus D_n$. From the construction of L_n we deduce that

$$U_n = \frac{1}{2^{1+n/2}} \max_{\tau \in M_n} |X_\tau|.$$

We deduce that for any $n > 0$ and any $c_n > 0$ we have

$$\mathbb{P}[U_n > c_n] \leq \sum_{\tau \in M_n} \mathbb{P}[|X_\tau| > 2^{1+n/2}c_n] = 2^{n+1}\mathbb{P}[Y > 2^{1+n/2}c_n],$$

$Y \sim N(0, 1)$. The Mills ratio inequalities (1.3.44) imply that

$$2^{n+1}\mathbb{P}[Y > 2^{1+n/2}c_n] \leq \frac{2^{n/2}}{\sqrt{2\pi}c_n} e^{-2^n c_n^2}.$$

When

$$c_n = \sqrt{rn2^{-n} \log 2}, \quad r > 1,$$

we have

$$\mathbb{P}[U_n > c_n] \leq \frac{2^{(1-r)n}}{\sqrt{2r\pi n \log 2}}.$$

Observing that the series

$$\sum_{n \geq 1} \frac{2^{(1-r)n}}{\sqrt{2r\pi n \log 2}}$$

is convergent we deduce from the Borel-Cantelli lemma that

$$\mathbb{P}[U_n > c_n \text{ i.o.}] = 0.$$

Hence $U_n \rightarrow 0$ a.s. since $c_n \rightarrow 0$. $\square$

Let us observe that if $(B(t))$ is a standard Brownian motion, then $B(0) = 0$ a.s.. For this reason, the standard Brownian motion is also referred to as the *Brownian motion started at 0*. For $x \in \mathbb{R}$ we set $B^x(t) = x + B(t)$. We will refer to $B^x(t)$ as the *Brownian motion started at x*.

Remark 2.5.16 (The Wiener measure). The space $\mathcal{C} := C([0, \infty))$ of continuous functions $[0, \infty) \rightarrow \mathbb{R}$ is equipped with a natural metric d ,

$$d(f, g) = \sum_{n \in \mathbb{N}} \frac{1}{2^n} \min(1, d_n(f, g)), \quad d_n(f, g) := \sup_{t \in [n-1, n]} |f(t) - g(t)|.$$

The topology induced by this metric is the topology of uniform convergence on the compact subsets of $[0, \infty)$. One can prove (see Exercise 1.4 and Exercise 2.79) that the Borel algebra of this metric space coincides with the sigma algebra generated by the functions

$$\mathbf{Ev}_t : \mathcal{C} \rightarrow \mathbb{R}, \quad \mathbf{Ev}_t(f) = f(t).$$

More generally, for any finite subset $I \subset [0, \infty)$ we have a measurable evaluation maps

$$\mathbf{Ev}_I : \mathcal{C} \rightarrow \mathbb{R}^I, \quad f \mapsto f|_I.$$

Proposition 1.2.4 shows that if μ_0, μ_1 are two probability measures on $\mathcal{C}$ such that

$$(\mathbf{Ev}_I)_\# \mu_0 = (\mathbf{Ev}_I)_\# \mu_1$$

for any finite subset $I \subset [0, \infty)$, then $\mu_0 = \mu_1$.

Note that if $(X_t)_{t \geq 0}$ is a stochastic process defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ whose paths are continuous, then it defines a map

$$X : \Omega \rightarrow \mathcal{C}, \quad \Omega \ni \omega \mapsto X(\omega) \in \mathcal{C}, \quad X(\omega)(t) = X_t(\omega).$$

The map X is measurable since its composition with all the evaluation maps $\mathbf{Ev}_I$ are measurable. Thus the stochastic process defines a probability measure

$$\mathbb{P}_X := X_\# \mathbb{P} \in \text{Prob}(\mathcal{C}, \mathcal{B}_{\mathcal{C}})$$

called the *distribution* of the process.

Suppose that B^0, B^1 are two Brownian motions defined on possibly different probability spaces. They have distributions

$$\mathbb{W}_0, \mathbb{W}_1 \in \text{Prob}(\mathcal{C}, \mathcal{B}_{\mathcal{C}}).$$

These distributions coincide since the finite dimensional distributions $\pi_I \mathbb{W}_j$, $i = 0, 1$ are centered Gaussian with identical covariances

$$\mathbb{E}[B_{t_1}^i B_{t_1}^i] = \min(t_1, t_2), \quad \forall t_1, t_2 \in I, \quad i = 0, 1.$$

Thus, the Brownian motions determine a probability measure $\mathbb{W}$ on $\mathcal{C}$ uniquely determined by the requirement that for any finite subset $\{t_1, \dots, t_n\} \subset [0, \infty)$ the random vector

$$(\mathbf{Ev}_{t_1}, \dots, \mathbf{Ev}_{t_n})$$

is centered Gaussian with covariances $\mathbb{E}[\mathbf{Ev}_{t_i} \mathbf{Ev}_{t_j}] = \min(t_i, t_j)$. This measure is known as the *Wiener measure*. We denote it by $\mathbb{W}$.

Note that $\mathbb{W}$ is unique probability measure on $\mathcal{C}$ such that the canonical process

$$B_t : (\mathcal{C}, \mathcal{B}_{\mathcal{C}}, \mathbb{W}) \rightarrow \mathbb{R}, \quad \mathcal{C} \ni f \mapsto \mathbf{E} \mathbf{v}_t(f) = f(t)$$

is itself a Brownian motion, i.e.,

$$\mathbb{E}_{\mathbb{W}}[B_s B_t] = \min(s, t), \quad \forall s, t \geq 0. \quad (2.5.16)$$

We have proved the existence of the Wiener measure by relying on the existence of Brownian motion. Conversely, if by some other method we can construct the Wiener measure on $\mathcal{C}$, then as a bonus we deduce the existence of Brownian motions. Here is one such alternate method.

Consider again the random functions defined by (2.5.2),

$$W^\nu(t) := \nu^{-1/2} S_{\lfloor \nu t \rfloor} + R_\nu(t), \quad R_\nu(t) := \nu^{-1/2} (\nu t - \lfloor \nu t \rfloor) X_{\lfloor \nu t \rfloor + 1}. \quad (2.5.17)$$

For each ν , the paths of the random process are continuous and piecewise linear. The above discussion shows that it defines a Borel probability measure $\mathbb{P}_\nu = \mathbb{P}_{W^\nu}$ on $\mathcal{C}$.

One can show that the sequence $\mathbb{P}_\nu$ converges weakly to a probability measure on $\mathbb{P}_\infty$ satisfying (2.5.16). In other words, $\mathbb{P}_\infty$ is the Wiener measure. Its proof requires an in depth investigation of the space of probability measures on Polish spaces¹⁴ and is beyond the scope of this text. For a most readable presentation of this theorem we refer to [18]. We will have more to say about this. $\square$

The next result suggests that the paths of a Brownian motion are very rough, i.e., they have poor differentiability properties.

Proposition 2.5.17 (The quadratic variation of Brownian paths). *Consider a Brownian motion $(B_t)_{t \geq 0}$ defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Fix $c > 0$ and let*

$$0 = t_0^n < t_1^n < \dots < t_{p_n}^n = c, \quad n \in \mathbb{N}$$

be a sequence of subdivisions of $[0, t]$ with mesh

$$\mu_n := \sup_{1 \leq k \leq p_n} (t_k^n - t_{k-1}^n)$$

tending to 0 as $n \rightarrow \infty$. Define the quadratic variations

$$Q_n(c) := \sum_{k=1}^{p_n} (B_{t_k^n} - B_{t_{k-1}^n})^2.$$

Then $\mathbb{E}[Q_n(c)] = c$, $\forall n$ and $Q_n(c) \rightarrow c$ in $L^2(\Omega, \mathcal{S}, \mathbb{P})$ as $n \rightarrow \infty$.

Proof. The Gaussian random variables $X_k^n = B_{t_k^n} - B_{t_{k-1}^n}$, $1 \leq k \leq p_n$, are independent, have mean zero and momenta

$$\mathbb{E}[(X_k^n)^2] = t_k^n - t_{k-1}^n, \quad \mathbb{E}[(X_k^n)^4] = 3(t_k^n - t_{k-1}^n)^2.$$

From the first equality we deduce $\mathbb{E}[Q_n(c)] = c$. Moreover

$$\sum_{k=1}^{p_n} (X_k^n)^2 - c = \sum_{k=1}^{p_n} \underbrace{((X_k^n)^2 - (t_k^n - t_{k-1}^n))}_{=: Y_k^n}.$$

¹⁴Recall that a Polish space is a complete separable metric space.

The random variables Y_k^n are independent and have mean zero so

$$\left\| \sum_{k=1}^{p_n} (X_k^n)^2 - c \right\|_{L^2}^2 = \sum_{k=1}^{p_n} \|Y_k^n\|_{L^2}^2.$$

Now observe that

$$\begin{aligned} \|Y_k^n\|_{L^2}^2 &= \mathbb{E}[(X_k^n)^4] - 2(t_k^n - t_{k-1}^n)\mathbb{E}[(X_k^n)^2] + (t_k^n - t_{k-1}^n)^2 \\ &= 2(t_k^n - t_{k-1}^n)^2. \end{aligned}$$

Hence

$$\begin{aligned} \left\| \sum_{k=1}^{p_n} (X_k^n)^2 - c \right\|_{L^2}^2 &= 2 \sum_{k=1}^{p_n} (t_k^n - t_{k-1}^n)^2 \\ &\leq 2\mu_n \sum_{k=1}^{p_n} (t_k^n - t_{k-1}^n) = 2\mu_n c \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

□

On a subsequence n_j we have $Q_{n_j}(c) \rightarrow c > 0$ a.s.. On the other hand, if for some $\omega \in \Omega$ the function $t \rightarrow B_t(\omega)$ where Hölder with exponent $\alpha > 1/2$ on $[0, c]$, then for some constant $C = C_\omega > 0$ independent of n we would have

$$0 \leq Q_n(t)(\omega) \leq C_\omega^2 \sum_k |t_k^n - t_{k-1}^n|^{2\alpha} \leq C_\omega^2 \mu_n^{2\alpha-1} c \rightarrow 0.$$

This prove that B_t is a.s. not α -Hölder on $[0, c]$, $\alpha > 1/2$.

On the other hand, we know that the paths of the Brownian motion are Hölder continuous for any exponent $< 1/2$. A 1933 result of Paley, Wiener, Zygmund [164] shows that they have very poor differentiability properties. First, let me offer some historical context.

One question raised in the 19th century was whether there exist continuous functions on an interval that are nowhere differentiable. Apparently Gauss believed that there are no such functions. K. Weierstrass explicitly produced in 1872 such examples defined by lacunary (or sparse) Fourier series.

In 1931 S. Banach [10] and S. Mazurkewicz [148] independently showed showed that the complement of the set of nowhere differentiable functions in the metric space of continuous functions on a compact interval is very small, meagre, in the Baire category sense.

The 1933 result of Paley, Wiener, Zygmund that I will discuss below is similar in nature. They prove that the complement set of continuous nowhere differentiable functions $f \in \mathcal{C}$ is negligible with respect to the Wiener measure.

Theorem 2.5.18 (Paley, Wiener, Zygmund). *The paths of a Brownian motion $(B_t)_{t \geq 0}$ are a.s. nowhere differentiable.*

Proof. I follow the very elegant argument of Dvoretzky, Erdős, Kakutani [69]. We will show that for any interval $I = [a, b) \subset [0, \infty)$ the paths of (B_t) are a.s. nowhere differentiable on I .

Assume the Brownian motion is defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. This probability space could be the space $\mathcal{C}$ equipped with the Wiener measure. For ease of presentation we assume that $I = [0, 1)$. Consider the set

$$S := \{ \omega \in \Omega; \text{ the path } B_t(\omega) \text{ is nowhere differentiable on } [0, 1) \}.$$

The set S may not be measurable¹⁵ but we will show that its complement is contained in a measurable subset of Ω of measure zero.

Let us observe that if $\omega \in \Omega \setminus S$, i.e.,

the path $t \mapsto B_t(\omega)$ is differentiable at a point $t_0 = t_0(\omega) \in [0, 1]$,

then there exist $M, N \in \mathbb{N}$ such that for any $n \geq N$ there exists $k \in \{1, \dots, n-2\}$ with the property that

$$|B_{(k-1+i)/n}(\omega) - B_{(k+i)/n}(\omega)| \leq \frac{M}{n}, \quad \forall i = 0, 1, 2.$$

To see this, set $f(t) := B_t(\omega)$, $m = |f'(t_0)|$, $M = \lfloor m \rfloor + 2$. Then there exists $\varepsilon > 0$ so that if $s, t \in (t_0 - \varepsilon, t_0 + \varepsilon)$, $s < t$ we have

$$|f(s) - f(t)| \leq M(t - s).$$

Now choose N such that $\frac{1}{N} < \frac{\varepsilon}{6}$ and, for $n \geq N$ choose $k \in \{1, 2, \dots, n\}$ such that

$$t_0 - \varepsilon < \frac{k-1}{n}, \frac{k}{n}, \frac{k+1}{n}, \frac{k+2}{n} < t_0 + \varepsilon. \quad (2.5.18)$$

We deduce that

$$\Omega \setminus S \subset \bigcup_{M \in \mathbb{N}} \bigcup_{N \in \mathbb{N}} \underbrace{\left(\bigcap_{n \geq N} \bigcup_{k=1}^n \bigcap_{i=0}^2 \{ |B_{(k-1+i)/n} - B_{(k+i)/n}| \leq M/n \} \right)}_{=: X_{M,N}}.$$

Clearly, the set $X_{M,N}$ is measurable and it suffices to show it is negligible. We have

$$\mathbb{P}[X_{M,N}] \leq \inf_{n \geq N} \sum_{k=1}^{n-2} \mathbb{P} \left[\max_{0 \leq i \leq 2} |B_{(k-1+i)/n} - B_{(k+i)/n}| \leq M/n \right]. \quad (2.5.19)$$

Now observe that the increments $B_{(k-1)/n} - B_{k/n}$ are independent Gaussians with mean zero and variance $1/n$. We deduce

$$\mathbb{P}[X_{M,N}] \leq \inf_{n \geq N} \sum_{k=1}^{n-2} \mathbb{P} \left[|B_{(k-1)/n} - B_{k/n}| \leq M/n \right]^3.$$

The exponent 3 above will make all the difference. It appears because of the constraint (2.5.18) on N . Since $\sqrt{n} |B_{(k-1)/n} - B_{k/n}|$ is standard normal, the random variable $|B_{(k-1)/n} - B_{k/n}|$ is normal with variance $\frac{1}{n}$ and we have

$$\mathbb{P} \left[|B_{(k-1)/n} - B_{k/n}| \leq M/n \right] = 2 \sqrt{\frac{n}{2\pi}} \int_0^{M/n} e^{-x^2 n/2} dx$$

($x = My/n$)

$$2 \sqrt{\frac{n}{2\pi}} \frac{M}{n} \int_0^1 e^{-\frac{M^2 y^2}{2n}} dy \leq \underbrace{\frac{2}{\sqrt{2\pi}}}_{=: C} M n^{-1/2}.$$

Hence, $\forall n \geq N$,

$$\sum_{k=1}^{n-2} \mathbb{P} \left[|B_{(k-1)/n} - B_{k/n}| \leq M/n \right]^3 \leq n C^3 M^3 n^{-3/2} = C^3 M^3 n^{-1/2},$$

¹⁵In 1936 S. Mazurkewicz proved that the set S is not a Borel subset of $\mathcal{C}$.

and (2.5.19) implies that $\mathbb{P}[X_{M,N}] = 0$.

□

2.6. Exercises

Exercise 2.1 (Ottaviani-Skorokhod). Suppose that $X_1, \dots, X_n$ are independent random variables. We set $S_0 = 0$, $S_k = X_1 + \dots + X_k$, $k = 1, \dots, n$. Let $\alpha > 0$ and set

$$c := \sup_{0 \leq j \leq n} \mathbb{P}[|S_n - S_j| > \alpha], \quad M_n := \sup_{1 \leq j \leq n} |S_j|.$$

Prove that if $c < 1$, then¹⁶

$$\mathbb{P}[M_n > 2\alpha] \leq \frac{1}{1-c} \mathbb{P}[|S_n| > \alpha] \leq \frac{c}{1-c}.$$

Hint. Denote by J the first j such that $|S_j| > 2\alpha$. Note that $\mathbb{P}[M_n > 2\alpha] = \mathbb{P}[J \leq n]$

$$\mathbb{P}[|S_n| > \alpha] \geq \mathbb{P}[|S_n| > \alpha, M_n > 2\alpha] = \sum_{j=1}^n \mathbb{P}[|S_n| > \alpha, J = j] \geq \sum_{j=1}^n \mathbb{P}[|S_n - S_j| \leq \alpha, J = j].$$

Observe that the event $\{J = j\}$ is independent of $S_n - S_j$ and $\mathbb{P}[|S_n - S_j| \leq \alpha] \geq 1 - c$. □

Exercise 2.2. Suppose that $(X_n)_{n \geq 1}$ is a sequence of independent random variables. Prove that the following statements are equivalent.

- (i) The series $\sum_{n \geq 1} X_n$ converges in probability.
- (ii) The series $\sum_{n \geq 1} X_n$ converges a.s.

Hint. Use Exercises 2.1 and 1.51. □

Exercise 2.3. A random variable X is called symmetric if X and $-X$ have identical distributions. Suppose that $X_1, \dots, X_n$ are independent, symmetric random variable. We set $S_n = X_1 + \dots + X_n$.

$$\frac{1}{2} \mathbb{P}\left[\max_{1 \leq k \leq n} |X_k| > u\right] \leq \mathbb{P}[|S_n| > u].$$

Hint. Set $T = \min \{k; 1 \leq j \leq n, |X_j| := \max_{1 \leq k \leq n} |X_k|\}$, $R_T := S_n - X_T$. □

Exercise 2.4. Let X be a real valued random variable. The *median set* of X is the collection

$$\text{med}(X) := \{c \in \mathbb{R}; \mathbb{P}[X < c] \leq 1/2 \leq \mathbb{P}[X \leq c]\}.$$

The numbers in $\text{med}(X)$ are called *medians* of X .

- (i) Prove that $\text{med}(X) \neq \emptyset$.
- (ii) Let $\bar{x} \in \text{med}(X)$. Suppose that X' is an independent copy of X , i.e., X, X' are i.i.d. and set $X^* = X - X'$. Prove that for any $\varepsilon > 0$, and any $a \in \mathbb{R}$

$$\frac{1}{2} \mathbb{P}[X - \bar{x} \geq \varepsilon] \leq \mathbb{P}[|X^*| \geq \varepsilon] \leq 2 \mathbb{P}[|X - a| \geq \varepsilon/2].$$

- (iii) Let $\bar{x} \in \text{med}(X)$ Prove that for any $a \in \mathbb{R}$ and any $p \in [1, \infty)$

$$\frac{1}{2} \mathbb{E}[|X - \bar{x}|^p] \leq \mathbb{E}[|X^*|^p] \leq 2^p \mathbb{E}[|X - a|^p].$$

Hint. For (iii) you need to use Proposition 1.3.43 and the integration by parts formula (1.3.52). □

¹⁶The weaker inequality, $\mathbb{P}[M_n > 2\alpha] \leq \frac{c}{1-c}$, is also known as *Lévy's inequality*.

Exercise 2.5. Suppose that $(X_n)_{n \geq 1}$ is a sequence of independent random variables. Fix another sequence $(X'_n)_{n \geq 1}$ of independent random variables, independent of $(X_n)_{n \geq 1}$, and such that X_n and X'_n have the same distributions for any n . Form the symmetrizations $X_n^* = X_n - X'_n$. Prove that the following statements are equivalent.

- (i) The series, and set

$$S_n^* = \sum_{n \geq 1}^n X_n^*$$

is a.s. convergent.

- (ii) There exists a sequence of real numbers $(a_n)_{n \geq 1}$ such that the series

$$\sum_{n \geq 1} (X_n - a_n)$$

is a.s. convergent.

Hint. Use Kolmogorov's three series theorem and Exercise 2.4. □

Exercise 2.6. Consider an infinite array of *nonnegative* numbers $P = (p_{n,k})_{k,n \geq 1}$ satisfying the following conditions.

- (i) The array is lower triangular, i.e., $p_{n,k} = 0, \forall k > n$.
(ii) For every n , the n -th row of P defines a probability distribution on $\mathbb{I}_n = \{1, 2, \dots, n\}$, i.e.,

$$\sum_{k=1}^n p_{n,k} = 1, \quad \forall n \geq 1.$$

- (iii) The sequence determined by each column of P converges to 0, i.e.,

$$\lim_{n \rightarrow \infty} p_{n,k} = 0, \quad \forall k \geq 1.$$

Show that if (x_n) is a sequence of real numbers that converges to a number x , then the sequence of weighted averages

$$y_n := \sum_{k=1}^n p_{n,k} x_k$$

converges to the same number x . □

Exercise 2.7 (J. von Neumann). In this exercise we describe the *acceptance-rejection method* frequently used in Monte-Carlo simulations. For any nonnegative function $f : \mathbb{R} \rightarrow [0, \infty)$ we denote by G_f the region below its graph

$$G_f := \{(x, y) \in \mathbb{R}^2; 0 \leq y \leq f(x)\}.$$

- (i) Suppose that we are given a probability density $p : \mathbb{R} \rightarrow [0, \infty)$

$$\int_{\mathbb{R}} p(x) dx = 1.$$

For any positive constant c we set

$$\mu_p^c = \frac{1}{c} \mathbf{I}_{G_{cp}}(x, y) dx dy.$$

Since $\text{area}(G_{cp}) = c$ we deduce that μ_p^c defines a Borel probability measure on $\mathbb{R}^2$. The natural projection $\mathbb{R}^2 \ni (x, y) \mapsto x \in \mathbb{R}$ is a random variable X defined on the probability space $(\mathbb{R}^2, \mathcal{B}_{\mathbb{R}^2}, \mu_p^c)$. Prove that the probability distribution of X is $p(x)dx$.

- (ii) Suppose that X is a random variable with probability distribution $p(x)dx$. Let U be a random variable independent of X and uniformly distributed over $[0, 1]$. Prove that the probability distribution of the random vector $(X, cp(X)U)$ is μ_p^c .
- (iii) Let $q : \mathbb{R} \rightarrow [0, \infty)$ be another probability density such that, there exists $c > 0$ with the property that

$$q(x) \leq cp(x), \quad \forall x \in \mathbb{R}.$$

Suppose that $(U_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables uniformly distributed on $[0, 1]$ and $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d., independent of the U_n 's and with common distribution $p(x)dx$. Denote by N the random variable

$$N = \inf \{ n \in \mathbb{N} : cp(X_n)U_n \leq q(X_n) \}.$$

Prove that

$$\mathbb{E}[N] = c.$$

Hint. Consider the random vector $V_n = (X_n, cp(X_n)U_n)$, observe that

$$N = \inf \{ n \in \mathbb{N} : V_n \in G_q \},$$

and use part (ii) to show that N is a geometric random variable.

- (iv) Define $Y = X_N$, i.e.,

$$Y(\omega) = X_{N(\omega)}(\omega).$$

From (iii) we know that $\mathbb{P}[N < \infty] = 1$ so Y is defined outside a probability zero set. Prove that the probability distribution of the random variable Y is $q(y)dy$.

□

Remark 2.6.1 (Acceptance-Rejection method). Suppose that a computer can sample the distribution $\text{Unif}(0, 1)$ and it can sample the distribution $p(x)dx$. We can then sample the distribution $q(y)dy$ as follows. Sample successively and independently $\text{Unif}(0, 1)$ and $p(x)dx$ and denote by U_n and respectively X_n the samples obtained at the n -th trial. Stop at the first trial N when the inequality $cU_n \leq \frac{q(X_n)}{p(X_n)}$ is observed. Set $Y = X_N$. The results in the above exercise show that the expected waiting time to observe this inequality is c and the random number Y samples the distribution $q(y)dy$.

□

Exercise 2.8 (Bernstein). For each $x \in [0, 1]$ we consider a sequence $(B_k^x)_{k \in \mathbb{N}}$ of i.i.d. Bernoulli random variables with probability of success x . We set

$$S_n^x = \sum_{k=1}^n B_k^x.$$

Note that $S_n^x/n \in [0, 1]$ and the SLLN shows that

$$S_n^x/n \rightarrow x \quad \text{a.s. as } n \rightarrow \infty.$$

The dominated converges theorem implies that for any continuous function $f : [0, 1] \rightarrow \mathbb{R}$ we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[f(S_n^x/n)] = f(x).$$

Set

$$B_n^f(x) := \mathbb{E}[f(S_n^x/n)].$$

(i) Show that

$$B_n^f(x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f(k/n).$$

(ii) Prove that as $n \rightarrow \infty$ the polynomials $B_n^f(x)$ converge *uniformly on* $[0, 1]$ to $f(x)$.

Hint. For (ii) imitate the argument in Step 2 of the proof of Theorem 2.2.31. □




Figure 2.5. The graph of $f(x) = \sin(4\pi x)$ (the continuous blue curve) and of the degree 50 Bernstein polynomial $B_{50}^f(x)$ (the dotted red curve).

Exercise 2.9. Suppose that $X_n \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ is a sequence of random variables with mean zero and variance one such that

$$\lim_{k \rightarrow \infty} \mathbb{E}[X_m X_{m+k}] = 0, \quad \text{uniformly in } m.$$

Prove that

$$\frac{1}{n}(X_1 + \cdots + X_n) \xrightarrow{p} 0 \quad \text{as } n \rightarrow \infty. \quad \square$$

Exercise 2.10. Suppose that a player rolls a die an indefinite amount of times. More formally, we are given a sequence independent random variables $(X_n)_{n \in \mathbb{N}}$, uniformly distributed on $\mathbb{I}_6 := \{1, 2, \dots, 6\}$. For $k \in \mathbb{N}$, we say that a k -run occurred at time n if $n \geq k$ and

$$X_n = X_{n-1} = \cdots = X_{n-k+1} = 6.$$

For $n \in \mathbb{N}$ we set

$$R_n = R_n^k := \#\{m \leq n; \text{ a } k\text{-run occurred at time } m\},$$

$$T = T_k = \min\{n \geq k; R_n > 0\}.$$

Thus T is the moment when the first k -run occurs. As shown in Example 1.4.13, $\mathbb{E}[T] < \infty$.

(i) Compute $\mathbb{E}[T]$.

(ii) Prove that $\frac{R_n}{n}$ converges in probability to $\frac{1}{6^k}$. **Hint.** For $n \geq k$ set

$$Y_n := \mathbf{I}_{\{X_n=6\}} \cdots \mathbf{I}_{\{X_{n-k+1}=6\}}.$$

Observe that $R_n = Y_k + \cdots + Y_n$.

□

Exercise 2.11 (A. Renyi). Suppose that $(A_n)_{n \geq 0}$ is a sequence of events in the sample space $(\Omega, \mathcal{S}, \mathbb{P})$ with the following properties.

- $A_0 = \Omega$.
- $\mathbb{P}[A_n] \neq 0, \forall n \geq 0$.
- There exists $\rho \in (0, 1]$ satisfying

$$\lim_{n \rightarrow \infty} \mathbb{P}[A_n | A_k] = \rho, \quad \forall k \geq 0. \quad (2.6.1)$$

Set $X_n := \mathbf{I}_{A_n} - \rho$.

- (i) Prove that

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n X_k] = 0, \quad \forall k \in \mathbb{N}.$$

- (ii) Prove that for any $X \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n X] = 0.$$

- (iii) Conclude that the sequence (A_n) satisfies the *mixing condition with density* ρ

$$\lim_{n \rightarrow \infty} \mathbb{P}[A_n \cap A] = \rho \mathbb{P}[A], \quad \forall A \in \mathcal{S}. \quad (2.6.2)$$

Thus, in the long run, the set A_n occupies the same proportion ρ of any measurable set A .

□

Exercise 2.12 (A. Renyi). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d., almost surely finite random variables. Set

$$M_n := \frac{X_1 + \cdots + X_n}{n}.$$

Assume that the empirical means M_n converge in probability to a random variable M . The goal of the exercise is to prove that M is a.s. constant. We argue by contradiction. Assume M is not a.s. constant. Let $F : \mathbb{R} \rightarrow [0, 1]$ the cdf of M , $F(m) = \mathbb{P}[M \leq m]$.

- (i) Prove that there exist two continuity points $a < b$ of $F(x)$ such that

$$p_0 := F(b) - F(a) = \mathbb{P}[a < M \leq b] \in (0, 1).$$

- (ii) Prove that there exists $\nu_0 \in \mathbb{N}$ such that

$$\mathbb{P}[a < M_n \leq b] > 0, \quad \forall n \geq \nu_0.$$

- (iii) Set $A_0 = \Omega$ and

$$A_n = \{a < M_{\nu_0+n} \leq b\}, \quad n \geq 1.$$

Prove that the sequence (A_n) satisfies the condition (2.6.1) with $\rho = p_0$.

- (iv) Set $B := \{a < M \leq b\}$. Prove that the restriction of M_n to $(B, \mathcal{S}|_B, \mathbb{P}[-|B])$ converges in probability to $M|_B$. Here

$$\mathcal{S}|_B = \{S \cap B; S \in \mathcal{S}\}.$$

- (v) Deduce that $p_0 = 1$, thus contradicting (i).

□

Exercise 2.13 (Vitali-Hahn-Saks). Suppose that $(\Omega, \mathcal{S}, \mu)$ is a probability space. Define an equivalence relation $\sim_\mu$ on $\mathcal{S}$ by setting $S \sim_\mu S'$ if $\mu[S \Delta S'] = 0$, where Δ denotes the symmetric difference $S \Delta S' = (S \setminus S') \cup (S' \setminus S)$. Define $d = d_\mu : \mathcal{S} \times \mathcal{S} \rightarrow [0, \infty)$

$$d(S_0, S_1) = \mu[S_0 \Delta S_1].$$

- (i) Prove that $\forall S_0, S_1, S_2 \in \mathcal{S}$ we have

$$d(S_0, S_1) = d(S_1, S_0), \quad d(S_0, S_2) \leq d(S_0, S_1) + d(S_1, S_2)$$

and $d(S_0, S_1) = 0$ iff $S_0 \sim_\mu S_1$.

- (ii) Prove that d defines a *complete* metric d on $\bar{\mathcal{S}} := \mathcal{S} / \sim_\mu$.

- (iii) Suppose that $\lambda : \mathcal{S} \rightarrow [0, \infty)$ is a finite measure that is absolutely continuous with respect to μ . Hence $\lambda[S_0] = \lambda[S_1] = 0$ if $S_0 \sim_\mu S_1$. Prove that the induced function

$$\lambda : \bar{\mathcal{S}} \rightarrow \mathbb{R}$$

is continuous with respect to the metric d .

- (iv) Suppose that (λ_n) is a sequence of finite measure such that $\lambda_n \ll \mu$ for any $n \in \mathbb{N}$ and, $\forall S \in \mathcal{S}$, the sequence $\lambda_n[S]$ has a finite limit $\lambda[S]$. Prove that $\lambda : \mathcal{S} \rightarrow \mathbb{R}$ is finitely additive and $\lambda[S] = 0$ if $\mu[S] = 0$.

- (v) For any $\varepsilon > 0$ and $k \in \mathbb{N}$ we set

$$\bar{\mathcal{S}}_{k,\varepsilon} := \{S \in \bar{\mathcal{S}}; \sup_{m \in \mathbb{N}} |\lambda_k[S] - \lambda_{k+m}[S]| \leq \varepsilon\}.$$

Prove that the sets $\bar{\mathcal{S}}_{k,\varepsilon} \subset \bar{\mathcal{S}}$ are closed with respect to the metric d and

$$\bar{\mathcal{S}} = \bigcup_{k \in \mathbb{N}} \bar{\mathcal{S}}_{k,\varepsilon}, \quad \forall \varepsilon > 0.$$

- (vi) Prove that $\bar{\lambda} : \bar{\mathcal{S}} \rightarrow [0, \infty]$ is continuous and deduce that $\bar{\lambda}$ is a finite measure. **Hint.** It suffices to show that for any decreasing sequence (S_n) in $\mathcal{S}$ with empty intersection we have $\lim \lambda[S_n] = 0$. Deduce this from (v) and Baire's theorem.

□

Exercise 2.14 (A. Renyi). Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and suppose that (A_n) is a *stable sequence of events*, i.e., for any $B \in \mathcal{S}$ the sequence $\mathbb{P}[A_n \cap B]$ has a finite limit $\lambda[B]$ and $\lambda[\Omega] \in (0, 1)$. Prove that $\lambda : \mathcal{S} \rightarrow [0, 1]$ is a finite measure absolutely continuous with respect to $\mathbb{P}$, $\lambda \ll \mathbb{P}$. Denote by ρ the density of λ with respect to $\mathbb{P}$, $\rho = \frac{d\lambda}{d\mathbb{P}}$. The function ρ is called the *density of the stable sequence of events*. **Hint.** Use Exercise 2.13. □

Exercise 2.15 (A. Renyi). Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and suppose that $(A_n)_{n \in \mathbb{N}}$ is a sequence of events such that the limits

$$\lambda_0 = \lim_{n \rightarrow \infty} \mathbb{P}[A_n], \quad \lambda_k := \lim_{n \rightarrow \infty} \mathbb{P}[A_k \cap A_n], \quad k \in \mathbb{N}$$

exist and $\lambda_0 \in (0, 1)$. Denote by X linear span of the indicators $\mathbf{I}_{A_n}$ and by $\bar{X}$ its closure in L^2 .

- (i) Prove that $\forall \xi \in \bar{X}$ there exists a limit

$$L(\xi) := \lim_{n \rightarrow \infty} \mathbb{E}[\xi \mathbf{I}_{A_n}] = \mathbb{E}[\rho \xi].$$

(ii) Prove that $\forall \xi \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ there exists a limit

$$L(\xi) = \lim_{n \rightarrow \infty} \mathbb{E}[\xi \mathbf{I}_{A_n}] = \mathbb{E}[\rho \xi].$$

(iii) Show that $\bar{X} = L^2(\Omega, \mathcal{S}, \mathbb{P})$ and $\exists \rho \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ such that $L(\xi) = \mathbb{E}[\rho \xi]$, $\forall \xi \in L^2(\Omega, \mathcal{S}, \mathbb{P})$.

(iv) Show that $(A_n)_{n \in \mathbb{N}}$ is a stable sequence with density ρ . (Note that when ρ is constant the sequence satisfies the mixing condition (2.6.1) with density $\rho = \lambda_0$.)

□

Exercise 2.16. Suppose that $f : [0, 1] \rightarrow [0, 1]$ is a continuous function that is not identically 0 or 1. For $n \in \mathbb{N}$ we set

$$A_n = \bigcup_{k=0}^{n-1} [k/n, k/n + f(k/n)].$$

Show that $(A_n)_{n \geq 1}$ is a stable sequence of events and compute its density.

□

Exercise 2.17. Let $(X_n)_{n \geq 1}$ be a sequence of i.i.d. random variables such $\mathbb{E}[|X_1|^r] < \infty$ that for some $r \in (0, 2)$. Set

$$Y_n := X_n \mathbf{I}_{\{|X_n| \leq n^{1/r}\}}$$

Prove that

$$\sum_{n \geq 1} \frac{1}{n^{2/r}} \text{Var}[Y_n] < \infty.$$

Hint. Have a look at the proof of (2.1.15).

□

Exercise 2.18 (Marcinkiewicz-Zygmund). Let $(X_n)_{n \geq 1}$ be a sequence of i.i.d. random variables such $\mathbb{E}[|X_1|^r] < \infty$ that for some $r \in (0, 1)$. Set

$$Y_n := X_n \mathbf{I}_{\{|X_n| \leq n^{1/r}\}}, \quad S_n = \sum_{k=1}^n X_k, \quad T_n = \sum_{k=1}^n Y_k.$$

(i) Show that $\mathbb{P}[X_n \neq Y_n \text{ i.o.}] = 0$.

(ii) Show that

$$\lim_{n \rightarrow \infty} \frac{1}{n^{1/r}} (T_n - \mathbb{E}[T_n]) = 0, \quad \text{a.s.}$$

(iii) Prove that

$$\lim_{n \rightarrow \infty} \frac{1}{n^{1/r}} \mathbb{E}[T_n] = 0.$$

(iv) Prove that

$$\lim_{n \rightarrow \infty} \frac{1}{n^{1/r}} S_n = 0, \quad \text{a.s.}$$

Hint. Use the proof of the SLLN as inspiration. To prove (ii) use Exercise 2.17. Part (iii) requires a bit more ingenuity. Note that

$$|\mathbb{E}[T_n]| \leq \sum_{k=1}^n \mathbb{E}[|X_k|^{1-r} \cdot |X_k|^r \mathbf{I}_{\{|X_k| \leq k^{1/r}\}}], \quad \mathbf{I}_{\{|X_k| \leq k^{1/r}\}} = \mathbf{I}_{\{|X_k| \leq k^{\frac{1}{2r}}\}} + \mathbf{I}_{\{k^{\frac{1}{2r}} < |X_k| \leq k^{1/r}\}},$$

and for any $\alpha > 0$

$$\sum_{k=1}^n k^\alpha = O(n^{\alpha+1}) \text{ as } n \rightarrow \infty.$$

□

Exercise 2.19. Suppose that π is a probability measure on $\mathbb{I}_n = \{1, 2, \dots, n\}$, $p_i = \pi[\{i\}]$. Consider a sequence $(X_n)_{n \in \mathbb{N}}$ of i.i.d. random variables uniformly distributed on $[0, 1]$. For $j \in \mathbb{I}_n$ and $m \in \mathbb{N}$ we set

$$Z_{m,j} := \# \left\{ 1 \leq k \leq m; \sum_{i=0}^{j-1} p_i \leq X_k < \sum_{i=0}^j p_i \right\}, \quad H_m := \frac{1}{m} \sum_{j=1}^n Z_{m,j} \log_2 p_j.$$

Prove that

$$\lim_{m \rightarrow \infty} H_m = -\text{Ent}_2[\pi] = \sum_{j=1}^n p_j \log_2 p_j, \text{ a.s..}$$

□

Exercise 2.20. Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of i.i.d. Bernoulli random variables with success probability $\frac{1}{2}$ and $(Y_n)_{n \in \mathbb{N}}$ a sequence of i.i.d. Bernoulli random variables with success probability $\frac{1}{3}$. (The sequences (X_n) and (Y_n) may not be independent of each other.) Set $\mathcal{B} = \{0, 1\}$ and denote by $\mathcal{F}_n$ the sigma-algebra of $\mathcal{B}^{\mathbb{N}}$ generated by the cylinders

$$C_\epsilon^k := \{x = (x_1, x_2, \dots) \in \mathcal{B}^{\mathbb{N}}; x_k = \epsilon\}, \quad k = 1, 2, \dots, n, \quad \epsilon = 0, 1.$$

We set

$$\mathcal{F} := \bigcup_{n \in \mathbb{N}} \mathcal{F}_n.$$

The sequence (X_n) (resp. Y_n) define a probability measures $\mathbb{P} = \text{Ber}(1/2)^{\otimes \mathbb{N}}$ (resp. $\mathbb{Q} = \text{Ber}(1/3)^{\otimes \mathbb{N}}$) on $\mathcal{B}^{\mathbb{N}}$; see Subsection 1.5.1. Denote by $\mathbb{P}_n$ (resp. $\mathbb{Q}_n$) the restrictions of $\mathbb{P}$ (resp. $\mathbb{Q}$) to $\mathcal{F}_n$.

- (i) Prove that for any $n \in \mathbb{N}$ the measure $\mathbb{Q}_n$ is absolutely continuous with respect to $\mathbb{P}_n$. Compute the density $\frac{d\mathbb{Q}_n}{d\mathbb{P}_n}$ of $\mathbb{Q}_n$ with respect to $\mathbb{P}_n$.
- (ii) Prove that $\mathbb{Q}$ is not absolutely continuous with respect to $\mathbb{P}$. **Hint.** Use the Law of Large Numbers.

□

Exercise 2.21. Suppose that $(\mu_n)_{n \in \mathbb{N}}$ is a sequence in $\text{Meas}(\mathbb{R}_{\geq 0})$. Denote by $\mathcal{L}_n$ the Laplace transform of μ_n

$$\mathcal{L}_n(\tau) = \int_{\mathbb{R}_{\geq 0}} e^{-\tau t} \mu_n[dt], \quad \tau \geq 0.$$

- (i) The sequence $(\mu_n)_{n \geq 0}$ converges vaguely to $\mu_\infty \in \text{Meas}(\mathbb{R}_{\geq 0})$ if and only if

$$\forall \tau > 0, \quad \lim_{n \rightarrow \infty} \mathcal{L}_n(\tau) = \mathcal{L}_\infty(\tau),$$

where $\mathcal{L}_\infty$ is the Laplace transform of μ_∞ .

- (ii) The sequence $(\mu_n)_{n \geq 0}$ converges weakly to μ_∞ if and only if

$$\forall \tau \geq 0, \quad \lim_{n \rightarrow \infty} \mathcal{L}_n(\tau) = \mathcal{L}_\infty(\tau).$$

□

Exercise 2.22. Let (X_0, d_0) and (X_1, d_1) be two metric spaces. Suppose that $\mu_n^i \in \text{Prob}(X_i)$ a sequence of Borel probability measures on X_i , $i = 0, 1$, $n \in \mathbb{N}$. Prove that following are equivalent.

- (i) The sequences $(\mu_n^0)_{n \in \mathbb{N}}$ and $(\mu_n^1)_{n \in \mathbb{N}}$ are weakly convergent.

(ii) The sequence $(\mu_n^0 \otimes \mu_n^1)_{n \in \mathbb{N}}$ is weakly convergent.

□

Exercise 2.23. For $a \geq 0$ we denote by $\text{Meas}_a(\mathbb{R}_{\geq 0})$ the set of Borel probability measures μ on $[0, \infty)$ such that

$$\int_{\mathbb{R}_{\geq 0}} e^{-\tau t} \mu[dt] < \infty, \quad \forall \tau > a.$$

The Laplace transform of $\mu \in \text{Meas}_a(\mathbb{R}_{\geq 0})$ is the nonincreasing function

$$\mathcal{L}_\mu : (a, \infty) \rightarrow [0, \infty), \quad \mathcal{L}_\mu(\tau) = \int_{\mathbb{R}_{\geq 0}} e^{-\tau t} \mu[dt].$$

- (i) Prove that a measure in $\text{Meas}_a(\mathbb{R}_{\geq 0})$ is uniquely determined by its Laplace transform.
- (ii) Suppose that if $(\mu_n)_{n \in \mathbb{N}}$ is a sequence in $\text{Meas}_a(\mathbb{R}_{\geq 0})$ such that their Laplace transforms converge pointwisely to a function $\mathcal{L}_\infty : (a, \infty) \rightarrow \mathbb{R}$. Then $\mathcal{L}_\infty$ is the Laplace transform of a measure $\mu_\infty \in \text{Meas}_a(\mathbb{R}_{\geq 0})$ and the measures μ_n converge vaguely to μ_∞ , i.e.

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}_{\geq 0}} f(t) \mu_n[dt] = \int_{\mathbb{R}_{\geq 0}} f(t) \mu_\infty[dt], \quad \forall f \in C_{\text{cpt}}(\mathbb{R}_{\geq 0}).$$

- (iii) Suppose that $(\mu_n)_{n \in \mathbb{N}}$ is a sequence in $\text{Meas}_a(\mathbb{R}_{\geq 0})$ that converges vaguely to a measure $\mu_\infty \in \text{Meas}_a(\mathbb{R}_{\geq 0})$. Prove that if

$$\sup_n \sup_{\tau > a} \mathcal{L}_{\mu_n}(\tau) < \infty,$$

then $\mathcal{L}_{\mu_n}$ converges pointwisely to $\mathcal{L}_{\mu_\infty}$ on (a, ∞) .

□

Exercise 2.24. For any Borel probability measure $\mu \in \text{Prob}(\mathbb{R})$ we denote by F_μ its cdf and by Q_μ its quantile; see Example 1.2.22. Prove that a sequence of Borel probability measures μ_n converges weakly to $\mu \in \text{Prob}(\mathbb{R})$ if and only if the sequence of quantiles $Q_{\mu_n} : [0, 1] \rightarrow \mathbb{R}$ converges almost everywhere to Q_μ .

□

Exercise 2.25. We say that Borel probability measure on $\mathbb{R}$ is *discrete* if $\mu[F] = 0$ for some finite subset of $\mathbb{R}$. Let $\rho \in C_b(\mathbb{R})$ be a nonnegative continuous function such that

$$\int_{\mathbb{R}} \rho(x) dx = 1.$$

Denote by λ_ρ the measure given by $\lambda_\rho[dx] = \rho(x) \lambda[dx]$. Prove that there exists a sequence of discrete probability measures converging weakly to λ_ρ .

□

Exercise 2.26. Suppose that $\rho \in C_b(\mathbb{R})$ is nonnegative and

$$\int_{\mathbb{R}} \rho(x) dx = 1.$$

For $\varepsilon > 0$ we set $\rho_\varepsilon(x) = \varepsilon^{-1} \rho(x/\varepsilon)$ and define as in Exercise 2.25

$$\lambda_{\rho_\varepsilon}[dx] = \rho_\varepsilon(x) \lambda[dx].$$

Fix $\mu \in \text{Prob}(\mathbb{R})$.

- (i) Prove that the convolutions $\lambda_{\rho_\varepsilon} * \mu$ converge weakly to μ as $\varepsilon \searrow 0$.

- (ii) Prove that there exists a sequence of discrete probability measures on $\mathbb{R}$ that converge weakly to μ .

□

Exercise 2.27. Let (X_n) be a sequence of geometric random variables $X_n \sim \text{Geom}(1/n)$. Prove that

$$Y_n := \frac{1}{n}X_n \Rightarrow X \sim \text{Exp}(1).$$

Hint. Show that $\mathbb{P}[Y_n > y] \rightarrow e^{-y}$ as $n \rightarrow \infty$, $\forall y \in \mathbb{R}$.

□

Remark 2.6.2. Let X be a geometric random variable with success probability p . In other words, X is the number of independent Bernoulli trials with success probability p until we record the first success. Suppose that we perform one trial per unit of time τ , where τ is measured in seconds. Then τX is the waiting time until we observe the first success. Suppose that $p = \frac{1}{n}$ we perform n trials per second so $\tau = \frac{1}{n}$. Then $\tau X = \frac{1}{n}X_n$. This exercise shows that, for n large, the distribution of the random time $\frac{1}{n}X_n$ is close to an exponential distribution. This partially explains the interpretation of exponential random variables as waiting times of rare (unlikely) events.

□

Exercise 2.28. Fix $\lambda > 0$. Show that as $n \rightarrow \infty$ we have $\text{Bin}(n, \lambda/n) \Rightarrow \text{Poi}(\lambda)$, where $\text{Bin}(n, \lambda/n)$ denotes the binomial probability distribution corresponding to n independent trials with success probability λ/n and $\text{Poi}(\lambda)$ denotes the Poisson distribution with parameter λ .

□

Exercise 2.29. When Bob gets bored, he goes to a nearby bus station with an urn containing balls of c colors in proportions $p_1, p_2, \dots, p_c$, $p_1 + \dots + p_c = 1$.

Each time a bus arrives at the bus station, Bob draws a ball at random from the urn, records its color, puts it back in the urn and waits for the next arrival. It is known that the waiting time for the next bus to arrive is exponential with rate $\lambda > 0$.

For each $i = 1, \dots, c$ and $t \geq 0$ denote by $N_i(t)$ the number of balls of color i the person has drawn from the urn during the interval $[0, t]$. Set $N(t) = N_1(t) + \dots + N_c(t)$ so $N(t)$ is a Poisson process with parameter λ ; see Example 1.3.7.

- (i) Prove that, for any $t \geq 0$, $N_i(t) = \text{Poi}(\lambda p_i t)$ **Hint.** Condition on $N(t)$.
(ii) Prove that for any $0 \leq s < t$, $N_i(t) - N_i(s)$ has the same distribution as $N_i(t - s)$. **Hint.** Use the memoryless property of the exponential distribution.
(iii) Prove that for any $0 \leq t_1 < t_2 < \dots < t_n$, the increments

$$N_i(t_1), N_i(t_2) - N_i(t_1), \dots, N_i(t_n) - N_i(t_{n-1})$$

are independent.

- (iv) Denote by T_i the waiting time until the first ball of color i extracted. Prove that $T_i \sim \text{Exp}(\lambda p_i)$.

Hint. Use (i), (ii) and Exercises 2.27.

□

Exercise 2.30 (P. Diaconis). Suppose that $(A_n)_{n \geq 1}$ is a sequence of independent events of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Set $p_n := \mathbb{P}[A_n]$.

- (i) Prove that if $\sum_{n \geq 1} p_n p_{n+1} < \infty$, then the random series

$$\sum_{n \geq 1} I_{A_n \cap A_{n+1}}$$

converges a.s..

- (ii) Prove that if $p_n = \frac{1}{n}$, $\forall n$, then the sum S of the above series is a Poisson random variable, $S \sim \text{Poi}(1)$.

□

Exercise 2.31 (Occupancy Problem). Suppose that b balls are successively and randomly placed in u urns. The urns are equally likely to be the destination of a given ball. Let $N_u = N_{u,b}$ the number of empty urns after all balls have been distributed.

- (i) Compute the expectation and variance of X_u . **Hint.** $X_r = \sum_{k=1}^u I_{U_k, u}$, $U_{k,u}$ = “urn k is empty after all the b balls have been randomly placed in the u urns.”
- (ii) Show that if $b/u \rightarrow c > 0$ as $u \rightarrow \infty$, then $\frac{X_u}{u} \rightarrow e^{-c}$ in probability.
- (iii) Compute $\mathbb{P}[X_{u,b} = m]$. **Hint.** Use the inclusion-exclusion equality (1.3.28).
- (iv) Show that if $ue^{-b/u} \rightarrow \lambda$ as $u \rightarrow \infty$, then $X_{u,b}$ converges in distribution to $\text{Poi}(\lambda)$. (Note that this means that $\frac{b}{u} - \log u + \log \lambda \rightarrow 0$ as $u \rightarrow \infty$.) **Hint.** Use the technique in Example 1.3.34.

□

Remark 2.6.3. Let me comment why the result in Exercise 2.31 is surprising. Consider the following concrete situation.

Assume $b = 2u$ and suppose that we want to distribute $2u$ gifts to u children. We want to do this in the “fairest” possible way since the gifts, of equal value, are different, and several kids may desire the same gift. To remove any bias, “common sense” suggests that each gift should be given to a child chosen uniformly at random. There are twice as many gifts as children so what can go wrong? Part (ii) of this exercise shows that for u large nearly surely $e^{-2}u \approx 0.13u$ children will receive no gifts!

□

Exercise 2.32 (Coupon collector problem). For $n \in \mathbb{N}$ denote by N_n the number of boxes of cereals one has to purchase in order to obtain all the n coupons of a collection; see Example 1.3.28. Recall that $\mathbb{E}[N_n] \sim n \log n$ as $n \rightarrow \infty$. Prove that¹⁷

$$\lim_{n \rightarrow \infty} \mathbb{P}[N_n - n \log n \leq nx] = \exp(-e^{-x}).$$

Hint. Reduce to Exercise 2.31(iv).

□

Exercise 2.33. For $N \in \mathbb{N}$ denote by B_N the birthday random variable defined in Exercise 1.29. Its range is $\{2, 3, \dots, N+1\}$. Prove that as $N \rightarrow \infty$, the sequence of random variables

$$X_N := \frac{1}{\sqrt{N}} B_N$$

converges in law to a *Raleigh random variable*, i.e., a random variable X with probability distribution

$$\mathbb{P}_X[dx] = xe^{-\frac{x^2}{2}} I_{[0, \infty)}(x) dx.$$

¹⁷The distribution with cdf $F(x) = \exp(-e^{-x})$ is called a *Gumbel distribution*.

Hint. Observe that $\mathbb{P}[X > x] = e^{-\frac{x^2}{2}}$. Show that

$$\lim_{N \rightarrow \infty} \log \mathbb{P}[X_N > x] = -\frac{x^2}{2}, \quad \forall x > 0.$$

Note. With considerably more effort one can prove that

$$\lim_{N \rightarrow \infty} \mathbb{E}[X_N] = \mathbb{E}[X] = \sqrt{\frac{\pi}{2}}.$$

□

Exercise 2.34 (P. Lévy). Consider the random variables L_n defined in Exercise 1.25. Prove that as $n \rightarrow \infty$ the random variables $\frac{L_n}{n}$ converge in distribution to the arcsine distribution $\text{Beta}(1/2, 1/2)$; see Example 1.3.39. **Hint.** You need to use Stirling formula (A.1.8) with error estimate (A.1.9). □

Exercise 2.35. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables uniformly distributed in $[0, L]$, $L > 0$. For $n \in \mathbb{N}$ we set

$$X_{(n)} := \max(X_1, X_2, \dots, X_n).$$

Prove that $\lim_{n \rightarrow \infty} \mathbb{E}[X_{(n)}] = L$ and $X_{(n)} \rightarrow L$ in probability. **Hint.** Have a look at Exercise 1.57. □

Exercise 2.36. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables uniformly distributed in $[0, 1]$. Denote by $X_{(1)}^n, X_{(2)}^n, \dots, X_{(n)}^n$ the order statistics of the first n of them; see Exercise 1.57. Prove that for any $k \in \mathbb{N}$ the random variable $nX_{(k)}^n$ converges in distribution to $\text{Gamma}(k, 1)$. □

Exercise 2.37. Suppose that $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ are two sequences of random vectors such that $X_n \rightarrow X$ and $|X_n - Y_n| \rightarrow 0$ in distribution. Then $Y_n \rightarrow X$ in distribution. □

Exercise 2.38. Suppose that $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ are two sequences of random variables such that X_n converges in distribution to X and Y_n converges in probability to the constant c . Prove that the random vector (X_n, Y_n) converges in distribution to (X, c) . **Hint.** Prove that (X_n, c) converges in probability to (X, c) and then use Exercise 2.37. □

Exercise 2.39. Suppose that $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ are two sequences of random variables such that

- X_n converges in distribution X .
- Y_n converges in distribution to Y .
- X_n is independent of Y_n for every n and X is independent of Y .

Prove the following.

- (i) The random vector (X_n, Y_n) converges in distribution to (X, Y) .
- (ii) The sum $X_n + Y_n$ converges in distribution to $X + Y$.

□

Exercise 2.40. Suppose that $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ are sequences of random variables with the following properties.

- (i) The random variables $(X_n)_{n \in \mathbb{N}}$ are identically distributed.
- (ii) The sequence of random vectors (X_n, Y_n) converges in distribution to the random vector (X, Y) .

Prove that for any Borel measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$ the sequence of random vectors $(f(X_n), Y_n)$ converges in distribution to $(f(X), Y)$. **Hint.** Fix a Borel measurable function f . It suffices to show that for any continuous and bounded functions $u, v : \mathbb{R} \rightarrow \mathbb{R}$ we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[u(f(X_n))v(Y_n)] = \mathbb{E}[u(f(X))v(Y)].$$

Consider the Borel measurable functions v_n defined by $v_n(X_n) = \mathbb{E}[v(Y_n) \mid X_n]$. □

Exercise 2.41. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. L^2 random variables with $\mu = \mathbb{E}[X_n]$, $\sigma^2 = \text{Var}[X_n]$. Set

$$\bar{X}_n = \frac{1}{n}(X_1 + \cdots + X_n), \quad Y_n = \frac{1}{n-1} \sum_{k=1}^{n-1} (X_k - \bar{X}_n)^2.$$

Prove that $\mathbb{E}[Y_n] = \sigma^2$ and $Y_n \rightarrow \sigma^2$ in probability. □

Exercise 2.42. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. Bernoulli random variables with success probability $p = \frac{1}{2}$. For each $n \in \mathbb{N}$ we set

$$S_n := \sum_{k=1}^n \frac{1}{2^k} X_k.$$

- (i) Find the probability distribution of S_n .
- (ii) Prove that for any $p \in [1, \infty]$ the sequence S_n converges a.s. and L^p to a random variable S uniformly distributed on $[0, 1]$.
- (iii) Compute the characteristic functions $F_n(\xi) = \mathbb{E}[\exp(i\xi S_n)]$ and deduce *Viète's formula*

$$\frac{\sin \xi}{\xi} = \prod_{n=1}^{\infty} \cos(\xi/2^n).$$

- (iv) Suppose that μ is a Borel probability measure on $\mathbb{R}$ with quantile $Q : [0, 1] \rightarrow \mathbb{R}$,

$$Q(p) = \inf \{x \in \mathbb{R}; \mu[(-\infty, x]) \geq p\}.$$

Prove that the sequence $Q(S_n)$ converges a.s. to a random variable with distribution μ . Have a look at Example 1.2.22. □

Remark 2.6.4. Part (iv) of the above exercise is essentially a universality property of the simplest random experiment: tossing a fair coin. If we are able to perform this experiment repeatedly and independently, then we can approximate any probability distribution. In other words, we can approximately sample any probability distribution by flipping fair coins. □

Exercise 2.43. Let $\mu \in \text{Prob}(\mathbb{R})$ be a Borel probability measure with characteristic function $\hat{\mu}$. Prove that for any $r > 1$ we have

$$\mu[\{|x| > r\}] \leq \frac{1}{C} \int_0^1 |1 - \text{Re} \hat{\mu}(t/r)| dt, \quad C := \inf_{|x| \geq 1} \left(1 - \frac{\sin x}{x}\right).$$

Hint. Set $h(x) = 1 - \frac{\sin x}{x}$, $x \in \mathbb{R}$, so, $h(x) > 0$ for $x \neq 0$, and $C\mu[\{|x| > r\}] \leq \int_{|x| \geq r} h(x/r)\mu[dx]$. □

Exercise 2.44. This exercise describes a strengthening of Levy's continuity theorem. Suppose that (μ_n) is a sequence of Borel probability measures on $\mathbb{R}$ with characteristic functions $\hat{\mu}_n(\xi)$. Assume that the functions $\hat{\mu}_n(\xi)$ converge pointwisely to a function $f : \mathbb{R} \rightarrow \mathbb{C}$ that is continuous at 0.

- (i) Prove that the sequence $(\mu_n)_{n \in \mathbb{N}}$ is tight, i.e.,

$$\lim_{r \rightarrow \infty} \sup_{n \geq 1} \mu_n[\{|x| > r\}] = 0.$$

Hint. Use Exercise 2.43.

- (ii) Show that f is the characteristic function of a Borel *probability* measure μ . **Hint.** Use Helly's Selection Theorem 2.2.22 and Proposition 2.2.23.
- (iii) Prove that μ_n converges weakly to μ .

□

Exercise 2.45. Suppose that X is a random variable and $\varphi(\xi)$ is its characteristic function

$$\varphi(\xi) = \mathbb{E}[e^{i\xi X}].$$

Prove that the following are equivalent.

- (i) X is a.s. constant.
- (ii) There exists $r > 0$ such that $|\varphi(\xi)| = 1, \forall \xi \in [-r, r]$.

Hint. Use an independent copy X' of X .

□

Exercise 2.46. Suppose that $(X_n)_{n \geq 1}$ is a sequence of random variables and $(a_n)_{n \geq 1}, (b_n)_{n \geq 1}$ are sequences of real numbers with the following properties

- $a_n > 0, \forall n \geq 1$.
- The random variables X_n and $Y_n = a_n X + b_n$ converge in distribution to the random variables X and respectively Y .
- The random variables X, Y are not a.s.-constant.

Prove that there exist real numbers a, b such that

$$a = \lim_{n \rightarrow \infty} a_n, \quad b = \lim_{n \rightarrow \infty} b_n.$$

and $\mathbb{P}_Y = \mathbb{P}_{aX+b}$. **Hint.** Use the characteristic functions of X_n , Proposition 2.2.32 and Exercise 2.45.

□

Exercise 2.47. A probability measure $\mu \in \text{Prob}(\mathbb{R})$ is said to be an *infinitely divisible distribution* if for any $n \in \mathbb{N}$, there exists $\mu_n \in \text{Prob}(\mathbb{R})$ such that

$$\mu = \mu_n^{*n} := \underbrace{\mu * \cdots * \mu}_n.$$

We denote by $\text{Prob}_\infty(\mathbb{R})$ the collection of infinitely divisible distributions. A random variable is called *infinitely divisible* if its distribution is such.

- (i) Prove that the $\text{Poi}(\lambda), N(0, \sigma^2) \in \text{Prob}_\infty, \forall \lambda, \sigma > 0$.
- (ii) Prove that any linear combination of independent infinitely divisible random variables is an infinitely divisible random variable. In particular, the convolution of two infinitely divisible distributions is infinitely divisible.

- (iii) Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables with common distribution $\nu \in \text{Prob}(\mathbb{R})$. Denote by $N(t), t \geq 0$ a Poisson process with intensity $\lambda > 0$; see Example 1.3.7. For $t \geq 0$ we set

$$Y(t) = \sum_{k=1}^{N(t)} X_k.$$

The distribution of $Y(t)$ denoted by Q_t , is called a *compound Poisson distribution*. The distribution ν is called the *compounding distribution*. Show that

$$Q_t = e^{-\lambda t} \sum_{n=0}^{\infty} \frac{(\lambda t)^n}{n!} \nu^{*n}$$

and deduce that $Q_t * Q_s = Q_{t+s}, \forall t, s \geq 0$. In particular, Q_t is infinitely divisible. We will denote Q_t by $\text{Poi}_{\nu}(\lambda t)$.

- (iv) Compute the characteristic function of Q_t .

□

Exercise 2.48. For any $\mu \in \text{Prob}(\mathbb{R})$ we denote by μ_- the measure defined by

$$\mu_-[B] = \mu[-B], \quad \forall B \in \mathcal{B}_{\mathbb{R}},$$

where $-B := \{y \in \mathbb{R}; -y \in B\}$ and we set $\mu_s := \mu * \mu_-$.

- (i) Prove that $\widehat{\mu}_s(\xi) = |\mu(\xi)|^2, \forall \xi$. Deduce that for any $n \in \mathbb{N}$ the function $|\widehat{\mu}(\xi)|^{2/n}$ is the characteristic function of a measure $\mu_n \in \text{Prob}(\mathbb{R})$ such that $\mu_n^{*n} = \mu_s$.
- (ii) Prove that $\widehat{\mu}(\xi) \neq 0, \forall \xi \neq 0$. **Hint.** Show that $|\mu(\xi)|^{2/n}$ converges as $n \rightarrow \infty$ and then conclude using Exercise 2.44.
- (iii) Deduce that there exists a continuous function $\psi : \mathbb{R} \rightarrow \mathbb{C}$ uniquely determined by the conditions $\psi(0) = 0$ and $\widehat{\mu}(\xi) = e^{\psi(\xi)}, \forall \xi \in \mathbb{R}$.
- (iv) Prove that for any $n \in \mathbb{N}$ there exists a unique measure $\nu \in \text{Prob}(\mathbb{R})$ such that $\nu^{*n} = \mu$. We will use the notation $\nu := \mu^{*1/n}$.
- (v) Prove that any weak limit of infinitely divisible distributions is also an infinitesimal distribution.

□

Exercise 2.49. Give an example of a sequence of random variables $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that X_n converge in distribution to 0 but

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n] = \infty.$$

□

Exercise 2.50 (Skhorokhod). Suppose that $\mu_n \in \text{Prob}(\mathbb{R}), n \in \mathbb{N}$, is a sequence converging weakly to μ . Denote by $F_n : \mathbb{R} \rightarrow [0, 1]$ the distribution function of μ_n ,

$$F_n(x) = \mu_n[(-\infty, x]],$$

and by Q_n the associated quantile function (see (1.2.5))

$$Q_n : [0, 1] \rightarrow \mathbb{R}, \quad Q_n(t) = \inf \{x; t \leq F_n(x)\}.$$

We can regard Q_n as random variables defined on the probability space

$$([0, 1], \mathcal{B}_{[0,1]}, \lambda_{[0,1]}),$$

where $\lambda_{[0,1]}$ denotes the Lebesgue measure on $[0, 1]$. As shown in Example 1.2.21,

$$\mu_n = (Q_n)_\# \lambda_{[0,1]},$$

so that μ_n is the probability distribution of Q_n . Prove that the sequence Q_n converges a.s. on $[0, 1]$ to a random variable with probability distribution μ .

In other words, given any sequence $\mu_n \in \text{Prob}(\mathbb{R})$ that converges weakly to $\mu \in \text{Prob}(\mathbb{R})$, we can find a sequence of random variables X_n , defined on the same probability space with $\mathbb{P}_{X_n} = \mu_n$ and such that X_n converges a.s. to a random variable X with distribution μ . $\square$

Exercise 2.51. Fix $n \in \mathbb{R}$ and denote by $C_b(\mathbb{R}^n, \mathbb{C})$ the space of continuous, bounded functions $\mathbb{R}^n \rightarrow \mathbb{C}$. Denote by $\mathcal{T}_n$ the complex subspace of $C_b(\mathbb{R}^n, \mathbb{C})$ spanned by the functions

$$\{e_\xi(x) = e^{i(\xi, x)}\}_{\xi \in \mathbb{R}^n},$$

where $(-, -)$ denotes the canonical inner product in $\mathbb{R}^n$.

- (i) Prove that $\mathcal{T}_n$ is a $\mathbb{C}$ -algebra of functions.
- (ii) Consider the continuous cut-off function $\eta : \mathbb{R} \rightarrow [0, \infty)$ defined by

$$\eta(x) = \begin{cases} 0, & |x| \geq 2, \\ 1, & |x| \leq 1, \\ \text{linear}, & 1 < |x| < 2. \end{cases}$$

For $L > 0$ define $\rho_L : \mathbb{R}^n \rightarrow \mathbb{R}$ $\rho_L(x_1, \dots, x_n) = \prod_{j=1}^n \eta(x_j/L)$. Note that ρ_L is supported in the cube $C_L = [-2L, 2L]^n$. Let $f \in C_b(\mathbb{R}, \mathbb{C})$. Prove that for any $\varepsilon > 0$ and any $L > 0$ there exists a trigonometric polynomial $T = T_{\varepsilon, L}$ such that

$$\sup_{x \in C_L} |\rho_L(x)f(x) - T(x)| < \varepsilon \quad \text{and} \quad \|T\| < \|f\| + 1.$$

- (iii) Prove that for any $f \in C_b(\mathbb{R}, \mathbb{C})$ there exists a sequence $(T_\nu)_{\nu \in \mathbb{N}}$ in $\mathcal{T}_n$ such that T_ν converges to F uniformly on compacts and $\|T_\nu\| < \|f\| + 1, \forall \nu$.
- (iv) Use (iii) to give a new proof that a probability measure on $\mathbb{R}^n$ is uniquely determined by its characteristic function.

$\square$

Exercise 2.52. Let μ be a Borel probability measure on $\mathbb{R}$ satisfying

$$\exists r_0 > 0 : \int_{\mathbb{R}} e^{tx} \mu[dx] < \infty, \quad \forall |t| < r_0.$$

- (i) Let $p \in [1, \infty)$. Prove that the map

$$L^p(\mathbb{R}, \mu) \ni f \mapsto Tf \in C_b(\mathbb{R}, \mathbb{C}), \quad (Tf)(\xi) = \int_{\mathbb{R}} e^{i\xi x} f(x) \mu[dx]$$

is injective. **Hint.** Reduce to Theorem 2.2.27 by writing $f = f_+ - f_-$.

- (ii) Let $f \in L^2(\mathbb{R}, \mu)$. Prove that there exists $r_1 > 0$ such that for any complex number such that $|\text{Im } z| < r_1$ the complex valued function $\mathbb{R} \ni x \mapsto e^{izx} f(x) \in \mathbb{C}$ is μ integrable and the resulting function

$$z \mapsto \hat{f}(z) = \int_{\mathbb{R}} e^{izx} f(x) \mu[dx]$$

is holomorphic in the strip $\{|\operatorname{Im} z| < r_1\}$.

- (iii) Prove that $\mathbb{R}[x]$, the space of polynomials with real coefficients, is dense in $L^2(\mathbb{R}, \mu)$. **Hint.** You have to show that if $f \in L^2(\mathbb{R}, \nu)$ satisfies

$$\int_{\mathbb{R}} f(x) x^n \mu[dx] = 0, \quad \forall n \geq 0,$$

the $f = 0$ μ -a.s. Prove that $\widehat{f}^{(n)}(0) = 0, \forall n = 0, 1, 2, \dots$

- (iv) Consider the Hermite polynomials $(H_n(x))_{n \geq 0}$ described in Exercise 1.33. Prove that the collection

$$\frac{1}{\sqrt{n!}} H_n, \quad n \geq 0$$

is a *complete* orthonormal basis of the Hilbert space $L^2(\mathbb{R}, \gamma_1)$, where γ_1 is the standard Gaussian measure on $\mathbb{R}$.

□

Exercise 2.53. Suppose that μ_0, μ_1 are two Borel probability measures such that $\exists t_0 > 0$

$$\int_{\mathbb{R}} e^{tx} \mu_0[dx] = \int_{\mathbb{R}} e^{tx} \mu_1[dx], \quad \forall |t| < t_0.$$

Fix $r > 0$ as in Exercise 2.52(ii) such that for any complex number the functions

$$z \mapsto F_k(z) = \int_{\mathbb{R}} e^{izx} f(x) \mu_k[dx], \quad k = 0, 1,$$

are well defined and holomorphic in the strip $\{|\operatorname{Im} z| < r\}$. Show that $F_0 = F_1$ and deduce that $\mu_0 = \mu_1$. **Hint.** Set $F = F_1 - F_0$. Use the Cauchy-Riemann equations to prove that $\frac{d^n F}{dz^n} \Big|_{z=t} = 0, \forall n \in \mathbb{N}, \forall t \in (-r, r)$. □

Exercise 2.54. Suppose that $(X_n)_{n \in \mathbb{N} \cup \infty}$ is a family of random variables such that there exists $T > 0$ with the following properties

(i)

$$\sup_{n \in \mathbb{N} \cup \infty} \mathbb{E}[e^{tX_n}] < \infty, \quad \forall |t| \leq T.$$

(ii)

$$\lim_{n \rightarrow \infty} \mathbb{E}[e^{tX_n}] = \mathbb{E}[e^{tX_\infty}], \quad \forall |t| \leq T.$$

Prove that X_n converge in distribution to X_∞ . **Hint.** Fix $t_0 \in (0, T)$ and consider the measures $\nu_n[dx] = \cosh(t_0 x) \mathbb{P}_{X_n}[dx]$ and argue as in the proof of Proposition 2.2.4 that $\nu_n \Rightarrow \nu_\infty$. □

Exercise 2.55. A function $F : \mathbb{R}^N \rightarrow \mathbb{C}$ is called *nonnegative definite* if it is continuous and for any $n \in \mathbb{N}$, any $\xi_1, \dots, \xi_n \in \mathbb{R}^N$ and any $z_1, \dots, z_n \in \mathbb{C}$ we have

$$\sum_{i,j=1}^n F(\xi_i - \xi_j) z_i \bar{z}_j \geq 0. \quad (2.6.3)$$

It is called *positive definite* if it is nonnegative definite and in (2.6.3) we have equality iff $z_1 = \dots = z_n = 0$.

Denote by $C_{\text{cpt}}(\mathbb{R}^N, \mathbb{C})$ the space of compactly supported continuous functions $\mathbb{R}^N \rightarrow \mathbb{C}$

(i) Let $\mu \in \text{Prob}(\mathbb{R}^n)$. Prove that its Fourier transform $\widehat{\mu}$ is a positive definite function.

(ii) Suppose $F : \mathbb{R}^N \rightarrow \mathbb{X}$ is nonnegative definite. Then the following hold.

(a) $F(0) \in [0, \infty)$.

- (b) $F(-x) = \overline{F(x)}$.
 (c) For any $\varphi : C_{\text{cpt}}(\mathbb{R}^N, \mathbb{C})$.

$$\int_{\mathbb{R}^N \times \mathbb{R}^N} F(\xi - \eta) \varphi(\xi) \overline{\varphi(\eta)} d\xi d\eta \geq 0. \quad (2.6.4)$$

- (iii) Conversely, prove that if the continuous function $F : \mathbb{R}^N \rightarrow \mathbb{C}$ satisfies (i)(c) then it is nonnegative definite.

Hint. (i)(c) Use Riemann sums to approximate the integral in (2.6.4). Fix a nonnegative continuous function $\rho : \mathbb{R}^N \rightarrow \mathbb{R}$ supported inside the unit ball of $\mathbb{R}^N$ and such that $\int_{\mathbb{R}^N} \rho(\xi) d\xi = 1$. For $t > 0$ we set $\rho_t(\xi) = t^{-N} \rho(\xi/t)$ (ii). In (2.6.4) choose φ of the form

$$\varphi(\xi) = \sum_{j=1}^n z_j \rho_\varepsilon(\xi - \xi_j),$$

and then let $\varepsilon \searrow 0$. □

Exercise 2.56 (De Moivre). Let $X_n \sim \text{Bin}(n, 1/2)$ and $Y \sim N(0, 1)$. Prove that

$$\lim_{n \rightarrow \infty} \frac{\mathbb{P}[|X_n - n/2| \leq \frac{r}{2}\sqrt{n}]}{\mathbb{P}[|Y| < r]} = 1, \quad \forall r > 0. \quad \square$$

Exercise 2.57 (t-statistic). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables such that $\mathbb{E}[X_n] = 0$, $\mathbb{E}[X_n^2] = \sigma^2 < \infty$, $\forall n$. We set

$$M_n = \frac{1}{n} \sum_{k=1}^n X_k, \quad V_n = \frac{1}{n-1} \sum_{k=1}^n (X_k - M_n)^2, \quad T_n = \sqrt{n} \frac{M_n}{\sqrt{V_n}}.$$

- (i) Prove that V_n converges in probability to σ^2 .
 (ii) Prove that T_n converges in distribution to a standard normal random variable. **Hint.** Use CLT and Slutsky's theorem. □

Exercise 2.58. Suppose that $X = X_\lambda$ is a $\text{Gamma}(1, \lambda)$ random variable (see Example 1.3.38) and $Y = Y_\lambda$ is a random variable such that

$$\mathbb{P}[Y = n | X] = \frac{X^n}{n!} e^{-nX}, \quad \forall n = 0, 1, 2, \dots$$

In other words, conditioned on $X = x$ the random variable Y is $\text{Poi}(x)$.

- (i) Compute the characteristic function of Y .
 (ii) Show that the random variable

$$\frac{1}{\sqrt{\text{Var}[Y_\lambda]}} (Y_\lambda - \mathbb{E}[Y_\lambda])$$

converges in distribution to $N(0, 1)$ as $\lambda \rightarrow \infty$. □

Exercise 2.59. Suppose that X, Y are independent random normal variables. Set $Z = XY$.

- (i) Show that

$$\mathbb{M}_Z(\lambda) = \frac{1}{\sqrt{1 - \lambda^2}}, \quad |\lambda| < 1,$$

and deduce that

$$\Psi_Z(\lambda) \leq \frac{\lambda^2}{2(1-\lambda^2)}, \quad \forall \lambda \in (-1, 1).$$

(ii) Prove that I_Z (see Proposition 2.3.1) satisfies

$$I_Z(z) \geq \frac{1}{3}(\sin z)^2 \geq \frac{1}{12}z^2, \quad \forall |z| < \frac{\pi}{6}.$$

□

Exercise 2.60. Given any zero mean random variable X the following are equivalent.

- (i) The random variable X is sub-gaussian (of some type σ^2).
- (ii) There exists $c \geq 0$ and a Gaussian random variable Z such that

$$\mathbb{P}[|X| > s] \leq c\mathbb{P}[|Z| > s], \quad \forall s \geq 0.$$

- (iii) There exists a constant $\theta \geq 0$ such that

$$\mathbb{E}[X^{2k}] \leq \frac{(2k)!}{2^k k!} \theta^{2k}, \quad \forall k \in \mathbb{N}.$$

There exists a constant $\sigma \geq 0$ such that

$$\mathbb{E}\left[\frac{\lambda}{2\sigma^2}X\right] \leq \frac{1}{\sqrt{1-\lambda}}, \quad \forall \lambda \in [0, 1).$$

□

Exercise 2.61. Let $\mathfrak{X}$ be a finite set. The *entropy* of a random variable $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}$ is

$$\text{Ent}_2[X] := \text{Ent}_2[\mathbb{P}_X] = - \sum_{x \in \mathfrak{X}} p_X(x) \log_2 p(x), \quad p_X(x) = \mathbb{P}[\{X = x\}].$$

Given two random variables $X_i : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}_i, i = 1, 2$, we define their *relative entropy* to be

$$\text{Ent}_2[X_2 | X_1] := \sum_{(x_1, x_2) \in \mathfrak{X}_1 \times \mathfrak{X}_2} p_{X_1, X_2}(x_1, x_2) \log_2 \left(\frac{p_{X_1, X_2}(x_1, x_2)}{p_{X_1}(x_1)} \right),$$

where $p_{X_1, X_2}(x_1, x_2) = \mathbb{P}[\{X_1 = x_1, X_2 = x_2\}]$.

- (i) Show that if $X_i : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}_i, i = 1, 2$, are random variables, then

$$\text{Ent}_2[X_2] - \text{Ent}_2[X_2 | X_1] = \mathbb{D}_{KL}[\mathbb{P}_{(X_1, X_2)}, \mathbb{P}_{X_1} \otimes \mathbb{P}_{X_2}],$$

where $\mathbb{D}_{KL}$ is the Kullback-Leibler divergence defined in (2.3.8).

- (ii) Suppose that we are given n finite sets $\mathfrak{X}_i, i = 1, \dots, n$ and n maps

$$X_i : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}_i.$$

We denote by $\text{Ent}_2(X_1, \dots, X_n)$ the entropy of the product random variable

$$(X_1, \dots, X_n) : \Omega \rightarrow \mathfrak{X}_1 \times \dots \times \mathfrak{X}_n.$$

Prove that

$$\text{Ent}_2[X_1, \dots, X_n] = \sum_{k=1}^n \text{Ent}_2[X_k | (X_{k-1}, \dots, X_1)].$$

□

Exercise 2.62. Suppose that X, Y are two discrete random variable. Denote by $\text{Ent}_2[X | Y]$ the relative entropy defined in Exercise 2.61.

(i) Prove that

$$\text{Ent}_2 [X|Y] \leq \text{Ent}_2 [X].$$

$$\text{Set } I[X, Y] := \text{Ent}_2 [X] - \text{Ent}_2 [X|Y]$$

(ii) Prove that

$$\begin{aligned} I[X, Y] &= I[Y, X] = \text{Ent}_2 [X] + \text{Ent}_2 [Y] - \text{Ent}_2 [(X, Y)] \\ &= \sum_{x,y} \mathbb{P}[X=x, Y=y] \log_2 \left(\frac{\mathbb{P}[X=x, Y=y]}{\mathbb{P}[X=x]\mathbb{P}[Y=y]} \right). \end{aligned}$$

(iii) Prove that if $Z \perp\!\!\!\perp_Y X$ then

$$I[X, Y] \geq I[X, Z].$$

Exercise 2.63 (Herbst). Let $\phi : [0, \infty) \rightarrow \mathbb{R}$, $\phi(x) = x \log x$, where $0 \cdot \log 0 := 0$. For any nonnegative random variable Z we set

$$\mathbb{H}[Z] = \text{Ent}_\phi [Z] = \mathbb{E}[\phi(Z)] - \phi(\mathbb{E}[Z]).$$

Suppose that X is a random variable such that $\mathbb{M}_X(\lambda) = \mathbb{E}[e^{\lambda X}] < \infty$, for all λ is an open interval J containing 0. We set $\mathbb{H}_X(\lambda) := \mathbb{H}[e^{\lambda X}]$. Prove that if

$$\mathbb{H}_X(\lambda) \leq \frac{\lambda^2 \sigma^2}{2} \mathbb{M}_X(\lambda),$$

then $X \in \mathbb{G}(\sigma^2)$. □

Exercise 2.64 (Poincaré phenomenon). Denote by S^n the unit sphere in $\mathbb{R}^{n+1}$,

$$S^n := \left\{ (x_0, x_1, \dots, x_n); \sum_{k=0}^n x_k^2 = 1 \right\}.$$

Suppose that $(X_0, \dots, X_n)$ is a random point uniformly distributed on S^n with respect to the canonical Euclidean volume on S^n .

(i) Prove that there exists $C > 0$ such that

$$\forall n \in \mathbb{N}, \quad r \in [0, 1]; \quad \mathbb{P}[|X_0| > r] \leq C e^{-\frac{nr^2}{2}}.$$

Thus, for spheres of large dimension n most of the volume is concentrated near the Equator $\{x_0 = 0\}$!

(ii) Prove that $\sqrt{n}X_0$ converges a.s. to a standard normal random variable.

Hint. Choose independent standard normal random variables $Y_0, \dots, Y_n$ set $Z_n = Y_0^2 + \dots + Y_n^2$. Show that the random vector

$$(X_0, \dots, X_n) = \frac{1}{\sqrt{Z}} (Y_0, \dots, Y_n)$$

is uniformly distributed on S^n . You can take for granted the fact that any finite Borel measure on S^n that is invariant under the action of $SO(n+1)$ on S^n is a multiple of the Euclidean volume measure.¹⁸ For (i) use Exercise 1.49 and Appendix A.1. Reduce (ii) to the SLLN. □

Exercise 2.65. Let X be a random variable such that $|X| < 1$, a.s.. Assume $\mathbb{E}[X] = 0$ and $\sigma^2 := \text{Var}[X] < \infty$.

¹⁸Can you prove this?

(i) Prove that

$$\mathbb{E}[e^{\lambda X}] \leq \exp\left(\sigma^2(e^\lambda - 1 - \lambda)\right), \quad \forall \lambda > 0.$$

(ii) Prove that

$$\mathbb{P}[X > x] \leq \exp\left(-\frac{x^2}{2\sigma^2 + 2x/3}\right).$$

□

Exercise 2.66. Let X, Y be random variables such that

$$\mathbb{E}[\Phi(X)] \leq \mathbb{E}[\Phi(Y)]$$

for every nondecreasing, convex function $\Phi : \mathbb{R} \rightarrow [0, \infty)$. If $\exists c_1, c_2, \alpha > 0$ such that

$$\mathbb{P}[Y > t] \leq c_1 e^{-c_2 t^\alpha}, \quad \forall t \geq 0,$$

then

$$\mathbb{P}[X > t] \leq c_1 e^{1-c_2 t^\alpha}, \quad \forall t \geq 0.$$

□

Exercise 2.67. Suppose that $\psi : [0, \infty) \rightarrow [0, \infty)$ is an *Young function*, i.e., it is convex, increasing, $\psi(0) = 0$, and $\psi(x)/x \rightarrow \infty$ as $x \rightarrow \infty$. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For any Young function ψ and any random variable $X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ we set

$$\|X\|_\psi = \inf \{ t > 0; \mathbb{E}[\psi(X/t)] \leq 1 \},$$

where $\inf \emptyset := \infty$. We set

$$\mathcal{L}_\psi(\Omega, \mathcal{S}, \mathbb{P}) = \{ X \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P}); \|X\|_\psi < \infty \}$$

and we denote by L_ψ the quotient of $\mathcal{L}_\psi$ modulo the a.s. equality.

(i) Prove that $L_\psi(\Omega, \mathcal{S}, \mathbb{P})$ is a normed space, called the *Orlicz space* determined by the young function ψ .

(ii) Show that when $\psi(x) = x^p$, $p \in [1, \infty)$ we have $L_\psi = L^p$.

(iii) Let $\Psi(x) = e^{x^2} - 1$. Prove that X is subgaussian if and only if $X \in L_\Psi$. □

Exercise 2.68. Suppose that Y_1, Y_2 are independent and uniformly distributed on $[0, 1]$. Prove that

$$X_1 = \sqrt{-2 \log(Y_1)} \cdot \cos(2\pi Y_2), \quad X_2 = \sqrt{-2 \log(Y_1)} \cdot \sin(2\pi Y_2)$$

are independent standard normal random variables. □

Exercise 2.69 (Cramér-Wold Device). Let V be a finite dimensional vector space. Fix a sequence of probability measure $\mu_n \in \text{Prob}(V)$. Prove that the following statements are equivalent.

(i) The sequence (μ_n) converges weakly to $\mu \in \text{Prob}(V)$.

(ii) For any $\xi \in V^*$, $\widehat{\mu_n}(\xi) \rightarrow \widehat{\mu}(\xi)$.

(iii) For any $\xi \in V^*$, $\xi_\#(\mu_n) \Rightarrow \xi_\#(\mu)$, where we recall that for any $\nu \in \text{Prob}(V)$ and any $\xi \in V^*$ we denote by $\xi_\#(\nu)$ the pushforward of ν via the function $\xi : V \rightarrow \mathbb{R}$. □

Exercise 2.70. Let V be an n -dimensional real vector space. We denote by V^* its dual, $V^* = \text{Hom}(V, \mathbb{R})$. We have a natural pairing

$$\langle -, - \rangle : V^* \times V \rightarrow \mathbb{R}, \quad \langle \xi, x \rangle := \xi(x), \quad \forall \xi \in V^*, \quad x \in V.$$

A Borel probability measure $\mu \in \text{Prob}(V)$ is called *Gaussian* if for every linear functional $\xi \in V^*$, the resulting random variable

$$\xi : (V, \mathcal{B}_V, \mu) \rightarrow \mathbb{R}$$

is Gaussian with mean $m[\xi]$ and variance $v[\xi]$, i.e., (see Example 1.3.37)

$$\mathbb{P}_\xi[dx] = \Gamma_{m[\xi], v[\xi]}[dx] = \frac{1}{(2\pi)^{n/2}} \cdot e^{-\frac{(x-m[\xi])^2}{2v[\xi]}} dx.$$

A random vector $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow V$ is called *Gaussian* if its probability distribution is a Gaussian measure on V

- (i) Show that the map $V^* \ni \xi \mapsto m[\xi] \in \mathbb{R}$ is linear and thus defines an element $m = m_\mu \in (V^*)^*$ called the *mean* of the Gaussian measure. Moreover, using the natural isomorphism¹⁹

$$J : V \rightarrow V^{**}$$

we have

$$J^{-1}(m_\mu) = \int_V x \mu[dx] \in V.$$

- (ii) Define $C = C_\mu : V^* \times V^* \rightarrow \mathbb{R}$

$$C(\xi, \eta) = \frac{1}{4} (v[\xi + \eta] - v[\xi - \eta]) = \mathbb{E}_\mu[(\xi - m[\xi])(\eta - m[\eta])].$$

Show that C is a bilinear form, it is symmetric and positive semidefinite. It is called the *covariance form* of the Gaussian measure μ .

- (iii) Show that if μ_0, μ_1 are Gaussian measures on V_0 and respectively V_1 , then the product $\mu_0 \otimes \mu_1$ is a Gaussian measure on $V_0 \oplus V_1$. Moreover,

$$m[\mu_0 \otimes \mu_1] = m_{\mu_0} \oplus m_{\mu_1}, \quad C_{\mu_0 \otimes \mu_1} = C_{\mu_0} \oplus C_{\mu_1}.$$

We set

$$\Gamma_{\mathbb{1}_n} := \underbrace{\Gamma_1 \otimes \cdots \otimes \Gamma_1}_n.$$

$\Gamma_{\mathbb{1}_n}$ is called the canonical Gaussian measure on $\mathbb{R}^n$. More explicitly

$$\Gamma_{\mathbb{1}_n}[dx] = \frac{1}{(2\pi)^{n/2}} e^{-\frac{|x|^2}{2}} dx,$$

where $|x|$ denotes the Euclidean norm of the vector $x \in \mathbb{R}^n$.

- (iv) Suppose that V_0, V_1 are real finite dimensional vector spaces, μ is a Gaussian measure on V_0 and $A : V_0 \rightarrow V_1$ is a linear map. Denote by μ_A the pushforward of μ via the map A , $\mu_A := A_\# \mu$. Prove that μ_A is a Gaussian measure on V_1 with mean $m_{\mu_A} = A m_\mu$ and covariance form

$$C_A : V_1^* \times V_1^* \rightarrow \mathbb{R}, \quad C_A(\xi_1, \eta_1) = C_\mu(A^* \xi_1, A^* \eta_1), \quad \forall \xi_1, \eta_1 \in V_1^*.$$

Above, $A^* : V_1 \rightarrow V_0$ is the dual (transpose) of the linear map A .

¹⁹For a vector $v \in V$, $J(v)$ is the linear functional on V^* that associated to $\xi \in V^*$ the number $\langle \xi, v \rangle$, i.e. $J(v)(\xi) = \langle \xi, v \rangle$. The map J is an isomorphism when V is finite dimensional.

- (v) Fix a basis $\{e_1, \dots, e_n\}$ of V so we can identify V and V^* with $\mathbb{R}^n$ and C with a symmetric positive semidefinite matrix. Denote by A its unique positive semidefinite square root. Show that the pushforward $A_{\#}\Gamma_{1_n}$ is a Gaussian measure on $\mathbb{R}^n$ with mean zero and covariance form $C = A^2$.
- (vi) Show that if μ is a Gaussian measure on V with mean m covariance form C , then its Fourier transform is

$$\hat{\mu}(\xi) : \int_V e^{i\langle \xi, v \rangle} \mu[dv] = e^{im[\xi]} e^{-\frac{1}{2}C(\xi, \xi)}, \quad \forall \xi \in V^*.$$

- (vii) Show that a Gaussian measure is uniquely determined by its mean and covariance form. We denote by $\Gamma_{m,C}$ the Gaussian measure with mean m and covariance C .
- (viii) Suppose that C is a symmetric positive definite $n \times n$ matrix. Prove that the Gaussian measure on $\mathbb{R}^n$ with mean 0 and covariance form C is

$$\Gamma_{0,C}[dx] = \frac{1}{(\det(2\pi C))^{n/2}} e^{-\frac{\langle C^{-1}x, x \rangle}{2}} dx$$

where $\langle -, - \rangle$ denotes the canonical inner product on $\mathbb{R}^n$. **Hint.** Analyze first the case when C is a diagonal matrix. $\square$

Exercise 2.71. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and E a finite dimensional real vector space. Recall that a Borel measurable map $X : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow E$ is called a *Gaussian random vector* if its distribution $\mathbb{P}_X = X_{\#}\mathbb{P}$ is a Gaussian measure on E ; see Exercise 2.70.

Suppose that $X_1, \dots, X_m \in \mathcal{L}^0(\Omega, \mathcal{S}, \mathbb{P})$ are *jointly Gaussian* random variables, i.e., the random vector $\vec{X} = (X_1, \dots, X_m) : \Omega \rightarrow \mathbb{R}^m$ is Gaussian.

- (i) Prove that each of the variables $X_1, \dots, X_m$ is Gaussian and the covariance form of the Gaussian measure $\mathbb{P}_{\vec{X}} \in \text{Prob}(\mathbb{R}^n)$ is given by the matrix $\text{Var}[\vec{X}] := (v_{ij})_{1 \leq i, j \leq m}$

$$v_{ij} = \text{Cov}[X_i, X_j], \quad \forall 1 \leq i, j \leq m$$

The matrix $\text{Var}[\vec{X}]$ is called the *variance* of the Gaussian vector $\vec{X}$

- (ii) Prove that $X_1, \dots, X_m$ are independent if and only if the matrix $\text{Var}[\vec{X}]$ is diagonal, i.e.,

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_i] \mathbb{E}[X_j], \quad \forall i \neq j.$$

- (iii) Suppose that $\vec{Y} = (Y_1, \dots, Y_n)$ is a Gaussian vector such that $(\vec{X}, \vec{Y})$ is also a Gaussian vector. Denote by $\text{Cov}[\vec{X}, \vec{Y}]$ the $m \times n$ -matrix with entries

$$\text{Cov}[X_i, Y_j], \quad 1 \leq i \leq m, \quad 1 \leq j \leq n.$$

Prove that the random vectors $\vec{X}$ and $\vec{Y}$ are independent iff $\text{Cov}[\vec{X}, \vec{Y}] = 0$.

Hint. Use the results in Exercise 2.70. $\square$

Exercise 2.72 (Gaussian regresion). Suppose that X_0, X_1 are mean zero, jointly Gaussian random vectors, i.e., the random vector (X_0, X_1) is Gaussian. Suppose that $\text{Var}[X_0]$ is invertible.

- (i) Prove that

$$\mathbb{E}[X_1 | X_0] = \text{Cov}[X_1, X_0] \text{Var}[X_0]^{-1} X_0.$$

In particular, $\mathbb{E}[X_0 | X_1]$.

- (ii) Consider the conditioned random vector $X_1 | X_0 = 0$ with distribution $\mathbb{P}_{X_1 | X_0=0}$; see Example 1.4.35. Prove that $X_1 | X_0 = 0$ is Gaussian random vectors with mean 0 and variance given by the *Gaussian regression formula*

$$\begin{aligned} \text{Var} [X_1 | X_0 = 0] &= \text{Var} [X_1] \\ &\quad - \text{Cov} [X_1, X_0] \cdot \text{Var} [X_0]^{-1} \cdot \text{Cov} [X_0, X_1]. \end{aligned}$$

Hint. (i) Use Exercise 2.71 to prove that the Gaussian vector

$$X_1 - \text{Cov} [X_1, X_0] \text{Var} [X_0]^{-1} X_0$$

is independent of X_0 . (ii) Use (i). □

Remark 2.6.5. The result in Exercise 2.72 is remarkable. Let us explain its typical use in statistics.

Suppose we want to understand the random vector X_1 , but all that we truly understand is the random vector X_0 . A quantity of the form $f(X_0)$, $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ measurable, is called a predictor. Predictors are guesses about X_1 using information about X_0 .

The simplest predictors are of the form LX_0 where $L : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is an affine operator. These are called *linear predictors*. The conditional expectation $\mathbb{E}[X_1 | X_0]$ is the predictor closest to X_1 . The linear predictor L^2 -closest to X_1 is called the *linear regression*. The linear regressions are obtained via the least squares approximation.

The result in the above exercise shows that, when the random vectors X_0, X_1 are jointly Gaussian, the best predictor of X_1 given X_0 is the linear regression. This is another reason why the Gaussian variables are extremely convenient to work with in practice. □

Exercise 2.73 (Maxwell). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of mean zero i.i.d. random variables. For each $n \in \mathbb{N}$ we denote by V_n the random vector $V_n := (X_1, \dots, X_n)$. Prove that the following are equivalent.

- (i) The random variables X_n are Gaussian.
- (ii) For any $n \in \mathbb{N}$ and for any orthogonal map $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ the random vectors V_n and RV_n have identical distributions. □

Exercise 2.74. Suppose that $X_1, \dots, X_n$ are independent standard normal random variables. Set

$$\bar{X} := \frac{1}{n} (X_1 + \dots + X_n), \quad S^2 := \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2.$$

- (i) Let $R = (r_{ij})_{1 \leq i, j \leq n}$ be an $n \times n$ orthogonal matrix such that

$$R_{1i} = \frac{1}{\sqrt{n}}, \quad \forall i = 1, \dots, n$$

and set

$$Y_i = \sum_{j=1}^n r_{ij} X_j$$

- (ii) Prove that $Y_1, \dots, Y_n$ are independent standard normal random variables.
- (iii) Prove that $(n-1)S^2 = Y_2^2 + \dots + Y_n^2 \sim \chi^2(n-1)$.

(iv) Set

$$T_n := \frac{\bar{X}}{\bar{S}}.$$

Prove that $\sqrt{n}T_n \sim \text{Stud}_{n-1}$, where Stud_p denotes the *Student t-distribution* with p degrees of freedom

$$\text{Stud}_p = \frac{1}{\sqrt{p\pi}} \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(1+t^2/p)^{(p+1)/2}} dt, \quad t \in \mathbb{R}, \quad p > 0.$$

Hint. Note that $\sqrt{n}\bar{X} = Y_1$ and $\bar{S}$ are independent. Conclude using (ii)

□

Exercise 2.75. Suppose that X is a standard normal random variable and Z is a Bernoulli random variable, independent of X , with success probability $p = \frac{1}{2}$.

- (i) Prove that $Y = XZ$ is also a standard normal random variable.
- (ii) Prove that $X + Y$ is not Gaussian.

□

Exercise 2.76. Suppose that X is a standard normal random variable defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For any $\theta > 0$ we set $Z_\theta := e^{-\theta X - \frac{1}{2}\theta^2}$.

- (i) Prove that $\mathbb{E}_{\mathbb{P}}[Z_\theta] = 1$.
- (ii) Define a new measure $\mathbb{P}_\theta$ on $\mathcal{S}$ by

$$\mathbb{P}_\theta[d\omega] = Z_\theta(\omega)\mathbb{P}[d\omega],$$

i.e.,

$$\mathbb{P}_\theta[S] = \int_S Z_\theta(\omega)\mathbb{P}[d\omega].$$

Prove that $\mathbb{P}_\theta$ is a probability measure and the random variable

$$X + \theta : (\Omega, \mathcal{S}, \mathbb{P}_\theta) \rightarrow \mathbb{R}$$

is standard normal.

□

Exercise 2.77. Suppose that $\mathbb{T}$ is a compact interval of the real axis, and $(X_t)_{t \in \mathbb{T}}, (Y_t)_{t \in \mathbb{T}}, (Z_t)_{t \in \mathbb{T}}$ are real valued stochastic processes such that (Y_t) and (Z_t) are modifications of (X_t) with a.s. continuous paths. Prove that the processes (Y_t) and (Z_t) are indistinguishable.

□

Exercise 2.78. Fix a Brownian motion $(B_t)_{t \geq 0}$ defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Denote by $\mathcal{E}$ the vector subspace of $L^2([0, 1], \lambda)$ spanned by the functions $\mathbf{I}_{(s, t]}$, $0 \leq s < t \leq 1$.

- (i) Prove that any function $f \in \mathcal{E}$ admits a convenient representation, i.e., a representation of the form

$$f = \sum_{k=1}^n c_k \mathbf{I}_{(s_k, t_k]}, \quad c_k \in \mathbb{R},$$

where the intervals $(s_j, t_j], (s_k, t_k]$ are disjoint for $j \neq k$.

- (ii) Prove that $\mathcal{E}$ is dense in $L^2([0, 1], \lambda)$.

(iii) Let $f \in \mathcal{E}$ and consider two convenient representations of f

$$\sum_{k=1}^n c_k \mathbf{I}_{(s_k, t_k]} = f = \sum_{i=1}^m c'_i \mathbf{I}_{(s'_i, t'_i]}.$$

Show that

$$\sum_{k=1}^n c_k (B_{t_k} - B_{s_k}) = \sum_{i=1}^m c'_i (B_{t'_i} - B_{s'_i}) =: \mathcal{W}(f).$$

(iv) Show that for any $f \in \mathcal{E}$ we have $\mathcal{W}(f) \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ and $\|\mathcal{W}(f)\|_{L^2(\Omega)} = \|f\|_{L^2([0,1])}$.

(v) Prove that the map $\mathcal{W} : \mathcal{E} \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$ is linear and extends to a linear isometry $\mathcal{W} : L^2([0,1], \lambda) \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$ whose image consists of Gaussian random variables. In other words, this isometry is a Gaussian white noise. The map $\mathcal{W}$ is called the *Wiener integral*. It is customary to write

$$\mathcal{W}(f) = \int_0^1 f(s) dB_s.$$

□

Exercise 2.79. The space $F := C([0, \infty))$ of continuous functions $[0, \infty) \rightarrow \mathbb{R}$ is equipped with a natural metric d ,

$$d(f, g) = \sum_{n \in \mathbb{N}} \frac{1}{2^n} \min(1, d_n(f, g)), \quad d_n(f, g) := \sup_{t \in [n-1, n]} |f(t) - g(t)|.$$

Denote by $\mathcal{B}_F$ the Borel algebra of F . For each $t \geq 0$ we define $E_t : F \rightarrow \mathbb{R}$, $E_t(f) = f(t)$. We set

$$\mathcal{S}_t = E_t^{-1}(\mathcal{B}_\mathbb{R}), \quad \forall t \geq 0, \quad \mathcal{S} = \bigcup_{t \geq 0} \mathcal{S}_t.$$

- (i) Prove that (F, d) is a complete separable metric space.
- (ii) Prove that E_t is a continuous function on F , $\forall t \geq 0$.
- (iii) Prove that $\mathcal{B}_F = \mathcal{S}$. **Hint.** Compare with Exercise 1.4.
- (iv) Suppose that $(\Omega, \mathcal{A})$ is a measurable space and $W : \Omega \rightarrow F$ is a map

$$\Omega \in \omega \mapsto W_\omega \in F.$$

Prove that W is $(\mathcal{A}, \mathcal{B}_F)$ -measurable if and only if for any $t \geq 0$ the function

$$W^t : (\Omega, \mathcal{A}) \rightarrow \mathbb{R}, \quad \omega \mapsto W_\omega(t)$$

is measurable. **Hint.** Use (iii).

- (v) For any $0 \leq t_1 < t_2 < \dots < t_n$ define

$$\mathbf{Ev}_{t_1, \dots, t_n} : F \rightarrow \mathbb{R}^n, \quad \mathbf{Ev}_{t_1, \dots, t_n}(f) = (f(t_1), \dots, f(t_n)).$$

Suppose that μ_0, μ_1 are two Borel probability measures on F such that

$$(\mathbf{Ev}_{t_1, \dots, t_n})_{\#} \mu_0 = (\mathbf{Ev}_{t_1, \dots, t_n})_{\#} \mu_1,$$

for any $n \in \mathbb{N}$ and any $0 \leq t_1 < \dots < t_n$. Prove that $\mu_0 = \mu_1$. **Hint.** Use (iii) and Proposition 1.2.4

□

Martingales

The usefulness of the martingale property was recognized by P. Lévy (condition (C) in [134, Chap. VIII]), but it was J. L. Doob [61] who realized its full potential by discovering its most important properties: optional stopping/sampling, existence of asymptotic limits, maximal inequalities.

I have to admit that when I was first introduced to martingales they looked alien to me. Why would anyone be interested in such things? What are really these martingales?

I can easily answer the first question. Martingales are ubiquitous, they appear in the most unexpected of situations, though not always in an obvious way, and they are “well behaved”. Since their appearance on the probabilistic scene these stochastic processes have found many applications.

As for the true meaning of this concept let me first remark that the name “martingale” itself is a bit unusual. It is a French word that has an equestrian meaning (harness) but, according to [146], the term was used among the French gamblers when referring to a gambling system. I personally cannot communicate clearly, beyond a formal definition, what is the true meaning of this concept. I believe it is a fundamental concept of probability theory and I subscribe to R. Feynman’s attitude: it is more useful to know how the electromagnetic waves behave than knowing what they look like. The same could be said about the concept of martingale and, why not, about the concept of probability. I hope that the large selection of examples discussed in this chapter will give the reader a sense of this concept.

This chapter is divided into two parts. The first part is devoted to discrete time martingales. The second part is devoted to continuous time martingales. I have included many and varied applications of martingales with the hope that they will allow the reader to see the many facets of this concept and convince him/her of its power and versatility. My presentation was inspired by many sources and I want to single out [94, 43, 61, 68, 131, 132, 180, 224] that influenced me the most.

3.1. Basic facts about martingales

We need to introduce some basic terminology.

3.1.1. Definition and examples. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $\mathbb{T} \subset \mathbb{R}$. Recall that a *random or stochastic process with parameter space* $\mathbb{T}$ is a family of random variables

$$X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad t \in \mathbb{T}.$$

A $\mathbb{T}$ -*filtration* of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$ is a family $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \in \mathbb{T}}$ of sub- σ -algebras of $\mathcal{S}$ such that

$$\mathcal{F}_s \subset \mathcal{F}_t, \quad \forall s \leq t.$$

We set

$$\mathcal{F}_\infty := \bigvee_{t \in \mathbb{T}} \mathcal{F}_t.$$

A $\mathbb{T}$ -*filtered probability space* is a quadruplet $(\omega, \mathcal{S}, \mathcal{F}_\bullet, \mathbb{P})$, where $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $\mathcal{F}_\bullet$ is $\mathbb{T}$ -filtration of sigma-subalgebras of $\mathcal{S}$. A family of random variables

$$X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad t \in \mathbb{T},$$

is said to be *adapted to the filtration* $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \in \mathbb{T}}$, if X_t is $\mathcal{F}_t$ -measurable for any t .

Remark 3.1.1. If we think of a σ -algebra as encoding all the measurable information in a given random experiment, then we can think of a $\mathbb{T}$ -filtration as an increasing flow of information. For example, if $\mathbb{T} = \mathbb{N}_0$, and $(X_n)_{n \geq 0}$ is a sequence of random variables, then the collection

$$\mathcal{F}_n = \sigma(X_0, X_1, \dots, X_n), \quad n \in \mathbb{N}_0,$$

is a filtration of σ -algebras. At epoch n , the information about X_n becomes available to us, on top of the information about $X_0, X_1, \dots, X_{n-1}$ that we have collected along the way. $\square$

Definition 3.1.2. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ equipped with a filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \in \mathbb{T}}$. An $\mathcal{F}_\bullet$ -*martingale* is a family of random variables $X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, t \in \mathbb{T}$, satisfying the following two conditions.

- (i) The family is adapted to the filtration $\mathcal{F}_\bullet$ and X_t is integrable for any $t \in \mathbb{T}$.
- (ii) For all $s, t \in \mathbb{T}, s < t$, we have $\mathbb{E}[X_t \mid \mathcal{F}_s] = X_s$.

The family of *integrable* random variables $(X_t)_{t \in \mathbb{T}}$ is called a $\mathcal{F}_\bullet$ -*submartingale* (resp. *supermartingale*) if it is adapted to the filtration and for any $s, t \in \mathbb{T}, s < t$, we have $X_s \leq \mathbb{E}[X_t \mid \mathcal{F}_s]$ (resp. $X_s \geq \mathbb{E}[X_t \mid \mathcal{F}_s]$).

When $\mathbb{T}$ is a discrete subset of $\mathbb{R}$ we say that the (sub- or super-)martingale is a *discrete time* (sub/super)martingale. $\square$

Note that a sequence of integrable random variables $(X_n)_{n \in \mathbb{N}_0}$ is a discrete time submartingale (resp. martingale) with respect to a filtration $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$ of $\mathcal{F}$ if

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] \geq X_n, \quad (\text{resp. } \mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = X_n), \quad \forall n \in \mathbb{N}_0.$$

Note that if $(X_n)_{n \in \mathbb{N}_0}$ is a martingale with respect to a filtration $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$,

$$\mathbb{E}[X_{n+1}] = \mathbb{E}[\mathbb{E}[X_{n+1} \mid \mathcal{F}_n]] = \mathbb{E}[X_n], \quad \forall n \geq 0.$$

Remark 3.1.3. Suppose that $(X_n)_{n \geq 0}$ is a sequence of integrable random variables and

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n).$$

Then $\mathbb{E}[X_{n+1} \mid \mathcal{F}_n]$ is a measurable function of the variables $X_0, \dots, X_n$,

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = f_{n+1}(X_0, X_1, \dots, X_n), \quad f_{n+1} : \mathbb{R}^{n+1} \rightarrow \mathbb{R}.$$

If the joint distribution of $(X_0, \dots, X_n)$ is described by a density $p_n(x_0, \dots, x_n)$, then

$$f_{n+1}(x_0, \dots, x_n) = \int_{\mathbb{R}} x_{n+1} \frac{p_{n+1}(x_0, \dots, x_n, x_{n+1})}{p_n(x_0, \dots, x_n)} dx_{n+1}.$$

The sequence $(X_n)_{n \geq 0}$ is a martingale if and only if $f_{n+1}(x_0, x_1, \dots, x_n) = x_n, \forall n \geq 0, \forall x_0, \dots, x_n \in \mathbb{R}$. $\square$

Example 3.1.4 (Closed martingales). Suppose that $\mathcal{F}_\bullet = (\mathcal{F}_n)_{n \in \mathbb{N}_0}$ is a filtration of $\mathcal{S}$ and $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then the sequence of random variables

$$X_n = \mathbb{E}[X \mid \mathcal{F}_n] \in L^1(\Omega, \mathcal{F}_n, \mathbb{P}), \quad n \in \mathbb{N}_0,$$

is a martingale since

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = \mathbb{E}[\mathbb{E}[X \mid \mathcal{F}_{n+1}] \mid \mathcal{F}_n] = \mathbb{E}[X \mid \mathcal{F}_n] = X_n.$$

Such a martingale is called *closed* or *Doob martingale*. $\square$

Example 3.1.5 (Unbiased random walk). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent integrable random variables such that $\mathbb{E}[X_n] = 0, \forall n \in \mathbb{N}_0$.

One should think that X_n is the size of the n -th step so that the location after n steps is

$$S_n = X_1 + \dots + X_n.$$

Set $\mathcal{F}_n := \sigma(X_1, \dots, X_n), S_0 := 0$. Then the sequence $(S_n)_{n \in \mathbb{N}_0}$ is a martingale adapted to the filtration $\mathcal{F}_n$. Indeed,

$$\begin{aligned} \mathbb{E}[S_{n+1} \mid \mathcal{F}_n] &= \mathbb{E}[X_{n+1} \mid X_1, \dots, X_n] + \mathbb{E}[X_1 + \dots + X_n \mid X_1, \dots, X_n] \\ &= \mathbb{E}[X_{n+1}] + X_1 + \dots + X_n = S_n. \end{aligned} \quad \square$$

Example 3.1.6 (Random products). Suppose that $(Y_n)_{n \in \mathbb{N}}$ are positive i.i.d. random variables such that $\mathbb{E}[Y_1] = 1$. Then the sequence of products

$$Z_n = Y_1 Y_2 \dots Y_n, \quad n \in \mathbb{N},$$

is a martingale adapted to the filtration $\mathcal{F}_n = \sigma(Y_1, \dots, Y_n)$. Indeed

$$\begin{aligned} \mathbb{E}[Z_{n+1} \mid Y_1, \dots, Y_n] &= \mathbb{E}[Y_1 \dots Y_n Y_{n+1} \mid Y_1, \dots, Y_n] \\ &= \mathbb{E}[Y_{n+1}] Y_1 \dots Y_n = Z_n. \end{aligned} \quad \square$$

Example 3.1.7 (Biased random walk). Suppose that $(X_n)_{n \in \mathbb{N}}$ are i.i.d. random variables such that the moment generating function

$$\mu(\lambda) := \mathbb{E}[e^{\lambda X_n}]$$

is well defined for λ in some interval Λ . We set

$$S_n := X_1 + \dots + X_n, \quad M_n = M_n(\lambda) := e^{\lambda S_n} \mu(\lambda)^{-n},$$

$$\mathcal{F}_n := \sigma(X_1, \dots, X_n).$$

If we define

$$Y_n := \frac{1}{\mu(\lambda)} e^{\lambda X_n},$$

then we deduce that

$$\mathbb{E}[Y_n] = 1, \quad M_n = Y_1 \cdots Y_n.$$

From the previous example we deduce that $(M_n(\lambda))_{n \in \mathbb{N}}$ is a martingale.

As a concrete example, suppose that the random variables X_n are all binomial type

$$\mathbb{P}[X_n = 1] = p, \quad \mathbb{P}[X_n = -1] = q = 1 - p.$$

In this case $\mu(\lambda) = pe^\lambda + qe^{-\lambda}$. Note that if $e^\lambda = \frac{q}{p}$, then $\mu(\lambda) = 1$ and we deduce that

$$M_n = \left(\frac{q}{p}\right)^{S_n}$$

is a martingale. This is sometimes referred to as the *De Moivre's martingale*. $\square$

Example 3.1.8 (Galton-Watson/branching processes). Fix a probability measure μ on $\mathbb{N}_0$ such that

$$m := \sum_{k \in \mathbb{N}_0} k\mu[k] < \infty, \quad \mu[k] := \mu[\{k\}],$$

and $\mu[k_0] > 0$ for some $k_0 > 0$. Consider next a sequence $(X_{n,j})_{j,n \in \mathbb{N}_0}$ of i.i.d. $\mathbb{N}_0$ -valued random variables with common probability distribution μ . Fix $\ell \in \mathbb{N}$, set $Z_0 := \ell$, and for any $n \in \mathbb{N}_0$ define

$$Z_{n+1} = \sum_{j=1}^{Z_n} X_{n,j}, \quad \mathcal{F}_n = \sigma(X_{k,j}; k \in \mathbb{N}_0, k < n).$$

The random variable Z_n can be interpreted as the population of the n -th generation of a species that had ℓ individuals at $n = 0$ and such that the number of offsprings of a given individuals is a random variable with distribution μ . The j -th individual of the n -th generators has $X_{n,j}$ offsprings. We will refer to μ as the *reproduction law*.

The sequence $(Z_n)_{n \geq 0}$ is known as the *Galton-Watson process* or the *branching process* with reproduction law μ .




Figure 3.1. Three generations of a Galton-Watson (random) tree. Here $Z_1 = 3$, $Z_2 = 2 + 1 + 3 = 6$, $Z_3 = 3 + 2 + 1 + 2 + 3 = 11$.

When $\ell = 1$ this process can be visualized as a random rooted tree. The root v_0 has $Z_1 = X_{0,1}$ successor vertices. $v_{1,1}, \dots, v_{1,Z_1}$. The vertex $v_{1,i}$ has $X_{1,i}$ successors etc.; see Figure 3.1. For any $n \in \mathbb{N}_0$ we have

$$Z_{n+1} = \sum_{k=1}^{\infty} \left(\sum_{j=1}^k X_{n,j} \right) \mathbf{I}_{\{Z_n=k\}}$$

so

$$\begin{aligned}\mathbb{E}[Z_{n+1} \mid \mathcal{F}_n] &= \sum_{k=1}^{\infty} \mathbb{E}\left[\left(\sum_{j=1}^k X_{n,j}\right) \mathbf{I}_{\{Z_n=j\}} \mid \mathcal{F}_n\right] \\ &= \sum_{k=0}^{\infty} \mathbb{E}\left[\left(\sum_{j=1}^k X_{n,j}\right) \mid \mathcal{F}_n\right] \mathbf{I}_{\{Z_n=k\}}\end{aligned}$$

$(X_{n,j} \perp \mathcal{F}_n, \forall n, j)$

$$= \sum_{k=1}^{\infty} \underbrace{\left(\sum_{j=1}^k \mathbb{E}[X_{n,j}]\right)}_{=km} \mathbf{I}_{\{Z_n=k\}} = m \sum_{k=0}^{\infty} k \mathbf{I}_{\{Z_n=k\}} = mZ_n.$$

This proves that the sequence $Y_n = m^{-n}Z_n$, $n \in \mathbb{N}_0$ defines a martingale.

The intuition behind the above algebraic manipulation can be easily explained: if on average an individual of this species has m successors, and the n -th generation consists of Z_n individuals, we expect that the population of the next generation to change by a factor of m , $\mathbb{E}[Z_{n+1} \mid Z_n] = mZ_n$. $\square$

Example 3.1.9 (Polya's urn). An urn contains $r > 0$ red balls and $g > 0$ green balls. At each moment of time we draw a ball uniformly likely from the balls existing at that moment, we replace it by $c + 1$ balls of the same color, $c \geq 0$. Denote by R_n and G_n the number of red and respectively green balls in the urn after the n -th draw. Note that $R_n + G_n = r + g + cn$. We denote by X_n the ratio of red balls after n draws, i.e.,

$$X_n := \frac{R_n}{R_n + G_n} = \frac{R_n}{r + g + cn}.$$

Note that when $c = 1$, the scheme can be alternatively described as randomly adding at each moment of time a red/green ball with probability equal to the fraction of red/green balls that exist at that moment in the urn.

We set

$$\mathcal{F}_n = \sigma(R_0, G_0, \dots, R_n, G_n) = \sigma(X_0, X_1, \dots, X_n).$$

We will show that (X_n) is an $\mathcal{F}_n$ -martingale. To see this observe that

$$X_n = \sum_{i,j \geq 0} \frac{i}{i+j} \mathbf{I}_{\{R_n=i, G_n=j\}}$$

so

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = \sum_{i,j \geq 0} \frac{i}{i+j} \mathbb{E}[\mathbf{I}_{\{R_{n+1}=i, G_{n+1}=j\}} \mid \mathcal{F}_n].$$

Now observe that

$$\begin{aligned}& \mathbb{E}[\mathbf{I}_{\{R_{n+1}=i, G_{n+1}=j\}} \mid \mathcal{F}_n] \\ &= \sum_{k, \ell \geq 0} \mathbb{P}[R_{n+1} = i, G_{n+1} = j \mid R_n = k, G_n = \ell] \mathbf{I}_{\{R_n=k, G_n=\ell\}} \\ &= \frac{i-c}{i+j-c} \mathbf{I}_{\{R_n=i-c, G_n=j-c\}} + \frac{j-c}{i+j-c} \mathbf{I}_{\{R_n=i, G_n=j-c\}}\end{aligned}$$

We deduce

$$\begin{aligned}
\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] &= \sum_{i,j} \frac{i}{i+j} \cdot \frac{i-c}{i+j-c} \mathbf{I}_{\{R_n=i-c, G_n=j\}} \\
&\quad + \sum_{i,j} \frac{i}{i+j} \cdot \frac{j-c}{i+j-c} \mathbf{I}_{\{R_n=i, G_n=j-c\}} \\
&= \sum_{u,v} \frac{u+c}{u+v+c} \cdot \frac{u}{u+v} \mathbf{I}_{\{R_n=u, G_n=v\}} + \sum_{u,v} \frac{u}{u+v+c} \cdot \frac{v}{u+v} \mathbf{I}_{\{R_n=u, G_n=v\}} \\
&= \sum_{u,v} \frac{u(u+v+c)}{(u+v)(u+v+c)} \mathbf{I}_{\{R_n=u, G_n=v\}} = \sum_{u,v} \frac{u}{u+v} \mathbf{I}_{\{R_n=u, G_n=v\}} = X_n. \quad \square
\end{aligned}$$

Example 3.1.10 (Random walks on graphs). Suppose that Γ is a connected simple graph with vertex set V_Γ and edges E_Γ . Assume that there are no multiple edges between two vertices $u, v \in V_\Gamma$. Assume that Γ is *locally finite* i.e., for any vertex $u \in V_\Gamma$, its set of neighbors $\mathcal{N}(u)$ is finite. We set $\deg(u) := |\mathcal{N}(u)|$.

A function $F : V_\Gamma \rightarrow \mathbb{R}$ is called *harmonic* if

$$F(u) = \frac{1}{\deg(u)} \sum_{v \in \mathcal{N}(u)} F(v).$$

Consider the simple random walk on Γ that starts at a given vertex v_0 and the probability of transitioning from a vertex u to a neighbor v is equal to $\frac{1}{\deg(u)}$. Denote by V_n the location after n steps of the walk. Suppose that $F : V_\Gamma \rightarrow \mathbb{R}$ is a harmonic function. Then the sequence of random variables

$$X_n = F(V_n), \quad n \in \mathbb{N}_0,$$

is a martingale with respect to the filtration $\mathcal{F}_n = \sigma(V_0, V_1, \dots, V_n)$. Moreover

$$\mathbb{E}[X_n] = F(v_0), \quad \forall n \in \mathbb{N}_0. \quad \square$$

Example 3.1.11 (New (sub)martingales from old). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is equipped with a filtration $\mathcal{F}_\bullet = (\mathcal{F}_n)_{n \in \mathbb{N}_0}$ and $X_n \in \mathcal{L}^1(\Omega, \mathcal{F}_n, \mathbb{P})$ is a sequence of random variables adapted to the above filtration.

- (i) If $(X_n)_{n \in \mathbb{N}_0}$ is a martingale and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function such that $\varphi(X_n)$ is integrable $\forall n \in \mathbb{N}_0$, then the conditional Jensen inequality implies that the sequence $\varphi(X_n)$ is a submartingale. Indeed, Jensen's inequality implies

$$\mathbb{E}[\varphi(X_{n+1}) \mid \mathcal{F}_n] \geq \varphi\left(\mathbb{E}[X_{n+1} \mid \mathcal{F}_n]\right) = \varphi(X_n).$$

- (ii) If $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a *nondecreasing* convex function such that $\varphi(X_n)$ is integrable $\forall n \in \mathbb{N}_0$, then the sequence $\varphi(X_n)$ is a submartingale. Indeed, follow the same argument as above where at the last step use the fact that φ is nondecreasing. In particular if $(X_n)_{n \geq 0}$ is a submartingale, then so is $(X_n^+)_{n \geq 0}$, $x^+ = \max(0, x)$.
- (iii) If $(X_n)_{n \in \mathbb{N}_0}$ is a supermartingale and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a nondecreasing concave function such that $\varphi(X_n)$ is integrable $\forall n \in \mathbb{N}_0$, then the sequence $\varphi(X_n)$ is a supermartingale. Indeed

$$\mathbb{E}[\varphi(X_{n+1}) \mid \mathcal{F}_n] \leq \varphi\left(\mathbb{E}[X_{n+1} \mid \mathcal{F}_n]\right) \leq \varphi(X_n).$$

In particular, if $(X_n)_{n \in \mathbb{N}_0}$ is a supermartingale, then so is $(\min(X_n, c))_{n \geq 0}$, $\forall c \in \mathbb{R}$.

□

3.1.2. Doob decomposition. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and an $\mathbb{N}_0$ -filtration $\mathcal{F}_\bullet$ of $\mathcal{S}$. If $C_\bullet$ is an increasing $\mathcal{F}_\bullet$ -adapted process, then obviously $C_\bullet$ is a submartingale. If we add to this process a martingale $M_\bullet$, then the resulting process $X_\bullet = M_\bullet + C_\bullet$ is a submartingale.

It turns out that all submartingales can be obtained in this fashion. In fact, the increasing process C_n can be chosen to be of a special type: the random variable C_{n+1} can be chosen to be $\mathcal{F}_n$ -measurable, i.e., the value of $C_\bullet$ at time $n+1$ is predictable at time n , i.e., can be determined from the information available to us at time n encoded in the σ -algebra $\mathcal{F}_n$.

Definition 3.1.12. A sequence of random variable $\{H_n : \Omega \rightarrow \mathbb{R}, n \in \mathbb{N}_0\}$ is called $\mathcal{F}_\bullet$ -previsible or predictable if H_0 is $\mathcal{F}_0$ -measurable, and H_n is $\mathcal{F}_{n-1}$ -measurable $\forall n \in \mathbb{N}$. $\square$

Note that if $\mathcal{F}_\bullet$ is the filtration determined by a stochastic process $(X_n)_{n \geq 0}$, $X_0 = \text{const}$,

$$; \mathcal{F}_n := \sigma(X_0, X_1, \dots, X_n), \quad \forall n \geq 0,$$

then the process $(H_n)_{n \in \mathbb{N}}$ is $\mathcal{F}_\bullet$ -predictable if there exist measurable functions $f_n : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$H_{n+1} = f_n(X_0, X_1, \dots, X_n), \quad ; \forall n \geq 0.$$

Thus, with the information available to us at moment $T = n$ we can predict the future value H_{n+1} .

The next result formalizes the discussion at the beginning of this subsection.

Proposition 3.1.13 (Doob decomposition of discrete submartingales). *Let $X_\bullet = (X_n)_{n \in \mathbb{N}_0}$ be an $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$ -adapted process such that $X_n \in L^1, \forall n \in \mathbb{N}_0$. Then the following statements are equivalent.*

- (i) *The process $X_\bullet$ is a submartingale.*
- (ii) *There exists an $\mathcal{F}_\bullet$ -martingale $M_\bullet$ and an $\mathcal{F}_\bullet$ -predictable nondecreasing process $C_\bullet$ such that*

$$M_0 = 0 = C_{\text{cpt}}, \quad X_n = X_0 + M_n + C_n, \quad \forall n \geq 0.$$

Moreover, when $X_\bullet$ is a submartingale, then the martingale $M_\bullet$ and the nondecreasing predictable process are uniquely determined by $X_\bullet$ up to indistinguishability; see Definition 2.5.11(ii). In this case $M_\bullet$ is called the martingale component of the submartingale $X_\bullet$ and $C_\bullet$ is called the compensator of $X_\bullet$. We denote it by $C(X_\bullet)$. The decomposition $X_n = X_0 + M_n + C_n$ is called the Doob decomposition of the submartingale $X_\bullet$.

Proof. Existence. We describe M_n and C_n in terms of their increments. More precisely

$$\begin{aligned} C_{n+1} - C_n &= \mathbb{E}[X_{n+1} - X_n \mid \mathcal{F}_n] - \mathbb{E}[M_{n+1} - M_n \mid \mathcal{F}_n] \\ &= \mathbb{E}[X_{n+1} \mid \mathcal{F}_n] - X_n = \mathbb{E}[(X_{n+1} - X_n) \mid \mathcal{F}_n], \quad \forall n \in \mathbb{N}_0, \end{aligned} \tag{3.1.1a}$$

$$M_{n+1} - M_n = (X_{n+1} - X_n) - (C_{n+1} - C_n), \quad \forall n \in \mathbb{N}_0. \tag{3.1.1b}$$

Note that $C_{n+1} - C_n$ is $\mathcal{F}_n$ measurable so (C_n) is predictable. By construction $M_\bullet$ is an $\mathcal{F}_\bullet$ -martingale. Clearly, if $X_\bullet$ is a submartingale then, tautologically, C_n is increasing.

Uniqueness. Suppose that $X_\bullet$ is a submartingale, $M'_\bullet$ is a martingale, and $C_\bullet$ is a nondecreasing predictable process such that

$$M_0 = C_{\text{cpt}} = 0, \quad X_n = X_0 + M'_n + C'_n, \quad \forall n \in \mathbb{N}_0.$$

We deduce

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] - X_n = \underbrace{\mathbb{E}[M'_{n+1} \mid \mathcal{F}_n] - M'_n}_{=0} + \underbrace{\mathbb{E}[C'_{n+1} \mid \mathcal{F}_n] - C'_n}_{=C'_{n+1}-C'_n}.$$

This shows that the increments of C'_n are given by (3.1.1a) so $C'_n = C_n$. In particular, $M'_n = M_n$, $\forall n \in \mathbb{N}$. $\square$

Example 3.1.14. Suppose that $(Y_n)_{n \geq 0}$ adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$. It has a Doob decomposition $Y_n = M_n + C_n$. We deduce from (3.1.1a) that

$$C_n = \sum_{k=0}^{n-1} \mathbb{E}[\Delta Y_k \mid \mathcal{F}_k], \quad \Delta Y_k = Y_{k+1} - Y_k.$$

Suppose that $(X_n)_{n \geq 0}$ is a sequence of independent integrable random variables such that $X_0 = 0$ and $\mathbb{E}[X_n] \geq 0, \forall n$. Then

$$S_n = X_1 + \cdots + X_n$$

is a submartingale with respect to the filtration $\mathcal{F}_n = \sigma(X_0, X_1, \dots, X_n), n \geq 0$. Indeed, for $n \geq 1$

$$\mathbb{E}[S_n \mid \mathcal{F}_{n-1}] = \mathbb{E}[X_n \mid \mathcal{F}_{n-1}] + S_{n-1} \geq S_{n-1}.$$

Consider the Doob decomposition $S_n = M_n + C_n$. The compensator C_n satisfies

$$C_n = \sum_{k=1}^n \mathbb{E}[X_k \mid \mathcal{F}_{k-1}] = \sum_{k=1}^n \mathbb{E}[X_k]$$

and

$$M_n = S_n - \sum_{k=1}^n \mathbb{E}[X_k] = \sum_{k=1}^n (X_k - \mathbb{E}[X_k]).$$

If the variables X_n are independent, then

$$M_n = \sum_{k=1}^n (X_k - \mathbb{E}[X_k]).$$

$\square$

Definition 3.1.15 (Quadratic variation). Suppose that $(M_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ such that $\mathbb{E}[M_n^2] < \infty, \forall n \geq 0$. The compensator of the submartingale $(M_n^2)_{n \geq 0}$ is called the *quadratic variation* and it is denoted by $\langle M_\bullet \rangle$. $\square$

Example 3.1.16. Suppose that $(M_n)_{n \geq 0}$ is an L^2 -martingale, $M_0 = 0$. Then $X_n = M_n^2$ is a submartingale. The compensator Q_n of X_n is found using the difference equation (3.1.1a)

$$\begin{aligned} Q_{k+1} - Q_k &= \mathbb{E}[X_{k+1} - X_k \mid \mathcal{F}_k] = \mathbb{E}[M_{k+1}^2 - M_k^2 \mid \mathcal{F}_k] \\ &= \mathbb{E}[(M_{k+1} - M_k)^2 \mid \mathcal{F}_k] + 2\mathbb{E}[M_k(X_{k+1} - M_k) \mid \mathcal{F}_k] \\ &= \mathbb{E}[(M_{k+1} - M_k)^2 \mid \mathcal{F}_k]. \end{aligned}$$

Hence

$$\langle M \rangle_n = Q_n = \sum_{k=1}^n \mathbb{E}[(M_k - M_{k-1})^2 \mid \mathcal{F}_{k-1}].$$

This explains why we refer to $\langle M_\bullet \rangle$ as quadratic *quadratic variation*. Note that

$$\mathbb{E}[M_{k+1}^2 - M_k^2] = \mathbb{E}[(M_{k+1} - M_k)^2]$$

and we deduce that

$$\forall n \in \mathbb{N}, \quad \mathbb{E}[M_n^2] = \sum_{k=1}^n \mathbb{E}[(M_{k+1} - M_k)^2]. \quad (3.1.2)$$

□

3.1.3. Discrete stochastic integrals. A very important method of producing a large supply of martingales *discrete stochastic integration*.

Theorem 3.1.17 (Discrete Stochastic Integral). *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ be an $\mathcal{F}_\bullet$ -adapted process and $(H_n)_{n \in \mathbb{N}}$ is a bounded predictable process. Define the process $(H \cdot X)_\bullet$ by setting*

$$(H \cdot X)_0 := 0, \quad (H \cdot X)_n = H_1(X_1 - X_0) + \cdots + H_n(X_n - X_{n-1}), \quad \forall n \in \mathbb{N}. \quad (3.1.3)$$

Then the following hold.

- (i) *If $(X_n)_{n \in \mathbb{N}_0}$ is a martingale, then the process $(H \cdot X)_n$, $n \in \mathbb{N}_0$ is also an $\mathcal{F}_\bullet$ -adapted martingale.*
- (ii) *If $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale and $H_n \geq 0$, $\forall n \in \mathbb{N}$, then the process $(H \cdot X)_n$, $n \in \mathbb{N}_0$ is also an $\mathcal{F}_\bullet$ -adapted submartingale.*

Proof. (i) Clearly $(H \cdot X)_n \in L^1(\Omega, \mathcal{F}_n)$. We have

$$\mathbb{E}[(H \cdot X)_{n+1} | \mathcal{F}_n] = \mathbb{E}[H_{n+1}(X_{n+1} - X_n) | \mathcal{F}_n] + (H \cdot X)_n$$

(H_{n+1} is $\mathcal{F}_n$ -measurable)

$$= H_{n+1} \mathbb{E}[X_{n+1} - X_n | \mathcal{F}_n] + (H \cdot X)_n$$

$$= H_{n+1} (\mathbb{E}[X_{n+1} | \mathcal{F}_n] - X_n) + (H \cdot X)_n$$

((X_n) is a martingale)

$$= (H \cdot X)_n.$$

The proof of (ii) is similar. □

Remark 3.1.18. (a) When $X_\bullet$ is a martingale the process $(H \cdot X)_\bullet$ is called the *discrete stochastic integral* of H with respect to X and it is alternatively denoted

$$\int_0^n H dX := (H \cdot X)_n.$$

One should think of X_n as a random signed measure assigning mass $X_n - X_{n-1}$ to the point n .

(b) The discrete stochastic integral has a stock-trading interpretation. Suppose that X_n represents the price of a stock *at the end* of the n -th trading day. A day trader buys H_n shares *at the beginning* of the n -th trading day, based on the information collected during the previous $(n-1)$ trading days. This information is encoded by the sigma-algebra $\mathcal{F}_{n-1}$ and the price of a share at the beginning of the n -th trading day is X_{n-1} . He sells these H_n shares at the end of the n -th trading day. The resulting profit at the end of day n is then $H_n(X_n - X_{n-1})$. We deduce that $(H \bullet X)_n$ represents the profit of the day trader after n trading days.

(c) The special case Theorem 3.1.17 when the variables H_n are Bernoulli random variables was discovered by P. Halmos and is classically known as the *impossibility of systems* theorem. In this case H_n represents the decision of a gambler to play or not the next game based on the information gathered during the games he observed so far. $\square$

The applicability of Theorem 3.1.17 depends on our ability of producing interesting predictable processes. We describe one very useful class of examples.

Example 3.1.19. Observe first that a discrete time process $(Y_n)_{n \in \mathbb{N}}$ on $(\Omega, \mathcal{S}, \mathbb{P})$ can be viewed as a map

$$Y : \mathbb{N} \times \Omega \rightarrow \mathbb{R}, \quad (n, \omega) \mapsto Y_n(\omega).$$

We equip $\mathbb{N} \times \Omega$ with the product σ -algebra. A measurable set $\mathcal{X} \subset \mathbb{N} \times \Omega$ defines a stochastic process

$$\mathbf{I}_{\mathcal{X}} : \mathbb{N} \times \Omega \rightarrow \{0, 1\}, \quad (\mathbf{I}_{\mathcal{X}})_n = \mathbf{I}_{\mathcal{X}_n}, \quad \mathcal{X}_n := \{ \omega \in \Omega; (n, \omega) \in \mathcal{X} \}.$$

The set $\mathcal{X}$ is called $\mathcal{F}_{\bullet}$ -predictable if the process $\mathbf{I}_{\mathcal{X}}$ is such. More precisely, this means that $\mathcal{X}_0 \in \mathcal{F}_0$ and, for any $n \in \mathbb{N}$, the set $\mathcal{X}_n$ is $\mathcal{F}_{n-1}$ -measurable. $\square$

3.1.4. Stopping and sampling: discrete time. We want to describe one technique that makes the martingales extremely useful in applications. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$.

Definition 3.1.20. A random variable $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{N}_0 \cup \{\infty\}$ is called a *stopping time* adapted to the filtration $\mathcal{F}_{\bullet} = (\mathcal{F}_n)_{n \geq 0}$, or an $\mathcal{F}_{\bullet}$ -stopping time if,

$$\{T \leq n\} \in \mathcal{F}_n, \quad \forall n \in \mathbb{N}_0 \cup \{\infty\}.$$

If $(X_n)_{n \in \mathbb{N}}$ is an $\mathcal{F}_{\bullet}$ -adapted process, and T is an $\mathcal{F}_{\bullet}$ -stopping time, then the T -sample of the process is the random variable

$$X_T := \sum_{n \in \mathbb{N}_0} X_n \mathbf{I}_{\{T=n\}}. \quad (3.1.4)$$

Observe that $X_T = 0$ on the set $\{T = \infty\}$. $\square$

Example 3.1.21. (a) For each $n \in \mathbb{N}_0$ the constant random variable equal to n is a stopping time.

(b) Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is $\mathcal{F}_{\bullet}$ -adapted and $C \subset \mathbb{R}$ is a Borel set. We define the *hitting time* of C to be the random variable

$$H_C : \Omega \rightarrow \mathbb{N}_0 \cup \{\infty\}, \quad H_C(\omega) := \min\{n \in \mathbb{N}_0; X_n(\omega) \in C\}.$$

This is a stopping time since the process (X_n) is $\mathcal{F}_{\bullet}$ -adapted and

$$\{H_C \leq n\} = \bigcup_{k \leq n} \{X_k \in C\}.$$

(c) If S, T are stopping times, then $S \wedge T = \min(S, T)$ and $S \vee T = \max(S, T)$ are also stopping times.

(d) If $(T_k)_{k \in \mathbb{N}}$ is a sequence of stopping times, then $\inf T_k$, $\sup T_k$, $\liminf T_k$ and $\limsup T_k$ are also stopping times. $\square$

Definition 3.1.22. Let $X_\bullet = (X_n)_{n \in \mathbb{N}}$ be a process adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$. For any stopping time T we denote by $X_\bullet^T$ the process stopped at T defined by

$$\begin{aligned} X_n^T &:= X_{T \wedge n}, \text{ where} \\ X_{T \wedge n} &= X_{\min(T(\omega), n)}(\omega) = \begin{cases} X_n(\omega), & n \leq T(\omega), \\ X_{T(\omega)}, & n > T(\omega). \end{cases} \end{aligned} \quad (3.1.5)$$

□

Note that the process stopped at T is also adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$.

Proposition 3.1.23. Suppose that S, T are stopping times such that $S \leq T$. Define

$$]]T, \infty[[:= \{ (n, \omega) \in \mathbb{N}_0 \times \Omega; T(\omega) < n \},$$

$$]]S, T]] := \{ (n, \omega) \in \mathbb{N}_0 \times \Omega; S(\omega) < n \leq T(\omega) \}.$$

Then $]]T, \infty[[$, $[[0, T]]$ and $]]S, T]]$ are predictable subsets of $\mathbb{N}_0 \times \Omega$.

Proof. We have $]]T, \infty[[_n = \{T < n\} = \{T \leq n-1\} \in \mathcal{F}_{n-1}$. Next observe that

$$I_{[[0, T]]} = 1 - I_{]]T, \infty[[}$$

so $I_{[[0, T]]}$ is a predictable process as a linear combination of predictable processes. Finally observe that since $S \leq T$ we have

$$I_{]]S, T]]} = I_{[[0, T]]} - I_{[[0, S]]},$$

so $I_{]]S, T]]}$ is predictable as a linear combination of predictable processes. Note that

$$(I_{]]S, T]]})_n = I_{\{n \leq T\}} - I_{\{S < n\}}.$$

□

Suppose now that $(X_n)_{n \in \mathbb{N}}$ is a $\mathcal{F}_\bullet$ -(sub)martingale and T is a stopping time. Then $S_0 = 0$ is also a stopping time, $S_0 \leq T$. Theorem 3.1.17 shows that the process $I_{]]S_0, T]]} = I_{[[0, T]]} \cdot X$ is also an $\mathcal{F}_\bullet$ -submartingale.

For every $n \in \mathbb{N}$ we have

$$\begin{aligned} (I_{[[0, T]]} \cdot X)_n &= (I_{[[0, T]]})_n (X_n - X_{n-1}) + \cdots + (I_{[[0, T]]})_1 (X_1 - X_0) \\ &= (I_{\{T \geq n\}})(X_n - X_{n-1}) + \cdots + (I_{\{T \geq 1\}})(X_1 - X_0) \\ &= X_{T \wedge n} - X_0 = X_n^T - X_0. \end{aligned}$$

Thus

$$\boxed{X_\bullet^T = X_0 + (I_{[[0, T]]} \cdot X)_\bullet}. \quad (3.1.6)$$

This proves the following result.

Theorem 3.1.24 (Optional Stopping Theorem). Suppose that $(X_n)_{n \geq 0}$ is a (sub)martingale adapted to the filtration $\mathcal{F}_\bullet$ and T is an $\mathcal{F}_\bullet$ -stopping time. Then $X_\bullet^T$, the process stopped at T , is also a (sub)martingale adapted to $\mathcal{F}_\bullet$. □

Remark 3.1.25. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale adapted to the filtration $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$ and T is an a.s. finite stopping time. Then

$$\lim_{n \rightarrow \infty} X_n^T = \lim_{n \rightarrow \infty} X_{n \wedge T} = X_T \text{ a.s.}$$

since $n \wedge T(\omega)$ for $n \geq T(\omega)$. If $\sup_n \mathbb{E}[|X_n|] < \infty$, then Fatou's lemma implies $X_T \in L^1$ and

$$\mathbb{E}[X_T] \leq \lim_{n \rightarrow \infty} \mathbb{E}[X_n].$$

If (X_n) is a martingale, then $\mathbb{E}[X_T] \leq \mathbb{E}[X_0]$. □

Suppose that $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{N}_0 \cup \{\infty\}$ is a stopping time adapted to the filtration $\mathcal{F}_\bullet$. We define

$$\begin{aligned} \mathcal{F}_T &:= \left\{ E \in \mathcal{F} : E \cap \{T \leq n\} \in \mathcal{F}_n, \forall n \in \mathbb{N}_0 \cup \{\infty\} \right\} \\ &= \left\{ E \in \mathcal{F} : E \cap \{T = n\} \in \mathcal{F}_n, \forall n \in \mathbb{N}_0 \cup \{\infty\} \right\}. \end{aligned} \quad (3.1.7)$$

Tautologically, the random variable T is $\mathcal{F}_T$ -measurable.

Example 3.1.26. Suppose that T is the hitting time of a Borel set $C \subset \mathbb{R}$. Then the event E belongs to $\mathcal{F}_T$ if, at any moment of time n , we can decide using the information $\mathcal{F}_n$ available to us at time n whether, up to that moment, we have visited C and the event E has occurred. □

A few remarks are in order.

- The collection $\mathcal{F}_T$ is a σ -subalgebra of $\mathcal{F}$. It is called the *past-until- T* σ -algebra.
- The random variable X_T is $\mathcal{F}_T$ -measurable. Indeed,

$$\{X_T \leq c\} \cap \{T = n\} = \{X_n \leq c\} \cap \{T = n\} \in \mathcal{F}_n, \quad \forall n.$$

- If S, T are stopping times such that $S \leq T$, then $\mathcal{F}_S \subset \mathcal{F}_T$.

Definition 3.1.27. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an $\mathcal{F}_\bullet$ -(sub)martingale and T is an $\mathcal{F}_\bullet$ -stopping time. We say that the stopping time T is a *Doob stopping time rel* (X_n) if it satisfies the *Doob conditions*¹ if the following hold.

$$\mathbb{P}[T < \infty] = 1. \quad (3.1.8a)$$

$$X_T \in L^1. \quad (3.1.8b)$$

$$\lim_{n \rightarrow \infty} \mathbb{E}[\mathbf{I}_{\{T > n\}} | X_n|] = 0. \quad (3.1.8c)$$

□

Roughly speaking, the condition (3.1.8c) states that the random process $(X_n)_{n \geq 0}$ is not sampled “too late” at time T . We want to emphasize that the Doob conditions are constraints of a *pair* (sub)martingale, stopping time) and not just of the stopping time alone. In Proposition 3.2.30 we provide another characterization of the Doob conditions in terms of the asymptotic behavior of the stopped process $X_\bullet^T$.

¹There is no consensus on terminology in the literature. We use the term *Doob conditions* since they were first spelled out by J.L. Doob in his influential monograph [61]

Example 3.1.28. Suppose that T is a bounded $\mathcal{F}_\bullet$ -stopping time. Then T satisfies the Doob conditions.

To see this, choose $N \in \mathbb{N}$ such that $T < N$ a.s.. The stopped process $X_\bullet^T$ is a submartingale so $X_T = X_{T \wedge N} = X_N^T \in L^1$. As for the second condition (3.1.8c), note that since T is a.s. bounded, then $\mathbf{I}_{\{T > n\}}|X_n|$ is a.s. 0 for $n > N$. $\square$

Theorem 3.1.29 (Optional Sampling Theorem). *Suppose that*

$$X_n : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow \mathbb{R}, \quad n \geq 0,$$

is a (sub)martingale adapted to the filtration $\mathcal{F}_\bullet$, and $S \leq T$ are stopping times adapted to the same filtration. If T satisfies Doob's conditions in Definition 3.1.27, then

$$\mathbb{E}[X_T \mid \mathcal{F}_S] \geq X_S$$

If $X_\bullet$ is a martingale, then

$$\mathbb{E}[X_T \mid \mathcal{F}_S] = X_S, \quad \mathbb{E}[X_T] = \mathbb{E}[X_S] = \mathbb{E}[X_0].$$

Proof. We follow the original approach in [61, VII.2]; see also [6, Thm. 6.7.4]. Suppose first that $(X_n)_{n \geq 0}$ is a martingale. We have to show that for any $A \in \mathcal{F}_S$ we have

$$\mathbb{E}[X_T \mathbf{I}_A] = \mathbb{E}[X_S \mathbf{I}_A].$$

Let $A \in \mathcal{F}_S$ and set $A_m := A \cap \{S = m\}$. Then

$$\mathbb{E}[X_S] = \sum_{m \geq 0} \mathbb{E}[X_S \mathbf{I}_{A_m}]$$

so it suffices to show that

$$\forall m \geq 0 : \quad \mathbb{E}[X_T \mathbf{I}_{A_m}] = \mathbb{E}[X_S \mathbf{I}_{A_m}].$$

We have

$$\begin{aligned} \mathbb{E}[X_S \mathbf{I}_{A_m}] &= \mathbb{E}[X_m \mathbf{I}_{\{T=m\}} \mathbf{I}_{A_m}] + \mathbb{E}[X_S \mathbf{I}_{A_m} \mathbf{I}_{\{T>m\}}] \\ &= \mathbb{E}[X_T \mathbf{I}_{\{T=m\}} \mathbf{I}_{A_m}] + \boxed{\mathbb{E}[X_m \mathbf{I}_{A_m} \mathbf{I}_{\{T>m\}}]}_* \end{aligned}$$

$(A_m \cap \{T > m\} \in \mathcal{F}_m, X_\bullet \text{ martingale})$

$$\begin{aligned} &= \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{T=m\}}] + \boxed{\mathbb{E}[X_{m+1} \mathbf{I}_{A_m} \mathbf{I}_{\{T>m\}}]}_* \\ &= \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{T=m\}}] + \mathbb{E}[X_{m+1} \mathbf{I}_{A_m} \mathbf{I}_{\{T=m+1\}}] + \mathbb{E}[X_{m+1} \mathbf{I}_{A_m} \mathbf{I}_{\{T>m+1\}}] \\ &= \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{m \leq T \leq m+1\}}] + \boxed{\mathbb{E}[X_{m+1} \mathbf{I}_{A_m} \mathbf{I}_{\{T>m+1\}}]}_\bullet \end{aligned}$$

$(A_m \cap \{T > m+1\} \in \mathcal{F}_{m+1}, X_\bullet \text{ martingale})$

$$\begin{aligned} &= \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{m \leq T \leq m+1\}}] + \boxed{\mathbb{E}[X_{m+2} \mathbf{I}_{A_m} \mathbf{I}_{\{T>m+1\}}]}_\bullet \\ &= \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{m \leq T \leq m+2\}}] + \mathbb{E}[X_{m+2} \mathbf{I}_{A_m} \mathbf{I}_{\{T>m+2\}}]. \end{aligned}$$

Iterating this procedure we deduce that that, $\forall n > 0$, we have

$$\mathbb{E}[X_S \mathbf{I}_{A_m}] = \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{m \leq T \leq m+n\}}] + \mathbb{E}[X_{m+n} \mathbf{I}_{\{T>m+n\}}].$$

The condition (3.1.8c) shows that

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_{m+n} \mathbf{I}_{\{T>m+n\}}] = 0,$$

so

$$\mathbb{E}[X_S \mathbf{I}_{A_m}] = \mathbb{E}[X_T \mathbf{I}_{A_m} \mathbf{I}_{\{0 \leq T \leq \infty\}}] = \mathbb{E}[X_T \mathbf{I}_{A_m}].$$

The submartingale situation is dealt with similarly. $\square$

Remark 3.1.30. Suppose that T is an a.s. finite $\mathcal{F}_\bullet$ -stopping time and $X_\bullet$ is an $\mathcal{F}_\bullet$ -submartingale such that $X_T \in L^1$. Then T satisfies Doob's conditions if and only if

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^+ \mathbf{I}_{\{T > n\}}] = 0. \quad (3.1.9)$$

Clearly (3.1.8c) implies (3.1.9). Let us show that (3.1.9) $\Rightarrow$ (3.1.8c). Assume first that $X_\bullet$ is a martingale.

Fix $m, n \in \mathbb{N}_0$, $m < n$. Observing that $\{T > m\} \in \mathcal{F}_m$ we deduce

$$\begin{aligned} \mathbb{E}[X_m \mathbf{I}_{T > n}] &= \mathbb{E}[X_{m+1} \mathbf{I}_{T > m}] = \mathbb{E}[X_{m+1} \mathbf{I}_{T=m+1}] + \mathbb{E}[X_{m+1} \mathbf{I}_{T > m+1}] \\ (\{T > m+1\} \in \mathcal{F}_{m+1}) &= \mathbb{E}[X_{m+1} \mathbf{I}_{T=m+1}] + \mathbb{E}[X_{m+2} \mathbf{I}_{T > m+1}] \\ &= \mathbb{E}[X_{m+1} \mathbf{I}_{T=m+1}] + \mathbb{E}[X_{m+2} \mathbf{I}_{T=m+2}] + \mathbb{E}[X_{m+2} \mathbf{I}_{T > m+2}] \\ &= \cdots = \mathbb{E}[X_{m+1} \mathbf{I}_{T=m+1}] + \cdots + \mathbb{E}[X_n \mathbf{I}_{T=n}] + \mathbb{E}[X_n \mathbf{I}_{T > n}] = \mathbb{E}[X_T \mathbf{I}_{m < T \leq n}]. \end{aligned}$$

We deduce

$$\mathbb{E}[X_m \mathbf{I}_{T > n}] - \mathbb{E}[X_n \mathbf{I}_{T > n}] = \mathbb{E}[X_T \mathbf{I}_{m < T \leq n}], \quad \forall n > m.$$

Using the equality $X_\bullet = X_\bullet^+ - X_\bullet^-$ we deduce that, $\forall n > m$.

$$\mathbb{E}[X_n^- \mathbf{I}_{T > n}] - \mathbb{E}[X_m^- \mathbf{I}_{T > m}] = \mathbb{E}[X_T \mathbf{I}_{m < T \leq n}] - (\mathbb{E}[X_m^+ \mathbf{I}_{T > m}] - \mathbb{E}[X_n^+ \mathbf{I}_{T > n}]).$$

If we let $n \rightarrow \infty$ in the above equality and recall that $T < \infty$ a.s., $X_T \in L^1$ and $X_\bullet^+$ satisfies (3.1.9) we deduce

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^- \mathbf{I}_{T > n}] - \mathbb{E}[X_m^- \mathbf{I}_{T > m}] = \mathbb{E}[X_T \mathbf{I}_{T > m}] - \mathbb{E}[X_m^+ \mathbf{I}_{T > m}].$$

Using the Optional Sampling Theorem 3.1.24 for the stopping times $S \equiv m$ and T we deduce

$$\mathbb{E}[X_T \mathbf{I}_{T > m}] - \mathbb{E}[X_m^+ \mathbf{I}_{T > m}] = \mathbb{E}[X_m \mathbf{I}_{T > m}] - \mathbb{E}[X_m^+ \mathbf{I}_{T > m}] = -\mathbb{E}[X_m^- \mathbf{I}_{T > m}].$$

Hence

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^- \mathbf{I}_{T > n}] - \mathbb{E}[X_m^- \mathbf{I}_{T > m}] = -\mathbb{E}[X_m^- \mathbf{I}_{T > m}]$$

so that

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X_n| \mathbf{I}_{T > n}] = \lim_{n \rightarrow \infty} \mathbb{E}[X_n^+ \mathbf{I}_{T > n}] + \lim_{n \rightarrow \infty} \mathbb{E}[X_n^- \mathbf{I}_{T > n}] = 0.$$

Suppose now that $(X_\bullet)$ is a submartingale. Consider its Doob decomposition $X_n = X_0 + M_n + C_n$. If $X_\bullet$ satisfies (3.1.9), then

$$0 \leq (X_0 + M_n)^+ \leq X_n^+$$

and we deduce that the martingale $Y_\bullet = X_0 + M_\bullet$ satisfies (3.1.9) and thus (3.1.9). Next, observe that

$$X_n^+ = (Y_n^+ + C_n^+) \mathbf{I}_{Y_n \geq 0} + (C_n - Y_n^-) \mathbf{I}_{0 < Y_n^- \leq C_n}.$$

This proves that $0 \leq C_n \leq X_n^+ + Y_n^-$, so

$$\lim_{n \rightarrow \infty} \mathbb{E}[C_n \mathbf{I}_{T > n}] = \lim_{n \rightarrow \infty} \mathbb{E}[(X_n^+ + Y_n^-) \mathbf{I}_{T > n}] = 0.$$

Hence

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X_n| \mathbf{I}_{T > n}] = \lim_{n \rightarrow \infty} \mathbb{E}[|Y_n| + C_n \mathbf{I}_{T > n}] = 0. \quad \square$$

Remark 3.1.31. In the existing literature the stopping and sampling theorems are often referred to by the same name, stopping theorems. We chose to stick with Doob's terminology in [61] because *stopping and sampling are qualitatively different operations*.

The operation of sampling associates to a process a *random variable*, whereas the operation of stopping associates to a process *another process*. $\square$

3.1.5. Applications of the optional sampling theorem. It is time to give the reader a first taste of the versatility of the optional sampling theorem. After we present more properties of martingales we will be able to extend the range of applications of this theorem.

Example 3.1.32 (The Ballot Problem). Let us consider again the *ballot problem* first discussed in Example 1.2.38. Recall the setup.

Two candidates A and B run for an election. Candidate A received a votes while candidate B received b votes, where $b < a$. The votes were counted in random order, so any permutation of the $a + b$ votes cast is equally likely. We have shown in Example 1.2.38 that the probability that A was ahead throughout the count is

$$p = \frac{a - b}{a + b}.$$

We want to describe an alternate proof using martingale methods. Our presentation is inspired from [152, Sec. 12.2].

Set $n := a + b$ and denote by D_k the number votes by which A was ahead when the k -th voted was tabulated. Note that $S_n = a - b$. Let X_k denote the random variable indicating the k -th vote. Thus, $X_k = 1$, if the vote went for A , and $X_k = -1$ if the vote went for B so that

$$D_0 = 0, \quad D_k = X_1 + \cdots + X_k.$$

For $k = 0, 1, \dots, n$ we denote by R_k the ratio

$$R_k := \frac{D_{n-k}}{n - k}.$$

In other words R_k is candidate's A the lead in percentages after the $(n - k)$ -th counted vote. Let us first show that R_k is a martingale with respect to the filtration

$$\mathcal{F}_k = \sigma(R_0, \dots, R_k) = \sigma(D_n, D_{n-1}, \dots, D_{n-k}).$$

Thus, conditioning on $\mathcal{F}_k$ corresponds to conditioning on the results of the last $k + 1$ votes. Observe that, given D_{n-k} the result D_{n-k-1} one vote earlier, is independent of the results at the later votes $D_{n-k+1}, \dots, D_n$. In other words,

$$\mathbb{E}[D_{n-k-1} \mid D_{n-k}, \dots, D_n] = \mathbb{E}[D_{n-k-1} \mid D_{n-k}].$$

One might be tempted to think of D_{n-k} as a random walk in reverse, but there is a silent trap: there is a condition at the n -th step in reverse namely $D_0 = 0$.

To compute the above conditional expectation denote by A_m (resp. B_m) the number of votes A (resp. B) has received after m votes. Thus

$$D_m = A_m - B_m, \quad m = A_m + B_m.$$

Note that A_m and B_m are determined by D_m via the equalities

$$A_m = \frac{D_m + m}{2}, \quad B_m = \frac{m - D_m}{2}.$$

Thus, if D_{n-k} is known, the $(n - k)$ -th vote could have been either a vote for A , and the probability of such a vote is $\frac{A_{n-k}}{n-k}$, or it could have been a vote for B , and the probability of such a vote is $\frac{B_{n-k}}{n-k}$. Hence

$$\begin{aligned} \mathbb{E}[D_{n-k-1} \mid D_{n-k}] &= (D_{n-k} - 1) \frac{A_{n-k}}{n-k} + (D_{n-k} + 1) \frac{B_{n-k}}{n-k} \\ &= D_{n-k} - \frac{D_{n-k}}{n-k} = \frac{n-k-1}{n-k} D_{n-k}. \end{aligned}$$

Dividing by $(n - k - 1)$ we deduce that $(R_k)_{0 \leq k \leq n-1}$ is indeed a martingale.

Now define the stopping times

$$S := \{ 0 \leq k \leq n - 1; R_k = 0 \},$$

where $\min \emptyset := \infty$ and $T := \min(S, n - 1)$. The stopping time T is bounded and the Optional Sampling Theorem 3.1.29 implies

$$\mathbb{E}[R_T] = \mathbb{E}[R_0] = \frac{D_n}{n} = \frac{a - b}{a + b}.$$

Now observe that

$$\mathbb{E}[R_T] = \mathbb{E}[R_T \mathbf{I}_{S=\infty}] + \mathbb{E}[R_T \mathbf{I}_{S<\infty}].$$

Note $R_T = 0$ on $\{S < \infty\}$. Observe that if $S = \infty$, then $D_k > 0$, for all $1 \leq k \leq n$. Hence $T = (n - 1)$ on $\{S = \infty\}$ so $R_T = D_1 = 1$ on $\{S = \infty\}$.

$$\frac{a - b}{a + b} = \mathbb{E}[R_T] = \mathbb{P}[S = \infty]$$

= the probability that candidate A lead throughout the vote count.

□

Example 3.1.33 (Expected time to observe a pattern). Suppose that we are given a finite set (alphabet) $\mathcal{A}$ and a probability distribution π on it so that

$$\pi(a) := \pi(\{a\}) > 0, \quad \forall a \in \mathcal{A}.$$

Define

$$f : \mathcal{A} \rightarrow (0, \infty), \quad f(a) = \frac{1}{\pi(a)}.$$

Fix a word (or pattern) of length $\ell > 0$ in this alphabet, $\mathbf{a} = (a_1, \dots, a_\ell) \in \mathcal{A}^\ell$.

Suppose that $(A_n)_{n \geq 1}$ is a sequence of independent $\mathcal{A}$ -valued random variables with common distribution π . We say that the *pattern $\mathbf{a}$ is observed at time n* if $n \geq \ell$ and

$$(A_{n-\ell+1}, A_{n-\ell+2}, \dots, A_n) = (a_1, a_2, \dots, a_\ell).$$

We let $T = T_{\mathbf{a}}$ denote the first time the pattern $\mathbf{a}$ is observed

$$T_{\mathbf{a}} := \min \{ n \geq \ell; (A_{n-\ell+1}, A_{n-\ell+2}, \dots, A_n) = (a_1, a_2, \dots, a_\ell) \}.$$

To visualize this, think that we have an urn with balls labeled by the letters in $\mathcal{A}$ in proportions given by π . We sample with replacement the urn and we record in succession the labels we draw. We are interested in the moment we first observe the labels $a_1, \dots, a_\ell$ in succession as we sample the urn. As a special case, think that we flip a fair coin and we stop the first we see T, H, T, H in succession. In this case $\mathcal{A} = \{H, T\}$, $\pi(H) = \pi(T) = \frac{1}{2}$, $\mathbf{a} = THTH$.

An amusing quote by Bertrand Russel comes to mind. “*There is a special department of Hell for students of probability. In this department there are many typewriters and many monkeys. Every time that a monkey walks on a typewriter, it types by chance one of Shakespeare’s sonnets.*”

We will compute $\mathbb{E}[T_{\mathbf{a}}]$ by using a clever martingale method due to Li [136]. The precise answer is contained in (3.1.12)

Let us first observe that $\mathbb{E}[T_{\mathbf{a}}] < \infty$. This follows from a very useful trick, [224, E10.5], generalizing the result in Example 1.4.13.

Lemma 3.1.34 (‘Sooner-rather-than-later’). *Suppose that T is a stopping time adapted to the filtration $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$ with the property that there exist $r_0 > 0$ and $N_0 \in \mathbb{N}$ such that*

$$\forall n \in \mathbb{N}_0, \quad \mathbb{P}[T \leq n + N_0 \mid \mathcal{F}_n] > r_0. \quad (3.1.10)$$

Then there exists $c \in (0, 1)$ such that $\mathbb{P}[T > n] < c^n, \forall n > N_0$. In particular,

$$\mathbb{E}[T] = \sum_{n \geq 0} \mathbb{P}[T > n] < \infty. \quad \square$$

In Exercise 3.6 we ask the reader to provide a proof of this result. It is a nice application of various properties of the conditional expectation.

In the case at hand the condition (3.1.10) is satisfied with $N_0 = \ell$ and $r = \left(\min_{a \in A} \pi(a) \right)^\ell$.

Following [136] we consider the following betting game involving the House (casino) and a random number of players. At each moment of time $n = 1, 2, \dots$ the House samples the alphabet $\mathcal{A}$ according to the probability distribution π . (The House runs a chance game with set of outcomes $\mathcal{A}$ and probability distribution π .) The outcome of this sampling is the sequence of i.i.d. random variables A_n .

The first player adopts the following α -based strategy.

- At time 0 he bets his fortune $F_0^1 = 1$ that the outcome of the first game is $A_1 = a_1$. If $A_1 = a_1$ his fortune will change to $F_1^1 = f(a_1) = \frac{1}{a_1}$. Otherwise, he will lose his fortune F_0^1 to the house, so $F_1^1 = 0$ in this case.
- At time 1 he bets his fortune that $A_2 = a_2$. If he wins, i.e., $A_2 = a_2$, his fortune at time 2 will grow to $F_2^1 = f(a_2)F_1^1$. If he loses, he will have to turn all its fortune to the House.
- In general, if $k \leq \ell$ and his fortune at time $k - 1$ is F_{k-1}^1 (the fortune could be 0 at that moment), the player bets all its fortune, $f(a_k)$ on a dollar, that $A_k = a_k$. If this happens, his fortune will grow to $F_k^1 = f(a_k)F_{k-1}^1$. Otherwise, he will surrender his fortune F_{k-1}^1 to the house, so $F_k^1 = 0$ in this case.
- At time ℓ the first gambler stops playing, so $F_n^1 = F_\ell^1, \forall n \geq \ell$
- We denote by X_n^1 the profit of the first player at time n , $X_n^1 = F_n^1 - F_0^1 = F_n^1 - 1$.

Concisely, if we define

$$M_k^1 = \begin{cases} f(a_k) \mathbf{I}_{\{A_k = a_k\}}, & 1 \leq k \leq \ell, \\ 1, & k < 1 \text{ or } k > \ell, \end{cases}$$

then

$$F_n^1 = \prod_{k=1}^n M_k^1.$$

Since $\mathbb{E}[M_k^1] = 1$ we deduce that $F_\bullet^1$ and $X_\bullet^1 = F_\bullet^1 - 1$ are martingales.

In general, for $m = 1, 2, \dots$, the m -th player also plays ℓ rounds using the same strategy as the first player, but with a delay of $m - 1$ units of times.

Thus, the second player skips game 1 and only starts betting before the 2nd game using the same betting strategy as if the game started when he began playing: at his j -th round he bets $f(a_j)$ on a dollar that the outcome is $A_{j+1} = a_j$. The third player skips the first two games etc.

In general, at his j -th round, the m -th player bets $f(a_j)$ on a dollar that the outcome is $A_{j+m-1} = a_j$. We denote by F_n^m the fortune of the m -th player at time n . More precisely, if we set

$$M_k^m := \begin{cases} f(a_{k-m+1}) \mathbf{I}_{\{A_k = a_{k-m+1}\}}, & m \leq k \leq m + \ell - 1, \\ 1, & k < m \text{ or } k \geq m + \ell, \end{cases}$$

then

$$F_n^m := \prod_{k=1}^n M_k^m, \quad X_n^m = F_n^m - 1 \quad n = 1, 2, \dots$$

Note that $F_n^m = 1$ for $n < m$ because the m -th player skips the games $n = 1, 2, \dots, m-1$. Define

$$S_n := \sum_{m \geq 1} X_n^m = \sum_{m=1}^n X_n^m = \sum_{m=1}^n F_n^m - n.$$

In other words, S_n is the sum of the profits of all the players after n games. The process $S_\bullet$ is obviously a martingale. Note that

$$S_T = \sum_{m \leq T} F_T^m - T, \quad T = T_a.$$

Recall that T is the first moment of time such that

$$A_{T-\ell+1} = a_1, \quad A_{T-\ell+2} = a_2, \dots, A_T = a_\ell. \quad (3.1.11)$$

Thus the player $(T - \ell + 1)$ will be the first player to hit the jackpot, i.e., observes the pattern $\mathbf{a}$ during the first ℓ games he plays. This proves $F_T^m = 0$ for $m \leq T - \ell$. Indeed, the minimality of T implies

$$(A_m, \dots, A_{m+\ell-1}) \neq (a_1, \dots, a_\ell)$$

and thus $\mathbf{I}_{\{A_m = a_1\}} \cdots \mathbf{I}_{\{A_{m+\ell-1} = a_\ell\}} = 0$.

The fortune of the player $T - \ell + 1$ at time T is

$$F_T^{T-\ell+1} = f(a_1) \cdots f(a_\ell).$$

Using the equalities (3.1.11) we deduce that the fortune of the next player, $T - \ell + 2$, at time T is nonzero if and only if

$$(a_2, \dots, a_\ell) = (a_1, \dots, a_{\ell-1}).$$

In this case the fortune is $f(a_1) \cdots f(a_{\ell-1})$. Similarly,

$$F_T^{T-\ell+3} = \begin{cases} f(a_1) \cdots f(a_{\ell-2}), & (a_1, \dots, a_{\ell-2}) = (a_3, \dots, a_\ell), \\ 0, & (a_1, \dots, a_{\ell-2}) \neq (a_3, \dots, a_\ell). \end{cases}$$

More generally, denote by $\delta_{\alpha, \beta}$ the Kronecker symbol

$$\delta_{\alpha, \beta} := \begin{cases} 1, & \alpha = \beta, \\ 0, & \alpha \neq \beta. \end{cases}$$

We deduce

$$\begin{aligned} S_T + T &= \underbrace{F_T^1 + \cdots + F_T^{T-\ell}}_{=0} + F_T^{T-\ell+1} + F_T^{T-\ell+2} + \cdots + F_T^T \\ &= F_T^{T-\ell+1} + F_T^{T-\ell+2} + \cdots + F_T^T \end{aligned}$$

$$\begin{aligned}
&= \underbrace{f(a_1) \cdots f(a_\ell)}_{F_T^T} + \underbrace{\prod_{j=1}^{\ell-1} f(a_j) \delta_{a_{j+1}, a_j}}_{F_T^{T-1}} + \underbrace{\prod_{j=1}^{\ell-2} f(a_j) \delta_{a_{j+2}, a_j}}_{F_T^{T-2}} + \cdots \\
&= \underbrace{\sum_{k=0}^{\ell-1} \prod_{j=1}^{\ell-k} f(a_j) \delta_{a_{j+k}, a_j}}_{=: \tau(\mathbf{a})}.
\end{aligned}$$

Hence $S_T = \tau(\mathbf{a}) - T$. If we could show that T satisfies Doob's conditions (Definition 3.1.27), then we could invoke the Optional Sampling Theorem 3.1.29 and conclude that

$$0 = \mathbb{E}[S_0] = \mathbb{E}[S_T] = \tau(\mathbf{a}) - \mathbb{E}[T].$$

Let us show that indeed the stopping time satisfies Doob's conditions.

Since $\mathbb{E}[T] < \infty$ and $S_T = \tau(\mathbf{a}) - T$ we deduce $S_T \in L^1$. Arguing as above we deduce that if $n < T$, then

$$\begin{aligned}
F_n^1 + \cdots + F_n^n &\leq F_n^{n-\ell+1} + \cdots + F_n^n \\
&\leq f(a_1) \cdots f(a_\ell) + \prod_{j=1}^{\ell-1} f(a_j) \delta_{a_{j+1}, a_j} + \prod_{j=1}^{\ell-2} f(a_j) \delta_{a_{j+2}, a_j} + \cdots = \tau(\mathbf{a}).
\end{aligned}$$

Hence

$$|S_n| \mathbf{I}_{\{T > n\}} \leq (\tau(\mathbf{a}) + n) \mathbf{I}_{\{T > n\}} \leq (\tau(\mathbf{a}) + T) \mathbf{I}_{\{T > n\}}.$$

Since $\mathbb{E}[T] < \infty$ we deduce

$$\lim_{n \rightarrow \infty} \mathbb{E}[|S_n| \mathbf{I}_{\{T > n\}}] = 0.$$

This shows that the stopping time $T_{\mathbf{a}}$ satisfies Doob's conditions so that

$$\mathbb{E}[T_{\mathbf{a}}] = \tau(\mathbf{a}) = \sum_{k=0}^{\ell-1} \prod_{j=1}^{\ell-k} \frac{\delta_{a_{j+k}, a_j}}{\pi(a_j)}. \quad (3.1.12)$$

Let us describe this equality using a more convenient notation. Denote by $V(\mathcal{A})$ the vocabulary of the alphabet $\mathcal{A}$

$$V(\mathcal{A}) = \bigsqcup_{\ell \geq 0} \mathcal{A}^\ell, \quad \mathcal{A}^0 := \{\emptyset\}.$$

We denote by $\ell(\mathbf{a})$ the length of a word $\mathbf{a}$.

We define a weight $w = w_\pi : V(\mathcal{A}) \rightarrow (0, \infty)$ by setting

$$w(a_1, \dots, a_\ell) = \prod_{k=1}^{\ell} f(a_k), \quad w(\emptyset) = 1.$$

For $\mathbf{a} = (a_1, \dots, a_\ell) \in \mathcal{A}^\ell$, $\ell \geq 1$, and $j = 1, \dots, \ell$ we define the left/right tail maps

$$L_j, R_j : \mathcal{A}^\ell \rightarrow \mathcal{A}^j, \quad L_j(\mathbf{a}) = (a_1, \dots, a_j), \quad R_j(\mathbf{a}) = (a_{\ell-j+1}, \dots, a_\ell).$$

Thus, R_j retains only the last j letters of a word while L_j retains the first j letters.

Given two words $\mathbf{a}, \mathbf{b} \in V(\mathcal{A})$ we set

$$\langle \mathbf{a}, \mathbf{b} \rangle := \begin{cases} 1, & \mathbf{a} = \mathbf{b}, \\ 0, & \mathbf{a} \neq \mathbf{b}. \end{cases}$$

Now define

$$\Phi : V(\mathcal{A}) \times V(\mathcal{A}) \rightarrow [0, \infty), \quad \Phi(\mathbf{a}, \mathbf{b}) = \sum_{j=1}^{\ell(\mathbf{a}) \wedge \ell(\mathbf{b})} \langle R_j \mathbf{a}, L_j \mathbf{b} \rangle w(L_j \mathbf{b}). \quad (3.1.13)$$

We can rewrite (3.1.12) as

$$\mathbb{E}[T_{\mathbf{a}}] = \Phi(\mathbf{a}, \mathbf{a}). \quad (3.1.14)$$

In the special case when $\mathcal{A} = \{1, 2, \dots, 6\}$, π is the uniform counting probability and

$$\mathbf{a} = \underbrace{6 \cdots 6}_k \in \mathcal{A}^k,$$

then the waiting time $\tau(\mathbf{a})$ coincides with the waiting time T to observe the first occurrence of a k -run of 6-s discussed in Example 1.4.13. In this case we have

$$\mathbb{E}[T] = \sum_{j=1}^k 6^j = \frac{6^{k+1} - 6}{5}.$$

We refer to Example A.3.20 for an R-code that simulates sampling an alphabet until a given pattern is observed.

Let us discuss in more detail the special case $\mathcal{A} = \{H, T\}$, $\pi(H) = \pi(T) = \frac{1}{2}$. Suppose that $\mathbf{a}$ is the pattern $\mathbf{a} = (TTHH)$, $\mathbf{b} = HHH$. Observe that $\langle L_j \mathbf{a}, R_j \mathbf{a} \rangle = 1$ for $j = 4$ and 0 otherwise. Hence $\mathbb{E}[T_{\mathbf{a}}] = 16$. A similar computation shows that $\mathbb{E}[T_{\mathbf{b}}] = 14$. Thus we have to wait a longer time for the pattern $\mathbf{a}$ to occur.

On the other hand, a formula of Conway (see Exercise 3.13) shows that

$$\frac{\mathbb{P}[T_{\mathbf{b}} < T_{\mathbf{a}}]}{\mathbb{P}[T_{\mathbf{a}} < T_{\mathbf{b}}]} = \frac{\Phi(\mathbf{a}, \mathbf{a}) - \Phi(\mathbf{a}, \mathbf{b})}{\Phi(\mathbf{b}, \mathbf{b}) - \Phi(\mathbf{b}, \mathbf{a})}.$$

We have $\langle L_i \mathbf{a}, R_j \mathbf{b} \rangle = 0, \forall j$ and $\langle R_j \mathbf{a}, L_j \mathbf{b} \rangle = 1$ for $j = 1, 2$ so that

$$\Phi(\mathbf{a}, \mathbf{b}) = 0, \quad \Phi(\mathbf{b}, \mathbf{a}) = 6, \quad \frac{\mathbb{P}[T_{\mathbf{b}} < T_{\mathbf{a}}]}{\mathbb{P}[T_{\mathbf{a}} < T_{\mathbf{b}}]} = \frac{5}{7}.$$

We have reached a somewhat surprising conclusion: although, on average, we have to wait a shorter amount of time to observe the pattern $\mathbf{b}$, it is less likely that we will observe $\mathbf{b}$ before $\mathbf{a}$. The odds that $\mathbf{b}$ will appear first versus that $\mathbf{a}$ will appear first are 5 : 7.

There are other strange phenomena. We should mention M. Gardner's even stranger nontransitivity paradox [83, Chap. 5]. More precisely, given any pattern $\mathbf{a} \in \mathcal{A}^k$ there exists a pattern $\mathbf{b} \in \mathcal{A}^k$ such that $\mathbf{b}$ is more likely to occur before $\mathbf{a}$, i.e., $\mathbb{P}[T_{\mathbf{b}} < T_{\mathbf{a}}] > \frac{1}{2}$. This seems counterintuitive since the set of patterns $\mathcal{A}^k$ is finite. As shown in by Guibas and Odlyzko [95], if $\mathbf{a} = (a_1, \dots, a_k)$ we can choose $\mathbf{b}$ to be of the form $\mathbf{b} = (b, a_1, \dots, a_{k-1})$. $\square$

3.1.6. Concentration inequalities: martingale techniques. Hoeffding's inequality (2.3.14) has a martingale counterpart usually referred to as Azuma's inequality.

Theorem 3.1.35 (Azuma). *Suppose that $(X_n)_{n \geq 0}$ is a martingale adapted to a filtration $\mathcal{F}_{\bullet} = (\mathcal{F}_n)_{n \geq 0}$ of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Assume that for any $n \in \mathbb{N}$ there exist constants $a_n < b_n$ such the differences $D_n = X_n - X_{n-1}$ satisfy*

$$a_n \leq D_n \leq b_n \quad \text{a.s..}$$

Then

$$\forall x > 0, \quad \mathbb{P}[|X_n - X_0| > x] \leq 2e^{-\frac{2x^2}{(s_1^2 + \dots + s_n^2)}}, \quad s_k = b_k - a_k. \quad (3.1.15)$$

Proof. The strategy is a variation on the Chernoff's method. Set

$$D_n := X_n - X_{n-1}, \quad \sigma_n^2 := s_1^2 + \dots + s_n^2, \quad \forall n \in \mathbb{N}.$$

We will prove inductively that

$$X_n - X_0 \in \mathbb{G}(\sigma_n^2/4), \quad \text{i.e.,} \quad \mathbb{E}[e^{\lambda(X_n - X_0)}] \leq e^{\frac{\lambda^2 \sigma_n^2}{8}}, \quad \forall n \in \mathbb{N}, \quad \lambda \in \mathbb{R}. \quad (3.1.16)$$

Assuming this, the inequality (3.1.15) follows from (2.3.13b).

To prove (3.1.16) note that since (X_n) is a martingale we have

$$\mathbb{E}[e^{\lambda(X_n - X_0)} \mid \mathcal{F}_{n-1}] = e^{\lambda(X_{n-1} - X_0)} \mathbb{E}[e^{\lambda D_n} \mid \mathcal{F}_{n-1}].$$

We set

$$Z_n(\lambda) := \mathbb{E}[e^{\lambda D_n} \mid \mathcal{F}_{n-1}], \quad \forall n \in \mathbb{N}, \quad \lambda \in \mathbb{R}.$$

We claim that

$$\forall n \in \mathbb{N}, \quad \forall \lambda \in \mathbb{R}, \quad Z_n(\lambda) \leq e^{\frac{\lambda^2 \sigma_n^2}{8}} \quad \text{a.s.} \quad (3.1.17)$$

Obviously this implies that

$$\mathbb{E}[e^{\lambda(X_n - X_0)}] \leq e^{\frac{\lambda^2 \sigma_n^2}{8}} \mathbb{E}[e^{\lambda(X_{n-1} - X_0)}],$$

from which we can conclude inductively that $X_n - X_0 \in \mathbb{G}(\sigma_n^2/4)$.

To prove (3.1.17) observe that, by construction, $Z_n(\lambda)$ is $\mathcal{F}_{n-1}$ -measurable. We have to show that for any $S \in \mathcal{F}_{n-1}$ such that $\mathbb{P}[S] \neq 0$

$$\mathbb{E}[Z_n(\lambda) \mathbf{I}_S] \leq \mathbb{P}[S] e^{\frac{\lambda^2 \sigma_n^2}{8}}.$$

Denote by D_n^S the random variable $D_n|_S$ defined on the probability space $(S, \mathcal{F}_{n-1}|_S, \mathbb{P}_S)$, where

$$\mathbb{P}_S[A] = \mathbb{P}[A|S] = \frac{\mathbb{P}[A \cap S]}{\mathbb{P}[S]}, \quad \forall A \in \mathcal{F}_{n-1}|_S.$$

We denote by $\mathbb{E}_S$ the expectation on $(S, \mathcal{F}_{n-1}|_S, \mathbb{P}_S)$. Since $\mathbb{E}[D_n \mid \mathcal{F}_{n-1}] = 0$ we deduce

$$\mathbb{E}_S[D_n^S] = \frac{1}{\mathbb{P}[S]} \int_{\Omega} \mathbf{I}_S D_n d\mathbb{P} = 0.$$

Clearly $a_n \leq D_n^S \leq b_n$. We deduce from Hoeffding's Lemma (Proposition 2.3.11) that

$$\frac{1}{\mathbb{P}[S]} \mathbb{E}[\mathbf{I}_S e^{\lambda D_n}] = \mathbb{E}_S[e^{\lambda D_n^S}] \leq e^{\frac{\lambda^2 \sigma_n^2}{8}},$$

and therefore, $\forall S \in \mathcal{F}_{n-1}$ such that, $\mathbb{P}[S] \neq 0$ we have

$$\mathbb{E}[Z_n(\lambda) \mathbf{I}_S] = \mathbb{E}[e^{\lambda D_n} \mathbf{I}_S] = \mathbb{P}[S] \mathbb{E}_S[e^{\lambda D_n^S}] \leq \mathbb{P}[S] e^{\frac{\lambda^2 \sigma_n^2}{8}}.$$

This concludes the proof of Azuma's inequality. $\square$

The strength of Azuma's inequality is best appreciated in concrete examples.

Example 3.1.36 (Longest common subsequence). We want to have another look at the problem of the longest common subsequence first discussed in Example 1.3.70. Let us briefly recall the set-up.

We are given a finite set (alphabet) $\mathcal{A}$, $|\mathcal{A}| = k$, and a family of independent $\mathcal{A}$ -valued random variables

$$\{X_n, Y_n; \quad m, n \in \mathbb{N}\}$$

all with the same distribution π . We denote by L_n the length of the longest common subsequence of two random words

$$(X_1, \dots, X_n) \quad \text{and} \quad (Y_1, \dots, Y_n).$$

We set

$$R_n := \frac{1}{n} L_n, \quad R := \sup_n R_n.$$

In Example 1.3.70 we have shown

$$\frac{1}{n} L_n \rightarrow R \quad \text{a.s.},$$

and

$$\lim_{n \rightarrow \infty} \mathbb{E}[R_n] = r(\pi) := \mathbb{E}[R].$$

We will to show that R_n is highly concentrated around its mean r_n . We follow the presentation in [198, Sec. 1.3].

Set $\ell_n := \mathbb{E}[L_n]$, $Z_n = (X_n, Y_n)$. Consider the finite filtration

$$\mathcal{F}_0 := \sigma(\emptyset), \quad \mathcal{F}_j = \sigma(Z_1, \dots, Z_j), \quad j = 1, \dots, n.$$

Form the Doob (closed) martingale $U_j := \mathbb{E}[L_n | \mathcal{F}_j]$. Note that $U_0 = \ell_n$. The random variable L_n is a function of the Z_j 's

$$L_n = L_n(Z_1, \dots, Z_n),$$

and U_j is a function of $Z_1, \dots, Z_j$, $U_j = F_j(Z_1, \dots, Z_j)$. More precisely, since the variables Z_n are independent, we have

$$\begin{aligned} F_j(z_1, \dots, z_j) &= \mathbb{E}[L_n(z_1, \dots, z_j, Z_{j+1}, \dots, Z_n)] \\ &= \int_{(\mathcal{A}^2)^{n-j}} L_n(z_1, \dots, z_j, z_{j+1}, \dots, z_n) \pi^{\otimes 2(n-j)}[dz_{j+1} \cdots dz_n]. \end{aligned}$$

Note that for any $z_1, \dots, z_{j-1}, z_j, z'_j, z_{j+1}, \dots, z_n \in \mathcal{A}^2$ we have

$$-1 \leq L_n(z_1, \dots, z_{j-1}, z'_j, z_{j+1}, \dots, z_n) - L_n(z_1, \dots, z_{j-1}, z_j, z_{j+1}, \dots, z_n) \leq 1$$

Integrating with respect to $z'_j, z_{j+1}, \dots, z_n$ we deduce

$$-1 \leq F_{j-1}(z_1, \dots, z_{j-1}) - F_j(z_1, \dots, z_n) \leq 1$$

Hence $|U_j - U_{j-1}| \leq 1$. From Azuma's inequality with $s_n = 2$ we deduce

$$\mathbb{P}[|L_n - \ell_n| \geq nx] \leq 2e^{-\frac{nx^2}{2}},$$

so that

$$\mathbb{P}[|R_n - r_n| \geq x] \leq 2e^{-\frac{nx^2}{2}}.$$

This proves that R_n is highly concentrated around its mean. Obviously

$$\forall \varepsilon > 0, \quad \sum_{n \geq 1} \mathbb{P}[|R_n - r_n| \geq \varepsilon] < \infty,$$

and Corollary 1.3.57 implies that $R_n - r_n \rightarrow 0$ a.s..

On the other hand, we know from Example 1.3.70 that

$$R_n \rightarrow R \text{ a.s. and } r_n \rightarrow r(\pi) = \mathbb{E}[R].$$

Hence $\frac{1}{n}L_n$ converges almost surely to a constant $r(\pi)$.

We write $r(k)$ instead of $r(\pi)$ when π is the uniform distribution on an alphabet of cardinality k . In this case one has additional information about the rate of convergence of r_n to $r(k)$. However, the exact value of $r(k)$ remains illusive, even for small k .

□

Example 3.1.37 (Bin packing). The bin packing problem has a short formulation: pack n items of sizes $x_1, \dots, x_n \in [0, 1]$ in as few bins of maximum capacity 1 each. We denote by $B_n(x_1, \dots, x_n)$ the lowest numbers of bins we can use to pack the items of sizes $x_1, \dots, x_n$.

As in the case of the longest common subsequence problem, the bin packing problem has a probabilistic counterpart. Consider independent random variables $X_n \sim \text{Unif}([0, 1])$, $n \in \mathbb{N}$ defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. We will describe the behavior of $b_n := \mathbb{E}[B_n(X_1, \dots, X_n)]$ as $n \rightarrow \infty$.

Note that

$$X_1 + \dots + X_n \leq B_n(X_1, \dots, X_n) \leq n.$$

By taking expectations we deduce

$$\frac{n}{2} \leq b_n \leq n, \quad \forall n \in \mathbb{N}, \quad (3.1.18)$$

showing that b_n has linear growth as $n \rightarrow \infty$. On the other hand,

$$\begin{aligned} B_{n+m}(X_1, \dots, X_n, X_{n+1}, \dots, X_{n+m}) \\ \leq B_n(X_1, \dots, X_n) + B_m(X_{n+1}, \dots, X_{n+m}), \end{aligned} \quad (3.1.19)$$

and thus

$$b_{n+m} \leq b_n + b_m, \quad \forall n, m \in \mathbb{N}.$$

Setting $r_n := \frac{b_n}{n}$, we deduce from Fekete's Lemma 1.3.71 that

$$\lim_{n \rightarrow \infty} r_n = r := \inf_n r_n.$$

The inequalities (3.1.18) show that $r \in [\frac{1}{2}, 1]$.

We set $R_n := \frac{B_n}{n}$. We deduce from (3.1.19) and Fekete's Lemma that

$$R_n \rightarrow R := \inf_n R_n \text{ a.s. and } r = \mathbb{E}[R].$$

We want to show that R_n is highly concentrated around its mean. We use the same approach as in Example 3.1.36.

We set

$$\mathcal{F}_j = \sigma(X_1, \dots, X_j), \quad \mathcal{F}_0 = \{\emptyset, \Omega\}$$

Fix $n \in \mathbb{N}$. For $j = 0, 1, \dots, n$ we set

$$U_j = U_{n,j} := \mathbb{E}[B_n \mid \mathcal{F}_j]$$

so the collection $(U_j)_{0 \leq j \leq n}$ is a martingale adapted to the filtration $(\mathcal{F}_j)_{0 \leq j \leq n}$.

There exist Borel measurable maps $F_j : [0, 1]^j \rightarrow \mathbb{N}$ such that $U_j = F_j(X_1, \dots, X_j)$. More precisely,

$$F_j(x_1, \dots, x_j) = \int_{[0,1]^{n-j}} B_n(x_1, \dots, x_j, x_{j+1}, \dots, x_n) dx_{j+1} \cdots dx_n.$$

For any $x_1, \dots, x_{j-1}, x_j, x'_j, x_{j+1}, \dots, x_n \in [0, 1]$ we have

$$-1 \leq B_n(x_1, \dots, x_{j-1}, x'_j, x_{j+1}, \dots, x_n) - B_n(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_n) \leq 1$$

Integrating with respect to $x'_j, x_{j+1}, \dots, x_n$ we deduce $|U_j - U_{j-1}| \leq 1$. Invoking Azuma's inequality we deduce as in Example 3.1.36 that

$$\mathbb{P}[|R_n - r_n| > x] \leq 2e^{-\frac{nx^2}{2}}.$$

This shows that R_n is highly concentrated around its mean and that $R_n \rightarrow r$ a.s..

In this case it is known that $r = \frac{1}{2}$. More precisely, there is an algorithm called MATCH which takes as input the sizes $x_1, \dots, x_n$ of the n items and packs them into $M_n = M_n(x_1, \dots, x_n)$ boxes where

$$\frac{n}{2} \leq \mathbb{E}[B_n] \leq \mathbb{E}[M_n] \leq \frac{n}{2} + O(\sqrt{n}),$$

This is the best one can hope for since it is also known

$$\mathbb{E}[B_n] \geq \frac{n}{2} + (\sqrt{3} - 1) \sqrt{\frac{n}{24\pi}} + o(\sqrt{n})$$

For details we refer to [44, Sec. 5.1]. □

The tricks used in the above examples are generalized and refined in McDiarmid's inequality.

Definition 3.1.38 (Bounded difference property). Suppose that S is a set. A function $f : S^n \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, is said to satisfy the *bounded difference property* if there exist $L_1, \dots, L_n > 0$ such that,

$$|f(s_1, \dots, s_{k-1}, s, s_{k+1}, \dots, s_n) - f(s_1, \dots, s_{k-1}, s', s_{k+1}, \dots, s_n)| \leq L_k, \quad (3.1.20)$$

$\forall k = 1, \dots, n, \forall s_1, \dots, s_{k-1}, s, s', s_{k+1}, \dots, s_n \in S$. □

Let us observe that the above condition is satisfied if and only if f is Lipschitz with respect to the *Hamming distance* on S

$$d_H : S^n \times S^n \rightarrow [0, \infty), \quad d_H(\underline{s}, \underline{t}) := \sum_{k=1}^n \mathbf{1}_{\mathbb{R} \setminus \{0\}}(s_k - t_k). \quad (3.1.21)$$

Theorem 3.1.39 (McDiarmid's inequality). Suppose that

$$X_1, \dots, X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$$

are independent random variables and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the bounded difference property with constants $L_1, \dots, L_n$. If $Z = f(X_1, \dots, X_n)$ is integrable, then

$$\mathbb{P}[Z - \mathbb{E}[Z] > t] \leq e^{-2t^2/L^2}, \quad L^2 = L_1^2 + \dots + L_n^2. \quad (3.1.22a)$$

$$\mathbb{P}[|Z - \mathbb{E}[Z]| > t] \leq 2e^{-2t^2/L^2}. \quad (3.1.22b)$$

Proof. Denote by $\mathbb{P}_k$ the distribution on X_k . Let $\mathcal{F}_0 = \{\emptyset, \Omega\}$, $\mathcal{F}_k := \sigma(X_1, \dots, X_k)$ and set

$$Z_k := \mathbb{E}[Z \mid \mathcal{F}_k], \quad k = 0, \dots, n$$

so that $Z_n = Z$ and $Z_0 = \mathbb{E}[Z]$. Since $X_1, \dots, X_n$ are independent we deduce that $\forall \omega \in \Omega$

$$Z_k(\omega) = g_k(X_1(\omega), \dots, X_k(\omega)),$$

where

$$\begin{aligned} & g_k(x_1, \dots, x_k) \\ &= \int_{\mathbb{R}^{n-k}} f(x_1, \dots, x_k, x_{k+1}, \dots, x_n) \mathbb{P}_{k+1}[dx_{k+1}] \cdots \mathbb{P}_n[dx_n]. \end{aligned}$$

Note that

$$g_{k-1}(x_1, \dots, x_{k-1}) = \mathbb{E}_k[g_k] := \int_{\mathbb{R}} g_k(x_1, \dots, x_{k-1}, x_k) \mathbb{P}_k[dx_k].$$

Hence

$$D_k = g_k - \mathbb{E}_k g_k, \quad \mathbb{E}[e^{\lambda D_k} \mid \mathcal{F}_{k-1}] = h_{k-1}(X_1, \dots, X_{k-1}),$$

where $h_{k-1}(x_1, \dots, x_{k-1}) := \mathbb{E}_k[e^{\lambda(g_k - \mathbb{E}_k g_k)}]$.

Fix $x_1, \dots, x_{k-1}$ and set

$$a_k = a_k(x_1, \dots, x_{k-1}) := \inf_{x_k} g(x_1, \dots, x_{k-1}, x_k),$$

$$b_k = b_k(x_1, \dots, x_{k-1}) := \sup_{y_k} g(x_1, \dots, x_{k-1}, y_k).$$

We deduce that

$$0 \leq b_k - a_k \leq \sup_{x_k, y_k} |g(x_1, \dots, x_{k-1}, y_k) - g(x_1, \dots, x_{k-1}, x_k)| \leq L_k.$$

We deduce from Hoeffding's inequality (2.3.16) that

$$\mathbb{E}_k[e^{\lambda(g_k - \mathbb{E}_k g_k)}] \leq e^{\frac{\lambda L_k^2}{8}}, \quad \forall x_1, \dots, x_{k-1}.$$

Hence

$$\mathbb{E}[e^{\lambda D_k} \mid \mathcal{F}_{k-1}] \leq e^{\frac{\lambda L_k^2}{8}} \quad \text{a.s., i.e.,} \quad \mathbb{E}[e^{\lambda(Z_n - Z_0)}] \leq e^{\frac{\lambda(L_1^2 + \dots + L_n^2)}{8}}.$$

□

Example 3.1.40 (Occupancy problem). Consider the occupancy problem discussed in Exercise 2.31. Suppose that b labeled balls are successively and randomly placed in u labeled urns. The urns are equally likely to be the destination of a given ball. Let $N = N_{u,b}$ the number of empty urns after all the balls have been distributed.

Denote by U_i the event i -th urn is empty. This is Bernoulli variable with success probability $(1 - 1/u)^b$. Then

$$N = \sum_{i=1}^u \mathbf{I}_{U_i}$$

so

$$m(u, b) := \mathbb{E}[N] = u \mathbb{E}[\mathbf{I}_{U_1}] = u \left(1 - \frac{1}{u}\right)^b.$$

For $j = 1, \dots, b$ denote by X_j the label of the urn in which the j -th ball is placed. Then

$$N_{u,b} = f(X_1, \dots, X_b),$$

for some Borel measurable function $f : \mathbb{R}^b \rightarrow \mathbb{R}$.

Then f satisfies the bounded difference property with $L_1 = \dots = L_b = 1$. Indeed if X_j changes by 1, the number of empty urns changes by at most 1. Using McDiarmid's inequality we deduce that for any $r \in (0, 1)$

$$\mathbb{P}[|N - m(u, b)| > ru] \leq 2e^{-2r^2u^2/b}.$$

For example, if $b = 2u$, then we deduce

$$\mathbb{P}[|N - m(u, b)| > ru] \leq 2e^{-r^2u/8}.$$

As $u \rightarrow \infty$ we have $\frac{m(u, 2u)}{u} = (1 - 1/u)^{2u} \rightarrow e^{-2} \approx 0.1335$.

If we denote by $F_u = \frac{N}{u}$ the fraction of empty urns we deduce that F_u is highly concentrated around its mean e^{-2} and $F_u \rightarrow e^{-2}$ a.s. as $u \rightarrow \infty$. $\square$

3.1.7. Uniform laws of large numbers revisited. Suppose that we are given

- a sequence of i.i.d. random vectors

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{X} := \mathbb{R}^N, \quad n \in \mathbb{N},$$

and

- a collection $\mathcal{F}$ of measurable functions $f : \mathbb{X} \rightarrow \mathbb{R}$ such that $f(X_n) \in L^1$, for all $f \in \mathcal{F}$.

In machine learning $\mathcal{F}$ is known as the hypothesis space. For $f \in \mathcal{F}$ we set

$$\bar{f}(x) := f(x) - \mathbb{E}[f(X_j)].$$

Note that the right-hand-side is indeed independent of j . The Strong Law of Large Numbers implies that

$$\forall f \in \mathcal{F}, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \bar{f}(X_k) = 0 \quad \text{a.s.}$$

We are interested if this happens almost surely uniformly in $f \in \mathcal{F}$. More precisely, for $n \in \mathbb{N}$, we set

$$D_n(\mathcal{F}) = D_n(\mathcal{F}, X) := \sup_{f \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(X_k) \right|,$$

Assume that $D_n(\mathcal{F})$ is measurable for every n .

We want to investigate if $D_n(\mathcal{F}) \rightarrow 0$ a.s.. In Section 2.4 we have investigated a special case of this problem, namely when $\mathcal{F}$ had the form

$$\mathcal{F} := \{ \mathbf{I}_C; \quad C \in \mathcal{C} \},$$

where $\mathcal{F}$ is a collection indicator functions of a collection $\mathcal{C}$ of subsets of $\mathbb{R}^N$. We showed that if the collection $\mathcal{C}$ has finite VC dimension, then $D_n(\mathcal{F}) \rightarrow 0$ in this case.

In this subsection we have a more limited goal. We want to provide an upper bound for $D_n(\mathcal{F})$ in terms of a probabilistic invariant of $\mathcal{F}$ that turns out to be relatable to the concept of VC-dimension. We follow the approach in [219, Sec. 4.2].

Let us first show that $D_n(\mathcal{F})$ is highly concentrated around its mean $\mathbb{E}[D_n(\mathcal{F})]$.

Lemma 3.1.41. For each $n \in \mathbb{N}$ the function the function $G_n : \mathbb{X}^n \rightarrow \mathbb{R}$

$$G_n(x_1, \dots, x_n) = \sup_{f \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(x_k) \right|$$

satisfies the bounded difference property (Definition 3.1.38) with $L_k = \frac{2C}{n}$, $\forall k = 1, \dots, n$.

Proof. Indeed, let $x, y \in \mathbb{R}^n$ such that there exists a single $j = 1, \dots, n$ such that $x_j \neq y_j$. Without loss of generality we can assume $x_1 \neq y_1$, $x_j = y_j$, $\forall j \geq 1$. for any $f \in \mathcal{F}$ we have

$$\begin{aligned} \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(x_k) \right| - G_n(y_1, \dots, y_n) &= \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(x_k) \right| - \sup_{h \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n \bar{h}(y_k) \right| \\ &\leq \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(x_k) \right| - \frac{1}{n} \left| \sum_{k=1}^n \bar{f}(y_k) \right| \leq \frac{1}{n} (|\bar{f}(x_1)| + |\bar{f}(y_1)|) \leq \frac{2C}{n}. \end{aligned}$$

Passing to sup over $f \in \mathcal{F}$ we deduce

$$G_n(x) - G_n(y) \leq \frac{2C}{n}, \quad \forall x, y \in \mathbb{R}^N.$$

□

Note that $D_n(\mathcal{F}) = G_n(X_1, \dots, X_n)$. We deduce from McDiarmid's inequality (3.1.22a) that

$$\mathbb{P} \left[D_n(\mathcal{F}) < \mathbb{E}[D_n(\mathcal{F})] + r \right] \geq 1 - e^{-\frac{nr^2}{2C^2}}, \quad \forall r > 0, \quad \forall n \in \mathbb{N}. \quad (3.1.23)$$

Fix a sequence of independent Rademacher random variables $(E_n)_{n \geq 1}$ that are also independent of $(X_n)_{n \geq 1}$. Define the *empirical Rademacher complexity*

$$R_n(\mathcal{F}, X) := \sup_{f \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n E_k f(X_k) \right|.$$

Assume that $R_n(\mathcal{F}, X)$ is measurable for every n . Hence, $R_n(\mathcal{F}, X)$ is a random variable. We define the *Rademacher complexity* to be the deterministic quantity

$$\begin{aligned} \mathcal{R}_n(\mathcal{F}) &= \mathbb{E}_X [R_n(\mathcal{F}, X)] = \frac{1}{2^n} \sum_{\epsilon \in \{-1, 1\}^n} \mathbb{E}_X \left[\sup_{f \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n \epsilon_k f(X_k) \right| \right] \\ &= \mathbb{E}_X \left[\frac{1}{n} \mathbb{E}_E \left[\sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n E_k f(X_k) \right| \right] \right]. \end{aligned}$$

A priori, $\mathcal{R}_n(\mathcal{F})$ depends on the common distribution of the random variables X_n of which we have no special information.

We want to show that the mean $\mathbb{E}[D_n(\mathcal{F})]$ can be controlled by the Rademacher complexity. More precisely, we have

$$\mathbb{E}[D_n(\mathcal{F})] \leq 2\mathcal{R}_n(\mathcal{F}), \quad \forall n. \quad (3.1.24)$$

The proof of the inequality (3.1.24) relies on a symmetrization trick similar to the one used in Section 2.4.

Let $(Y_1, \dots, Y_n)$ be an independent copy of $(X_1, \dots, X_n)$. Then

$$\begin{aligned}
 \mathbb{E}[D_n(\mathcal{F})] &= \mathbb{E}_X \left[\sup_{f \in \mathcal{F}} \frac{1}{n} \left| \sum_{k=1}^n (f(X_k) - \mathbb{E}_{Y_k}[f(Y_k)]) \right| \right] \\
 &= \mathbb{E}_X \left[\sup_{f \in \mathcal{F}} \frac{1}{n} \left| \mathbb{E}_Y \left[\sum_{k=1}^n (f(X_k) - f(Y_k)) \right] \right| \right] \\
 &\leq \mathbb{E}_X \left[\sup_{f \in \mathcal{F}} \frac{1}{n} \mathbb{E}_Y \left[\left| \sum_{k=1}^n (f(X_k) - f(Y_k)) \right| \right] \right] \\
 &\leq \mathbb{E}_X \left[\frac{1}{n} \mathbb{E}_Y \left[\sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n (f(X_k) - f(Y_k)) \right| \right] \right] \\
 &= \mathbb{E}_{X,Y} \left[\frac{1}{n} \sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n (f(X_k) - f(Y_k)) \right| \right].
 \end{aligned}$$

Now observe that the random vectors $(f(X_k) - f(Y_k))$ and $E_k(f(X_k) - f(Y_k))$ have the same distribution so

$$\begin{aligned}
 \mathbb{E}_{X,Y} \left[\frac{1}{n} \sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n (f(X_k) - f(Y_k)) \right| \right] &= \mathbb{E}_{X,Y,E} \left[\frac{1}{n} \sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n E_k(f(X_k) - f(Y_k)) \right| \right] \\
 &\leq 2\mathbb{E}_{X,E} \left[\frac{1}{n} \sup_{f \in \mathcal{F}} \left| \sum_{k=1}^n E_k f(X_k) \right| \right] = 2\mathcal{R}_n(\mathcal{F}).
 \end{aligned}$$

We deduce that

$$\mathbb{P}[D_n(\mathcal{F}) < \mathcal{R}_n(\mathcal{F}) + r] \geq \mathbb{P}[D_n(\mathcal{F}) < \mathbb{E}[D_n(\mathcal{F})] + r] \geq 1 - e^{-\frac{nr^2}{2C^2}}.$$

Hence, $\forall n \in \mathbb{N}$

$$\sum_n \mathbb{P}[D_n(\mathcal{F}) \geq \mathcal{R}_n(\mathcal{F}) + r] \leq \sum_n e^{-\frac{nr^2}{2C^2}} < \infty.$$

Invoking Borel-Cantelli we deduce that for any $r > 0$

$$\mathbb{P}[D_n(\mathcal{F}) \geq \mathcal{R}_n(\mathcal{F}) + r \text{ i.o.}] = 0.$$

In particular, we deduce that $D_n(\mathcal{F}) \rightarrow 0$ if $\mathcal{R}_n(\mathcal{F}) \rightarrow 0$.

Remark 3.1.42. (a) Using McDiarmid inequality as above we can deduce that $R_n(\mathcal{F}, X)$ is also highly concentrated near its mean $\mathcal{R}_n(\mathcal{F})$. We can conclude that there exists $c. > 0$ such that for any $\varepsilon > 0$

$$\mathbb{P}[\mathcal{D}_n(X) < 2\mathcal{R}_n(\mathcal{F}, X) + \varepsilon] < e^{-c n \varepsilon^2}.$$

(b) Note that $0 \leq D_n(\mathcal{F}) \leq C, \forall n$. Hence, if $D_n(\mathcal{F}) \rightarrow 0$ a.s., then $\mathbb{E}[D_n(\mathcal{F})] \rightarrow 0$. Conversely, if $\mathbb{E}[D_n(\mathcal{F})] \rightarrow 0$, then the inequality (3.1.23) implies that $D_n(\mathcal{F}) \rightarrow 0$ a.s.. Hence

$$D_n(\mathcal{F}) \text{ a.s.} \iff \mathbb{E}[D_n(\mathcal{F})] \rightarrow 0.$$

We have shown that $\mathcal{R}_n(\mathcal{F}) \rightarrow 0$ implies $\mathbb{E}[D_n(\mathcal{F})] \rightarrow 0$ and thus $D_n(\mathcal{F}) \rightarrow 0$ a.s..

One can prove that (see [219, Prop. 4.11])

$$\mathbb{E}[D_n(\mathcal{F})] \geq \frac{1}{2} \mathcal{R}_n(\mathcal{F}).$$

Thus, we have a uniform law of large numbers if and only if the Rademacher complexity goes to 0 as $n \rightarrow \infty$. This requires a better understanding of the Rademacher complexity.

A lot is known nowadays. The empirical Rademacher complexity of a family of functions can be estimated in terms of certain entropies. To the family of functions $\mathcal{F}$ we can associate a collection $\mathcal{G}$ of subsets of $\mathbb{R}^N \times \mathbb{R}$, the subgraphs of the functions in $\mathcal{F}$, namely the sets

$$G_f := \{ (x, t) \in \mathbb{R}^N \times \mathbb{R}; \ t \leq f(x) \}, \quad f \in \mathcal{F}.$$

The family $\mathcal{F}$ is said to be a VC-family if the family $\mathcal{G}$ of subgraphs is a VC collection of subsets. These metric entropies of $\mathcal{F}$ can be estimated in terms of the VC dimension and one obtains uniform limits in this fashion. and one concludes that $\mathcal{R}_n(\mathcal{F}) = O(n^{-1/2})$ as $n \rightarrow \infty$. For the rather involved details I refer to [65, 89, 153, 212, 219]. $\square$

3.2. Limit theorems: discrete time

We have seen in the previous section how the Optional Stopping Theorem combined with quite a bit of ingenuity can produce miraculous results. This section is devoted to another miraculous property of martingales, namely, their rather nice asymptotic behavior. The foundational results in this section are all due to J. L. Doob. To convince the reader of the amazing versatility of martingales we have included a large eclectic collection of concrete applications.

3.2.1. Almost sure convergence. Fix a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and a $\mathbb{N}_0$ -filtration $\mathcal{F}_\bullet$ of $\mathcal{F}$. We will investigate the behavior of an $\mathcal{F}_\bullet$ -submartingale $X = (X_n)_{n \in \mathbb{N}_0}$ as $n \rightarrow \infty$. The key to this investigation is *Doob's upcrossing inequality*.

Given real numbers $a < b$ and a sequence of real numbers $\alpha = (\alpha_n)_{n \geq 0}$ we define inductively the sequences

$$(S_k(\alpha) = S_k(\alpha; a, b))_{k \geq 1} \text{ and } (T_k(\alpha) = T_k(\alpha; a, b))_{k \geq 1}$$

in $\mathbb{N}_0 \cup \{\infty\}$ as follows. We set

$$S_1(\alpha) := \inf_{n \geq 0} \{ \alpha_n \leq a \}, \quad T_1(\alpha) := \inf_{n \geq S_1} \{ \alpha_n \geq b \}.$$

Thus, S_1 is the first moment the sequence α drops below a , and T_1 is the first moment after S_1 when the sequence α crosses the upper level b . We then define inductively

$$S_{k+1}(\alpha) := \inf_{n \geq T_k} \{ \alpha_n \leq a \}, \quad T_{k+1}(\alpha) := \inf_{n \geq S_{k+1}} \{ \alpha_n \geq b \},$$

where we set $\inf \emptyset = \infty$; see Figure 3.2.




Figure 3.2. Up/downcrossing of the interval $[a, b]$.

The terms S_k are called *downcrossing times* while the terms T_k are called the *upcrossing times* of the sequence $(\alpha_n)_{n \geq 0}$. We define the *upcrossing numbers*

$$U_n([a, b], \alpha) := \#\{k \in \mathbb{N}; T_k(\alpha) \leq n\}, \quad n \in \mathbb{N}. \quad (3.2.1)$$

The sequence of nonnegative integers $U_n([a, b], \alpha)$ is nonincreasing and thus it has a, possibly infinite, limit

$$U_\infty([a, b], \alpha) := \lim_{n \rightarrow \infty} U_n([a, b], \alpha).$$

The importance of the upcrossing numbers in convergence problems is explained by the following elementary but rather clever result. In Exercise 3.8 we ask the reader to provide a proof.

Lemma 3.2.1. *Suppose that $\alpha = (\alpha_n)_{n \geq 0}$ is a sequence of real numbers. Then the following statements are equivalent.*

- (i) *The sequence α has a limit (possibly infinite).*
- (ii) *For any rational numbers $a < b$ the total number of upcrossings $U_\infty([a, b], \alpha)$ is finite.* $\square$

Suppose now that $(X_n)_{n \in \mathbb{N}_0}$ is a process adapted to the filtration $\mathcal{F}_\bullet$. Then, for any $k \in \mathbb{N}$, the down/up-crossing times $S_k(X)$ and $T_k(X)$ are stopping times.

Theorem 3.2.2 (Doob's upcrossing inequality). *Assume that $X = (X_n)_{n \in \mathbb{N}_0}$ is a submartingale. Then for any real numbers $a < b$ we have*

$$(b - a)\mathbb{E}[U_n([a, b], X)] \leq \mathbb{E}[(X_n - a)^+] - \mathbb{E}[(X_0 - a)^+], \quad x^+ := \max(x, 0). \quad (3.2.2)$$

Proof. Since $(X - a)^+$ is a submartingale and

$$U_n([a, b], X) = U_n([0, b - a], (X - a)^+),$$

we see that it suffices to prove the result in the special case $X \geq 0$ and $a = 0 < b$. In other words, it suffices to prove that if $X \geq 0$, then

$$b\mathbb{E}[U_n([0, b], X)] \leq \mathbb{E}[(X_n - X_0)]. \quad (3.2.3)$$

The key fact underlying this inequality is the existence of a submartingale Y that lies above the random process $U_n([0, b], X)$ and, in the mean, below the process X .

Consider the predictable process

$$H = \sum_{k=1}^{\infty} \mathbf{I}_{[S_k(X), T_k(X)]},$$

i.e.,

$$H_n = \sum_{k=1}^{\infty} \mathbf{I}_{\{S_k(X) < n \leq T_k(X)\}}.$$

Since the intervals

$$(S_1(X), T_1(X)], (S_2(X), T_2(X)], \dots,$$

are pairwise disjoint (when finite) we have $H_n \in \{0, 1\}, \forall n$. Set $Y_n := (H \cdot X)_n$.

In stock market terms, think of the following investing strategy. Start buying a stock when its price cost hits zero, and sell it at the end the trading day. Continue buying (and selling) the stock as long as its price at the start of the trading day is below b . Once the price crosses b stop buying and wait until the price hits 0 again. The price of the stock at the beginning of the n -th trading day is X_{n-1} and changes to X_n at the end of the n -th trading day. Then Y_n is the profit following this

strategy at the end of n days. Clearly the profit will be at least as big as $b \times$ the number of upcrossings of the interval $(0, b)$. This is the content of the following fundamental inequality.

$$\boxed{Y_n \geq bU_n([0, b], X)}. \quad (3.2.4)$$

Here is a formal proof of this inequality. Let $M := U_n([0, b], X)$. Then

$$\begin{aligned} Y_n &= \sum_{j=1}^n H_j \cdot (X_j - X_{j-1}) = \sum_{k=1}^M \sum_{j=S_k(X)+1}^{T_k(X)} (X_j - X_{j-1}) + \sum_{j=S_{M+1}+1}^n (X_j - X_{j-1}) \\ &= \sum_{k=1}^M (X_{T_k} - X_{S_k}) + \mathbf{I}_{\{S_{M+1} < n\}} (X_n - X_{S_{M+1}}) \\ &\quad \text{(use the fact that } X_{S_{M+1}} = 0 \text{ and } X_n \geq 0) \\ &\geq \sum_{k=1}^M \underbrace{(X_{T_k} - X_{S_k})}_{\geq b} \geq bM = bU_n([0, b], X). \end{aligned}$$

Hence

$$b\mathbb{E}[U_n([0, b], X)] \leq \mathbb{E}[Y_n], \quad \forall n \in \mathbb{N}.$$

Note that the inequality (3.2.4) does not rely on the fact that X is a submartingale.

The process (H_n) is predictable and thus

$$\begin{aligned} \mathbb{E}[Y_k - Y_{k-1} | \mathcal{F}_{k-1}] &= \mathbb{E}[H_k(X_k - X_{k-1}) | \mathcal{F}_{k-1}] \\ &= H_k \mathbb{E}[X_k - X_{k-1} | \mathcal{F}_{k-1}]. \end{aligned}$$

Since X is a *submartingale* we deduce

$$\mathbb{E}[X_k - X_{k-1} | \mathcal{F}_{k-1}] \geq 0.$$

On the other hand $0 \leq H_k \leq 1$ so that

$$H_k \mathbb{E}[X_k - X_{k-1} | \mathcal{F}_{k-1}] \leq \mathbb{E}[X_k - X_{k-1} | \mathcal{F}_{k-1}].$$

Hence

$$\mathbb{E}[Y_k - Y_{k-1}] \leq \mathbb{E}[X_k - X_{k-1}]. \quad (3.2.5)$$

We deduce

$$b\mathbb{E}[U_n([0, b], X)] \leq \mathbb{E}[Y_n] = \sum_{k=1}^n \mathbb{E}[Y_k - Y_{k-1}] \leq \mathbb{E}[X_n - X_0].$$

□

Remark 3.2.3. We should ponder why the inequality (3.2.5) is miraculous. We know that $H_k \in [0, 1]$ so, whenever $X_k \leq X_{k-1}$, i.e., the price of stock goes down, we have

$$X_k - X_{k-1} \leq H_k(X_k - X_{k-1}) = Y_k - Y_{k-1}.$$

The inequality (3.2.5) shows that *this is not the expected behavior*. The fact that $X_\bullet$ is a *submartingale* biases the price in favor of increase. That is the reason why (3.2.5) holds. □

Theorem 3.2.4 (Submartingale Convergence Theorem). *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale satisfying*

$$\sup_{n \in \mathbb{N}_0} \mathbb{E}[X_n^+] < \infty. \quad (3.2.6)$$

Then X_n converges almost surely to an integrable random variable X_∞ .

Remark 3.2.5. Observe that since X_n is a submartingale we have

$$\mathbb{E}[X_0] \leq \mathbb{E}[X_n] = \mathbb{E}[X_n^+] - \mathbb{E}[X_n^-], \quad x^- = \max(-x, 0),$$

so that

$$\sup_{n \in \mathbb{N}_0} \mathbb{E}[X_n^-] < \infty$$

showing that (3.2.6) is equivalent to

$$\sup_{n \in \mathbb{N}_0} \mathbb{E}[|X_n|] < \infty. \quad (3.2.7)$$

□

Proof. Set

$$M := \sup_{n \in \mathbb{N}_0} \mathbb{E}[|X_n|].$$

Now let $a, b \in \mathbb{Q}$, $a < b$. Doob's upcrossing inequality shows that, for all $n \geq 1$, we have

$$(b - a)\mathbb{E}[N_n(a, b, X_\bullet)] \leq \mathbb{E}[(X_n - a)^+] \leq |a| + \mathbb{E}[|X_n|] \leq |a| + M.$$

Letting $n \rightarrow \infty$ we deduce $\mathbb{E}[N_\infty([a, b], X_\bullet)] < \infty$, and thus $N_\infty([a, b], X_\bullet) < \infty$ a.s.. By removing a countable family of negligible sets (one for each pair of rational numbers a, b , $a < b$) we deduce that there exists a negligible set $\mathcal{N} \subset \Omega$ such that $\forall \omega \in \Omega \setminus \mathcal{N}$ we have

$$N_\infty([a, b], X_\bullet(\omega)) < \infty, \quad \forall a, b \in \mathbb{Q}, \quad a < b.$$

Lemma 3.2.1 implies that the sequence $X_\bullet$ converges a.s. to a random variable X_∞ . The integrability of X_∞ follows from Fatou's lemma

$$\mathbb{E}[|X_\infty|] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[|X_n|] < \infty.$$

□

Corollary 3.2.6. *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a nonnegative supermartingale. Then X_n converges a.s. to an integrable random variable X_∞ . In particular, any nonnegative martingale has an integrable a.s. limit.*

Proof. Observe that $Y_n = -X_n$ is a submartingale and $Y_n^+ = 0$. The result now follows from the Submartingale Convergence Theorem. □

Example 3.2.7 (Galton-Watson/branching processes). ² Consider again the branching process in Example 3.1.8 with reproduction law $\mu \in \text{Prob}(\mathbb{N}_0)$ and mean m ,

$$0 < m := \sum_{n \geq 0} n\mu[n] < \infty.$$

²**To the post pandemic reader.** I wrote a substantial part of this book during the great covid pandemic. I even taught this example to a group of masked students that were numbed by the news about the R -factor. The mean m is a close relative of this R -factor. This example explains the desirability of $R < 1$.

As explained in Example 3.1.8, the sequence

$$W_n = \frac{1}{m^n} Z_n, \quad n \in \mathbb{N}_0,$$

is a nonnegative martingale so, according to Corollary 3.2.6, it converges a.s. to an integrable random variable W_∞ .

If $m < 1$, the original sequence $Z_n = m^n W_n$ converges a.s. and in mean to 0. Moreover

$$\mathbb{E}[Z_n] = m^n \mathbb{E}[Z_0] = m^n \ell.$$

Thus, the expected population decays *exponentially* to zero. Something more dramatic holds.

Since $Z_n \geq 1$ if $Z_n > 0$ we deduce

$$\mathbb{P}[Z_n > 0] = \mathbb{P}[Z_n \geq 1] \leq \mathbb{E}[Z_n] = \ell m^n.$$

Hence

$$\sum_{n \geq 0} \mathbb{P}[Z_n > 0] < \infty.$$

The Borell-Cantelli Lemma implies that $\mathbb{P}[Z_n > 0 \text{ i.o.}] = 0$.

Thus, a population of bacteria that have on average less than one successor will die out, i.e., with probability 1 there exists $n \in \mathbb{N}$ such that $Z_n = 0$. If we set

$$E_n := \{Z_k = 0, \quad \forall k \geq n\} = \{Z_n = 0\},$$

then the event

$$E = \bigcup_{n \geq 0} E_n$$

is called the *extinction event*. Note that

$$E_0 \subset E_1 \subset \cdots \subset E_n \subset \cdots.$$

The probability of E is called *extinction probability*. We see that when $m < 1$, the extinction probability is 1. $\square$

Example 3.2.8 (Lévy's equivalence theorem). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent, mean zero random variables defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that the sums

$$S_n := X_1 + \cdots + X_n$$

converge in probability to an a.s.-finite random variable S_∞ . Assume that

$$|S_n(\omega)| < \infty, \quad \forall n \in \mathbb{N} \cup \{\infty\}, \quad \forall \omega \in \Omega.$$

We want to use the martingale convergence theorem to prove the *Lévy's equivalence theorem*, namely, that the convergence in distribution of S_n implies a.s. convergence. I will first prove that S_n converges in probability following the strategy in [98, Sec.1.3].

Lévy's Continuity Theorem 2.2.31 implies that

$$\Phi_{S_n}(t) = \mathbb{E}[e^{itS_n}]$$

converges for any $t \in \mathbb{R}$ to $\Phi_{S_\infty}(t) = \mathbb{E}[e^{itS_\infty}]$. Moreover, Proposition 2.2.32 implies that there exists δ, c such that

$$|\Phi_{S_n}(t)| > \frac{1}{c}, \quad \forall |t| \leq \delta, n \in \mathbb{N}.$$

Arguing as in Example 3.1.6 we deduce that for any $|t| \leq \delta$ we have a martingale (see

$$Y_n(t) = \frac{1}{\Phi_{S_n}(t)} e^{itS_n}$$

This is obviously bounded in L^1 and thus, for any $t \in [-\delta, \delta]$, it converges a.s.. Note that

$$\Phi_{S_n}(t) \rightarrow \Phi_{S_\infty}(t), \quad \forall t$$

since S_n converges to S_∞ in distribution. We deduce that for any $t \in [-\delta, \delta]$ there exists a negligible subset $\mathcal{N}_t \in \mathcal{S}$ such that

$$\forall \omega \in \Omega \setminus \mathcal{N}_t : e^{itS_n(\omega)} \rightarrow e^{itS_\infty(\omega)} \quad \text{as } n \rightarrow \infty. \quad (3.2.8)$$

We want to prove that this implies that $S_n \rightarrow S_\infty$ in probability.

Since $e^{i\delta S_n} \rightarrow e^{i\delta S_\infty}$ on $\Omega \setminus \mathcal{N}_\delta$, we deduce that there exists a sequence of $\mathbb{Z}$ -valued random variables $(R_n)_{n \in \mathbb{N}}$ such that

$$\delta S_n(\omega) + 2\pi R_n(\omega) \rightarrow \delta S_\infty(\omega), \quad \forall \omega \in \Omega \setminus \mathcal{N}_\delta. \quad (3.2.9)$$

For example, the random variables

$$R_n = \left\lfloor \frac{\delta}{2\pi} (S_\infty - S_n) \right\rfloor$$

have this property. We deduce that for any $t \in [-1, 1]$, and any $\omega \in \Omega \setminus \mathcal{N}_{t\delta}$ we have

$$\Psi_n(t, \omega) := e^{2\pi i t R_n(\omega)} \rightarrow 1. \quad (3.2.10)$$

Hence

$$\lim_{n \rightarrow \infty} \Phi_{R_n}(2\pi t) = 1, \quad \forall t \in [-1, 1].$$

Since $t \mapsto \Phi_{R_n}(2\pi t)$ is 1-periodic we deduce that the above equality holds for any $t \in \mathbb{R}$. Lévy's continuity theorem implies that the random variables R_n converge in distribution to 0 as $n \rightarrow \infty$.

The variables R_n are $\mathbb{Z}$ -valued so $\mathbb{P}[|R_n| > \varepsilon] = 1 - \mathbb{P}[R_n = 0] \rightarrow 0$ for any $\varepsilon \in (0, 1)$, i.e., R_n converges in probability to 0. Since $\delta S_n + 2\pi R_n$ converges a.s. to δS_∞ we deduce from Corollary 1.3.63 that

$$\delta S_n = (\delta S_n + 2\pi R_n) - 2\pi R_n$$

converges in probability to δS_∞ . Let us prove that $\delta S_n \rightarrow \delta S_\infty$ a.s..

Given that the random variables R_n are integer valued we can say a bit more. For $n \in \mathbb{N}$ and $m \in \mathbb{Z}$ we set

$$\Omega_{n,m} := \{R_n = m\}, \quad p_{n,m} := \mathbb{P}[\Omega_{n,m}].$$

If $p_{n,m} = 0$, then we modify R_n on $\Omega_{n,m}$ by setting it equal to 0 on this set. Thus we can assume that for $m \neq 0$ we have either $p_{n,m} > 0$ or $\Omega_{n,m} = \emptyset$. Denote by $\mathcal{F}$ the sigma-algebra generated by the (modified) R_n 's. Thus a set is $\mathcal{F}$ -measurable iff it is a union of sets of type $\Omega_{n,m}$. This shows that if $\mathcal{N} \in \mathcal{F}$ is a nonempty negligible, then $R_n(\omega) = 0, \forall n$.

The negligible sets $\mathcal{N}_{t\delta}, t \in [-1, 1]$ in (3.2.8) and (3.2.10) belong to $\mathcal{F}$. Hence $R_n = 0$ on $\mathcal{N}_{t\delta}$ so for $\omega \in \mathcal{N}_{t\delta}$ we have

$$e^{2\pi i t R_n(\omega)} = 1, \quad \forall n.$$

We deduce that for any $t \in [-1, 1]$ and any $\omega \in \Omega$ we have

$$\lim_{n \rightarrow \infty} e^{2\pi i t R_n(\omega)} = 1.$$

Clearly the above equality holds for any $t \in \mathbb{R}$. Fix $\omega \in \Omega$ and set

$$x_n = x_n(\omega) := \delta R_n(\omega).$$

At this point I want to invoke the following technical result.

Lemma 3.2.9. *Suppose that (x_n) is a sequence of integers such that for any $t \in \mathbb{R}$ the sequence e^{itx_n} converges to 1. Then the sequence (x_n) converges to 0.*

Proof. I will present two proofs.

1st Proof. Set $L_n(t) = e^{itx_n}$. The functions $L_n(t)$ converge to 1 for any $t \in \mathbb{R}$ and Egorov's theorem [45, Prop. 3.1.4] implies that there exists a Borel measurable set $B \subset [-1, 1]$ such that $\lambda[B] > 1$ and L_n converges to 1 uniformly on B . Since $L_n(t)/L_n(s) = L_n(t-s)$ we deduce that $L_n(t)$ converges uniformly to 1 on the set

$$B - B = \{t - s; t, s \in B\}.$$

A 1920 result of Hugo Steinhaus [200]³ shows that there exists $\varepsilon > 0$ such that

$$[-\varepsilon, \varepsilon] \subset B - B.$$

Hence $L_n(t)$ converges to 1 uniformly on $[-\varepsilon, \varepsilon]$. This implies that the sequence (x_n) is bounded.

We argue by contradiction. If this were not the case the sequence (x_n) would contain a monotone subsequence with limit $\pm\infty$. For ease of notation I will assume that $x_n \rightarrow \infty$ and (x_n) is positive and strictly increasing. Then

$$\bigcup_n [-\varepsilon x_n, \varepsilon x_n] = \mathbb{R}.$$

Set $t_n := \frac{\pi}{x_n}$. Then $t_n \in [-\varepsilon, \varepsilon]$ for n sufficiently large and $t_n \rightarrow 0$ as $n \rightarrow \infty$. Since L_n converges uniformly to 1 on $[-\varepsilon, \varepsilon]$ we deduce that

$$-1 = e^{\pi i} = L_n(t_n) \rightarrow 1.$$

Hence the sequence (x_n) is bounded.

If (x_n) had a subsequence (x_{n_k}) that converged to $N \in \mathbb{N} \setminus 0$, then $x_{n_k} = N$ for all k sufficiently large and, if we set $t = \frac{\pi}{N}$ then we deduce that

$$-1 = e^{\pi i} = L_{n_k}(t).$$

for all k sufficiently large. We deduce that if a subsequence of (x_n) converges, then its limit has to be zero.

2nd Proof. This is based on an idea in [186, Lemma 45.8]. The Dominated Convergence Theorem implies that for any $r > 0$ we have

$$1 = \frac{1}{2r} \lim_{n \rightarrow \infty} \int_{-r}^r e^{itx_n} dt = \lim_{n \rightarrow \infty} \frac{\sin(rx_n)}{rx_n},$$

where we set $\frac{\sin 0}{0} := 1$.

No subsequence (x_{n_k}) can have a nonzero limit because if that were the case we would have

$$\lim_{k \rightarrow \infty} \frac{\sin(rx_{n_k})}{rx_{n_k}} < 1.$$

Hence x_n has to converge to 0. □

³I refer to [64, Prop.3.4.3] for a more modern proof.

We deduce that $R_n(\omega) \rightarrow 0$, for any ω . Using this in (3.2.9) we deduce that $S_n \rightarrow S_\infty$ a.s.. $\square$

Remark 3.2.10. Lévy's equivalence theorem shows that a random series with mean zero *independent* terms converges a.s. if and only if it converges in distribution. The partial sums S_n form a martingale. However, there exist martingales that converge in probability, but not a.s..

Here is one such example, [68, Example 4.2.14]. Consider the following random walk $(X_n)_{n \geq 0}$ on $\mathbb{Z}$ where you should think of X_n as the location at time n . We set $X_0 = 0$. If X_{n-1} is known, then

$$\mathbb{P}[X_n = \pm 1 \mid X_{n-1} = 0] = \frac{1}{2n}, \quad \mathbb{P}[X_n = 0 \mid X_{n-1} = 0] = 1 - \frac{1}{n},$$

$$\mathbb{P}[X_n = 0 \mid X_{n-1} \neq 0] = 1 - \frac{1}{n}, \quad \mathbb{P}[X_n = nX_{n-1} \mid X_{n-1} \neq 0] = \frac{1}{n}.$$

The existence of such a process is guaranteed by Kolmogorov's theorem.

Denote by $\mathcal{F}_n$ the sigma-algebra generated by the random variables $X_0, X_1, \dots, X_n$. From the construction we deduce that $\mathbb{E}[X_n \mid \mathcal{F}_{n-1}] = X_{n-1}$ so (X_n) is a martingale with respect to the filtration $\mathcal{F}_n$. Let $p_n := \mathbb{P}[X_n \neq 0]$. Note that

$$\begin{aligned} p_n &= \mathbb{P}[X_n \neq 0 \mid X_{n-1} = 0] \mathbb{P}[X_{n-1} = 0] + \mathbb{P}[X_n \neq 0 \mid X_{n-1} \neq 0] \mathbb{P}[X_{n-1} \neq 0] \\ &= \frac{1}{n} (1 - p_{n-1}) + \frac{1}{n} p_{n-1} = \frac{1}{n}. \end{aligned}$$

Hence

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_n \neq 0] = 0,$$

so that X_n converges in probability to 0. To show it does not converge a.s. it suffices to show that it does not converge a.s. to 0.

Denote by F_n the event $\{X_n \neq 0\}$. The random variables X_n have integer values so $F_n = \{|X_n| \geq 1\}$. Note that $F_n \in \mathcal{F}_n$ and

$$\mathbb{E}[\mathbf{I}_{F_n} \mid \mathcal{F}_{n-1}] = \frac{1}{n} \mathbf{I}_{\{X_{n-1}=0\}} + \frac{1}{n} \mathbf{I}_{\{X_{n-1} \neq 0\}} = \frac{1}{n}.$$

Hence

$$\sum_{n \geq 1} \mathbb{E}[\mathbf{I}_{F_n} \mid \mathcal{F}_{n-1}] = \sum_{n \geq 1} \frac{1}{n} = \infty.$$

The conditional Borel-Cantelli result in Exercise 3.12 implies that

$$\mathbb{P}[|X_n| \geq 1 \text{ i.o.}] = \mathbb{P}[F_n \text{ i.o.}] = 1.$$

Thus (X_n) does not converge a.s. to 0.

Recently (2021) Iosif Pinelis gave another beautiful example of martingale converging in probability but not a.s.. Here is briefly the construction.

Choose a sequence of independent geometric random variables

$$(T_n)_{n \geq 1}, \quad T_n \sim \text{Geom}(p_n).$$

We perform the following delayed and frequently stopped random walk on $\mathbb{Z}$. We start at $X_0 = 0$ and we wait for T_1 moments and we begin a standard random walk on $\mathbb{Z}$ until we first return to the origin. At that moment take a brake lasting T_2 moments and begin the standard walk until we return

back to the origin etc. Denote by X_n the location after n moments. Then (X_n) is a martingale (with respect to an appropriate filtration). Moreover, if

$$\sum_{n \geq 1} \sqrt{p_n} < \infty,$$

then X_n converges in probability to 0 but not a.s.. For details I refer to [170]. $\square$

Example 3.2.11. The assumptions in the (sub)martingale convergence theorem are not strong enough to guarantee L^1 -convergence. The following example shows what can happen.

Consider the standard random walk on $\mathbb{Z}$ that starts at 1. Each second the traveler takes a size 1 step forward or back with equal probability. More precisely, consider a sequence of i.i.d. Rademacher random variables $(X_n)_{n \in \mathbb{N}}$, $\mathbb{P}[X_1 = 1] = \mathbb{P}[X_n = -1] = \frac{1}{2}$. Then the sequence

$$S_0 = 1, \quad S_n = 1 + X_1 + \cdots + X_n, \quad n \in \mathbb{N}$$

is a martingale describing the evolution of the walk. Denote by N the first moment the walk reaches the origin, i.e.,

$$N := \inf \{ n \in \mathbb{N}; \quad S_n = 0 \}.$$

Observe that $N < \infty$ a.s.; see Exercise 3.23. Consider the random walk stopped at N

$$Y_n := S_n^N = S_{n \wedge N}.$$

From the Optional Stopping Theorem 3.1.24 we deduce that Y_n is a martingale which, by construction, is also nonnegative. Clearly $Y_n \rightarrow 0$ a.s. since $N < \infty$ a.s.. This convergence is not L^1 since

$$\mathbb{E}[Y_n] = \mathbb{E}[Y_0] = 1, \quad \forall n \in \mathbb{N}. \quad \square$$

3.2.2. Uniform integrability. We will describe in this subsection necessary and sufficient conditions guaranteeing that a sequence that converges in probability also converges in the mean. We begin with a basic fact.

Lemma 3.2.12. Let $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X| \mathbf{I}_{\{|X| \geq n\}}] = 0.$$

Proof. The sequence $Z_n := |X| \mathbf{I}_{\{|X| > n\}}$ converges a.s. to 0 and $|Z_n| \leq |X|$, $\forall n$. The desired conclusion now follows from the Dominated Convergence theorem. $\square$

Definition 3.2.13 (Uniform integrability). A collection $\mathfrak{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$ is called *uniformly integrable* (or UI for brevity) if

$$\lim_{r \rightarrow \infty} \mathbb{E}[|X| \mathbf{I}_{\{|X| \geq r\}}] = 0 \quad \text{uniformly in } X \in \mathfrak{X}. \quad (\text{UI}_1) \quad \square$$

Example 3.2.14. Let $\mathfrak{X} \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ be a family of random variables such that there exists $Z_0 \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ with the property

$$|X| \leq |Z_0| \quad \text{a.s.,} \quad \forall X \in \mathfrak{X}.$$

Then $\mathfrak{X}$ is UI.

Indeed, for any $X \in \mathfrak{X}$ we have $|X| \mathbf{I}_{\{|X| > r\}} \leq |Z_0| \mathbf{I}_{\{|Z_0| > r\}}$. The conclusion now follows from Lemma 3.2.12. $\square$

Remark 3.2.15. (a) Let $\mathfrak{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$\chi(r) = \chi(r, \mathfrak{X}) := \sup_{X \in \mathfrak{X}} \mathbb{E}[|X| \mathbf{I}_{\{|X| \geq r\}}]. \quad (3.2.11)$$

Then $\mathfrak{X}$ is uniformly integrable iff $\lim_{r \rightarrow \infty} \chi(r) = 0$.

(b) A uniformly integrable family $\mathfrak{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$ is bounded in the L^1 -norm, i.e.,

$$\sup_{X \in \mathfrak{X}} \mathbb{E}[|X|] < \infty.$$

Indeed, $\forall X \in \mathfrak{X}$ and r is sufficiently large so that $\chi(r) < 1$, we have

$$\mathbb{E}[|X|] = \mathbb{E}[|X| \mathbf{I}_{\{|X| < r\}}] + \mathbb{E}[|X| \mathbf{I}_{\{|X| \geq r\}}] \leq r + \chi(r) < \infty.$$

□

Theorem 3.2.16. Let $\mathfrak{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then the following statements are equivalent.

- (i) $\mathfrak{X}$ is uniformly integrable.
- (ii) The family $\mathfrak{X}$ is L^1 -bounded and, for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that, for any $X \in \mathfrak{X}$ and any $S \in \mathcal{S}$, we have

$$\mathbb{P}[S] \leq \delta \Rightarrow \mathbb{E}[|X| \mathbf{I}_S] = \int_S |X(\omega)| \mathbb{P}[d\omega] < \varepsilon. \quad (\mathbf{UI}_2)$$

Proof. (i) $\Rightarrow$ (ii) Fix $\varepsilon > 0$. There exists $r_\varepsilon > 0$ such that $\chi(r_\varepsilon) < \varepsilon/2$, with $\chi(r)$ defined by (3.2.11). Next, fix $\delta > 0$ such that $\delta r_\varepsilon < \frac{\varepsilon}{2}$. Then, for any $X \in \mathfrak{X}$ and any $S \in \mathcal{S}$ such that $\mathbb{P}[S] < \delta$, we have

$$\begin{aligned} \mathbb{E}[|X| \mathbf{I}_S] &= \mathbb{E}[|X| \mathbf{I}_{S \cap \{|X| < r_\varepsilon\}}] + \mathbb{E}[|X| \mathbf{I}_{S \cap \{|X| \geq r_\varepsilon\}}] \\ &\leq r_\varepsilon \mathbb{P}[S] + \mathbb{E}[|X| \mathbf{I}_{\{|X| \geq r_\varepsilon\}}] \leq \delta r_\varepsilon + \chi(r_\varepsilon) < \varepsilon. \end{aligned}$$

(ii) $\Rightarrow$ (i) Set

$$B := \sup_{X \in \mathfrak{X}} \mathbb{E}[|X|] < \infty.$$

Markov's inequality implies that for $r > 0$ we have

$$\mathbb{P}[|X| > r] \leq \frac{B}{r}, \quad \forall X \in \mathfrak{X}.$$

Fix $\varepsilon > 0$ and $r_\varepsilon > 0$ such that $\frac{B}{r_\varepsilon} < \delta(\varepsilon)$. Then $\mathbb{P}[|X| > r_\varepsilon] < \delta(\varepsilon)$, $\forall X \in \mathfrak{X}$. Assumption (ii) implies

$$\chi(r_\varepsilon) = \sup_{X \in \mathfrak{X}} \mathbb{E}[|X| \mathbf{I}_{\{|X| > r_\varepsilon\}}] < \varepsilon.$$

□

Remark 3.2.17. (a) We should draw attention to the qualitatively different conditions $(\mathbf{UI}_1)$ and $(\mathbf{UI}_2)$.

Condition $(\mathbf{UI}_1)$ involves only the probability distributions of the random variables $X \in \mathfrak{X}$ with no mention of the probability space on which they are defined. On the other hand, condition $(\mathbf{UI}_2)$ makes explicit reference to their domain of definition $(\Omega, \mathcal{S}, \mathbb{P})$. Condition $(\mathbf{UI}_2)$ is usually referred to as *uniform absolute continuity*.

(b) An *atom* of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$ is a measurable set $S \in \mathcal{S}$ such that $\mathbb{P}[S] > 0$ and for every measurable subset $S' \subset S$, $\mathbb{P}[S'] = 0$ or $\mathbb{P}[S'] = \mathbb{P}[S]$. The L^1 -boundedness condition

follows from (UI₂) alone if the probability measure $\mathbb{P}$ has *no atoms*. For a proof of this fact we refer to [21, Prop.4.5.3]. $\square$

Recall (see Exercise 2.67) that a *Young function* is a continuous, nondecreasing convex function $f : [0, \infty) \rightarrow [0, \infty)$ such that

$$f(0) = 0, \quad \lim_{x \rightarrow \infty} f(x) = \infty.$$

The Young function f is called *superlinear* if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{x} = \infty.$$

Theorem 3.2.18. *Let $\mathfrak{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$. Then the following statements are equivalent.*

(i) $\mathfrak{X}$ is UI.

(ii)

$$\lim_{r \rightarrow \infty} \sup_{X \in \mathfrak{X}} \int_r^\infty \mathbb{P}[|X| > x] dx = 0.$$

(iii) *There exists a superlinear Young function $f : [0, \infty) \rightarrow [0, \infty)$ such that $f(0) = 0$ such that*

$$\sup_{X \in \mathfrak{X}} \mathbb{E}[f(|X|)] < \infty. \quad (3.2.12)$$

Proof. (i) $\iff$ (ii) Proposition 1.3.43 shows that

$$\int_r^\infty \mathbb{P}[|X| > x] dx = \mathbb{E}[|X| \mathbf{I}_{\{|X| > r\}}], \quad \forall X \in \mathfrak{X}.$$

(ii) $\Rightarrow$ (iii) Set

$$h(r) := \sup_{X \in \mathfrak{X}} \int_r^\infty \mathbb{P}[|X| > x] dx.$$

Note that $h(0) \leq r + h(r) < \infty$. Since $h(r) = o(1)$ as $r \rightarrow \infty$ we can find

$$0 = r_0 < r_1 < r_2 < \dots$$

such that

$$h(r_n) \leq \frac{h(0)}{2^n}, \quad \forall n \in \mathbb{N}.$$

Now define

$$g(r) := \sum_{n \geq 0} \mathbf{I}_{[r_n, \infty)}(r), \quad f(x) = \int_0^x g(r) dr.$$

Note that $g(r)$ is nondecreasing and $\lim_{r \rightarrow \infty} g(r) = \infty$. This shows that f is increasing convex and superlinear. Using the Fubini-Tonelli theorem as in the proof of Proposition 1.3.43 we deduce that for any $X \in \mathfrak{X}$ we have

$$\begin{aligned} \mathbb{E}[f(|X|)] &= \mathbb{E}\left[\int_0^{|X|} g(r) dr\right] = \mathbb{E}\left[\sum_{n \geq 0} \int_{r_n}^\infty \mathbf{I}_{|X| > r_n}(x) dx\right] \\ &\leq \sum_{n \geq 0} h(r_n) \leq h(0) \sum_{n \geq 0} \frac{1}{2^n}. \end{aligned}$$

(iii) $\Rightarrow$ (i) For every $n \in \mathbb{N}$ there exists $r_n > 0$ such that

$$\forall x : x > r_n \Rightarrow x < \frac{f(x)}{n}.$$

We deduce that for any $X \in \mathfrak{X}$ we have

$$\mathbb{E}[|X|\mathbf{I}_{|X|>r_n}] \leq \frac{1}{n}\mathbb{E}[f(|X|)\mathbf{I}_{|X|>r_n}] \leq \frac{1}{n}\mathbb{E}[f(|X|)].$$

The conclusion now follows from (3.2.12). $\square$

The equivalence (i) $\Longleftrightarrow$ (iii) is sometimes referred to as the *de la Vallée-Poussin theorem*.

If in the above theorem we choose $f(r) = r^p$, $p > 1$, we obtain the following result.

Corollary 3.2.19. *Let $\mathfrak{X} \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ be a family of random variables such that there exist $p \in (1, \infty)$ with the property*

$$\sup_{X \in \mathfrak{X}} \mathbb{E}[|X|^p] < \infty.$$

Then $\mathfrak{X}$ is UI. $\square$

Corollary 3.2.20. *Let $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and suppose that $(\mathcal{F}_i)_{i \in I}$ is a family of sigma-subalgebras of $\mathcal{S}$. Set $X_i := \mathbb{E}[X \mid \mathcal{F}_i]$, $i \in I$. Then the family $(X_i)_{i \in I}$ is UI.*

Proof. First Method. Lemma 3.2.12 shows that the family $\{X\}$ consisting of the integrable random variable X is uniformly integrable. Hence

$$|X_i| = |\mathbb{E}[X \mid \mathcal{F}_i]| \leq \mathbb{E}[|X| \mid \mathcal{F}_i].$$

Theorem 3.2.18 implies that there exists a superlinear Young function $f : [0, \infty) \rightarrow [0, \infty)$ such that $\mathbb{E}[f(X)] < \infty$. Since f is increasing and convex we deduce from the conditional Jensen inequality in Theorem 1.4.12(ix) that

$$f(|X_i|) \leq f(\mathbb{E}[|X| \mid \mathcal{F}_i]) \leq \mathbb{E}[f(|X|) \mid \mathcal{F}_i].$$

Taking the expectations of both sides of this inequality we deduce

$$\mathbb{E}[f(|X_i|)] \leq \mathbb{E}[f(|X|)], \quad \forall i \in I.$$

Using Theorem 3.2.18 again we deduce that the family $(X_i)_{i \in I}$ is UI.

Second Method. Fix $\varepsilon > 0$ and then choose $\delta = \delta(\varepsilon) > 0$ such that

$$\forall S \in \mathcal{S}, \mathbb{P}[S] < \delta \Rightarrow \mathbb{E}[|X|\mathbf{I}_S] < \varepsilon. \quad (3.2.13)$$

Choose $C > 0$ sufficiently large so $C^{-1}\mathbb{E}[|X|] < \delta$.

For each i choose a version X_i of $\mathbb{E}[X \mid \mathcal{F}_i]$. Then

$$|X_i| \leq \mathbb{E}[|X| \mid \mathcal{F}_i] \text{ a.s. and } \mathbb{E}[|X_i|] \leq \mathbb{E}[|X|].$$

Hence

$$\mathbb{P}[|X_i| > C] \leq C^{-1}\mathbb{E}[|X_i|] \leq C^{-1}\mathbb{E}[|X|] < \delta.$$

Then

$$\mathbb{E}[|X_i|\mathbf{I}_{\{|X_i|>C\}}] \leq \mathbb{E}[\mathbb{E}[|X| \mid \mathcal{F}_i]\mathbf{I}_{|X_i|>C}] = \mathbb{E}[|X|\mathbf{I}_{\{|X_i|>C\}}] \stackrel{(3.2.13)}{\leq} \varepsilon.$$

$\square$

The next result clarifies the importance of the uniform integrability condition.

Theorem 3.2.21 (Lebesgue-Vitali). *Consider a sequence (X_n) in $L^1(\Omega, \mathcal{S}, \mathbb{P})$ that converges in probability to X . Then the following statements are equivalent.*

- (i) *The sequence (X_n) is UI.*
- (ii) *The limit X is integrable and the sequence (X_n) converges to X in the L^1 -norm.*
- (iii) *The limit X is integrable and*

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X_n|] = \mathbb{E}[|X|].$$

Proof. We follow the approach in [68, Thm. 5.5.2].

(i) $\Rightarrow$ (ii). For every $M > 0$ we define

$$\Phi_M : \mathbb{R} \rightarrow \mathbb{R}, \quad \Phi_M(x) = \begin{cases} M, & x \geq M, \\ x, & |x| < M, \\ -M, & x \leq -M. \end{cases}$$

We have

$$\mathbb{E}[|X_n - X|] \leq \mathbb{E}[|X_n - \Phi_M(X_n)|] + \mathbb{E}[|\Phi_M(X_n) - \Phi_M(X)|] + \mathbb{E}[|\Phi_M(X) - X|]$$

$$\leq 2\mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M\}}] + \mathbb{E}[|\Phi_M(X_n) - \Phi_M(X)|] + 2\mathbb{E}[|X| \mathbf{I}_{\{|X| > M\}}].$$

The sequence (X_n) is uniformly integrable and Theorem 3.2.16 implies that $\sup_n \mathbb{E}[|X_n|] < \infty$. Fatou's Lemma applied to an a.s. convergent subsequence of X_n implies that $X \in L^1$. We conclude that for any $\varepsilon > 0$ there exists $M = M(\varepsilon) > 0$ such that

$$2\mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M\}}] + 2\mathbb{E}[|X| \mathbf{I}_{\{|X| > M\}}] < \frac{\varepsilon}{2}, \quad \forall n \in \mathbb{N}. \quad (3.2.14)$$

From Corollary 1.3.62 we deduce that $\Phi_{M(\varepsilon)}(X_n)$ converges to $\Phi_{M(\varepsilon)}(X)$ in probability. Moreover,

$$|\Phi_{M(\varepsilon)}(X_n)| < M(\varepsilon), \quad \forall n \in \mathbb{N}.$$

The Bounded Convergence Theorem 1.3.73 implies that there exists $n = n(\varepsilon) > 0$ such that for any $n \geq n(\varepsilon)$ we have

$$\mathbb{E}[|\Phi_{M(\varepsilon)}(X_n) - \Phi_{M(\varepsilon)}(X)|] < \frac{\varepsilon}{2}.$$

Form (3.2.14) we deduce that $\mathbb{E}[|X_n - X|] < \varepsilon$ for $n > n(\varepsilon)$.

Clearly (ii) $\Rightarrow$ (iii) since $X_n \rightarrow X$ in L^1 implies $\|X_n\|_{L^1} \rightarrow \|X\|_{L^1}$.

(iii) $\Rightarrow$ (i) For any $M > 0$ consider the *continuous* function

$$\Psi_M : [0, \infty) \rightarrow \mathbb{R}, \quad \Psi_M(x) = \begin{cases} x, & x \in [0, M-1], \\ 0, & x \geq M, \\ \text{linear}, & x \in (M-1, M). \end{cases}$$

The Dominated Convergence Theorem implies that $\Psi_M(|X|)$ converges to $|X|$ in L^1 as $M \rightarrow \infty$. Thus, there exists $M = M(\varepsilon)$ such that

$$\mathbb{E}[|X|] - \mathbb{E}[\Psi_M(|X|)] < \frac{\varepsilon}{2}, \quad \forall M \geq M(\varepsilon). \quad (3.2.15)$$

Using the Bounded Convergence Theorem as in the proof of the implication (i) $\Rightarrow$ (ii) we deduce that

$$\mathbb{E}[\Psi_M(|X_n|)] \rightarrow \mathbb{E}[\Psi_M(|X|)], \quad \forall M > 0. \quad (3.2.16)$$

Thus, for any $n \in \mathbb{N}$ we have

$$\begin{aligned} \mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M(\varepsilon)\}}] &\leq \mathbb{E}[|X_n|] - \mathbb{E}[\Psi_{M(\varepsilon)}(|X_n|)] \\ &= \left(\mathbb{E}[|X_n|] - \mathbb{E}[|X|] \right) \\ &\quad + \left(\mathbb{E}[|X|] - \mathbb{E}[\Psi_{M(\varepsilon)}(|X|)] \right) + \left(\mathbb{E}[\Psi_{M(\varepsilon)}(|X|)] - \mathbb{E}[\Psi_{M(\varepsilon)}(|X_n|)] \right) \\ &\stackrel{(3.2.15)}{<} \left(\mathbb{E}[|X_n|] - \mathbb{E}[|X|] \right) + \left(\mathbb{E}[\Psi_{M(\varepsilon)}(|X|)] - \mathbb{E}[\Psi_{M(\varepsilon)}(|X_n|)] \right) + \frac{\varepsilon}{2}. \end{aligned}$$

We can choose $N = N(\varepsilon, M(\varepsilon))$ so that for $n > N(\varepsilon)$ we have

$$\left(\mathbb{E}[|X_n|] - \mathbb{E}[|X|] \right) + \left(\mathbb{E}[\Psi_{M(\varepsilon)}(|X|)] - \mathbb{E}[\Psi_{M(\varepsilon)}(|X_n|)] \right) < \frac{\varepsilon}{2}.$$

Hence for any $M \geq M(\varepsilon)$

$$\sup_{n > N(\varepsilon)} \mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M\}}] \leq \sup_{n > N(\varepsilon)} \mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M(\varepsilon)\}}] < \varepsilon.$$

Now choose $M_1 > M(\varepsilon)$ such that

$$\mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M_1\}}] < \varepsilon, \quad \forall n = 1, 2, \dots, N(\varepsilon).$$

Hence for $M \geq M_1$ we have

$$\sup_{n \in \mathbb{N}} \mathbb{E}[|X_n| \mathbf{I}_{\{|X_n| > M\}}] < \varepsilon.$$

Thus (X_n) is uniformly integrable. $\square$

Remark 3.2.22. (a) The UI condition is a compactness condition. More precisely, the *Dunford-Pettis theorem* states that a collection $\mathcal{X} \subset L^1(\Omega, \mathcal{S}, \mathbb{P})$ is UI if and only if its closure in the weak topology of $L^1(\Omega, \mathcal{S}, \mathbb{P})$ is compact. For a proof we refer to [21, Thm. 4.7.18] or [66, Sec. IV.8.11].

(b) The implication (iii) $\Rightarrow$ (ii) is sometimes referred to as *Scheffé's Lemma*. It goes back to earlier work of Vitali and Riesz. Here is a direct proof. Suppose that $\mathbb{E}[|X_n|] \rightarrow \mathbb{E}[|X|]$.

Assume first that $X_n \rightarrow X$ a.s.. Set $Y_n := |X_n| + |X| - |X_n - X|$. Then $Y_n \geq 0$ and $Y_n \rightarrow 2|X|$ a.s.. Fatou's lemma implies that

$$\begin{aligned} 2\mathbb{E}[|X|] &\leq \liminf_{n \rightarrow \infty} \mathbb{E}[Y_n] = \liminf_{n \rightarrow \infty} \mathbb{E}[|X_n| + |X| - |X_n - X|] \\ &= 2\mathbb{E}[|X|] - \limsup_{n \rightarrow \infty} \mathbb{E}[|X_n - X|]. \end{aligned}$$

We conclude that

$$\limsup_{n \rightarrow \infty} \mathbb{E}[|X_n - X|] = 0 \iff \mathbb{E}[|X_n - X|] = 0.$$

Suppose that $X_n \rightarrow X$ in probability. Then any subsequence of (X_n) has a sub-subsequence that converges a.s. to X , and thus in L^1 . Hence $X_n \rightarrow X$ in L^1 if $X_n \rightarrow X$ in probability.

(c) We used the Bounded Convergence Theorem to prove the implication (i) $\Rightarrow$ (ii). Obviously, the Bounded Convergence Theorem is a special case of this implication. One can prove the equivalence (i) $\iff$ (ii) without relying on the Bounded Convergence Theorem; see [64, Thm. 10.3.6].

(d) The sequence in Example 2.2.10 converges in law, it is uniformly integrable yet it does not converge in probability.

On the other hand, if a sequence of random variables (X_n) is uniformly integrable and converges in law to a random variable X_∞ , then $X_\infty \in L^1$

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n] = \mathbb{E}[X_\infty].$$

Indeed, Skorokhod's Representation Theorem (Exercise 2.24) implies that there exists a sequence of random variables $Y_n : ([0, 1], \lambda) \rightarrow \mathbb{R}$ that converge a.s. to a random variable $Y_\infty : [0, 1] \rightarrow \mathbb{R}$ and such that

$$\mathbb{P}_{Y_n} = \mathbb{P}_{X_n}, \quad \forall n \in \mathbb{N} \cup \{\infty\}.$$

Then (Y_n) is UI and $\mathbb{E}[Y_n] = \mathbb{E}[X_n]$. □

3.2.3. Uniformly integrable martingales. We can now formulate and prove a refinement of Theorem 3.2.4.

Theorem 3.2.23. *If $(X_n)_{n \in \mathbb{N}_0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ of $(\Omega, \mathcal{S}, \mathbb{P})$. Set*

$$\mathcal{F}_\infty := \bigvee_{n \geq 0} \mathcal{F}_n = \sigma(\mathcal{F}_n, n \geq 0).$$

The following are equivalent.

- (i) *The collection $(X_n)_{n \in \mathbb{N}_0}$ is UI.*
- (ii) *The sequence $(X_n)_{n \in \mathbb{N}_0}$ converges a.s. and L^1 to a random variable X_∞ .*
- (iii) *The sequence $(X_n)_{n \in \mathbb{N}_0}$ converges L^1 to a random variable X_∞ .*
- (iv) *There exists an integrable random variable $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that*

$$X_n = \mathbb{E}[X \mid \mathcal{F}_n], \quad \text{a.s.,} \quad \forall n \in \mathbb{N}_0$$

If the above conditions are satisfied, then the limiting random variable X_∞ in (ii) and (iii) is related to the random variable X in (iv) via the equality $X_\infty = \mathbb{E}[X \mid \mathcal{F}_\infty]$, i.e.,

$$\lim_{n \rightarrow \infty} \mathbb{E}[X \mid \mathcal{F}_n] = \mathbb{E}[X \mid \mathcal{F}_\infty]$$

a.s. and L^1 . In particular, $\mathbb{E}[X_\infty] = \mathbb{E}[X_0]$.

Proof. Note that if a martingale (X_n) is UI, then it is bounded in L^1 and, according to Theorem 3.2.4, converges a.s. to an integrable random variable X_∞ . In view of the previous discussion the statements (i)-(iii) are equivalent. The implication (iv) $\Rightarrow$ (i) follows from Corollary 3.2.20. The only thing left to prove is (iii) $\Rightarrow$ (iv).

More precisely, we will show that if $X_n \rightarrow X_\infty$ in L^1 , then

$$X_n = \mathbb{E}[X_\infty \mid \mathcal{F}_n], \quad \text{a.s.,} \quad \forall n \in \mathbb{N}_0$$

In other words, we have to show that, for all $m \in \mathbb{N}_0$, and all $A \in \mathcal{F}_m$ we have

$$\mathbb{E}[X_m \mathbf{I}_A] = \mathbb{E}[X_\infty \mathbf{I}_A].$$

Since (X_n) is a martingale we deduce that, for $n > m$, we have

$$\mathbb{E}[X_m \mathbf{I}_A] = \mathbb{E}[\mathbb{E}[X_n \mid \mathcal{F}_m] \mathbf{I}_A] = \mathbb{E}[\mathbb{E}[X_n \mathbf{I}_A \mid \mathcal{F}_m]] = \mathbb{E}[X_n \mathbf{I}_A].$$

Now let $n \rightarrow \infty$.

Suppose now that for some integrable random variable X we have $X_n = \mathbb{E}[X \mid \mathcal{F}_n]$. We want to show that

$$\lim_n X_n = X_\infty := \mathbb{E}[X \mid \mathcal{F}_\infty]$$

i.e., for any $F \in \mathcal{F}_\infty$ we have

$$\mathbb{E}[X_\infty \mathbf{I}_F] = \mathbb{E}[X \mathbf{I}_F].$$

Denote by $\mathcal{Z} \subset \mathcal{F}_\infty$ the collection of $F \in \mathcal{F}_\infty$ for which the above holds. Clearly $\mathcal{F}_n \subset \mathcal{Z}$. Moreover, $\mathcal{Z}$ is a λ -system and contains the π -system

$$\bigcup_{n \geq 0} \mathcal{F}_n.$$

Thus it contains $\mathcal{F}_\infty$, the σ -algebra generated by this system. $\square$

Theorem 3.2.23 implies that

$$\lim_{n \rightarrow \infty} \mathbb{E}[X \mid \mathcal{F}_n] = \mathbb{E}[X \mid \mathcal{F}_\infty] \text{ a.s. and } L^1, \quad \forall X \in L^1(\Omega, \mathcal{F}, \mathbb{P}). \quad (3.2.17)$$

In particular, we deduce

Corollary 3.2.24 (Lévy's 0-1 law). *For any set $A \in \mathcal{F}_\infty$, the random variables*

$$\mathbb{P}[A \mid \mathcal{F}_n] = \mathbb{E}[\mathbf{I}_A \mid \mathcal{F}_n], \quad n \in \mathbb{N},$$

converge a.s. and L^1 to $\mathbf{I}_A$ as $n \rightarrow \infty$. $\square$

Corollary 3.2.25 (Kolmogorov's 0-1 law). *Suppose that $\mathcal{G}_1, \mathcal{G}_2, \dots$ are independent σ -subalgebras of $\mathcal{F}$. We set*

$$\mathcal{T}_n := \sigma(\mathcal{G}_{n+1}, \mathcal{G}_{n+2}, \dots)$$

and form the tail σ -algebra

$$\mathcal{T}_\infty = \bigcap_{n \geq 1} \mathcal{T}_n.$$

Then $\mathcal{T}_\infty$ is a zero-one sigma-algebra,

$$H \in \mathcal{T}_\infty \Rightarrow \mathbb{P}[H] \in \{0, 1\}.$$

Proof. Define $\mathcal{F}_n := \sigma(\mathcal{G}_1, \dots, \mathcal{G}_n)$. Let $H \in \mathcal{F}_\infty$. By Levy's 0 – 1 law we have

$$\mathbb{E}[\mathbf{I}_H \mid \mathcal{F}_n] \rightarrow \mathbf{I}_H \text{ a.s..}$$

On the other hand, if $H \in \mathcal{T}$, then since $\mathcal{T} \perp \mathcal{F}_n$ we deduce $\mathbb{E}[\mathbf{I}_H \mid \mathcal{F}_n] = \mathbb{P}[H]$, so that $\mathbb{P}[H] = \mathbf{I}_H$ a.s.. In other words, $\mathbf{I}_H$ is a.s. constant and this constant can only be 0 or 1. $\square$

Example 3.2.26. Consider again the Galton-Watson branching process in Example 3.1.8. Suppose that the reproduction law μ satisfies

$$m := \sum_{n \geq 0} n\mu[n] < \infty.$$

Assume $m \geq 1$. Consider the *extinction event* defined in Example 3.2.7,

$$E := \bigcup_{n \geq 0} E_n, \quad E_n = \{Z_k = 0, \quad \forall k \geq n\}.$$

Consider next the event

$$U := \left\{ \sup_n Z_n = \infty \right\}.$$

We want to prove that if the probability that an individual has no successor is positive then, with probability 1, either the population extinguishes in finite time, or explodes. In particular it cannot stabilize to a finite nonzero limit. More precisely, we have the following dichotomy result.

If $p_0 = \mu[0] > 0$, then, with probability 1, the population either becomes extinct or explodes, i.e.,

$$E = U^c, \quad \mathbb{P}[E \cup U] = 1. \quad (3.2.18)$$

In particular

$$E = \left\{ \lim_n Z_n = 0 \right\}. \quad (3.2.19)$$

Note that

$$\begin{aligned} \forall \nu \in \mathbb{N}_0, \quad \exists \delta(\nu) \in (0, 1) : \forall n \in \mathbb{N}, \\ \mathbb{P}[E \mid Z_1, \dots, Z_n] \geq \delta(\nu) \text{ on } \{Z_n \leq \nu\}. \end{aligned} \quad (3.2.20)$$

Indeed, if the population of the n -th generation has at most ν individuals then the probability that there will be no $(n+1)$ -th generation is at least p'_0 . More formally,

$$\mathbb{P}[E \mid Z_1, \dots, Z_n] \geq \mathbb{P}[E_{n+1} \mid Z_1, \dots, Z_n] = \mathbb{P}[E_{n+1} \mid Z_n].$$

We have

$$\begin{aligned} \mathbb{P}[E_{n+1} \mid Z_n] \mathbf{I}_{\{Z_n \leq \nu\}} &= \sum_{k=0}^{\nu} \mathbb{P}[E_{n+1} \mid Z_n = k] \mathbf{I}_{\{Z_n = k\}} \\ &= \sum_{k=0}^{\nu} p_0^k \mathbf{I}_{\{Z_n = k\}} \geq p'_0 \mathbf{I}_{\{Z_n \leq \nu\}}. \end{aligned}$$

This proves (3.2.20) with $\delta(\nu) = p'_0$.

Since Z_n are integer valued we deduce that

Lemma 3.2.27. *Suppose that $(Z_n)_{n \geq 1}$ is a sequence of nonnegative random variables. Set*

$$E = \{Z_n = 0 \text{ for some } n\}, \quad B := \left\{ \sup_n Z_n < \infty \right\}.$$

If (Z_n) satisfies (3.2.20), then

$$E \supset B \quad (3.2.21)$$

Proof. Set $\mathcal{F}_n := \sigma(Z_1, \dots, Z_n)$, $B_\nu := \left\{ \sup_n Z_n \leq \nu \right\}$, so that

$$B_1 \subset B_2 \subset \dots, \quad \bigcup_{\nu} B_\nu = B.$$

We have $\mathbb{E}[E \mid \mathcal{F}_n] \geq \delta(\nu)$ on B_ν . Letting $n \rightarrow \infty$ we deduce from Lévy's 0-1 theorem (Corollary 3.2.24) that

$$\lim_{n \rightarrow \infty} \mathbb{E}[\mathbf{I}_E \mid \mathcal{F}_n] = \mathbf{I}_E.$$

Hence $B_\nu \subset E$ for any ν . Hence $B \subset E$. □

In our special case, $B = U^c$. Note also that if the population dies at a time n_0 , then $Z_n = 0, \forall n \geq n_0$. Hence $E \subset B$ or, in view of (3.2.21), $E = B = U^c$. This proves the claimed dichotomy (3.2.18).

When $m = 1$, then $W_n = Z_n$ converges almost surely to an integrable random variable and we see that

$$\left\{ \lim_n Z_n < \infty \right\} \subset \left\{ \sup Z_n < \infty \right\} \subset E$$

and we deduce that

$$1 \geq \mathbb{P}[E] \geq \mathbb{P}\left[\lim_n Z_n < \infty\right] = 1.$$

Thus, when $m = 1$ and the probability having *no* successors is positive, i.e., $\mu[0] > 0$, then the extinction probability is also 1. One can show (see [8, Sec. I.9] or [115]) that if $m = 1$ and

$$\sigma^2 := \text{Var}[X_{n,j}] = \sum_k k(k-1)\mu[k] < \infty,$$

then

$$\lim_{n \rightarrow \infty} n\mathbb{P}[Z_n > 0] = \frac{2}{\sigma^2}.$$

Thus, the probability of the population surviving more than n generations, given that individuals have on average 1 successor is $O(1/n)$.

When $m > 1$ the extinction probability is still positive but < 1 . Exercise 3.29 describes this probability and gives additional information about the distribution of W . For more details about branching processes we refer to [8, 101]. $\square$

Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a process adapted to a filtration $\mathcal{F}_\bullet$ such that X_n converges a.s. to a random variable X_∞ as $n \rightarrow \infty$. If T is a stopping time adapted to the same filtration, finite or not, we set

$$\hat{X}_T := \sum_{n \in \mathbb{N}_0} X_n \mathbf{I}_{\{T=n\}} + X_\infty \mathbf{I}_{\{T=\infty\}} = X_T + X_\infty \mathbf{I}_{\{T=\infty\}}. \quad (3.2.22)$$

Note that

$$\begin{aligned} \mathbb{P}[T = \infty] = 0 &\Rightarrow \hat{X}_T = X_T \text{ a.s.,} \\ \hat{X}_T = X_\infty^T &:= \lim_{n \rightarrow \infty} X_{T \wedge n} = \lim_{n \rightarrow \infty} X_n^T. \end{aligned} \quad (3.2.23)$$

Theorem 3.2.28 (UI Optional sampling: martingales). *Suppose that $X_\bullet = (X_n)_{n \in \mathbb{N}_0}$ is a UI martingale and T is a stopping time, not necessarily a.s. finite. Then $\hat{X}_T \in L^1$ and*

$$\hat{X}_T = \mathbb{E}[X_\infty | \mathcal{F}_T]. \quad (3.2.24)$$

Moreover, if S, T are stopping times such that $S \leq T$, then

$$\mathbb{E}[X_\infty | \mathcal{F}_S] = \mathbb{E}[\hat{X}_T | \mathcal{F}_S] = \hat{X}_S. \quad (3.2.25)$$

In particular

$$\mathbb{E}[\hat{X}_T] = \mathbb{E}[X_0].$$

Proof. Let us first prove that $\hat{X}_T \in L^1$. We have

$$\mathbb{E}[|\hat{X}_T|] = \sum_{n \geq 0} \mathbb{E}[\mathbf{I}_{\{T=n\}} |X_n|] + \mathbb{E}[\mathbf{I}_{\{T=\infty\}} |X_\infty|]$$

$$\begin{aligned}
&= \sum_{n \geq 0} \mathbb{E} \left[\mathbf{I}_{\{T=n\}} \left| \mathbb{E}[X_\infty \mid \mathcal{F}_n] \right| \right] + \mathbb{E} \left[\mathbf{I}_{\{T=\infty\}} |X_\infty| \right] \\
& \left(\left| \mathbb{E}[X \mid \mathcal{F}] \right| \leq \mathbb{E}[|X| \mid \mathcal{F}] \right) \\
&\leq \sum_{n \geq 0} \mathbb{E} \left[\mathbf{I}_{\{T=n\}} \mathbb{E}[|X_\infty| \mid \mathcal{F}_n] \right] + \mathbb{E} \left[\mathbf{I}_{\{T=\infty\}} |X_\infty| \right]
\end{aligned}$$

(use the definition of conditional expectation)

$$= \sum_{n \geq 0} \mathbb{E} \left[\mathbf{I}_{\{T=n\}} |X_\infty| \right] + \mathbb{E} \left[\mathbf{I}_{\{T=\infty\}} |X_\infty| \right] = \mathbb{E}[|X_\infty|] < \infty.$$

Moreover, for $A \in \mathcal{F}_T$ we have

$$\begin{aligned}
\mathbb{E}[\mathbf{I}_A \hat{X}_T] &= \sum_{n \in \mathbb{N}_0} \mathbb{E}[\mathbf{I}_{A \cap \{T=n\}} X_n] + \mathbb{E}[\mathbf{I}_{A \cap \{T=\infty\}} X_\infty] \\
& \left(\mathbf{I}_{A \cap \{T=n\}} X_n = \mathbb{E}[\mathbf{I}_{A \cap \{T=n\}} X_\infty \mid \mathcal{F}_n] \right) \\
&= \sum_{n \in \mathbb{N}_0} \mathbb{E}[\mathbf{I}_{A \cap \{T=n\}} X_\infty] + \mathbb{E}[\mathbf{I}_{A \cap \{T=\infty\}} X_\infty] = \mathbb{E}[\mathbf{I}_A X_\infty],
\end{aligned}$$

and thus $\hat{X}_T = \mathbb{E}[X_\infty \mid \mathcal{F}_T]$. The since $\mathcal{F}_S \subset \mathcal{F}_T$, the equality (3.2.25) follows immediately from (3.2.24) and the properties of conditional expectation. $\square$

Corollary 3.2.29 (Optional Stopping). *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a UI martingale and T is any stopping time. Then the stopped martingale $X_n^T = X_{T \wedge n}$ is also a uniformly integrable martingale with respect to the filtration $\mathcal{F}_{T \wedge n}$.*

Proof. From Theorem 3.2.28 we deduce that $X_{T \wedge n} = \mathbb{E}[X_\infty \mid \mathcal{F}_{T \wedge n}]$ and Corollary 3.2.20 implies it is UI. $\square$

Doob's conditions in Definition 3.1.27 are closely related to uniform integrability.

Proposition 3.2.30. *Suppose that $(X_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ and T is an a.s. finite stopping time adapted to the same filtration. Then the following statements are equivalent.*

- (i) *The stopping time satisfies Doob's conditions (3.1.8b) and (3.1.8c).*
- (ii) *The stopped martingale $X_n^T = X_{T \wedge n}$ is UI.*

Proof. (i) $\Rightarrow$ (ii) Consider the submartingale $|X_n|$. Since T satisfies Doob's conditions we deduce from Theorem 3.1.29 that

$$\mathbb{E}[|X_T|] \geq \mathbb{E}[|X_{T \wedge n}|] \quad \forall n \geq 0.$$

Thus

$$\limsup_{n \rightarrow \infty} \mathbb{E}[|X_{T \wedge n}|] \leq \mathbb{E}[|X_T|]$$

Since $\lim_{n \rightarrow \infty} X_{T \wedge n} = X_T$, a.s., we deduce from Fatou's Lemma that

$$\mathbb{E}[|X_T|] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[|X_{T \wedge n}|]$$

so that

$$\limsup_{n \rightarrow \infty} \mathbb{E}[|X_{T \wedge n}|] = \mathbb{E}[|X_T|].$$

The desired conclusion now follows from Theorem 3.2.21.

(ii) $\Rightarrow$ (i) Observe first that $\lim_{n \rightarrow \infty} X_{T \wedge n} = X_T$ and since X_n^T is UI we deduce X^T is integrable. Now observe that

$$\mathbb{E}[|X_n| \mathbf{I}_{T > n}] = \mathbb{E}[|X_{T \wedge n}| \mathbf{I}_{T > n}].$$

Since $\mathbb{P}[T < \infty] = 1$ we deduce $\lim_{n \rightarrow \infty} \mathbb{P}[T > n] = 0$. Finally, using the fact that the stopped martingale X_n^T is UI we deduce from (UI₁)

$$\lim_{n \rightarrow \infty} \mathbb{E}[|X_{T \wedge n}| \mathbf{I}_{T > n}] = 0.$$

□

Definition 3.2.31. Suppose that $(X_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ and T is an a.s. finite stopping time adapted to the same filtration. We say that T *reduces* X if the stopped process $X_\bullet^T = X_{n \wedge T}$ is a UI martingale. □

Thus, a stopping time satisfies the Doob conditions iff it is reducing. The bounded stopping times are reducing. If T is an a.s. finite stopping time such that for some $p > 1$

$$\sup_{n \in \mathbb{N}} \|X_{n \wedge T}\|_{L^p} < \infty,$$

then T is a reducing stopping time.

Let us observe that the above discussion yields an alternate proof of the Optional Sampling Theorem 3.1.29.

Corollary 3.2.32 (Optional Sampling Theorem). *Suppose that $(X_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)$, $S \leq T$ are stopping times adapted to the same filtration and T is reducing. Then*

$$\mathbb{E}[X_T \mid \mathcal{F}_S] = X_S.$$

In particular, $\mathbb{E}[X_T] = \mathbb{E}[X_S]$.

Proof. Note that X^T is UI and, since $X^S = (X^T)^S$, we deduce from Theorem 3.2.28 that

$$\mathbb{E}[X_T \mid \mathcal{F}_S] = \mathbb{E}[X_\infty^T \mid \mathcal{F}_S] = X_S^T = X_S.$$

□

3.2.4. Applications of the optional sampling theorem. The Optional Sampling Theorem is a very versatile tool for computing expectations. We restate below the special case of this theorem frequently used in applications.

Corollary 3.2.33 (Optional Sampling Theorem). *Suppose that $(X_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ and T is an a.s. finite reducing stopping time, i.e., the stopped martingale X^T is UI. Then $\mathbb{E}[X_T] = \mathbb{E}[X_0]$.* □

The applicability of the above result is greatly enhanced once we have simple criteria for recognizing when a stopped martingale is UI. We have the following result of J. L. Doob, [61, Thm. VII.2.2].

Proposition 3.2.34. Suppose that $(M_n)_{n \geq 0}$ is a random process adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ such that

$$\mathbb{E}[M_n] < \infty, \quad \forall n,$$

and T is a stopping time adapted to the same filtration. Suppose that

$$\mathbb{E}[T] < \infty, \quad (3.2.26a)$$

$$\exists C > 0 : \quad \forall n \in \mathbb{N}, \quad \mathbb{E}[|M_n - M_{n-1}| \mid \mathcal{F}_{n-1}] \leq C. \quad (3.2.26b)$$

Then the stopped process $M_n^T = M_{T \wedge n}$ is UI.

Proof. We will show that there exists $Y \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ such that

$$|M_{T \wedge n}| \leq Y, \quad \forall n \in \mathbb{N}.$$

Note that

$$\begin{aligned} M_{T \wedge n} &= \sum_{k=0}^{n-1} M_k \mathbf{I}_{\{T \geq k\}} + M_n \mathbf{I}_{\{T \geq n\}} \\ &= \sum_{k=0}^{n-1} M_k (\mathbf{I}_{\{T \geq k\}} - \mathbf{I}_{\{T \geq k+1\}}) + M_n \mathbf{I}_{\{T \geq n\}} = M_0 + \sum_{k=1}^n (M_k - M_{k-1}) \mathbf{I}_{\{T \geq k\}} \end{aligned}$$

so

$$|M_{T \wedge n}| \leq |M_0| + \sum_{k=1}^n |M_k - M_{k-1}| \mathbf{I}_{\{T \geq k\}}.$$

Set

$$Y := |M_0| + \sum_{k=1}^{\infty} |M_k - M_{k-1}| \mathbf{I}_{\{T \geq k\}}.$$

Clearly $|M_{T \wedge n}| \leq Y, \forall n \geq 0$. We will show that $\mathbb{E}[Y] < \infty$. We have

$$\mathbb{E}[|M_k - M_{k-1}| \mathbf{I}_{\{T \geq k\}}] = \mathbb{E}[\mathbb{E}[|M_k - M_{k-1}| \mathbf{I}_{\{T \geq k\}} \mid \mathcal{F}_{k-1}]],$$

($\{T \geq k\} \in \mathcal{F}_{k-1}$)

$$\mathbb{E}[\mathbf{I}_{\{T \geq k\}} \mathbb{E}[|M_k - M_{k-1}| \mid \mathcal{F}_{k-1}]] \stackrel{(3.2.26b)}{\leq} C \mathbb{E}[\mathbf{I}_{\{T \geq k\}}] = C \mathbb{P}[T \geq k].$$

Thus

$$\mathbb{E}[Y] \leq \mathbb{E}[|M_0|] + C \sum_{k=1}^{\infty} \mathbb{P}[T \geq k] = \mathbb{E}[|M_0|] + C \mathbb{E}[T] \stackrel{(3.2.26a)}{<} \infty.$$

□

Theorem 3.2.35 (Wald's formula). Suppose that $(Y_n)_{n \geq 0}$ is a sequence of i.i.d. integrable random variables with finite mean μ . Set

$$S_n := \sum_{k=0}^n Y_k.$$

Let T be a stopping time adapted to the filtration $\mathcal{F}_n = \sigma(Y_0, \dots, Y_n)$ and such that $\mathbb{E}[T] < \infty$. The following hold.

$$(i) \quad \mathbb{E}[S_T] = \mu \mathbb{E}[T].$$

(ii) Suppose additionally that

$$Y_n \in L^2, \quad \mu = 0, \quad \sigma^2 = \text{Var}[Y_n].$$

$$\text{Then } \text{Var}[S_T] = \sigma^2 \mathbb{E}[T].$$

Proof. (i) Set $\bar{Y}_n = Y_n - \mu$,

$$M_n := S_n - n\mu = \sum_{k=1}^n \bar{Y}_k.$$

Then

$$\begin{aligned} \mathbb{E}[M_n \mid \mathcal{F}_{n-1}] &= \mathbb{E}[\bar{Y}_n + M_{n-1} \mid \mathcal{F}_{n-1}] \\ &= \mathbb{E}[\bar{Y}_n \mid \mathcal{F}_{n-1}] + \mathbb{E}[M_{n-1} \mid \mathcal{F}_{n-1}] = \mathbb{E}[\bar{Y}_n] + M_{n-1} = M_{n-1}. \end{aligned}$$

Observe that

$$\mathbb{E}[|M_n - M_{n-1}| \mid \mathcal{F}_{n-1}] = \mathbb{E}[|\bar{Y}_n| \mid \mathcal{F}_{n-1}] = \mathbb{E}[|\bar{Y}_n|] = \mathbb{E}[|\bar{Y}_0|]$$

so that (3.2.26b) is satisfied. We deduce from Proposition 3.2.34 that the stopped martingale M_n^T is UI and the Optional Sampling Theorem implies

$$0 = \mathbb{E}[M_0] = \mathbb{E}[M_T] = \mathbb{E}[S_T] - \mu \mathbb{E}[T].$$

(ii) From (i) we deduce $\mathbb{E}[S_T] = 0$ so $\text{Var}[S_T] = \mathbb{E}[S_T^2]$. Set

$$Q_n := \sum_{k=1}^n Y_k^2.$$

We have

$$\mathbb{E}[S_n^2] = \sum_{k=1}^n \mathbb{E}[Y_k^2] + 2 \sum_{1 \leq i < j \leq n} \mathbb{E}[Y_i Y_j] = \mathbb{E}[Q_n].$$

As in (i) we observe that $Z_n = Q_n - n\sigma^2$ is a martingale adapted to the filtration $\mathcal{F}_n$, the increments $Q_n - Q_{n-1}$ are independent of $\mathcal{F}_n$ and

$$\mathbb{E}[|Z_n - Z_{n-1}| \mid \mathcal{F}_{n-1}] = \mathbb{E}[|Q_n - Q_{n-1}|] \leq \mathbb{E}[Y_n^2] + \sigma^2 = 2\sigma^2.$$

We deduce from Proposition 3.2.34 that the stopped martingale Z_n^T is UI and the Optional Sampling Theorem implies

$$\begin{aligned} 0 &= \mathbb{E}[Z_0] = \mathbb{E}[Z_T] = \mathbb{E}[Q_T] - \sigma^2 \mathbb{E}[T] \\ &= \mathbb{E}[S_T^2] - \sigma^2 \mathbb{E}[T] = \text{Var}[S_T] - \sigma^2 \mathbb{E}[T]. \end{aligned}$$

□

Remark 3.2.36. In Exercise 1.21 we described a version (1.6.1) of Wald's formula that has a different nature than the one presented in Theorem 3.2.35. The random time T in (1.6.1) is independent of the variables X_n and the proof of (1.6.1) is a simple exercise in conditioning.

In Theorem 3.2.35 the random time T is quite dependent of these variables given that it is adapted to the filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$ and the proof of the corresponding version of Wald's formula requires the machinery of martingale theory.

We want to point out that, without some assumptions on T , we cannot expect the equality $\mathbb{E}[S_T] = \mu \mathbb{E}[T]$ to hold. Here is an example.

Suppose that the random variables X_n are exponential with parameter λ . For fix $t > 0$ we set

$$N(t) := \max \{n \geq 0; S_n \leq t\}.$$

The collection $(N(t))_{t \geq 0}$ is the Poisson process introduced in Example 1.3.7. Thus, $N(t) \sim \text{Poi}(\lambda t)$ so that

$$\mathbb{E}[N(t)] = \lambda t.$$

In this case

$$\mu = \mathbb{E}[\text{Exp}(\lambda)] = \frac{1}{\lambda}.$$

For fixed t , the random variable $N(t)$ is *not adapted* to the filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$. Indeed, knowing $S_1, \dots, S_n$, we cannot conclude that $S_{n+1} > t$, i.e., that n is the largest index k such that $S_k \leq t$. If Wald's formula were true in this case it would predict $\mathbb{E}[S_{N(t)}] = t$. However, we know from (1.3.57) that

$$\mathbb{E}[S_{N(t)}] = t - \frac{1}{\lambda} + \frac{e^{-\lambda t}}{\lambda}.$$

Let us observe that $T = N(t) + 1$ is adapted to the filtration $\mathcal{F}_n$. Indeed

$$T = n \iff N(t) = n - 1$$

$$\iff X_1 + \dots + X_{n-1} \leq t \text{ and } X_1 + \dots + X_n + X_{n+1} > t,$$

so that $\{T = n\} \in \sigma(X_1, \dots, X_n)$. Wald's formula implies

$$\mathbb{E}[S_{N(t)+1}] = \mathbb{E}[N(t) + 1] \cdot \mathbb{E}[X_1] = \frac{\lambda t + 1}{\lambda} = t + \frac{1}{\lambda}.$$

This agrees with our earlier conclusion (1.3.58). □

Example 3.2.37 (Gambler's Ruin). Suppose that

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \{-1, 1\}, \quad n \in \mathbb{N},$$

is a sequence of i.i.d. random variables with common probability distribution $\mathbb{P}[X_n = 1] = p$, $\mathbb{P}[X_n = -1] = q = 1 - p$, $p \in (0, 1)$. Fix $k \in \mathbb{N}$ and set

$$S_0 =: k, \quad S_n := k + X_1 + \dots + X_n, \quad \forall n \in \mathbb{N},$$

Intuitively, k is the initial fortune of a gambler that plays a sequence of independent games where he wins \$1 with probability p and loses \$1 with probability q . Then S_n is the fortune of the gambler after n games. The game stops when the gambler is out of money, or his fortune reaches a prescribed threshold $N > k$.

The sequence $(S_n)_{n \in \mathbb{N}}$ is a random process adapted to the filtration

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n).$$

The random variable

$$T = T_k := \min \{n \in \mathbb{N}; S_n \in \{0, N\}\}. \quad (3.2.27)$$

is a stopping time adapted to this filtration. It is the moment the gambler stops playing. The 'sooner-rather-than-later' Lemma 3.1.34 implies that $\mathbb{E}[T] < \infty$ since T satisfies (3.1.10)

$$\forall n \in \mathbb{N}_0, \quad \mathbb{P}[T \leq n + N | \mathcal{F}_n] > r, \quad r = (\min(p, q))^N.$$

In particular, $\mathbb{P}[T < \infty] = 1$. We want to compute $p_k(N) := \mathbb{P}[S_T = N]$. We distinguish two cases.

A. $p = 1/2$ so that the game is fair. Then S_n is a martingale. Consider now the stopped process S^T . It is a UI martingale since it is uniformly bounded. We deduce from the Optional Sampling Theorem that

$$k = \mathbb{E}[S_0] = \mathbb{E}[S_T] = p_k(N)N \Rightarrow p_k(N) = \frac{k}{N}.$$

Thus, the ruin probability is $1 - p_k(N) = \frac{N-k}{N} = 1 - \frac{k}{N}$.

B. $p \neq 1/2$ so the game is biased. Consider the De Moivre's martingale M_n defined Example 3.1.7, i.e.,

$$M_n = \left(\frac{q}{p}\right)^{S_n}.$$

The stopped martingale M^T is UI since it is bounded. Hence

$$\mathbb{E}[M_T] = \mathbb{E}[M_0] = \left(\frac{q}{p}\right)^k.$$

If we set $p_k(N) := \mathbb{P}[S_T = N]$, then we deduce

$$\left(\frac{q}{p}\right)^k = \mathbb{P}[S_T = 0] \left(\frac{q}{p}\right)^0 + \mathbb{P}[S_T = N] \left(\frac{q}{p}\right)^N = (1 - p_k(N)) + p_k(N) \left(\frac{q}{p}\right)^N.$$

Hence

$$p_k(N) = \frac{\left(\frac{q}{p}\right)^k - 1}{\left(\frac{q}{p}\right)^N - 1} = p^{N-k} \frac{q^k - p^k}{q^N - p^N}. \quad \square$$

Example 3.2.38 (The coupon collector problem revisited). Let us recall the coupon collector problem we discussed in Example 1.3.28.

Suppose that each box of cereal contains one of m different coupons. Once you obtain one of every type of coupons, you can send in for a prize. Ann wants that prize and, for that reason, she buys one box of cereals everyday. Assuming that the coupon in each box is chosen independently and uniformly at random from the m possibilities and that Ann does not collaborate with others to collect coupons, how many boxes of cereal is she expected to buy before she obtain at least one of every type of coupon?

Let N denote the number of boxes bought until Ann has at least one of every coupon. We have shown in Example 1.3.28 that

$$\mathbb{E}[N] = mH_m, \quad H_m := \left(1 + \frac{1}{2} + \cdots + \frac{1}{m-1} + \frac{1}{m}\right).$$

Suppose now that Ann has a little brother, Bob and, every time she collects a coupon she already has, she gives it to Bob. At the moment when she completed her collection, Bob is missing B coupons. What is the expectation of B ?

To answer this question we follow the approach in [78, Sec. 12.5.1]. Assume that the coupons are labelled $1, \dots, m$. We denote by C_k the label of the coupon Ann found in the k -th box she bought. Thus $(C_k)_{k \geq 1}$ are i.i.d., uniformly distributed in $\{1, \dots, m\}$. Let $\mathcal{F}_0$ denote the trivial sigma-subalgebra and set

$$\mathcal{F}_n := \sigma(C_1, \dots, C_n), \quad n \in \mathbb{N}.$$

We introduce two new random variables.

- X_n is the number of coupons Ann is missing after she bought n cereal boxes, $n \geq 0$.

- Y_n the number of coupons that have appeared exactly one time in the first n boxes Ann bought, $n \geq 0$.

Note that $X_0 = m$, $Y_0 = 0$. From the equality

$$N = \min \{ n \in \mathbb{N}; X_n = 0 \},$$

we deduce that N is a stopping time adapted to the filtration $\mathcal{F}_\bullet$. Observe that $Y_N = B$, the number of coupons Bob is missing the moment Ann completed her collection.

Fix a function $f : \mathbb{N}_0^2 \rightarrow \mathbb{N}_0$ satisfying the difference equation

$$x(f(x-1, y+1) - f(x, y)) + yf(x, y-1) = 0, \quad \forall x, y \geq 1 \quad (3.2.28)$$

and form the processes

$$W_n := f(X_n, Y_n), \quad Z_n = W_n^N = W_{N \wedge n}, \quad n \geq 0.$$

Lemma 3.2.39. *The process $(Z_n)_{n \geq 0}$ is a martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$.*

Proof. We set $\Delta Z_n := Z_{n+1} - Z_n$. Note that Z_n is $\mathcal{F}_n$ -measurable so we have to show that

$$\mathbb{E}[\Delta Z_n \mid \mathcal{F}_n] = 0. \quad (3.2.29)$$

Observe first that

$$\Delta Z_n = \mathbf{I}_{\{X_n > 0\}} \Delta W_n.$$

Let us observe that, when $X_n > 0$ and Ann buys a new cereal box there are only three, mutually exclusive possibilities,

$$\Delta X_n = -1, \quad \Delta Y_n = -1, \quad \Delta X_n = \Delta Y_n = 0.$$

The first possibility corresponds to Ann obtaining a new coupon. In this case $Y_{n+1} = Y_n + 1$. The second possibility corresponds to Bob obtaining a new coupon. In this case $X_{n+1} = X_n$. The third possibility occurs when the $(n+1)$ -th coupon is owned by both Ann and Bob. Hence

$$\begin{aligned} \Delta Z_n &= \mathbf{I}_{\{\Delta X_n = -1\}} (f(X_n - 1, Y_n + 1) - f(X_n, Y_n)) \mathbf{I}_{\{X_n > 0\}} \\ &\quad + \mathbf{I}_{\{\Delta Y_n = -1\}} (f(X_n, Y_n - 1) - f(X_n, Y_n)) \mathbf{I}_{\{X_n > 0\}}. \end{aligned}$$

Now observe that

$$\mathbb{E}[\mathbf{I}_{\{\Delta X_n = -1\}} \mid \mathcal{F}_n] = \frac{X_n}{m} \quad \text{and} \quad \mathbb{E}[\mathbf{I}_{\{\Delta Y_n = -1\}} \mid \mathcal{F}_n] = \frac{Y_n}{m}.$$

To understand the first equality observe that if Ann is missing X_n coupons at time n , then the probability of getting a new one in the new box is $\frac{X_n}{m}$. The second equality is proved in a similar fashion. Hence

$$\begin{aligned} \mathbb{E}[\Delta Z_n \mid \mathcal{F}_n] &= \frac{X_n}{m} (f(X_n - 1, Y_n + 1) - f(X_n, Y_n)) \mathbf{I}_{\{X_n > 0\}} \\ &\quad + \frac{Y_n}{m} (f(X_n, Y_n - 1) - f(X_n, Y_n)) \mathbf{I}_{\{X_n > 0\}} \stackrel{(3.2.28)}{=} 0. \end{aligned}$$

□

The martingale $(Z_n)_{n \geq 0}$ is bounded so it is uniformly integrable and we deduce from the Optional Sampling Theorem that

$$\mathbb{E}[f(0, Y_N)] = \mathbb{E}[Z_N] = \mathbb{E}[Z_0] = \mathbb{E}[f(m, 0)].$$

This holds for any function f satisfying (3.2.28). Now observe that the function

$$f : \mathbb{N}_0 \rightarrow \mathbb{N}_0, \quad f(x, y) = H_x + \frac{y}{1+x},$$

where

$$H_0 = 0, \quad H_x = 1 + \frac{1}{2} + \cdots + \frac{1}{x}, \quad \forall x > 0,$$

satisfies (3.2.28), and we conclude

$$\mathbb{E}[Y_N] = \mathbb{E}[f(0, Y_N)] = \mathbb{E}[f(m, 0)] = H_m \sim \log m \quad \text{as } m \rightarrow \infty.$$

For example, if $m = 30$, then $H_m \approx 3.99$ so at the moment Ann has all the complete collection of 30 coupons, we expect that her little brother is missing only about 4 of them. Nearly there. $\square$

3.2.5. Uniformly integrable submartingales. The proof of Theorem 3.2.23 yields the following submartingale counterpart.

Theorem 3.2.40. *If $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale, then the following are equivalent.*

- (i) *The collection $(X_n)_{n \in \mathbb{N}_0}$ is uniformly integrable.*
- (ii) *The sequence $(X_n)_{n \in \mathbb{N}_0}$ converges a.s. and L^1 to a random variable X_∞ .*
- (iii) *The sequence $(X_n)_{n \in \mathbb{N}_0}$ converges L^1 to a random variable X_∞ .* $\square$

Corollary 3.2.41. *Suppose that $X_\bullet = (X_n)_{n \in \mathbb{N}_0}$ is a submartingale with Doob decomposition $X_n = X_0 + M_n + C_n$, where $(M_n)_{n \in \mathbb{N}_0}$ is the martingale component and $(C_n)_{n \in \mathbb{N}_0}$ is the predictable compensator. Then the following are equivalent.*

- (i) *The submartingale $(X_n)_{n \in \mathbb{N}_0}$ is uniformly integrable.*
- (ii) *The martingale $(M_n)_{n \in \mathbb{N}_0}$ and the compensator $(C_n)_{n \in \mathbb{N}_0}$ are uniformly integrable.*

Proof. Clearly (ii) $\Rightarrow$ (i). To prove the converse note that

$$\mathbb{E}[|C_n|] = \mathbb{E}[C_n] = \mathbb{E}[X_n] - \mathbb{E}[X_0]$$

and since (X_n) is uniformly integrable we deduce

$$\sup_n \mathbb{E}[|C_n|] \leq \sup_n \mathbb{E}[|X_n|] - \mathbb{E}[X_0] < \infty.$$

The limit $C_\infty := \lim_{n \rightarrow \infty} C_n$ exists because (C_n) is nondecreasing. The Monotone Convergence theorem implies that C_∞ is integrable. Since $|C_n| = C_n \leq C_\infty$, $\forall n$, we deduce that the family (C_n) is UI. $\square$

We can use Doob's decomposition to prove a submartingale version of Theorem 3.2.28.

Corollary 3.2.42. *If $X_\bullet = (X_n)_{n \in \mathbb{N}_0}$ is a uniformly integrable submartingale, then for any stopping time T the stopped submartingale $X_n^T = X_{T \wedge n}$ is a uniformly integrable submartingale.*

Proof. Consider the Doob decomposition of $X_\bullet$, $X_n = X_0 + M_n + C_n$. From Corollary 3.2.41 we deduce that $M_\bullet$ and $C_\bullet$ are UI. Moreover, the Doob decomposition of X^T is $X^T = M^T + C^T$. Corollary 3.2.29 shows that M^T is UI and C^T is UI since $0 \leq C_n^T \leq C_\infty \in L^1$. $\square$

Theorem 3.2.43 (UI Optional Sampling: submartingales). *Suppose that*

$$X_\bullet = (X_n)_{n \in \mathbb{N}_0}$$

is a UI submartingale. Then for any stopping times S, T such that $S \leq T$ we have

$$\hat{X}_S \leq \mathbb{E}[\hat{X}_T \mid \mathcal{F}_T]. \quad (3.2.30)$$

In particular, if we let $T = \infty$,

$$\hat{X}_S \leq \mathbb{E}[X_\infty \mid \mathcal{F}_S]. \quad (3.2.31)$$

Proof. If $X_\bullet = M_\bullet + C_\bullet$ is the Doob decomposition of X , then

$$X_\bullet^S = M_\bullet^S + C_\bullet^S, \quad X_\bullet^T = M_\bullet^T + C_\bullet^T$$

In this case $X^S = (X^T)^S$ we deduce that $\hat{X}_S = \widehat{X^T}_S$. Then, since $\hat{C}_S$ is $\mathcal{F}_S$ -measurable,

$$\begin{aligned} \hat{X}_S &= \hat{M}_S + \hat{C}_S \stackrel{(3.2.24)}{=} \mathbb{E}[M_\infty^T \mid \mathcal{F}_S] + \mathbb{E}[\hat{C}_S \mid \mathcal{F}_S] \\ &\leq \mathbb{E}[M_\infty^T \mid \mathcal{F}_S] + \mathbb{E}[\hat{C}_T \mid \mathcal{F}_S] = \mathbb{E}[M_\infty^T \mid \mathcal{F}_S] + \mathbb{E}[C_\infty^T \mid \mathcal{F}_S] = \mathbb{E}[\hat{X}_T \mid \mathcal{F}_S]. \end{aligned}$$

□

Corollary 3.2.44 (Optional Sampling). *Suppose that $Y_\bullet = (Y_n)_{n \in \mathbb{N}_0}$ is a uniformly integrable submartingale and S, T are a.s. finite stopping times such that $S \leq T$. Then $Y_S \leq \mathbb{E}[Y_T \mid \mathcal{F}_S]$.*

Proof. Use (3.2.31) with the UI submartingale $X = Y^T$ and observe that

$$X_\infty = Y_\infty^T = \hat{Y}_T = Y_T.$$

□

Example 3.2.45 (Optimal Gambling Strategies). Consider a game of chance where the winning probability is $p < \frac{1}{2}$. For example, in the red-and-black roulette game one bets on black with winning probability $p = \frac{18}{38} \approx 0.473$. (The fair case $p = \frac{1}{2}$ is discussed in Exercise 3.21.)

Before each game, the player bets a sum s , called *stake*, that cannot be larger than his fortune at that moment. If he wins, his fortune increases by the amount that he bet. Otherwise he loses his stake.

The player starts with a sum of money x and decides that he will play until the first moment his fortune goes above a set sum, the goal, say 1. His strategy is based on a function $\sigma(x)$. If his fortune after n games is X_n , then the amount he wagers for the next game depends on his current fortune X_n and is $\sigma(X_n)$. The player stops playing when, either he is broke, or he has reached (or surpassed) his goal. The function σ is known as the *strategy* of the gambler.

We denote by $\pi(x, \sigma)$ the probability that the gambler will reach his goal using the strategy σ , given that his initial fortune is x .

We want to show that the strategy that maximizes the winning probability $\pi(x, \sigma)$ is the “*go-bold*” strategy: if your fortune is less than half the goal, bet it all, and if your fortune is more than half the goal, bet as much as you need to reach your goal. Our presentation follows [81, §24.8]. To find out about gambling strategies for more complex games we refer to [63].

First let us introduce the appropriate formalism. The strategies will be chosen from a space $\mathcal{S}$, the collection of measurable functions $\sigma : [0, \infty) \rightarrow [0, \infty)$ such that

$$\sigma(0) = 0, \quad 0 < \sigma(x) \leq x, \quad \forall x \in (0, 1] \quad \text{and} \quad \sigma(x) = 0, \quad \forall x > 1.$$

Note that the stopping rule is built in the definition of $\mathcal{S}$. Also, a strategy σ forces the player to play if he is not bankrupt and has not reached his goal.

The sequence of games encoded by the sequence of i.i.d. random variables $(Y_n)_{n \in \mathbb{N}}$ such that

$$\mathbb{P}[Y_n = 1] = p, \quad \mathbb{P}[Y_n = -1] = 1 - p, \quad 0 < p < \frac{1}{2}.$$

For each $x \geq 0$ and each $\sigma \in \mathcal{S}$ define inductively a sequence of random variables $X_n = X_n^{x, \sigma}$,

$$X_0^x = x, \quad X_{n+1} = X_n + \sigma(X_n)Y_{n+1}, \quad n \geq 0. \quad (3.2.32)$$

We denote by $(\Omega, \mathcal{S}, \mathbb{P})$ the probability space where the random variables X_n and Y_n are defined. Thus $X_n^{x, \sigma}$ is the fortune of the player after n games starting with initial fortune x and using the strategy σ . Note that $\sigma(X_n)$ is the amount of money the player bets before the $(n+1)$ -th game. It depends only on its fortune X_n at that time. If $Y_n = 1$ the player gains $\sigma(X_n)$ and if $Y_n = -1$, the player loses this amount. His strategy σ stays the same for the duration of the game.

Let us observe first that

$$X_\infty^{x, \sigma} := \lim_{n \rightarrow \infty} X_n^{x, \sigma}$$

exists a.s. and L^1 . We will prove this by showing that $X_n^{x, \sigma}$ is a bounded supermartingale.

Since $\sigma(x) \leq x$ we deduce $x - \sigma(x) \geq 0$ and we deduce inductively that $X_n \geq 0$, a.s., $\forall n$. Next, we observe that if $x \leq 1 + x$ then $x + \sigma(x) \leq x + 1$. We deduce inductively that $X_n \leq x + 1$, a.s., $\forall n$.

We have $\mathbb{E}[Y_n] = 2p - 1 < 0$ and thus

$$\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = X_n + \sigma(X_n)\mathbb{E}[Y_{n+1}] \leq X_n.$$

Hence (X_n) is a uniformly bounded supermartingale and thus UI. Set

$$h(x, \sigma) := \mathbb{E}[\min(X_\infty^{x, \sigma}, 1)] \quad \text{and} \quad \pi(x, \sigma) := \mathbb{P}[X_\infty^{x, \sigma} \geq 1].$$

I claim that

$$x \geq h(x, \sigma) \geq \pi(x, \sigma), \quad \forall x \in [0, 1], \quad \sigma \in \mathcal{S}. \quad (3.2.33)$$

Indeed, (X_n) is a supermartingale and the function $x \mapsto \min(x, 1)$ is concave and nondecreasing, so the sequence $\min(X_n, 1)$ is also a supermartingale. Using the continuity of $x \mapsto \min(x, 1)$ we deduce from (i) that

$$\min(X_n^{x, \sigma}, 1) \rightarrow \min(X_\infty^{x, \sigma}, 1) \quad \text{a.s.}$$

Since $0 \leq \min(X_n, 1) \leq 1$ we deduce from the Dominated Convergence theorem that

$$\begin{aligned} x &= \mathbb{E}[\min(X_0, 1)] \geq \mathbb{E}[\min(X_\infty^{x, \sigma}, 1)] \\ &\geq \mathbb{E}[\mathbf{I}_{X_\infty \geq 1}] = \mathbb{P}[X_\infty \geq 1] = \pi(x, \sigma). \end{aligned}$$

Lemma 3.2.46. *If σ is continuous, then $\mathbb{P}[0 < X_\infty < 1] = 0$.*

Proof. We argue by contradiction and assume

$$\mathbb{P}[0 < X_\infty < 1] > 0.$$

Thus, assume there exists $\omega \in \Omega$ such that $X_\infty(\omega) \in (0, 1)$ and

$$\lim_{n \rightarrow \infty} X_n(\omega) = X_\infty(\omega).$$

Using the continuity of σ we deduce

$$\lim_{n \rightarrow \infty} \sigma(X_n(\omega)) = \sigma(X_\infty(\omega)) > 0.$$

On the other hand,

$$\sigma(X_n) = |X_{n+1}(\omega) - X_n(\omega)| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Hence $\mathbb{P}[0 < X_\infty < 1] = 0$ so

$$\mathbb{E}[\min(X_\infty, 1)] = \mathbb{P}[X_\infty \geq 1].$$

□

We have the following *optimality criterion*.

Lemma 3.2.47. *Let $\sigma_0 \in \mathcal{S}$ and set $h_0(x) := h(x, \sigma_0)$, $\pi_0(x) = \pi(x, \sigma_0)$. If $h_0(x)$ is continuous and satisfies,*

$$h_0(x) \geq ph_0(x+s) + (1-p)h_0(x-s), \quad (3.2.34)$$

then, for any $\sigma \in \mathcal{S}$, and any $x \in [0, 1]$ we have $\pi(x, \sigma_0) = h_0(x) \geq \pi(x, \sigma)$.

Proof. Note $h_0(x) = 1$, for $x \geq 1$. Indeed, if the initial fortune x is greater than the goal then

$$h_0(x, \sigma_0) = \mathbb{E}[\min(X_\infty^{x, \sigma_0}, 1)] = \mathbb{E}[\min(X_0, 1)] = 1.$$

Let $\sigma \in \mathcal{S}$ and set $\bar{X}_n := X_n^{x, \sigma}$. Observe that the random process $Z_n = h_0(\bar{X}_n)$ is a supermartingale. Indeed,

$$\begin{aligned} \mathbb{E}[h_0(\bar{X}_{n+1}) \mid \mathcal{F}_n] &= \mathbb{E}[h_0(\bar{X}_n + \sigma(\bar{X}_n)Y_{n+1}) \mid \mathcal{F}_n] \\ &= \mathbb{E}[h_0(\bar{X}_n + \sigma(\bar{X}_n))\mathbf{I}_{\{Y_{n+1}=1\}} + h_0(\bar{X}_n - \sigma(\bar{X}_n))\mathbf{I}_{\{Y_{n+1}=-1\}} \mid \mathcal{F}_n] \\ &= ph_0(\bar{X}_n + \sigma(\bar{X}_n)) + (1-p)h_0(\bar{X}_n - \sigma(\bar{X}_n)) \stackrel{(3.2.34)}{\leq} h_0(\bar{X}_n). \end{aligned}$$

Hence, Z_n is a bounded supermartingale and thus

$$h_0(x) = \mathbb{E}[h_0(X_0^{x, \sigma})] = \mathbb{E}[Z_0] \geq \mathbb{E}[Z_\infty].$$

On the other hand, since $h_0(x)$ is continuous and bounded we deduce that $h_0(X_n)$ converges a.s. and L^1 to $h_0(X_\infty)$. Thus

$$\mathbb{E}[Z_\infty] = \mathbb{E}[h_0(X_\infty^{x, \sigma})] \geq \mathbb{P}[X_\infty^{x, \sigma} \geq 1] \geq \pi(x, \sigma).$$

□

Define $\sigma_0 \in \mathcal{S}$

$$\sigma_0(x) := \begin{cases} \min(x, 1-x), & x \in [0, 1], \\ 0, & x \geq 1, \end{cases}$$

and set $h_0(x) := h(x, \sigma_0)$, $\pi_0(x) = \pi(x, \sigma_0)$. We want to show that σ_0 satisfies all the conditions of Lemma 3.2.47.

Clearly σ_0 is a continuous strategy. By construction, for any $x \in [0, 1]$ we have $0 \leq X_n^{x, \sigma_0} \leq 1$ a.s.. Moreover, Lemma 3.2.46 shows that

$$X_\infty^{x, \sigma_0} \in \{0, 1\} \text{ a.s.}$$

so

$$\pi_0(x) = h_0(x) = \mathbb{E}[\min(X_\infty^{x, \sigma_0}, 1)], \quad \forall x \in [0, 1].$$

The functions

$$[0, 1] \ni x \mapsto x + \sigma_0(x) \in [0, 1], \quad [0, 1] \ni x \mapsto x - \sigma_0(x) \in [0, 1]$$

are non-decreasing. We deduce inductively that if $x \leq y$ then

$$\mathbb{E}[X_n^{x, \sigma_0}] \leq \mathbb{E}[X_n^{y, \sigma_0}]$$

and, by letting $n \rightarrow \infty$ we deduce that $h_0(x) \leq h_0(y)$ so that h_0 is non-decreasing.

By conditioning on Y_1 we deduce that

$$h_0(x) = \begin{cases} ph_0(2x), & x \leq 1/2, \\ p + (1-p)h_0(2x-1), & 1/2 \leq x \leq 1, \\ 1, & x > 1. \end{cases} \quad (3.2.35)$$

Set

$$\mathcal{D} := \left\{ \frac{k}{2^n}; \quad n \in \mathbb{N}_0, \quad 0 \leq k \leq 2^n \right\}.$$

We will prove by induction on n that the optimality condition (3.2.34) holds for x of the form $x = \frac{k}{2^n}$. Start with $n = 1$ so $x = \frac{1}{2}$. We have

$$\begin{aligned} & h_0(1/2) - ph_0(1/2 + s) - (1-p)h_0(1/2 - s) \\ & \stackrel{(3.2.35)}{=} p - p\left(p + (1-p)h_0(2s)\right) - (1-p)ph_0(1-2s) \\ & = p(1-p)\left(1 - (h_0(2s) + h_0(1-2s))\right) \geq 0, \end{aligned}$$

where at the last step we used the fact that $h_0(x) \leq x, \forall x \in [0, 1]$.

For the inductive step, assume that $n > 1$ and $x = \frac{k}{2^n}, k < 2^n$. Choose $s \in [0, x]$. We consider several cases.

Case 1. $x + s \leq \frac{1}{2}$. Using (3.2.35) and the induction hypothesis we deduce

$$\begin{aligned} ph_0(x+s) + (1-p)h_0(x-s) &= p(ph_0(2x+2s) + (1-p)ph_0(2x-2s)) \\ &\leq ph_0(2x) = h_0(x). \end{aligned}$$

Case 2. $x - s \geq \frac{1}{2}$. Similar to **Case 1**.

Case 3. $x \leq \frac{1}{2}$ and $x + s \geq \frac{1}{2}$. Using (3.2.35) we have

$$\begin{aligned} A &:= h_0(x) - ph_0(x+s) - (1-p)h_0(x-s) \\ &= ph_0(2x) - p\left(p + (1-p)h_0(2x+2s-1)\right) - (1-p)ph_0(2x-2s) \\ &= p\left(h_0(2x) - p - (1-p)h_0(2x+2s-1) - (1-p)h_0(2x-2s)\right). \end{aligned}$$

Observe that $\frac{1}{2} \leq x+s \leq 2x \leq 1$. Using (3.2.35) we deduce

$$h_0(2x) = p + (1-p)h_0(4x-1)$$

so that

$$\begin{aligned} A &= p\left(p + (1-p)h_0(4x-1) - p - (1-p)h_0(2x+2s-1) - (1-p)h_0(2x-2s)\right) \\ &= p(1-p)\left(h_0(4x-1) - h_0(x+s-1/2) - h_0(x-s)\right) \end{aligned}$$

$$\begin{aligned}
&= (1-p) \left(h_0(2x-1/2) - ph_0(2x+2s-1) - ph_0(2x-2s) \right) \\
(p \leq 1-p) \\
&\geq (1-p) \left(h_0(2x-1/2) - ph_0(2x+2s-1) - (1-p)h_0(2x-2s) \right).
\end{aligned}$$

The induction hypothesis implies

$$\begin{aligned}
&h_0(2x-1/2) - ph_0(2x+2s-1) - (1-p)h_0(x-2s) \\
(x' = 2x-1/2, s' = 2s-1/2) \\
&= h_0(x') - ph(x' + s') + (1-p)h(x' - s') \geq 0.
\end{aligned}$$

Case 4. $x \geq \frac{1}{2}$ and $x-s \leq \frac{1}{2}$. This is similar to the previous case.

We can now prove that h_0 is continuous. Since h is nondecreasing we deduce that the right/left limits $h(x \pm 0)$ exist at each $x \in [0, 1]$. Since (3.2.34) holds for every x in a dense set we deduce

$$h(x-0) \geq ph_0((x+s)-0) + (1-p)h_0((x-s)-0)$$

$\forall 0 \leq s < x \leq 1$. Now let $s \searrow 0$ to conclude

$$h_0(x-0) \geq ph(x+0) + (1-p)h_0(x-0) \Rightarrow ph_0(x-0) \geq ph_0(x+0)$$

so that $h_0(x-0) = h_0(x+0)$, i.e., h_0 is continuous. Since $\mathcal{D}$ is dense in $[0, 1]$ we deduce that h_0 satisfies (3.2.34) on $[0, 1]$. We can now invoke Lemma 3.2.47 to deduce that

$$\pi(x, \sigma_0) = h_0(x) \geq \pi(x, \sigma), \quad \forall x \in [0, 1], \quad \sigma \in \mathcal{S},$$

i.e., σ_0 is an optimal gambling strategy.

Let us explain how to compute $h_0(x)$, $x \in \mathcal{D}$. Every number $x \in \mathcal{D}$ has a binary expansion

$$x = 0.\epsilon_1\epsilon_2\cdots = \sum_{n \geq 1} \frac{\epsilon_n}{2^n}$$

where $\epsilon_n \in \{0, 1\}$, and $\epsilon_n = 0$ for $n \gg 0$. Note that

$$x < \frac{1}{2} \iff \epsilon_1 = 0.$$

The first equation in (3.2.35) reads

$$h_0(0.0\epsilon_2\cdots) = p \cdot h_0(0.\epsilon_2\cdots).$$

In particular

$$h_0(0.\underbrace{0\cdots 0}_k 1\epsilon_{k+2}\cdots) = p^k(0.1\epsilon_{k+2}\cdots).$$

The second equation in (3.2.35) reads

$$h_0(0.1\epsilon_2\cdots) = p + (1-p)h_0(0.\epsilon_2\cdots).$$

We define $f_0, f_1 : [0, 1] \rightarrow [0, 1]$ by

$$f_0(x) = px, \quad f_1(x) = p + (1-p)x.$$

The above discussion shows that

$$h_0(0.\epsilon_1\cdots\epsilon_n) = f_{\epsilon_1}(h_0(0.\epsilon_2\cdots\epsilon_n)).$$

Since $h_0(0) = 0$ and $h_0(1/2) = p$ we deduce by iteration that if,

$$x = 0.\epsilon_1\epsilon_2\cdots\epsilon_n 1,$$

then

$$h_0(x) = f_{\epsilon_1} \circ f_{\epsilon_2} \circ \cdots \circ f_{\epsilon_n}(p).$$

Thus h is uniquely determined on $\mathcal{D}$ and, since $\mathcal{D}$ is dense on $[0, 1]$, the function h is uniquely determined on $[0, 1]$. Let us emphasize that $h_0(x)$ depends on the winning probability p .

As an illustration let us compute $h_0(21/32)$. Note that $\frac{21}{32}$ has the binary expansion

$$\frac{21}{32} = 0.10101$$

so that

$$\begin{aligned} h_0(21/32) &= f_1 \circ f_0 \circ f_1 \circ f_0(p) = f_1 \circ f_0 \circ f_1(p^2) \\ &= f_1 \circ f_0(p + p^2 - p^3) = f_1(p^2 + p^3 - p^4) = p + (1 - p)(p^2 + p^3 - p^4) \\ &= p + p^2 + p^3 - p^4 - p^3 - p^4 + p^5 = p + p^2 - 2p^4 + p^5. \end{aligned}$$

For example, if $p = 0.4$, then $h_0(21/32) = 0.519 > 0.5$. Thus, although the winning probability $p = 0.4 < 0.5$, using this strategy with an initial fortune $21/32$, the odds of increasing the fortune to 1 are better than 50 : 50.

If the initial fortune is $x = \frac{1}{4}$, then using its binary expansion $\frac{1}{4} = 0.01$ we deduce

$$h_0(1/4) = ph_0(1/2) = \frac{p}{2}.$$

In this case, if $p = 0.4$, the probability of reaching his goal is 0.2, substantially smaller, in fact smaller than p . Thus, with a small initial fortune the probability of achieving is smaller than the probability of winning the first game.

In the R-generated Figure 3.3 we have depicted the graphs of $x \mapsto h_0(x, p)$ for different values of p .

In the top left corner of Figure 3.3 we depicted the optimal winning probability for the game of roulette, i.e., $p = 18/38$. For example

$$h_0(33/64, 18/38) \approx 0.4862, \quad h_0(34/64, 18/38) \approx 0.5001,$$

$$h_0(35/64, 18/38) \approx 0.5141.$$

Noting that $35/64 = 0.546875$ we conclude that if the initial fortune is bigger than 55% of the goal, then the go-bold strategy will get you to your goal with probability 51%. So, if you have money to burn, going bold will beat the house in the long run.

In Figure 3.4 we have depicted the graph of the function $p \mapsto h(63/64, p)$. Since the binary expansion of $63/64$ is $0.\underbrace{111111}_6$ we deduce that

$$h_0(63/64, p) = \underbrace{f_1 \circ \cdots \circ f_1}_5(p) = 1 - (1 - p)^6.$$

If $p = 1 - 2^{-1/6} \approx 0.109$, then $h(63/64, p) = 1/2$. Thus, if the probability p of winning one game is > 0.11 and your initial fortune is $> 63/64 \approx 0.98$ your chance of reaching your goal is using go-bold strategy is $> 1/2$ although the probability of winning a single game is small. $\square$




Figure 3.3. The plot of $x \mapsto h_0(x, p)$, for $p \in \{0.1, 0.25, 0.4, 18/38\}$.




Figure 3.4. The plot of $p \mapsto h_0(63/64, p) = 1 - (1 - p)^6$, $p \in [0, 1/2)$.

3.2.6. Maximal inequalities and L^p -convergence. The results in this subsection are wide ranging generalizations of Kolmogorov's one series theorem. They depend on Doob's maximal inequality which generalizes Kolmogorov's inequality (2.1.3).

Theorem 3.2.48 (Doob's maximal inequality). *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a submartingale. Set*

$$\tilde{X}_n := \sup_{k \leq n} X_k.$$

Then, for any $a > 0$, we have

$$a\mathbb{P}[\tilde{X}_n \geq a] \leq \mathbb{E}[X_n \mathbf{I}_{\{\tilde{X}_n \geq a\}}] \leq \mathbb{E}[X_n^+]. \quad (3.2.36)$$

Proof. Let us introduce the stopping time

$$T := \inf\{n \geq 0; X_n \geq a\}.$$

Then

$$A := \{\tilde{X}_n \geq a\} = \left\{ \sup_{k \leq n} X_k \geq a \right\} = \{T \leq n\}.$$

Applying the Optional Sampling Theorem 3.1.29 to the bounded stopping times $T \wedge n$ and n we deduce $\mathbb{E}[X_{T \wedge n}] \leq \mathbb{E}[X_n]$.

On the other hand,

$$X_{T \wedge n}(\omega) = X_{T(\omega)} \mathbf{I}_A(\omega) + X_n(\omega) \mathbf{I}_{A^c}(\omega) \geq a \mathbf{I}_A(\omega) + X_n(\omega) \mathbf{I}_{A^c}(\omega),$$

so $X_{T \wedge n} \geq a \mathbf{I}_A + X_n \mathbf{I}_{A^c}$. We deduce

$$a\mathbb{P}[A] + \mathbb{E}[X_n \mathbf{I}_{A^c}] \leq \mathbb{E}[X_{T \wedge n}] \leq \mathbb{E}[X_n] = \mathbb{E}[X_n \mathbf{I}_A] + \mathbb{E}[X_n \mathbf{I}_{A^c}]$$

so that

$$a\mathbb{P}[\tilde{X}_n \geq a] = a\mathbb{P}[A] \leq \mathbb{E}[X_n \mathbf{I}_{\{\tilde{X}_n \geq a\}}] \leq \mathbb{E}[X_n^+].$$

□

Corollary 3.2.49. *Suppose that (Y_n) is a martingale. We set*

$$Y_n^* = \max_{k \leq n} |Y_k|.$$

Then for every $c > 0$ and any $p \in [1, \infty)$ we have

$$\mathbb{P}[Y_n^* > c] \leq \frac{1}{c^p} \mathbb{E}[|Y_n|^p].$$

Proof. Doob's maximal inequality applied to the submartingale $X_n = |Y_n|^p$ yields

$$\mathbb{P}[Y_n^* > c] = \mathbb{P}\left[\max_{0 \leq k \leq n} |Y_k|^p > c^p\right] \leq \frac{1}{c^p} \mathbb{E}[|Y_n|^p]$$

□

Theorem 3.2.50 (Doob's L^p -inequality). *Let $p > 1$ and suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a positive submartingale such that $X_n \in L^p, \forall n \geq 0$. Set*

$$\tilde{X}_n := \sup_{k \leq n} X_k.$$

Then for any $n \geq 0$ we have

$$\mathbb{E}[(\tilde{X}_n)^p]^{\frac{1}{p}} \leq q \mathbb{E}[X_n^p]^{\frac{1}{p}}, \quad (3.2.37)$$

where

$$\frac{1}{p} + \frac{1}{q} = 1 \quad \text{or} \quad q = \frac{p}{p-1}.$$

In particular, if $(Y_n)_{n \in \mathbb{N}_0}$ is a martingale and if

$$Y_n^* := \max_{k \leq n} |Y_k|,$$

then, for any $n \geq 0$, we have

$$\|Y_n^*\|_{L^p} \leq q \|Y_n\|_{L^p}. \quad (3.2.38)$$

Proof. Clearly (3.2.37) $\Rightarrow$ (3.2.38). Note that $(X_n^p)_{n \geq 0}$ is also a submartingale and $\tilde{X}_n \in L^p$. From Doob's maximal inequality we deduce

$$a \mathbb{P}[\tilde{X}_n \geq a] \leq \mathbb{E}[X_n \mathbf{I}_{\{\tilde{X}_n \geq a\}}]$$

so

$$\frac{1}{p} \mathbb{E}[\tilde{X}_n^p] \stackrel{(1.3.48)}{=} \int_0^\infty a^{p-1} \mathbb{P}[\tilde{X}_n \geq a] da \stackrel{(3.2.36)}{\leq} \int_0^\infty a^{p-2} \mathbb{E}[X_n \mathbf{I}_{\{\tilde{X}_n \geq a\}}] da.$$

Switching the order of integration we deduce

$$\int_0^\infty a^{p-2} \mathbb{E}[X_n \mathbf{I}_{\{\tilde{X}_n \geq a\}}] da = \mathbb{E}\left[X_n \int_0^{\tilde{X}_n} a^{p-2} da\right] = \frac{1}{p-1} \mathbb{E}[X_n \tilde{X}_n^{p-1}]$$

(use Hölder's inequality with $\frac{1}{q} = 1 - \frac{1}{p}$)

$$\leq \frac{1}{p-1} \mathbb{E}[X_n^p]^{\frac{1}{p}} \mathbb{E}[\tilde{X}_n^p]^{\frac{p-1}{p}}.$$

Hence

$$\frac{1}{p} \mathbb{E}[\tilde{X}_n^p] \leq \frac{1}{p-1} \mathbb{E}[X_n^p]^{\frac{1}{p}} \mathbb{E}[\tilde{X}_n^p]^{\frac{p-1}{p}}.$$

This proves (3.2.37). $\square$

Definition 3.2.51. Let $p \in [1, \infty)$. A martingale $(X_n)_{n \in \mathbb{N}_0}$ is called an L^p -martingale if

$$\mathbb{E}[|X_n|^p] < \infty, \quad \forall n \in \mathbb{N}_0.$$

A *bounded L^p -martingale* is a martingale $(X_n)_{n \in \mathbb{N}_0}$ such that

$$\sup_{n \in \mathbb{N}_0} \mathbb{E}[|X_n|^p] < \infty. \quad \square$$

Corollary 3.2.52 (L^p -martingale convergence theorem). *Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is a bounded L^p -martingale for some $p > 1$. Set*

$$X_n^* := \max_{k \leq n} |X_k|, \quad X_\infty^* = \sup_{k \geq 0} |X_k| = \lim_{n \rightarrow \infty} X_n^*.$$

Then $(X_n)_{n \in \mathbb{N}_0}$ is a UI martingale and X_n converges a.s. and L^p to a random variable

$$X_\infty \in L^p(\Omega, \mathcal{F}_\infty, \mathbb{P}).$$

Moreover

$$\mathbb{E}[(X_\infty^*)^p] \leq \left(\frac{p}{p-1}\right)^p \mathbb{E}[|X_\infty|^p].$$

Proof. From the Monotone Convergence Theorem we deduce

$$\mathbb{E}[(X_\infty^*)^p] = \lim_{n \rightarrow \infty} \mathbb{E}[(X_n^*)^p] \leq \left(\frac{p}{p-1}\right)^p \sup_{n \geq 0} \mathbb{E}[|X_n|^p] < \infty.$$

so $X_\infty^* \in L^p$ and $|X_n| \leq X_\infty^*, \forall n \geq 0$. The desired conclusions now follow from the martingale convergence theorem and the Dominated Convergence Theorem. $\square$

Example 3.2.53 (Kolmogorov's one series theorem). Suppose that $(X_n)_{n \geq 0}$ is a sequence of independent random variables such that

$$\mathbb{E}[X_n] = 0, \quad \forall n \geq 0, \quad \sum_{n \geq 0} \text{Var}[X_n] < \infty.$$

Then the random series $X_0 + X_1 + \dots$ is a.s. and L^2 -convergent. Indeed, the sequence of partial sums

$$S_n = X_0 + \dots + X_n$$

is a bounded L^2 -martingale and so it converges a.s. and L^2 . $\square$

Example 3.2.54 (Likelihood ratio). This example has origin in statistics. Suppose that we have a random quantity and we have reasons to believe that its probability distribution is either of the form $p(x)dx$ or $q(x)dx$ where $p, q : \mathbb{R} \rightarrow [0, \infty)$ are mutually absolutely continuous probability densities on $\mathbb{R}$

$$\int_{\mathbb{R}} p(x)dx = \int_{\mathbb{R}} q(x)dx = 1.$$

We want to describe a statistical test that helps deciding which is the real distribution. Our presentation follows [94, Sec.12.8].

We take a large number of samples of the random quantity, or equivalently, suppose that we are given a sequence of i.i.d. random variables $(X_n)_{n \geq 1}$ with common probability density f , where f is one of the two densities p or q . Assume for simplicity that $q(x) = 0 \Rightarrow p(x) = 0$ and agree to set $\frac{0}{0} = 0$. The products

$$Y_n := \prod_{k=1}^n \frac{p(X_k)}{q(X_k)}.$$

are called *likelihood ratios*. Note that if $f = q$, then $\mathbb{E}[Y_n] = 1, \forall n$.

To decide whether $f = q$ or $f = p$ we fix a (large) positive number a and a large $n \in \mathbb{N}$ and adopt the prediction strategy

$$\bar{f}_n := \begin{cases} p, & Y_n \geq a, \\ q, & Y_n < a. \end{cases}$$

We want to show that this strategy picks the correct density with high confidence, i.e., $\mathbb{P}[f = \bar{f}_n]$ is very close to 1 for large n and a .

If $f = q$, then Y_n is a product of i.i.d. nonnegative random variables with mean 1 and, as shown in Example 3.1.6, it is a martingale with respect to the filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$.

The function $\log$ is *strictly* concave and we deduce from Jensen's inequality

$$\mathbb{E}\left[\log \frac{p(X_n)}{q(X_n)}\right] < \log \mathbb{E}\left[\frac{p(X_n)}{q(X_n)}\right] = 0.$$

The Strong Law of Large Numbers shows that

$$\frac{1}{n} \sum_{k=1}^n \log \frac{p(X_k)}{q(X_k)} \rightarrow \mathbb{E}[\log Y_1] < 0, \quad \text{a.s.}$$

Thus

$$\log Y_n = \sum_{k=1}^n \log \frac{p(X_k)}{q(X_k)} \rightarrow -\infty \quad \text{a.s.}$$

Thus, if $f = q$, then $Y_n \rightarrow 0$ a.s.. If $f = p$, then a similar argument shows that $\frac{1}{Y_n} \rightarrow 0$ a.s.. We deduce that

$$Y_n \rightarrow \begin{cases} 0, & f = q, \\ \infty, & f = p. \end{cases}$$

In particular,

$$\lim_{n \rightarrow \infty} \mathbb{P}[\bar{f}_n = q \mid f = q] = \lim_{n \rightarrow \infty} \mathbb{P}[Y_n < a] = 1,$$

and

$$\lim_{n \rightarrow \infty} \mathbb{P}[\bar{f}_n = p \mid f = p] = \lim_{n \rightarrow \infty} \mathbb{P}[Y_n \geq a] = 1.$$

Moreover, Doob's maximal inequality (3.2.36) shows that if $f = q$, so Y_n is a martingale, we have

$$\mathbb{P}\left[\max_{1 \leq k \leq n} Y_k \geq a\right] \leq \frac{1}{a}, \quad \forall n.$$

Thus, $\forall n$,

$$\mathbb{P}[\bar{f}_n \neq q \mid f = q] = \mathbb{P}[Y_n \geq a \mid f = q] \leq \frac{1}{a} \rightarrow 0 \quad \text{as } a \rightarrow \infty,$$

and this statistical test makes the right decision with high confidence if $a \gg 1$. $\square$

Example 3.2.55. Consider again the branching process in Example 3.1.8. Suppose that the reproduction law μ satisfies

$$m = \sum_{k=0}^{\infty} k\mu(k) < \infty, \quad \sum_{k=0}^{\infty} k^2\mu(k) < \infty.$$

We set

$$\sigma^2 := \text{Var}[\mu] = \sum_{k=0}^{\infty} k^2\mu(k) - m^2.$$

Note that

$$\begin{aligned} Z_{n+1} &= \sum_{k=1}^{\infty} \left(\sum_{j=1}^k X_{n,j} \right) \mathbf{I}_{\{Z_n=k\}} \\ &= \sum_{j=0}^{\infty} X_{n,j} \sum_{k \geq j} \mathbf{I}_{\{Z_n=k\}} = \sum_{j=0}^{\infty} X_{n,j} \mathbf{I}_{\{Z_n \geq j\}}, \end{aligned}$$

$$\mathbb{E}[Z_{n+1}^2 \mid \mathcal{F}_n] = \mathbb{E}\left[\sum_{k,j=1}^{\infty} \mathbf{I}_{\{Z_n \geq j, Z_n \geq k\}} X_{n,j} X_{n,k} \mid \mathcal{F}_n\right]$$

$(X_{n,j}, X_{n,k} \perp \mathcal{F}_n)$

$$= \sum_{k,j=1}^{\infty} \mathbf{I}_{\{Z_n \geq j, Z_n \geq k\}} \mathbb{E}[X_{n,j} X_{n,k}] = \sum_{k,j=1}^{\infty} \mathbf{I}_{\{Z_n \geq j, Z_n \geq k\}} (m^2 + \delta_{jk} \sigma^2)$$

$$\begin{aligned}
&= m^2 \sum_{j,k=j=1}^{\infty} I_{\{Z_n \geq j\}} I_{\{Z_n \geq k\}} + \sigma^2 \sum_{k=1}^{\infty} I_{\{Z_n \geq k\}} \\
(\mathbb{E}[Z_n] = \sum_{k \geq 1} \mathbb{P}(Z_n \geq k)) \\
&= m^2 \left(\sum_{k=1}^{\infty} I_{\{Z_n \geq k\}} \right)^2 + \sigma^2 \sum_{k=1}^{\infty} I_{\{Z_n \geq k\}} = m^2 Z_n^2 + \sigma^2 Z_n.
\end{aligned}$$

Hence

$$\mathbb{E}[Z_{n+1}^2] = m^2 \mathbb{E}[Z_n^2] + m^2 \mathbb{E}[Z_n] = \sigma^2 \mathbb{E}[Z_n^2] + \sigma^2 m^n \mathbb{E}[Z_0] = m^2 \mathbb{E}[Z_n^2] + \sigma^2 m^n \ell.$$

We set

$$q_{n+1} := m^{-2n} \mathbb{E}[Z_n^2]$$

and we get from the above that

$$q_{n+1} = q_n + m^{-n-2} \sigma^2 \ell.$$

This shows that if $m > 1$, then the sequence (q_n) converges, so the martingale $W_n := m^{-n} Z_n$ converges in L^2 and a.s. The limit W_∞ is nonzero if $\ell = \mathbb{E}[Z_0] > 0$, because

$$\mathbb{E}[W_\infty] = \mathbb{E}[W_0] = \mathbb{E}[Z_0] = \ell.$$

We refer to Exercise 3.29 for more details about W_∞ . □

3.2.7. Backwards martingales. Suppose that the parameter set $\mathbb{T}$ is

$$\mathbb{T} = -\mathbb{N}_0 = \{0, -1, -2, \dots\}$$

In this case a $\mathbb{T}$ -filtration $\mathcal{F}_n$, $n \in -\mathbb{N}_0$ is called a *backwards filtration*. We set

$$\mathcal{F}_{-\infty} := \bigcap_{n \leq 0} \mathcal{F}_n.$$

A *backwards martingale* (submartingale, supermartingale) is a martingale (resp. submartingale, supermartingale) adapted to a backwards filtration.

Theorem 3.2.56 (Convergence of backwards submartingales). *Suppose that $\mathcal{F}_\bullet = (\mathcal{F}_n)_{n \in -\mathbb{N}_0}$ is a backwards filtration of $(\Omega, \mathcal{S}, \mathbb{P})$ and $X_\bullet = (X_n)_{n \in -\mathbb{N}_0}$ is $\mathcal{F}_\bullet$ -submartingale, i.e.,*

$$X_n \leq \mathbb{E}[X_m | \mathcal{F}_n], \quad \forall n, m \in -\mathbb{N}_0, \quad n \leq m,$$

and

$$C := \inf_{n \leq 0} \mathbb{E}[X_n] > -\infty.$$

Then the following hold.

- (i) The family $(X_n)_{n \in -\mathbb{N}_0}$ is UI.
- (ii) There exists $X_{-\infty} \in L^1(\Omega, \mathcal{F}_{-\infty}, \mathbb{P})$ such that $X_n \rightarrow X_{-\infty}$ a.s. and L^1 as $n \rightarrow -\infty$.

Moreover

$$X_{-\infty} \leq \mathbb{E}[X_n | \mathcal{F}_{-\infty}], \tag{3.2.39}$$

with equality if $(X_n)_{n \in -\mathbb{N}_0}$ is a martingale.

Proof. Step 1. Boundedness in L^1 . Observe that (X_n^+) is a submartingale and thus

$$\mathbb{E}[X_n^+] \leq \mathbb{E}[X_0^+], \quad \forall n \leq 0.$$

On the other hand, there exists $C \in \mathbb{R}$ such that

$$\mathbb{E}[X_n] = \mathbb{E}[X_n^+] - \mathbb{E}[X_n^-] \geq C, \quad \forall n \geq 0.$$

Hence

$$\mathbb{E}[X_n^-] \leq C + \mathbb{E}[X_n^+] \leq C + \mathbb{E}[X_0^+], \quad \forall n \leq 0,$$

and consequently,

$$Z := \sup_{n \leq 0} \mathbb{E}[|X_n|] < \infty. \quad (3.2.40)$$

Step 2. Almost sure convergence. For $K \in \mathbb{N}$ consider the (increasing) filtration

$$\mathcal{G}_n^K := \mathcal{F}_{(-K+n) \wedge 0}, \quad n \in \mathbb{N}_0,$$

and the $\mathcal{G}_n^K$ -submartingale $Y_n^K = X_{(-K+n) \wedge 0}$. Thus

$$Y_0^K = X_{-K}, Y_1^K = X_{-K+1}, \dots, Y_K^K = X_0, Y_{K+1}^K = X_0, \dots$$

Doob's upcrossing inequality applied to the submartingale Y_n^K shows that, for any rational numbers $a < b$ we have

$$\begin{aligned} (b-a)\mathbb{E}[N_K([a, b], Y^K)] &\leq \mathbb{E}[(X_0 - a)^+] - \mathbb{E}[(X_{-K} - a)^+] \\ &\leq \mathbb{E}[(X_0 - a)^+] \leq |a| + \mathbb{E}[|X_0|]. \end{aligned}$$

This proves that, for any rational numbers $a < b$, the nondecreasing sequence

$$K \mapsto N_K([a, b], Y^K)$$

is also bounded, and thus it has a finite limit $N_\infty([a, b], X)$ as $K \rightarrow \infty$. An obvious version of Lemma 3.2.1 shows that X_n has an a.s. limit a.s. $n \rightarrow -\infty$. The limit is a $\mathcal{F}_{-\infty}$ -measurable random variable $X_{-\infty}$. Fatou's Lemma shows that

$$\mathbb{E}[|X_{-\infty}|] < \infty.$$

Step 3. Uniform integrability. This is obvious if $(X_n)_{n \leq 0}$ is a martingale since

$$X_n = \mathbb{E}[X_0 | \mathcal{F}_n]$$

and the conclusion follows from Corollary 3.2.20.

In general, if $(X_n)_{n \leq 0}$ is a submartingale, we have

$$\mathbb{E}[X_n] \leq \mathbb{E}[X_m], \quad \forall n \leq m \leq 0.$$

Since the sequence $\mathbb{E}[X_n]$ is bounded below we deduce that it has a finite limit. Thus, for any $\varepsilon > 0$, there exists $K = K(\varepsilon) > 0$ such that

$$\mathbb{E}[X_{-n}] \geq \mathbb{E}[X_{-K}] - \frac{\varepsilon}{2}, \quad \forall n \geq K.$$

For $n > K$ and $a > 0$ we have

$$\begin{aligned} \mathbb{E}[|X_{-n}| \mathbf{I}_{\{|X_{-n}| > a\}}] &= \mathbb{E}[(-X_{-n}) \mathbf{I}_{\{X_{-n} < -a\}}] + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} > a\}}] \\ &= \boxed{-\mathbb{E}[X_{-n}]} + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} \geq -a\}}] + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} > a\}}] \\ &\leq \boxed{-\mathbb{E}[X_{-K}] + \frac{\varepsilon}{2}} + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} \geq -a\}}] + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} > a\}}]. \end{aligned}$$

Now observe that, for any $H \in \mathcal{F}_n$, we have

$$X_{-n} \mathbf{I}_H \leq \mathbb{E}[X_{-K} | \mathcal{F}_{-n}] \mathbf{I}_H = \mathbb{E}[X_{-K} \mathbf{I}_H | \mathcal{F}_{-n}],$$

so

$$\mathbb{E}[X_{-n} \mathbf{I}_H] \leq \mathbb{E}[X_{-K} \mathbf{I}_H].$$

Hence, if $H = \{X_{-n} \geq -a\}$, or $H = \{X_{-n} > a\}$, then

$$\mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} \geq -a\}}] + \mathbb{E}[X_{-n} \mathbf{I}_{\{X_{-n} > a\}}] \leq \mathbb{E}[X_{-K} \mathbf{I}_{\{X_{-n} \geq -a\}}] + \mathbb{E}[X_{-K} \mathbf{I}_{\{X_{-n} > a\}}],$$

and

$$\begin{aligned} \mathbb{E}[|X_{-n}| \mathbf{I}_{\{|X_{-n}| > a\}}] &\leq -\mathbb{E}[X_{-K}] + \mathbb{E}[X_{-K} \mathbf{I}_{\{X_{-n} \geq -a\}}] + \mathbb{E}[X_{-K} \mathbf{I}_{\{X_{-n} > a\}}] + \frac{\varepsilon}{2} \\ &= \mathbb{E}[|X_{-K}| \mathbf{I}_{\{|X_{-n}| \geq a\}}] + \frac{\varepsilon}{2}. \end{aligned}$$

From Markov's inequality and (3.2.40) we deduce

$$\mathbb{P}[|X_{-m}| > a] \leq \frac{Z}{a}, \quad \forall m \in \mathbb{N}_0.$$

Since the family consisting of the single random variable X_{-K} is uniformly integrable, we deduce that there exists $\delta = \delta(\varepsilon) > 0$ such that, for any $A \in \mathcal{F}_K$ satisfying $\mathbb{P}[A] < \delta$ we have

$$\mathbb{E}[|X_{-K}| \mathbf{I}_A] < \frac{\varepsilon}{2}.$$

We deduce that for any $a > 0$ such that $\frac{Z}{a} < \delta(\varepsilon)$ we have

$$\mathbb{E}[|X_{-n}| \mathbf{I}_{\{|X_{-n}| > a\}}] \leq \mathbb{E}[|X_{-K}| \mathbf{I}_{\{|X_{-n}| \geq a\}}] < \frac{\varepsilon}{2}.$$

This proves that the family $(X_{-n})_{n \in \mathbb{N}_0}$ is UI.

Step 4. Conclusion. Finally, observe that for any $A \in \mathcal{F}_{-\infty}$ and any $n \leq m \leq 0$ we have $\mathbb{E}[X_n \mathbf{I}_A] \leq \mathbb{E}[X_m \mathbf{I}_A]$. If we let $n \rightarrow -\infty$ we deduce

$$\mathbb{E}[X_{-\infty} \mathbf{I}_A] \leq \mathbb{E}[X_m \mathbf{I}_A], \quad \forall m \leq 0, \quad A \in \mathcal{F}_{-\infty}$$

This is precisely the inequality (3.2.39). When (X_n) is a martingale all the above inequalities are equalities. $\square$

Corollary 3.2.57 (Backwards Martingale Convergence). *Suppose that $(\mathcal{G}_n)_{n \in \mathbb{N}_0}$ is a decreasing family of σ -subalgebras of $\mathcal{F}$ and*

$$Z \in L^1(\Omega, \mathcal{S}, \mathbb{P}).$$

Then the sequence $\mathbb{E}[Z | \mathcal{G}_n]$ converges a.s. and L^1 to $\mathbb{E}[Z | \mathcal{G}_{\infty}]$, where

$$\mathcal{G}_{\infty} = \bigcap_{n \geq 0} \mathcal{G}_n.$$

Proof. Apply the previous theorem to the backwards filtration $\mathcal{F}_n := \mathcal{G}_{-n}$, $n \leq 0$, and the martingale $Z_n := \mathbb{E}[Z | \mathcal{F}_n]$, $n \leq 0$. $\square$

3.2.8. Exchangeable sequences of random variables. An n -dimensional random vector

$$\mathbf{X} = (X_1, \dots, X_n)$$

is called *exchangeable* if, for any permutation π of $\llbracket 1, n \rrbracket$, the random vectors $(X_1, \dots, X_n)$ and $(X_{\pi(1)}, \dots, X_{\pi(n)})$ have identical distributions.

A sequence of random variables $(X_k)_{k \in \mathbb{N}}$ is called *exchangeable* if for any $n \in \mathbb{N}$ the random vector $(X_1, \dots, X_n)$ is exchangeable. One also refers to an exchangeable sequence as an *exchangeable process*.

Equivalently, if we denote by $\mathfrak{S}_n$ the subgroup of permutations φ of $\mathbb{N}$ such that $\varphi(r) = r$, $\forall r > n$, then the sequence $(X_n)_{n \geq 1}$ is exchangeable if for any $n \in \mathbb{N}$ and any $\varphi \in \mathfrak{S}_n$ the sequences $(X_n)_{n \in \mathbb{N}}$ and $(X_{\varphi(n)})_{n \in \mathbb{N}}$ are identically distributed.

Example 3.2.58. (a) A sequence of i.i.d. random variables $(X_n)_{n \geq 1}$ is exchangeable.

(b) Suppose that $(\mu_\lambda)_{\lambda \in \Lambda}$ is a family of Borel probability measures on $\mathbb{R}$ parametrized by a probability space $(\Lambda, \mathcal{S}, \mathbb{P}_\Lambda)$ such that, for any Borel subset $B \subset \mathbb{R}$, the function

$$\Lambda \ni \lambda \mapsto \mu_\lambda[B]$$

is measurable. In other words, $\mu_\bullet$ is a *random probability measure*. In the language of kernels, the function $\mu_\bullet : \Lambda \rightarrow \mathcal{B}_\mathbb{R}$ is a Markov kernel $(\Lambda, \mathcal{S}) \rightsquigarrow (\mathbb{R}, \mathcal{B}_\mathbb{R})$.

For each $\lambda \in \Lambda$ we have a product measure $\mu_\lambda^{\otimes n}$ on $\mathbb{R}^n$ equipped with its natural σ -algebra, $\mathcal{B}_n = \mathcal{B}_\mathbb{R}^{\otimes n}$. The *mixture* of the family (μ_λ^n) directed by $\mathbb{P}^\Lambda$ is the measure μ_Λ^n defined by the averaging formula

$$\mu_\Lambda^n[S] := \int_\Lambda \mu_\lambda^n[B] \mathbb{P}_\Lambda[d\lambda], \quad \forall B \in \mathcal{B}_n.$$

The collection $(\mu_\Lambda^n)_{n \in \mathbb{N}}$ forms a projective family. Kolmogorov's existence theorem shows that this family induces a unique probability measure μ_Λ^∞ on $\mathbb{R}^\mathbb{N}$. The random variables

$$X_n : \mathbb{R}^\infty \rightarrow \mathbb{R}, \quad X_n(x_1, x_2, \dots) = x_n$$

form an exchangeable sequence. The measure μ_Λ^∞ is called a *mixture of i.i.d. directed by the random measure μ* .

For example, suppose that ν is a Borel probability measure on $\Lambda = [0, 1]$. For any $p > 0$ define

$$\mu_p = \text{Bin}(p) = (1-p)\delta_0 + p\delta_1 \in \text{Prob}(\{0, 1\}).$$

Then we obtain the mixtures $\mu_\nu^n \in \text{Prob}(\{0, 1\}^n)$ defined by,

$$\mu_\nu^n[\{\epsilon_1, \dots, \epsilon_n\}] = \binom{n}{k} \int_{[0,1]} (1-p)^{n-k} p^k \nu[dp], \quad k = \epsilon_1 + \dots + \epsilon_n.$$

The collection $\mu_\nu^n \in \text{Prob}(\{0, 1\}^n)$, $n \in \mathbb{N}$ is a projective family and thus it defines a measure μ_ν^∞ on $\{0, 1\}^\mathbb{N}$.

The random vector $X = (X_1, X_2, \dots)$ with distribution μ^∞ defines an exchangeable sequence of Bernoulli random variables. Observe that their common success probability is

$$\bar{p} := \mathbb{P}[X_n = 1] = \int_{[0,1]} p \nu[dp], \quad \forall n \in \mathbb{N}.$$

□

Denote by $\mathcal{B}$ the Borel σ -algebra of $\mathbb{R}$. The groups $\mathfrak{S}_n$ act on $\mathbb{R}^\mathbb{N}$ by permuting the first n coordinates and we say that a function $\Phi : \mathbb{R}^\mathbb{N} \rightarrow \mathbb{R}$ is n -symmetric if it is $\mathfrak{S}_n$ -invariant. We denote by $\mathcal{S}_n \subset \mathcal{B}^\mathbb{N}$ the sigma-subalgebra generated by the n -symmetric measurable functions $\Phi : \mathbb{R}^\mathbb{N} \rightarrow \mathbb{R}$. Equivalently,

$$S \subset \mathcal{S}_n \iff \sigma(S) = S, \quad \forall \sigma \in \mathfrak{S}_n.$$

We set

$$\mathcal{S}_\infty := \bigcap_{n \geq 1} \mathcal{S}_n \subset \mathcal{B}^\mathbb{N}.$$

We will refer to $\mathcal{S}_\infty$ as the σ -algebra of *permutable or exchangeable events* associated to the exchangeable sequence $(X_n)_{n \in \mathbb{N}}$.

Lemma 3.2.59. We have $\mathcal{B}_{\mathbb{R}}^{\mathbb{N}} \supset \mathcal{S}_{\infty} \supset \mathcal{T}_{\infty}$, where $\mathcal{T}_{\infty}$ denotes the tail σ -algebra of the coordinate sequence $X_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}$, $X_n(x_1, x_2, \dots) = x_n$, $n \in \mathbb{N}$,

$$\mathcal{T}_{\infty} = \bigcap_{m \geq 1} \underbrace{\sigma\{X_n; n > m\}}_{\mathcal{T}_m}.$$

Proof. Note that for any $S \in \mathcal{T}_m$ and any $\varphi \in \mathfrak{S}_m$, $\varphi(S) = S$. Indeed, a set in $\mathcal{T}_m$ is determined by conditions on the variables $X_{m+1}, \dots$ while a permutation in the first m variables does not affect these conditions. Hence $\mathcal{T}_m \subset \mathcal{S}_m$, $\forall m$. $\square$

It turns out that exchangeable sequences have a rather nice structure.

Theorem 3.2.60 (de Finetti). Suppose that $\underline{X} := (X_n)_{n \in \mathbb{N}}$ is an exchangeable sequence of integrable random variables defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$\underline{\mathcal{S}}_n := \underline{X}^{-1}\mathcal{S}_n, \quad \forall n \in \mathbb{N} \cup \{\infty\}.$$

Then the following hold.

- (i) The random variables $(X_n)_{n \geq 1}$ are conditionally independent given $\underline{\mathcal{S}}_{\infty}$.
- (ii) The random variables $(X_n)_{n \geq 1}$ are identically distributed given $\underline{\mathcal{S}}_{\infty}$, i.e., there exists a negligible subset $\mathcal{N} \in \mathcal{F}$ such that, on $\Omega \setminus \mathcal{N}$

$$\mathbb{P}[X_i \leq x \mid \underline{\mathcal{S}}_{\infty}] = \mathbb{P}[X_j \leq x \mid \underline{\mathcal{S}}_{\infty}], \quad \forall i, j \in \mathbb{N}, \quad \forall x \in \mathbb{R}.$$

- (iii) The empirical means

$$\frac{X_1 + \dots + X_n}{n}$$

converge a.s. and L^1 to $\mathbb{E}[X_1 \mid \underline{\mathcal{S}}_{\infty}]$.

Proof. We follow the presentation in [117]. Without any loss of generality we can assume that $(\Omega, \mathcal{S}) = (\mathbb{R}^{\mathbb{N}}, \mathcal{B}^{\mathbb{N}})$ and $X_n(x_1, x_2, \dots) = x_n$. In this case $\underline{\mathcal{S}}_n = \mathcal{S}_n$. Observe that the exchangeability condition implies that the random variables X_n are identically distributed. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is a measurable function such that $f(X_1) \in L^1$. We claim that

$$\forall k \in \mathbb{N}, \quad \frac{1}{n} \left(f(X_1) + \dots + f(X_n) \right) = \mathbb{E}[f(X_k) \mid \mathcal{S}_n] \quad (3.2.41)$$

Note that $\mathcal{S}_n = \underline{X}^{-1}(\mathcal{B}_n)$, where $\mathcal{B}_n$ is the σ -subalgebra of $\mathcal{B}^{\mathbb{N}}$ consisting of $\mathfrak{S}_n$ -invariant subsets. In particular, a function $g : \Omega \rightarrow \mathbb{R}$ is $\mathcal{S}_n$ -measurable iff there exists an n -symmetric function Φ such that $g = \Phi(\underline{X})$.

Let $A \in \mathcal{S}_n$ and choose an n -symmetric function Φ such that $\mathbf{I}_A = \Phi(\underline{X})$. Then, for $1 \leq j \leq n$ we have

$$\begin{aligned} \mathbb{E}[f(X_j)\Phi(\underline{X})] &= \mathbb{E}[f(X_j)\Phi(X_j, X_2, \dots, X_{j-1}, X_1, X_{j+1}, \dots)] \\ &= \mathbb{E}[f(X_1)\Phi(\underline{X})], \end{aligned}$$

so that

$$\mathbb{E}[f(X_1)\mathbf{I}_A] = \mathbb{E}\left[\frac{f(X_1) + \dots + f(X_n)}{n}\mathbf{I}_A\right] = \mathbb{E}[f(X_j)\Phi(\underline{X})].$$

The equality (3.2.41) follows by observing that $f(X_1) + \dots + f(X_n)$ is $\mathcal{S}_n$ -measurable.

The convergence theorem for backwards martingales (Corollary 3.2.57) shows that the empirical mean

$$\frac{f(X_1) + \cdots + f(X_n)}{n}$$

converges a.s. and L^1 to $\mathbb{E}[f(X_1) \mid \mathcal{S}_\infty]$. By choosing $f(x) = x$ we obtain the statement (iii) of Theorem 3.2.60.

By choosing $f(x) = \mathbf{I}_{(-\infty, x]}$ we deduce

$$\lim_{n \rightarrow \infty} \frac{\#\{j \leq n; X_j \leq x\}}{n} = F(x) := \mathbb{P}[X_1 \leq x \mid \mathcal{S}_\infty], \quad (3.2.42)$$

a.s. and L^1 .

Let $k \in \mathbb{N}$. For $n \geq k$ we set $(n)_k := n(n-1) \cdots (n-k+1)$. Suppose that $f_1, \dots, f_k : \mathbb{R} \rightarrow \mathbb{R}$ are bounded and measurable. The above argument generalizes to prove that for $n \geq k$ we have

$$A_{k,n} := \frac{1}{(n)_k} \sum_{\substack{j_1, \dots, j_k \\ j_i \text{ distinct}}} f_1(X_{j_1}) \cdots f_k(X_{j_k}) = \mathbb{E}[f_1(X_1) \cdots f_k(X_k) \mid \mathcal{S}_n].$$

Using the backwards martingale convergence theorem we deduce

$$\lim_{n \rightarrow \infty} A_{k,n} = \mathbb{E}[f_1(X_1) \cdots f_k(X_k) \mid \mathcal{S}_\infty]. \quad (3.2.43)$$

Consider now

$$B_{k,n} := \frac{1}{n^k} \sum_{j_1, \dots, j_k=1}^n f_1(X_{j_1}) \cdots f_k(X_{j_k}) = \prod_{i=1}^k \frac{f_i(X_1) + \cdots + f_i(X_n)}{n},$$

where the functions $f_i : \mathbb{R} \rightarrow \mathbb{R}$ are bounded and Borel measurable. We deduce from (3.2.41) that

$$\lim_{n \rightarrow \infty} B_{k,n} = \prod_{i=1}^k \mathbb{E}[f_i(X_i) \mid \mathcal{S}_\infty].$$

Now observe that

$$A_{k,n} - B_{k,n} = O(1/n) \text{ as } n \rightarrow \infty,$$

since the contribution to $B_{k,n}$ corresponding to k -tuples with j_i non-distinct is $O(n^{k-1})$ and $n^k \sim (n)_k$ as $n \rightarrow \infty$. If we choose

$$f_i = \mathbf{I}_{(-\infty, x_i]}, \quad 1 \leq i \leq k,$$

we deduce from (3.2.42) that

$$\mathbb{P}[X_1 \leq x_1, \dots, X_k \leq x_k \mid \mathcal{S}_\infty] = \prod_{i=1}^k \mathbb{P}[X_i \leq x_i \mid \mathcal{S}_\infty].$$

This proves (i) and (ii) of the theorem. $\square$

Remark 3.2.61. Suppose that $(X_n)_{n \in \mathbb{N}}$ is an exchangeable sequence of random variables defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Denote by $\mathcal{S}_\infty$ the sigma-algebra of exchangeable events. Suppose that

$$Q : \Omega \times \mathcal{B}_{\mathbb{R}} \rightarrow [0, 1], \quad (\omega, B) \mapsto Q_\omega[B].$$

is a regular version of the conditional distribution $\mathbb{P}_{X_1}[dx \mid \mathcal{S}_\infty]$, i.e.,

$$\forall B \in \mathcal{B}_{\mathbb{R}}, \quad \mathbb{P}[X_1 \in B \mid \mathcal{S}_\infty] = Q_\square[B], \text{ a.s..}$$

De Finetti's theorem implies that

$$\mathbb{P}[X_1 \in B \mid \mathcal{S}_\infty] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \mathbf{I}_B(X_k) = \mathbb{P}[X_m \in B \mid \mathcal{S}_\infty], \quad \forall m \in \mathbb{N}.$$

Thus the random variables (X_n) are equidistributed, conditional on $\mathcal{S}_\infty$.

Let us show that the distribution of the sequence $(X_n)_{n \in \mathbb{N}}$ is a mixture directed by the random measure $\omega \mapsto Q_\omega[-]$ as in Example 3.2.58(b).

Indeed, for any Borel subsets $B_1, \dots, B_n \subset \mathbb{R}$ we have

$$\mathbb{P}[X_1 \in B_1, \dots, X_n \in B_n] = \mathbb{E}\left[\mathbb{E}[\mathbf{I}_{B_1}(X_1) \cdots \mathbf{I}_{B_n}(X_n) \mid \mathcal{S}_\infty]\right]$$

(use the conditional independence given $\mathcal{S}_\infty$)

$$\begin{aligned} &= \mathbb{E}\left[\mathbb{E}[\mathbf{I}_{B_1}(X_1) \mid \mathcal{S}_\infty] \cdots \mathbb{E}[\mathbf{I}_{B_n}(X_n) \mid \mathcal{S}_\infty]\right] \\ &= \mathbb{E}\left[Q[B_1] \cdots Q[B_n]\right] = \int_{\Omega} Q_\omega^{\otimes n}[B_1 \times \cdots \times B_n] \mathbb{P}[d\omega]. \end{aligned}$$

Thus the distribution of the sequence (X_n) is a mixture of i.i.d. driven by the random distribution Q . $\square$

According to Lemma 3.2.59, the σ -algebra $\mathcal{S}_\infty \subset \mathcal{B}^{\mathbb{N}}$ of permutable events of an exchangeable sequence $(X_n)_{n \in \mathbb{N}}$ contains its tail σ -algebra $\mathcal{T}_\infty$. It turns out that they are not so different. We have the following general result.

Theorem 3.2.62 (Hewitt-Savage). *Suppose that $(X_n)_{n \in \mathbb{N}}$ is an exchangeable sequence of random variables. Then the $\mathbb{P}$ -completion of $\mathcal{S}_\infty$ coincides with the completion of the tail $\mathcal{T}_\infty$.*

Proof. We follow the approach in [37, Sec.7.3, Thm. 4]. Denote by $\mathcal{S}_\infty^*$ and $\mathcal{T}_\infty^*$ the completions of $\mathcal{S}_\infty$ and respectively $\mathcal{T}_\infty$. We have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \mathbb{1}_{\{X_k \leq x\}} = \mathbb{P}[X_1 \leq x \mid \mathcal{S}_\infty].$$

Clearly the limit in the left-hand side is $\mathcal{T}_\infty$ -measurable since it is not affected by changing finitely many of the random variables. We know from Lemma 3.2.59 that $\mathcal{S}_\infty \supset \mathcal{T}_\infty$ and we deduce

$$\mathbb{P}[X_1 \leq x \mid \mathcal{S}_\infty] = \mathbb{P}[X_1 \leq x \mid \mathcal{T}_\infty] = \mathbb{P}[X_1 \leq x \mid \mathcal{T}_\infty^*]. \quad (3.2.44)$$

Similarly, for any $x_1, \dots, x_n \in \mathbb{R}$, the random variable

$$\prod_{k=1}^n \mathbb{P}[X_k \leq x_k \mid \mathcal{S}_\infty]$$

is $\mathcal{T}_\infty$ -measurable. Hence, for any $S \in \mathcal{S}_\infty$ we have

$$\begin{aligned} \mathbb{P}\left[S \cap \bigcap_{k=1}^n \{X_k \leq x_k\} \mid \mathcal{T}_\infty\right] &= \mathbb{E}\left[\mathbf{I}_S \cdot \prod_{k=1}^n \mathbb{P}[X_k \leq x_k \mid \mathcal{S}_\infty] \mid \mathcal{T}_\infty\right] \\ &= \mathbb{E}\left[\prod_{k=1}^n \mathbb{P}[X_k \leq x_k \mid \mathcal{S}_\infty] \mid \mathcal{T}_\infty\right] \mathbb{P}[S \mid \mathcal{T}_\infty] \end{aligned}$$

$$\stackrel{(3.2.44)}{=} \prod_{k=1}^n \mathbb{P}[X_k \leq x_k \mid \mathcal{T}_\infty] \mathbb{P}[S \mid \mathcal{T}_\infty].$$

Thus $\mathcal{S}_\infty$ and $X_1, \dots, X_n$ are conditionally independent given $\mathcal{T}_\infty$ so $\mathcal{S}_\infty$ and $(X_n)_{n \in \mathbb{N}}$ are conditionally independent given $\mathcal{T}_\infty$. Since $\mathcal{S}_\infty \subset \sigma(X_n, n \in \mathbb{N})$ we deduce that for any $S \in \mathcal{S}_\infty$ is conditionally independent of itself given $\mathcal{T}_\infty$, i.e.,

$$\mathbb{P}[S \mid \mathcal{T}_\infty]^2 = \mathbb{P}[S \mid \mathcal{T}_\infty].$$

Hence $\mathbb{P}[S \mid \mathcal{T}_\infty] \in \{0, 1\}$, $\forall S \in \mathcal{S}_\infty$. Set $F_S := \mathbb{P}[S \mid \mathcal{T}_\infty]$. This 0 – 1 valued random variable is $\mathcal{T}_\infty$ -measurable so there exists $T = T(S) \in \mathcal{T}_\infty$ such that $F_S = \mathbf{I}_T$. We have

$$\mathbb{P}[S \cap T] = \mathbb{E}[\mathbf{I}_S \mathbf{I}_T] = \mathbb{E}[F_S \mathbf{I}_T] = \mathbb{E}[\mathbf{I}_T] = \mathbb{P}[T].$$

On the other hand,

$$\mathbb{P}[S] = \mathbb{E}[F_S] = \mathbb{E}[\mathbf{I}_T] = \mathbb{P}[T].$$

Hence $T \setminus S = T \setminus (T \cap S)$ is negligible. This concludes the proof. $\square$

Remark 3.2.63. For different proofs of Theorem 3.2.62 we refer to [1, Cor.(3.10)] or [151, Thm. VIII.T-3]. In [163], the completion of $\mathcal{S}_\infty$ is shown to coincide with the sigma-algebra of shift-invariant events; see Definition 5.1.3 and Remark 5.1.4(b) and Example 5.1.9 $\square$

Observe that a sequence of i.i.d. random variables $(X_n)_{n \geq 1}$ is exchangeable. The Kolmogorov zero-one law and the above proposition imply the following result.

Theorem 3.2.64 (Hewitt-Savage 0-1 Law). *If $(X_n)_{n \geq 1}$ is a sequence of iid random variables and $A \in \mathcal{S}_\infty$, then $\mathbb{P}[A] \in \{0, 1\}$.* $\square$

For a brief and elementary proof of the above result we refer to [76, Sec. IV.6].

Corollary 3.2.65 (The Strong Law of Large Numbers). *Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. integrable random variables. Then*

$$\bar{X}_n := \frac{1}{n} (X_1 + \dots + X_n)$$

converges a.s. and L^1 to $\mathbb{E}[X_1]$.

Proof. From de Finetti's Theorem 3.2.60 we deduce that $\bar{X}_n$ converges a.s. and L^1 to $\mathbb{E}[X_1 \mid \mathcal{S}_\infty]$. Theorem 3.2.62 implies that $\mathbb{E}[X_1 \mid \mathcal{S}_\infty] = \mathbb{E}[X_1 \mid \mathcal{T}_\infty]$ and Kolmogorov's zero-one theorem shows that $\mathbb{E}[X_1 \mid \mathcal{T}_\infty] = \mathbb{E}[X_1]$. $\square$

Theorem 3.2.66 (de Finetti). *Suppose that $(X_n : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow \{0, 1\})_{n \in \mathbb{N}}$ is an exchangeable sequence of Bernoulli random variables. Set*

$$S := \lim_{n \rightarrow \infty} \frac{1}{n} (X_1 + \dots + X_n).$$

Then

$$S = \mathbb{P}[X_1 = 1 \mid \mathcal{S}_\infty], \tag{3.2.45a}$$

$$\mathbb{P}[X_1 = \dots = X_k = 1, X_{k+1} = \dots = X_n = 0 \mid S] = S^k (1 - S)^{n-k}, \tag{3.2.45b}$$

$$\mathbb{P}[X_1 = \dots = X_k = 1, X_{k+1} = \dots = X_n = 0] = \mathbb{E}[S^k (1 - S)^{n-k}]. \tag{3.2.45c}$$

In particular, the moment generating function of S is

$$\mathbb{E}[e^{tS}] = \sum_{n \geq 0} \mathbb{P}[X_1 = \dots = X_n = 1] \frac{t^n}{n!}.$$

Proof. Using de Finetti's theorem 3.2.60 we deduce that $S = \mathbb{E}[X_1 \mid \mathcal{S}_\infty]$. Observe that $X_1 = \mathbf{I}_{\{X_1=1\}}$ so that $S = \mathbb{E}[X_1 \mid \mathcal{S}_\infty] = \mathbb{E}[\mathbf{I}_{\{X_1=1\}} \mid \mathcal{S}_\infty] = \mathbb{P}[X_1 = 1 \mid \mathcal{S}_\infty]$.

Note that $0 \leq S \leq 1$ a.s and

$$1 - S = \mathbb{E}[1 - \mathbf{I}_{\{X_1=1\}} \mid \mathcal{S}_\infty] = \mathbb{P}[X_1 = 0 \mid \mathcal{S}_\infty].$$

Then, since $X_1, \dots, X_n$ are conditionally i.i.d. given $\mathcal{S}_\infty$, we have

$$\begin{aligned} & \mathbb{P}[X_1 = 1, \dots, X_k = 1, X_{k+1} = 0, \dots, X_n = 0 \mid \mathcal{S}_\infty] \\ &= \mathbb{P}[X_1 = 1 \mid \mathcal{S}_\infty]^k \mathbb{P}[X_1 = 0 \mid \mathcal{S}_\infty]^{n-k} = S^k (1 - S)^{n-k}. \end{aligned}$$

Since S is $\mathcal{S}_\infty$ -measurable we have

$$\begin{aligned} & \mathbb{P}[X_1 = 1, \dots, X_k = 1, X_{k+1} = 0, \dots, X_n = 0 \mid S] \\ &= \mathbb{E}[\mathbb{P}[X_1 = 1, \dots, X_k = 1, X_{k+1} = 0, \dots, X_n = 0 \mid \mathcal{S}_\infty] \mid S] \\ &= \mathbb{E}[S^k (1 - S)^{n-k} \mid S] = S^k (1 - S)^{n-k}. \end{aligned}$$

Clearly,

$$\mathbb{P}[X_1 = 1, \dots, X_k = 1, X_{k+1} = 0, \dots, X_n = 0] = \mathbb{E}[S^k (1 - S)^{n-k}].$$

□

Remark 3.2.67. If we denote by $\mathbb{P}_S$ the distribution of S we deduce

$$\mathbb{P}[X_1 = 1, \dots, X_k = 1, X_{k+1} = 0, \dots, X_n = 0] = \int_0^1 s^k (1 - s)^{n-k} \mathbb{P}_S[ds].$$

For a more elementary proof of this equality we refer to [76, Sec. VII.4].

□

Example 3.2.68 (Polya's urn revisited). We want to conclude this introduction to exchangeability with an application to Polya's urn problem introduced in Example 3.1.9. We recall this process.

We start with an urn containing $r > 0$ red balls and $g > 0$ green balls. At each moment of time we draw a ball uniformly likely from the balls existing at that moment, we replace it by $c + 1$ balls of the same color, $c \geq 0$. Denote by R_n and G_n the number of red and respectively green balls in the urn after the n th draw. As we have seen in Example 3.1.9 the ratio of red balls

$$Z_n = \frac{R_n}{R_n + G_n} = \frac{R_n}{r + g + cn}$$

is a bounded martingale and thus it has an a.s. and L^1 limit Z_∞ . We will determine this limit using de Finetti's theorem. We discuss only the nontrivial case $c > 0$.

Introduce the $\{0, 1\}$ -valued random variables $(X_n)_{n \geq 1}$ where $X_n = 1$ if the n -drawn ball is red and $X_n = 0$ if it is green. Then

$$R_n = r + cS_n, \quad S_n := X_1 + \dots + X_n,$$

and we deduce that

$$\lim_{n \rightarrow \infty} \frac{cS_n}{cn} = \lim_{n \rightarrow \infty} \frac{R_n}{R_n + G_n} = Z_\infty.$$

Let us observe that the sequence $(X_n)_{n \geq 1}$ is exchangeable. We prove by induction that $(X_1, \dots, X_n)$ is exchangeable. For $n = 1$ the result is trivial.

Let $n > 1$ and $\epsilon_1, \dots, \epsilon_n \in \{0, 1\}$. We denote by r_k and g_k the number of red balls and respectively green balls after the k -th draw. We deduce

$$\mathbb{P}[X_1 = \epsilon_1, \dots, X_n = \epsilon_n] = \begin{cases} \frac{\prod_{k=1}^n (\epsilon_k r_{k-1} + (1 - \epsilon_k) g_{k-1})}{\prod_{k=1}^n (r + g + (k-1)c)}, & c > 0, \\ z_0^k (1 - z_0)^{n-k}, & c = 0, \end{cases},$$

where $z_0 = Z_0 = \frac{r}{r+g}$. When $c > 0$ the denominator above is independent of $\{\epsilon_1, \dots, \epsilon_n\}$. We set $S_n := \epsilon_1 + \dots + \epsilon_n$ and we rewrite the numerator in the form

$$\prod_{k=1}^n (\epsilon_k r_{k-1} + (1 - \epsilon_k) g_{k-1}) = \prod_{i=1}^{S_n} (r + c(i-1)) \prod_{j=1}^{n-S_n} (g + c(j-1)).$$

The last expression only depends on S_n which is obviously a symmetric function in the variables $\epsilon_1, \dots, \epsilon_n$. If $c = 0$, then this expression is equal to 0.

When $c > 0$, we set

$$\rho := \frac{r}{c}, \quad \gamma := \frac{g}{c},$$

and we deduce

$$\begin{aligned} \mathbb{P}[X_1 = \dots = X_k = 1, X_{k+1} = \dots = X_n = 0] &= \frac{\prod_{i=0}^{k-1} (\rho + i) \prod_{j=0}^{n-k-1} (\gamma + j)}{\prod_{k=0}^{n-1} (r + \gamma + k)} \\ &= \frac{\Gamma(\rho + \gamma)}{\Gamma(\rho)\Gamma(\gamma)} \cdot \frac{\Gamma(\rho + k)\Gamma(\gamma + n - k)}{\Gamma(\rho + \gamma + n)} = \frac{B(\rho + k, \gamma + n - k)}{B(\rho, \gamma)}, \end{aligned} \quad (3.2.46)$$

where $B(x, y)$ denotes the Beta function

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

We now invoke Theorem 3.2.66. Note that

$$Z_\infty = \lim_{n \rightarrow \infty} \frac{1}{n} (X_1 + \dots + X_n) = S = \mathbb{P}[X_1 = 1 \mid \mathcal{S}_\infty]$$

is a $[0, 1]$ -valued random variable and (3.2.45c) with $k = n$ shows that, for any $n \geq 0$, we have

$$\begin{aligned} \int_0^1 z^n \mathbb{P}_{Z_\infty}[dz] &= \mathbb{E}[Z_\infty^n] = \mathbb{P}[X_1 = \dots = X_n = 1] \\ &\stackrel{(3.2.46)}{=} \begin{cases} \frac{B(\rho+n, \gamma)}{B(\rho, \gamma)}, & c > 0, \\ z_0^n, & c = 0 \end{cases} = \begin{cases} \int_0^1 z^n \frac{z^{\rho-1} (1-z)^{\gamma-1}}{B(\rho, \gamma)} dz, & c > 0, \\ \int_0^1 s^n \delta_{z_0}[dz] & c = 0, \end{cases} \end{aligned}$$

where δ_{z_0} is the Dirac measure concentrated at z_0 . Hence

$$\int_0^1 z^n \mathbb{P}_{Z_\infty}[dz] = \begin{cases} \int_0^1 z^n \frac{z^{\rho-1} (1-z)^{\gamma-1}}{B(\rho, \gamma)} dz, & c > 0, \\ \int_0^1 z^n \delta_{z_0}[dz], & c = 0, \quad \forall n \geq 0. \end{cases}$$

Since the probability measures on $[0, 1]$ are uniquely determined by their momenta (see Corollary 1.3.24) we deduce

$$\mathbb{P}_{Z_\infty}[dz] = \begin{cases} \frac{z^{\rho-1}(1-z)^{\gamma-1}}{B(\rho, \gamma)} dz, & c > 0, \\ \delta_{z_0}[dz], & c = 0. \end{cases}$$

The distribution in the case $c > 0$ is the *Beta distribution* with parameters ρ, γ discussed in Example 1.3.39. $\square$

3.3. Generalities about filtered processes

The study of martingales parametrized by $\mathbb{T} = [0, \infty)$ faces several fundamental technical difficulties stemming from the fact that the space of parameters is not countable. To deal with these issues we need to introduce several new concepts. I believe that for a first encounter with continuous-time martingales it may be more productive to skip the proofs of many (and nontrivial) measure theoretic results and discuss only a more limited set-up yet general enough to cover many interesting applications. My presentation is heavily influenced by [132, 180] to which I refer for more details and additional references.

3.3.1. Continuous parameter filtrations. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and

$$\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$$

is a filtration of sigma-subalgebras of $\mathcal{S}$. We set

$$\mathcal{F}_\infty := \bigvee_{t \geq 0} \mathcal{F}_t,$$

and denote by $\text{Proc}(\mathcal{F}_\bullet)$ the collection of stochastic processes (parametrized by $\mathbb{T}$) that are adapted to the filtration $\mathcal{F}_\bullet$. If no confusion is possible, we will use the simpler notation Proc when referring to adapted processes.

A function $f : [0, \infty) \rightarrow \mathbb{R}$ is called an *R-function*⁴ if it is *right*-continuous with left limits. It is called an *L-function*⁵ if it is *left*-continuous with right limits.

Definition 3.3.1. Let $X_\bullet = \{X_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}\}_{t \in [0, \infty)}$ be a random process, not necessarily adapted to the filtration $\mathcal{F}_\bullet$.

- (i) We say that the random process $X_\bullet$ is *measurable* if the map

$$X : [0, \infty) \times \Omega \rightarrow \mathbb{R}, \quad (t, \omega) \mapsto X_t(\omega)$$

is measurable with respect to the σ -algebra $\mathcal{B}_{[0, \infty)} \otimes \mathcal{S}$.

- (ii) We say that the random process $X_\bullet$ is *progressively measurable* or *progressive* (with respect to the filtration $\mathcal{F}_\bullet$) if for any $t > 0$, the map

$$[0, t] \times \Omega \ni (s, \omega) \mapsto X_t(\omega) \in \mathbb{R}$$

is $\mathcal{B}_{[0, t]} \otimes \mathcal{F}_t$ measurable, where $\mathcal{B}_{[0, t]}$ denotes the σ -algebra of Borel subsets of $[0, t]$.

⁴A.k.a. *cadlag* function, *continue à droite limite à gauche*.

⁵A.k.a. *caglad* function, *continue à gauche limite à droite*.

(iii) A subset $A \subset [0, \infty) \times \Omega$ is called *progressive* if the associated process

$$I_A : [0, \infty) \times \Omega \rightarrow \mathbb{R}$$

is progressive.

(iv) We say that the adapted random process $X_\bullet$ is an *R-process* (resp. *L-process*) if there exists a negligible subset $\mathcal{N} \subset \Omega$ such that, for any $\omega \in \Omega \setminus \mathcal{N}$, the function $\mathbb{T} \ni t \mapsto X_t(\omega)$ is and *R-function* (resp. *L-function*).

□

Remark 3.3.2. The progressive subsets of $[0, \infty) \times \Omega$ form a σ -subalgebra of $\mathcal{B}(\mathbb{R}) \otimes \mathcal{S}$ that we denote by $\mathcal{F}_{\text{prog}}$. Observe that a process is progressively measurable if and only if it is $\mathcal{F}_{\text{prog}}$ -measurable. For this reason we will denote by $\text{Proc}(\mathcal{F}_{\text{prog}})$ or $\text{Proc}_{\text{prog}}$ the collection of progressive processes.

An $\mathcal{F}_\bullet$ -progressive process is also adapted to the filtration $\mathcal{F}_\bullet$ so

$$\text{Proc}(\mathcal{F}_{\text{prog}}) \subset \text{Proc}(\mathcal{F}_\bullet).$$

□

Proposition 3.3.3. Suppose that $X_\bullet \in \text{Proc}(\mathcal{F}_\bullet)$ is either an *R-process* or an *L-process*. Then $X_\bullet$ is a *progressive process*.

Proof. Assume X is an *R-process*. The case of *L-processes* is similar. Fix $t \geq 0$. For each $n \in \mathbb{N}$, we subdivide the interval $[0, t]$ into n intervals of the same size. For $n \in \mathbb{N}$, define $X^n : [0, t] \times \Omega \rightarrow \mathbb{R}$

$$X_s^n(\omega) = \begin{cases} X_{kt/n}(\omega), & s \in [(k-1)t/n, kt/n), \quad 1 \leq k \leq n, \\ X_t(\omega), & s = t. \end{cases}$$

Since $X_\bullet$ is an *R-process* we deduce that there exists a negligible subset $\mathcal{N} \subset \Omega$ such that

$$\lim_{n \rightarrow \infty} X_s^n(\omega) = X_s(\omega), \quad \forall s \in [0, t], \quad \omega \in \Omega \setminus \mathcal{N}.$$

Clearly the function $X^n : [0, t] \times \Omega \rightarrow \mathbb{R}$ is $\mathcal{B}_{[0,t]} \otimes \mathcal{F}_t$ -measurable. It follows that the a.s. limit $X : [0, t] \times \Omega \rightarrow \mathbb{R}$ is also $\mathcal{B}_{[0,t]} \otimes \mathcal{F}_t$ -measurable, $\forall t \geq 0$. □

We have the following nontrivial result, [38], [151, IV T46].

Theorem 3.3.4 (Chung-Doob). Suppose that

$$X_\bullet = \{ X_t : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow \mathbb{R} \}_{t \in [0, \infty)}$$

is a measurable process adapted to the filtration $\mathcal{F}_\bullet$. Then $X_\bullet$ admits a *progressive modification*. □

Definition 3.3.5. Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ of $\mathcal{F}$. We set

$$\mathcal{F}_{t+} := \bigcap_{s > t} \mathcal{F}_s, \quad t \geq 0.$$

(i) We say that the filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ is *right-continuous* if

$$\mathcal{F}_t = \mathcal{F}_{t+}, \quad \forall t \geq 0.$$

(ii) We say that the filtration $\mathcal{F}_t$ is $\mathbb{P}$ -complete if the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is $\mathbb{P}$ -complete and the collection $\mathcal{N} \subset \mathcal{F}$ of $\mathbb{P}$ -negligible events is contained in $\mathcal{F}_t, \forall t \geq 0$

- (iii) We say that the filtration $\mathcal{F}_t$ satisfies the *usual conditions* (or that it is *usual*) if it is both right-continuous and $\mathbb{P}$ -complete. Following [180], I will refer to filtrations satisfying the usual conditions as *R-filtrations*.

□

Remark 3.3.6. If $(\mathcal{F}_t)_{t \geq 0}$ is a filtration of the *complete* probability space $(\Omega, \mathcal{F}, \mathbb{P})$, then the *usual augmentation* of $(\mathcal{F}_t)$ is the minimal filtration $(\hat{\mathcal{F}}_t)$ containing $(\mathcal{F}_t)$ and satisfying the usual conditions. More precisely if $\mathcal{N} \subset \mathcal{F}$ is the collection of probability zero events, then

$$\hat{\mathcal{F}}_t = \bigcap_{s > t} \sigma(\mathcal{N}, \mathcal{F}_s). \quad \square$$

3.3.2. Stopping and optional times.

Definition 3.3.7. Fix a filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ of the probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

- (i) An $\mathcal{F}_\bullet$ -*stopping time* is a random variable $T : \Omega \rightarrow [0, \infty]$ such that

$$\{T \leq t\} \in \mathcal{F}_t, \quad \forall t \geq 0.$$

- (ii) An $\mathcal{F}_\bullet$ -*optional time* is a random variable $T : \Omega \rightarrow [0, \infty]$ such that

$$\{T < t\} \in \mathcal{F}_t, \quad \forall t > 0.$$

- (iii) If $T : \Omega \rightarrow [0, \infty]$ is a stopping time, then the *past before* T is collection $\mathcal{F}_T \subset \mathcal{F}_\infty$ consisting of the sets $F \in \mathcal{F}$ satisfying the property $F \cap \{T \leq t\} \in \mathcal{F}_t, \forall t \geq 0$.

□

Lemma 3.3.8. For any stopping time T adapted to the filtration $\mathcal{F}_\bullet$ the collection $\mathcal{F}_T$ is a σ -algebra. □

The proof is left to the reader as an exercise.

Lemma 3.3.9. Any stopping time T is an optional time.

Proof. Indeed,

$$\{T < t\} = \bigcup_{n \geq 0} \{T \leq t - 1/n\},$$

and $\{T \leq t - 1/n\} \in \mathcal{F}_{t-1/n} \subset \mathcal{F}_t$. □

Proposition 3.3.10. Consider a random variable $T : \Omega \rightarrow [0, \infty]$. Then the following statements are equivalent.

- (i) T is an optional time for $(\mathcal{F}_t)$.
- (ii) T is a stopping time for $(\mathcal{F}_{t+})$.
- (iii) $T \wedge t = \min(T, t)$ is $\mathcal{F}_t$ -measurable for any $t > 0$.

In particular, if $\mathcal{F}_t$ is right-continuous, then T is a stopping time if and only if it is an optional time.⁶ □

⁶This settles an inconsistency in the existence literature. Many authors refer to stopping times as optional times, while our optional times are sometimes referred to as weakly optional times. When the filtration is right continuous all these terms refer to the same concept, that of stopping time.

Example 3.3.11. Suppose that $(X_t)_{t \geq 0}$ is a process adapted to $\mathcal{F}_\bullet$ and $\Gamma \subset \mathbb{R}$. The (Γ) -*début time* or *hitting time* of (X_t) is the function

$$D_\Gamma : \Omega \rightarrow [0, \infty], \quad D_\Gamma(\omega) = \inf \{ t \geq 0; X_t(\omega) \in \Gamma \},$$

and the (Γ) -*return time* of (X_t) is the function

$$T_\Gamma : \Omega \rightarrow [0, \infty], \quad T_\Gamma(\omega) = \inf \{ t > 0; X_t(\omega) \in \Gamma \}.$$

The following facts are not hard to prove; see [109, Lemma 9.6], [132, Prop. 3.9].

- (i) If Γ is *open*, and the paths of X_t are right continuous, then the *début time* D_Γ is a *stopping time* of (X_t) , while the *return time* T_Γ is an *optional time*.
- (ii) If Γ is *closed*, and the paths of X_t are continuous, then the *début time* D_Γ is a *stopping time* of (X_t) , while the *return time* T_Γ is an *optional time*.

We deduce from the above that if the filtration $\mathcal{F}_t$ is *right-continuous* and the paths of (X_t) are continuous, then both D_Γ and T_Γ are *stopping times* if Γ is either open or closed. $\square$

If the filtration $\mathcal{F}_\bullet$ satisfies the usual conditions, then a much more general result is true. More precisely, we have the following highly nontrivial result of Dellacherie and Meyer [50, Thm. IV.50].

Theorem 3.3.12 (Début Theorem). *Suppose that the filtration $\mathcal{F}_\bullet$ satisfies the usual conditions and $(X_t)_{t \geq 0}$ is an $\mathcal{F}_\bullet$ -progressive process. Then, for any Borel subset $\Gamma \subset \mathbb{R}$, the *début time* D_Γ is a *stopping time*.* $\square$

We list below a few elementary properties of stopping times.

Proposition 3.3.13. *Fix a filtered probability space $(\Omega, \mathcal{F}_\bullet, \mathbb{P})$.*

- (i) *If T is a stopping time, then T is also $\mathcal{F}_T$ -measurable.*
- (ii) *If S is a stopping time and T is an $\mathcal{F}_S$ -measurable random variable such that $T \geq S$, then T is also a stopping time and $\mathcal{F}_S \subset \mathcal{F}_T$.*
- (iii) *Suppose that S, T are stopping times. Then $S \wedge T$ and $S \vee T$ are also stopping times and*

$$\mathcal{F}_{S \vee T} = \mathcal{F}_S \cap \mathcal{F}_T.$$

- (iv) *An increasing limit of stopping times is a stopping time while a decreasing limit of stopping times is an optional time.*
- (v) *Suppose that T is a stopping time. A function*

$$\{T < \infty\} \ni \omega \mapsto Y(\omega) \in \mathbb{R}$$

is $\mathcal{F}_T$ -measurable if and only if, $\forall t \geq 0$, the restriction of Y to $\{T \leq t\}$ is $\mathcal{F}_t$ -measurable.

Proof. We prove only (i). The rest are left to the reader as an exercise. To prove that the sublevel set $\{T \leq c\}$ is measurable we have to show that for any $t \geq 0$ the intersection

$$\{T \leq c\} \cap \{T \leq t\} = \{T \leq t \wedge c\}$$

is $\mathcal{F}_t$ -measurable. This is a consequence of the fact that T is compatible with the filtration $\mathcal{F}_t$. $\square$

Definition 3.3.14. Fix a filtered probability space $(\Omega, \mathcal{F}_\bullet, \mathbb{P})$. Given a random process $(X_t)_{t \geq 0}$ and an $\mathcal{F}_\bullet$ -stopping time $T : \Omega \rightarrow [0, \infty]$ we denote by X_T the random variable

$$X_T(\omega) = \begin{cases} X_{T(\omega)}(\omega), & T(\omega) < \infty \\ 0, & T(\omega) = \infty \end{cases}.$$

We will sometimes refer to X_T as the T -sample of the process. The process *stopped at T* is the random process $(X_t^T)_{t \geq 0}$ defined by

$$X_t^T(\omega) = X_{t \wedge T}(\omega) = X_{t \wedge T(\omega)}(\omega), \quad t \wedge T = \min(t, T)$$

□

The proof of the following result is left to the reader as an exercise.

Proposition 3.3.15. Suppose that $(X_t)_{t \geq 0}$ is a progressive random process and T is a stopping time. Then the following hold.

- (i) The random variable X_T is $\mathcal{F}_T$ -measurable.
- (ii) The stopped process $(X_t^T)_{t \geq 0}$ is progressive.

□

3.4. The Brownian motion as a filtered process

Let us illustrate the concepts introduced in the previous section on the stochastic process defined by the Brownian motion.

3.4.1. Some fundamental properties of the Brownian motion. Recall that the Brownian motion is a Gaussian process

$$B_t : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad t \geq 0,$$

with continuous paths such that, $B_0 = 0$ and, for any $0 \leq t_0 < t_1$, the increment $B_{t_1} - B_{t_0}$ is Gaussian random variable with mean zero, variance $t_1 - t_0$ and independent of $\{B_s, 0 \leq s \leq t_0\}$.

Equivalently, the Brownian motion is the unique Gaussian process $(B_t)_{t \geq 0}$ with continuous paths, mean zero and covariance kernel

$$K(s, t) = \mathbb{E}[B_s B_t] = \min(s, t).$$

Let us observe that if $(B_t)_{t \geq 0}$ is a Brownian motion, then for any $t_0 \geq 0$, the process

$$(B_{t+t_0} - B_{t_0})_{t \geq 0}$$

is also a Brownian motion, independent of $\sigma(B_s, 0 \leq s \leq t_0)$. We will refer to this elementary fact as the *simple Markov property*.

Proposition 3.4.1. Suppose that B is a Brownian motion. Then the following hold.

Symmetry. The stochastic process $-B$ is also a Brownian motion.

Time rescaling. For any $c > 0$ the rescaled Brownian motion

$$B_t^c := \frac{1}{c} B_{c^2 t}$$

is another Brownian motion.

Time inversion. *The stochastic process*

$$X_t := \begin{cases} tB_{1/t}, & t > 0, \\ 0, & t = 0, \end{cases}$$

is another standard Brownian motion.

Proof. The statements (i) and (ii) are immediate. The last statement concerning time inversion requires a bit more work. We follow the approach in the proof of [43, Thm. VIII.1.6].

Observe first that X_t is a Gaussian process with mean zero and covariances

$$\mathbb{E}[X_s X_t] = \min(s, t), \quad \forall s, t \geq 0.$$

Indeed,

$$\mathbb{E}[X_t X_s] = st \frac{1}{\max(s, t)} = \min(s, t).$$

Thus it suffices to show that (X_t) is a.s. continuous, i.e.,

$$\lim_{t \searrow 0} X_t = 0 \quad \text{a.s.}$$

Equivalently, we will show that

$$\lim_{t \rightarrow \infty} \frac{1}{t} B_t = 0 \quad \text{a.s.}$$

Note that for $n \in \mathbb{N}$ and $t \in (n, n+1]$ we have

$$\frac{1}{t} |B_t| \leq \frac{1}{n} |B_n + (B_t - B_n)| \leq \frac{1}{n} |B_n| + \frac{1}{n} \sup_{s \in [0,1]} |B_{n+s} - B_n|$$

(the process (B_t) is a.s. continuous on $(0, \infty)$)

$$= \frac{1}{n} |B_n| + \frac{1}{n} \sup_{s \in [0,1] \cap \mathbb{Q}} |B_{n+s} - B_n|.$$

The Strong Law of Large Numbers shows that

$$\lim_{n \rightarrow \infty} \frac{1}{n} B_n = 0 \quad \text{a.s.}$$

For each $m \in \mathbb{N}$, the process $(D_m^k)_{0 \leq k \leq m}$

$$D_m^0 = 0, \quad D_m^k = B_{n+k/m} - B_n = \sum_{j=1}^k (B_{n+\frac{j}{m}} - B_{n+\frac{j-1}{m}}), \quad 1 \leq k \leq m,$$

is a martingale since the above summands have mean zero and are independent. Applying Doob's maximal inequalities (3.2.36) to the discrete submartingales

$$Y^k + m = \{|D_m^k|^2; \quad 0 \leq k \leq m\}, \quad m \in \mathbb{N},$$

we deduce that, for any $\varepsilon > 0$,

$$\begin{aligned} \mathbb{P} \left[\sup_{s \in [0,1]} |B_{n+s} - B_n| > n\varepsilon \right] &= \mathbb{P} \left[\sup_{s \in [0,1] \cap \mathbb{Q}} |B_{n+s} - B_n|^2 > n^2 \varepsilon^2 \right] \\ &\leq \frac{1}{n^2 \varepsilon^2} \mathbb{E} \left[|B_{n+1} - B_n|^2 \right] = \frac{1}{n^2 \varepsilon^2}. \end{aligned}$$

Since $\sum_{n \geq 1} \frac{1}{n^2} < \infty$ we deduce from the Borel-Cantelli Lemma that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sup_{s \in [0,1]} |B_{n+s} - B_n| = 0, \text{ a.s.}$$

□

Theorem 3.4.2. *Suppose that*

$$B : [0, \infty) \times (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$$

is a Brownian motion and $(\Omega, \mathcal{S}, \mathbb{P})$ is a complete probability space. Let $\mathcal{N}$ denote the collection of $\mathbb{P}$ -negligible events. We set

$$\mathcal{F}_t := \sigma(B_s, 0 \leq s \leq t), \quad \bar{\mathcal{F}}_t = \sigma(\mathcal{N}) \vee \mathcal{F}_t.$$

Then the following hold

- (i) *For any $0 \leq s < t$, the increment $B_t - B_s$ is independent of $\bar{\mathcal{F}}_s$.*
- (ii) *The filtration $(\bar{\mathcal{F}}_t)_{t \geq 0}$ is right-continuous and thus it is the usual augmentation of the canonical filtration $\bar{\mathcal{F}}_t$.*

Proof. We follow the approach in the proof of [43, Thm. VII.3.20]. Part (i) is immediate since any $F \in \bar{\mathcal{F}}_s$ can be written as a disjoint union $F_0 \cup N$, $F_0 \in \mathcal{F}_s$, $N \in \mathcal{N}$.

(ii) We set

$$\mathcal{G}_n = \sigma(B_t - B_s, 2^{-n-1} \leq s, t \leq 2^{-n}), \quad n \in \mathbb{N}.$$

Clearly the σ -algebras $\mathcal{G}_1, \dots$, are independent. Set

$$\mathcal{T}_n := \sigma(\mathcal{G}_{n+1}, \mathcal{G}_{n+2}, \dots), \quad \mathcal{T}_\infty := \bigcap_{n \in \mathbb{N}} \mathcal{T}_n.$$

From Kolmogorov's zero-one law we deduce that $\mathcal{T}_\infty$ is a zero-one sigma-algebra so

$$\mathcal{F}_{0+} = \mathcal{T}_\infty \subset \sigma(\mathcal{N}).$$

Hence

$$\bar{\mathcal{F}}_{0+} = \sigma(\mathcal{N}) \vee \mathcal{F}_{0+} \subset \sigma(\mathcal{N}) = \bar{\mathcal{F}}_0.$$

Fix $t_0 \geq 0$ and set $\hat{B}_t = B_{t_0+t} - B_{t_0}$. Then $\hat{B}_t$ is a Brownian motion. Set

$$\hat{\mathcal{F}}_t = \sigma(\hat{B}_s, 0 \leq s \leq t) \vee \sigma(\mathcal{N}).$$

The argument above shows that

$$\bigcap_{t > 0} \hat{\mathcal{F}}_t = \hat{\mathcal{F}}_0 = \sigma(\mathcal{N}).$$

On the other hand, $\bar{\mathcal{F}}_{t_0+t} = \bar{\mathcal{F}}_{t_0} \vee \hat{\mathcal{F}}_t$, so

$$\bar{\mathcal{F}}_{t_0+} = \bar{\mathcal{F}}_{t_0} \vee \mathcal{N} = \bar{\mathcal{F}}_{t_0}.$$

□

Corollary 3.4.3 (Blumenthal's 0-1 law). *If $H \in \mathcal{F}_{0+}$ then, $\mathbb{P}[H] \in \{0, 1\}$.*

□

Fix a Brownian motion $(B(t))_{t \geq 0}$ defined on a complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Denote by $\mathcal{F}_\bullet^B = (\mathcal{F}_t)_{t \geq 0}$ the completed Brownian filtration

$$\mathcal{F}_t^B = \sigma(B(s), 0 \leq s \leq t) \vee \sigma(\mathcal{N}), \quad (3.4.1)$$

where $\mathcal{N} \subset \mathcal{S}$ denotes the collection of $\mathbb{P}$ -negligible events. As shown in Theorem 3.4.2, this filtration is also right-continuous.

Proposition 3.4.4. *Suppose that $(B_t)_{t \geq 0}$ is a Brownian motion. Then the following hold.*

(i) *For any $\varepsilon > 0$ we have*

$$\mathbb{P}\left[\sup_{s \in [0, \varepsilon]} B_s > 0\right] = \mathbb{P}\left[\inf_{s \in [0, \varepsilon]} B_s < 0\right] = 1.$$

(ii) *For any $a \in \mathbb{R}$ we set*

$$T_a := \inf_{t \geq 0} B_t = a.$$

Then

$$\mathbb{P}[T_a < \infty] = 1, \quad \forall a \in \mathbb{R}.$$

In particular, a.s.,

$$\limsup_{t \rightarrow \infty} B_t = \infty, \quad \liminf_{t \rightarrow \infty} B_t = -\infty.$$

Proof. (i) For any $c \neq 0$, the rescaled process

$$B^c(t) := \frac{1}{c} B_{c^2 t}, \quad t \geq 0$$

is also a Brownian motion. Note that since the paths of B_t are continuous we have

$$\sup_{t \in [0, 1]} B_t = \sup_{t \in \mathbb{Q} \cap [0, 1]} B_t.$$

Thus the set

$$\left\{ \omega; \sup_{t \in [0, 1]} B_t(\omega) > 0 \right\}$$

is a Brownian event. The discussion in Remark 2.5.7 shows that

$$\mathbb{P}\left[\sup_{t \in [0, 1]} B_t > 0\right] = \mathbb{P}\left[\sup_{t \in [0, 1]} B_t^c > 0\right], \quad \forall c \neq 0. \quad (3.4.2)$$

If we let $c = -1$ in the above equality we deduce,

$$\mathbb{P}\left[\sup_{t \in [0, 1]} B_t > 0\right] = \mathbb{P}\left[\inf_{t \in [0, 1]} B_t < 0\right]. \quad (3.4.3)$$

If we let $c_n = n^{-1/2}$, $n \in \mathbb{N}$, we deduce that, $\forall n \in \mathbb{N}$,

$$\mathbb{P}\left[\sup_{t \in [0, 1]} B_t > 0\right] = \mathbb{P}\left[\sup_{t \in [0, 1]} B_t^{c_n} > 0\right] = \mathbb{P}\left[\sup_{t \in [0, 1/n]} B_t > 0\right]. \quad (3.4.4)$$

We denote by E_n the Brownian event $\sup_{t \in [0, 1/n]} B_t > 0$. Clearly

$$E_1 \supset E_2 \supset \cdots \supset E_n \supset \cdots$$

and $E_n \in \mathcal{F}_{1/n}$. We deduce from (3.4.4) that $\mathbb{P}[E_n] = \mathbb{P}[E_1]$, $\forall n$. If we set

$$E_\infty := \bigcap_n E_n,$$

then we deduce that $E_\infty \in \mathcal{F}_{0+}$ and $\mathbb{P}[E_\infty] = \mathbb{P}[E_1]$. Blumenthal's 0-1 theorem implies that

$$\mathbb{P}[E_n] = \mathbb{P}[E_\infty] \in \{0, 1\}.$$

Now observe that

$$\mathbb{P}[E_1] \geq \mathbb{P}[B_{1/2} > 0] = \frac{1}{2} > 0.$$

Hence

$$\mathbb{P}\left[\sup_{t \in [0, 1/n]} B_t > 0\right] = \mathbb{P}\left[\inf_{t \in [0, 1/n]} B_t < 0\right] = 1, \quad \forall n \in \mathbb{N}. \quad (3.4.5)$$

This shows that the paths of the Brownian motion oscillate wildly.

(ii) We have

$$1 = \mathbb{P}\left[\sup_{0 \leq s \leq 1} B_s > 0\right] = \lim_{\delta \searrow 0} \mathbb{P}\left[\sup_{0 \leq s \leq 1} B_s > \delta\right],$$

where the second is an increasing limit. The rescaling invariance of the Brownian motion implies

$$\mathbb{P}\left[\sup_{0 \leq s \leq 1} B_s > \delta\right] = \mathbb{P}\left[\sup_{0 \leq s \leq 1/\delta^2} B_s^\delta > 1\right] = \mathbb{P}\left[\sup_{0 \leq s \leq 1/\delta^2} B_s > 1\right].$$

We deduce

$$\mathbb{P}\left[\sup_{s \geq 0} B_s > 1\right] = \lim_{\delta \searrow 0} \mathbb{P}\left[\sup_{0 \leq s \leq 1/\delta^2} B_s > 1\right] = \lim_{\delta \searrow 1} \mathbb{P}\left[\sup_{0 \leq s \leq 1} B_s > \delta\right] = 1,$$

so

$$\mathbb{P}\left[\sup_{s \geq 0} B_s > c\right] = \mathbb{P}\left[\sup_{s > 0} B_s^c > 1\right] = 1, \quad \forall c > 0.$$

Replacing B by $-B$ we deduce

$$\mathbb{P}\left[\inf_{s \geq 0} B_s < -c\right] = 1, \quad \forall c > 0.$$

The conclusion (ii) is now obvious. $\square$

Remark 3.4.5. The above result shows that, with probability 1 the Brownian motion has a zero on any arbitrarily small interval $[0, \varepsilon]$. As a matter of fact, the set of zeros of a Brownian motion is a large set: its Hausdorff dimension is a.s. $\frac{1}{2}$, [154, Thm. 4.24]. $\square$

In many applications, we will need to view a Brownian motion as a filtered process where the filtration $\mathcal{F}_\bullet$ is bigger than the canonical filtration $\mathcal{F}_t^B = \sigma(B_s, 0 \leq s \leq t)$ but still maintains many of the desirable features of the canonical filtration.

Definition 3.4.6. Consider a Brownian motion B defined on the complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$. A filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ of $(\Omega, \mathcal{S}, \mathbb{P})$ is called *B-admissible* or *B-non-anticipating* if

- $\mathcal{F}_\bullet$ satisfies the usual conditions,
- $B(t)$ is $\mathcal{F}_t$ -measurable, $\forall t \geq 0$, and
- $\forall t > s \geq 0$, the increment $B(t) - B(s)$ is independent of $\mathcal{F}_s$.

In this case we also say that B is a $\mathcal{F}_\bullet$ -Brownian motion. $\square$

Example 3.4.7. Suppose that B_t is a Brownian motion defined on the complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and X is a random variable independent of the Brownian motion. If $\mathcal{F}_\bullet^B$ is the completed Brownian filtration defined in (3.4.1), then the filtration $\mathcal{F}_t = \mathcal{F}_t^B \vee \sigma(X)$ is *B-admissible*. $\square$

Observe that if $\mathcal{F}_\bullet$ is a filtration admissible with respect to a Brownian motion B , then for any $s < t$ we have

$$\mathbb{E}[B_t \mid \mathcal{F}_s] = \mathbb{E}[B_t \mid B_s] = B_s.$$

The next result is a very powerful generalization of the above simple Markov property.

Theorem 3.4.8 (The strong Markov property). *Suppose that $(B_t)_{t \geq 0}$ is a Brownian motion, $\mathcal{F}_\bullet$ is a B -admissible filtration, and T is a $\mathcal{F}_\bullet$ -stopping time such that $\mathbb{P}[T < \infty] > 0$. For every $t \geq 0$ we set*

$$B_t^{(T)} := \mathbf{I}_{\{T < \infty\}}(B_{T+t} - B_T).$$

Then, with respect to the probability measure $\mathbb{P}[- \mid T < \infty]$, the process $B_t^{(T)}$ is a standard Brownian motion, independent of $\mathcal{F}_T$.

Proof. I will follow the approach in [132, Thm. 2.20]. We have the following result whose proof we postpone.

Lemma 3.4.9. *Fix $A \in \mathcal{F}_T$. Let $F : \mathbb{R}^p \rightarrow \mathbb{R}$ be a bounded continuous function. Then, $\forall t_1, \dots, t_p \geq 0$, we have*

$$\mathbb{E}[\mathbf{I}_A \mathbf{I}_{T < \infty} F(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)})] = \mathbb{P}[A \cap \{T < \infty\}] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})] \quad (3.4.6)$$

□

Let us first observe that the conclusions of theorem follow from the above lemma. Set

$$S_\infty := \{T < \infty\}.$$

Assume first that $\mathbb{P}[S_\infty] = 1$. Then (3.4.6) reads

$$\mathbb{E}[\mathbf{I}_A F(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)})] = \mathbb{P}[A] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})]. \quad (3.4.7)$$

If we set $A = \Omega$ in (3.4.7) we deduce from Proposition 1.2.63 that for every choice of $t_1, \dots, t_p \geq 0$, the vectors

$$(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)}) \text{ and } (B_{t_1}, \dots, B_{t_p})$$

have the same distribution. Thus the process $t \rightarrow B_t^{(T)}$ is also a Brownian motion.

Next, (3.4.7) implies that for every choice of $t_1, \dots, t_p \geq 0$ the vector $(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)})$ is independent of $\mathcal{F}_T$.

If $\mathbb{P}[S_\infty] < 1$, we denote by $\mathbb{E}_{S_\infty}$ the expectation with respect to the conditional probability measure $\mathbb{P}[- \mid S_\infty]$, then (3.4.6) implies

$$\mathbb{E}_{S_\infty}[\mathbf{I}_A F(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)})] = \mathbb{P}[A \mid S_\infty] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})].$$

Arguing as before we reach the conclusions of Theorem 3.4.8 assuming the validity of Lemma 3.4.9. □

Proof of Lemma 3.4.9. For the clarity of exposition we discuss only the case $\mathbb{P}[S_\infty] = 1$. The case $\mathbb{P}[S_\infty] < 1$ requires no new ideas. The details can be safely left to the reader.

For every $t \geq 0$ and any $n \in \mathbb{N}$ we denote by $\lceil t \rceil_n$ the smallest rational number of the form $k/2^n$ and $\geq t$. More explicitly, $\lceil t \rceil_n = 2^{-n} \lceil 2^n t \rceil$.

Note that the quantities $\lceil T \rceil_n$ are stopping times since

$$\{ \lceil T \rceil_n \leq t \} = \bigcup_{k/2^n \leq t} \{ T \leq k/2^n \} \in \mathcal{F}_t$$

Stopping the process at $\lceil T \rceil_n$ corresponds to stopping the process at the first time of the form $k/2^n$ after T . Then

$$\lim_{n \rightarrow \infty} \lceil T \rceil_n = T$$

and

$$F(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)}) = \lim_{n \rightarrow \infty} F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)}).$$

From the Dominated Convergence theorem we deduce that

$$\begin{aligned} \mathbb{E}[I_A F(B_{t_1}^{(T)}, \dots, B_{t_p}^{(T)})] &= \lim_{n \rightarrow \infty} \mathbb{E}[I_A F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)})] \\ &= \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} \mathbb{E}[I_A I_{(k-1)2^{-n} < T \leq k2^{-n}} F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)})]. \end{aligned}$$

Observe now that if $A \in \mathcal{F}_T$, then the event

$$\begin{aligned} A_{k,n} &:= A \cap \{ (k-1)2^{-n} < T \leq k2^{-n} \} \\ &= A \cap \{ T \leq k2^{-n} \} \cap \{ T > (k-1)2^{-n} \} \end{aligned}$$

is $\mathcal{F}_{k2^{-n}}$ -measurable.

From the simple Markov property of the Brownian motion we deduce

$$\begin{aligned} &\mathbb{E}[I_{A_{k,n}} F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)})] \\ &= \mathbb{E}[I_{A_{k,n}} F(B_{t_1+k2^{-n}} - B_{k2^{-n}}, \dots, B_{t_p+k2^{-n}} - B_{k2^{-n}})] \\ &= \mathbb{P}[A_{k,n}] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})]. \end{aligned}$$

Observing that

$$\sum_{k=0}^n \mathbb{P}[A_{k,n}] = \mathbb{P}[A]$$

we deduce

$$\begin{aligned} &\sum_{k=0}^{\infty} \mathbb{E}[I_A I_{(k-1)2^{-n} < T \leq k2^{-n}} F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)})] \\ &= \sum_{k=0}^{\infty} \mathbb{E}[I_{A_{k,n}} F(B_{t_1}^{(\lceil T \rceil_n)}, \dots, B_{t_p}^{(\lceil T \rceil_n)})] \\ &= \sum_{k=0}^n \mathbb{P}[A_{k,n}] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})] = \mathbb{P}[A] \mathbb{E}[F(B_{t_1}, \dots, B_{t_p})]. \end{aligned}$$

□

3.4.2. Some applications. The strong Markov property has many consequences. I want to present a few of them and some of their applications.

Theorem 3.4.10 (Reflection Principle). *If $(B_t)_{t \geq 0}$ is a standard Brownian motion,*

$$\mathcal{F}_t = \sigma(B_s, s \leq t),$$

and T is an a.s. finite $\mathcal{F}_\bullet$ -stopping time, then the process

$$\tilde{B}_t = \begin{cases} B_t, & t < T, \\ 2B_T - B_t, & t \geq T \end{cases} \quad (3.4.8)$$

is also a standard Brownian motion.

Proof. I follow the approach in [180, I.13]. Consider the processes

$$Y_t = B_t \mathbf{I}_{[0, T]}, \quad Z_s = B_{s+T} - B_T, \quad s \geq 0.$$

By the strong Markov property, Z is a standard Brownian motion, independent of Y . The process $-Z$ is also a Brownian motion independent of Y . Thus, the processes (Y, Z) and $(Y, -Z)$ have the same distribution. The map

$$(Y, Z) \mapsto \varphi(Y, Z) := Y_t \mathbf{I}_{[0, T]} + (B_T + Z_{t-T}) \mathbf{I}_{[T, \infty[}$$

produces the a continuous process which will therefore have the same law as $\varphi(Y, -Z)$. Now observe that $\varphi(Y, Z) = B$ and $\varphi(Y, -Z) = \tilde{B}$. $\square$

Remark 3.4.11. The above result is called the reflection principle for a simple reason. In the region $t \geq T$ the graph of the function $t \rightarrow \tilde{B}_t$, viewed as a curve in the Cartesian plane with coordinates (t, x) , is the reflection of the graph of B_t in the horizontal line $y = B_T$; see Figure 3.5. This reflection principle is intimately related to André's reflection trick we used in Example 1.2.38 to solve the ballot problem. $\square$

Corollary 3.4.12. *Define*

$$S_t := \sup_{u \leq t} B_u.$$

Then, for any $a, y, t \geq 0$ we have

$$\mathbb{P}[S_t \geq a, B_t \leq a - y] = \mathbb{P}[B_t \geq a + y]. \quad (3.4.9)$$

In particular, S_t has the same distribution as $|B_t|$.

Proof. Consider the return time

$$T_a := \inf \{t > 0; B_t = a\}.$$

Proposition 3.4.4(ii) shows that $\mathbb{P}[T_a < \infty] = 1$. Note that $S_t \geq a$ if and only if $T_a \leq t$. We have

$$\mathbb{P}[S_t \geq a, B_t \leq a - y] = \mathbb{P}[T_a \leq t, B_t \leq a - y] \stackrel{(3.4.8)}{=} \mathbb{P}[\tilde{B}_t \geq a + y]$$

(use the Reflection Principle)

$$= \mathbb{P}[B_t \geq a + y].$$




Figure 3.5. A brownian path and its (dashed) reflection at $y = B_{60}$.

Now observe that

$$\begin{aligned} \mathbb{P}[S_t \geq a] &= \underbrace{\mathbb{P}[S_t \geq a, B_t \geq a]}_{=\mathbb{P}[B_t \geq a]} + \mathbb{P}[S_t \geq a, B_t \leq a] \\ &\stackrel{(3.4.9)}{=} 2\mathbb{P}[B_t \geq a] = \mathbb{P}[B_t \geq a] + \mathbb{P}[B_t \leq -a] = \mathbb{P}[|B_t| \geq a]. \end{aligned}$$

□

Corollary 3.4.13. For every $a > 0$ the stopping time T_a has the same distribution as $\frac{a^2}{B_1^2}$ and has density

$$f_a(t) = \frac{a}{\sqrt{2\pi t^3}} \exp\left(-\frac{a^2}{2t}\right) \mathbf{I}_{\{t>0\}}.$$

Proof. Note that

$$\begin{aligned} \mathbb{P}[T_a \leq t] &= \mathbb{P}[S_t \geq a] = \mathbb{P}[|B_t| \geq a] = \mathbb{P}[B_t^2 \geq a^2] \\ &= \mathbb{P}[tB_1^2 \geq a^2] = \mathbb{P}\left[\frac{a^2}{B_1^2} \leq t\right]. \end{aligned}$$

The statement about f_a now follows from the fact that B_1 is a standard normal random variable. □

3.4.3. The support theorem. We want to describe another nice application of the reflection principle. It is called the support theorem since it proves that the Wiener measure of any open set $U \subset C([0, 1])$ is positive. Equivalently, this means that for any continuous function $f : [0, 1] \rightarrow \mathbb{R}$ such that $f(0) = 0$, and any $\varepsilon > 0$

$$\mathbb{P} \left[\sup_{0 \leq t \leq 1} |B(t) - f(t)| < \varepsilon \right] > 0.$$

Thus, during the first second of its existence, the Brownian motion, with positive probability, closely traces any given continuous function. $f : [0, 1] \rightarrow \mathbb{R}$ such that $f(0) = 0$. $\square$

The following proof is essentially P. Lévy's original approach, based on clever usage of the reflection principle.

Proposition 3.4.14. *Suppose that $(B(t))_{t \geq 0}$ is a Brownian motion. Then, for any $a_0 < 0 < a_1$ and any nontrivial interval $I \subset (a_0, a_1)$*

$$\begin{aligned} p(a_0, a_1, I) &:= \mathbb{P} \left[a_0 \leq \min_{s \in [0, 1]} B(s) \leq \max_{s \in [0, 1]} B(s) < a_1, \ B(1) \in I \right] \\ &= \sum_{k \in \mathbb{Z}} \mathbb{P} [B(1) \in 2k\ell + I] - \sum_{k \in \mathbb{Z}} \mathbb{P} [B(1) \in 2a_1 + 2k\ell - I] \\ &= \frac{1}{\sqrt{2\pi}} \int_I \sum_{k \in \mathbb{Z}} \left(e^{-\frac{(x-2k\ell)^2}{2}} - e^{-\frac{(x-2a_1+2k\ell)^2}{2}} \right) dx, \end{aligned} \quad (3.4.10)$$

where $\ell = a_1 - a_0$ and $a + bI = \{a + bx; x \in I\}$, $\forall a, b \in \mathbb{R}$. Moreover

$$p(a_0, a_1, I) > 0. \quad (3.4.11)$$

Proof. The approach I use, based on the reflection principle, is inspired from [80, Sec.I.3] and [185, Sec. 6.5]. For a different but related approach I refer to [135, Sec. 20°].

Set

$$\begin{aligned} T_a &:= \inf \{ t \geq 0; B_t = a \}, \\ T_i &:= T_{a_i}, \ i = 0, 1, \ T_* = \min(T_0, T_1), \ J = \mathbf{1}_{\{T_* = T_1\}}. \end{aligned}$$

Note that $J = 0$ if $T_* = T_0$ and $J = 1$ if $T_* = T_1$. For each $c \in \mathbb{R}$ we denote by r_c the reflection about c

$$r_c : \mathbb{R} \rightarrow \mathbb{R}, \ r_c(x) = 2c - x.$$

Lemma 3.4.15. *Let $S \subset \mathbb{R}$ be a Borel set.*

(i) *If $S \subset (-\infty, a_0]$, then*

$$\begin{aligned} \mathbb{P} [T_1 \leq T_0, B(1) \in S] &= \mathbb{P} [B(1) \in r_{a_1}(S)] \\ &\quad - \mathbb{P} [T_0 \leq T_1, B(1) \in r_{a_1}(S)]. \end{aligned} \quad (3.4.12)$$

(ii) *If $S \subset [a_1, \infty)$, then*

$$\begin{aligned} \mathbb{P} [T_0 \leq T_1, B(1) \in S] &= \mathbb{P} [B(1) \in r_{a_0}(S)] \\ &\quad - \mathbb{P} [T_1 \leq T_0, B(1) \in r_{a_0}(S)]. \end{aligned} \quad (3.4.13)$$

Proof. We prove only (i). Statement (ii) can be dealt with in a similar fashion. Alternatively, (ii) follows from (i) applied to the Brownian motion $-B(t)$ and the interval $[-a_1, -a_0]$.

Since $S \subset (-\infty, a_0]$ we have $\{B(1) \in S\} \subset \{T_0 \leq 1\}$ so

$$\{T_1 \leq T_0, B(1) \in S\} \subset \{T_1 \leq 1\}.$$

Note that $\mathbf{r}_{a_1}S \subset [a_1, \infty)$ so $\{B(1) \in \mathbf{r}_{a_1}S\} \subset \{T_1 \leq 1\}$. The reflection principle shows that

$$\begin{aligned} \mathbb{P}[T_1 \leq T_0, B(1) \in S] &= \mathbb{P}[T_1 \leq T_0, B(1) \in \mathbf{r}_{a_1}(S)] \\ &= \mathbb{P}[B(1) \in \mathbf{r}_{a_1}(S)] - \mathbb{P}[T_0 \leq T_1, B(1) \in \mathbf{r}_{a_1}(S)]. \end{aligned}$$

□

For ease of notation we set

$$M_* := \min_{s \in [0,1]} B(s), \quad M^* := \max_{s \in [0,1]} B(s),$$

Then

$$p(a_0, a_1, I) = \mathbb{P}[B(1) \in I] - \mathbb{P}[\{B(1) \in I\} \cap \{[M_*, M^*] \not\subset (a_0, a_1)\}].$$

Note that $[M_*, M^*] \not\subset [a_0, a_1]$ iff $B(t)$ hits a_0 or a_1 for some $t \leq 1$. Hence

$$\begin{aligned} &\mathbb{P}[\{B(1) \in I\} \cap \{[M_*, M^*] \not\subset [a_0, a_1]\}] \\ &= \underbrace{\mathbb{P}[B(1) \in I, T_* = T_0 \leq 1]}_{=: p_0(a_0, a_1, I)} + \underbrace{\mathbb{P}[B(1) \in I, T_* = T_1 \leq 1]}_{=: p_1(a_0, a_1, I)}. \end{aligned}$$

Hence

$$p(a_0, a_1, I) = \mathbb{P}[B(1) \in I] - p_0(a_0, a_1, I) - p_1(a_0, a_1, I). \quad (3.4.14)$$

Using the reflection principle we deduce

$$\begin{aligned} p_0(a_0, a_1, I) &= \mathbb{P}[T_0 \leq T_1, B(1) \in \mathbf{r}_{a_0}(I)] \\ &= \mathbb{P}[B(1) \in \mathbf{r}_{a_0}(I)] - \mathbb{P}[T_1 \leq T_0, B(1) \in \mathbf{r}_{a_0}(I)] \\ &\stackrel{(3.4.12)}{=} \mathbb{P}[B(1) \in \mathbf{r}_{a_0}(I)] - \mathbb{P}[B(1) \in \mathbf{r}_{a_1}\mathbf{r}_{a_0}(I)] \\ &\quad + \mathbb{P}[T_0 \leq T_1, B(1) \in \mathbf{r}_{a_1}\mathbf{r}_{a_0}(I)] \\ &\stackrel{(3.4.13)}{=} \mathbb{P}[B(1) \in \mathbf{r}_{a_0}(I)] - \mathbb{P}[B(1) \in \mathbf{r}_{a_1}\mathbf{r}_{a_0}(I)] + \mathbb{P}[B(1) \in \mathbf{r}_{a_0}\mathbf{r}_{a_1}\mathbf{r}_{a_0}(I)] \\ &\quad - \mathbb{P}[T_1 \leq T_0, B(1) \in \mathbf{r}_{a_0}\mathbf{r}_{a_1}\mathbf{r}_{a_0}(I)]. \end{aligned}$$

Define

$$R_{2k,0} := (\mathbf{r}_{a_1}\mathbf{r}_{a_0})^k, \quad R_{2k+1,0} := \mathbf{r}_{a_0}R_{2k,0}.$$

Iterating we deduce that for any $n \in \mathbb{B}$ we have

$$\begin{aligned} p_0(a_0, a_1, I) &= \sum_{k=1}^n (-1)^{k+1} \mathbb{P}[B(1) \in R_{k,0}(I)] \\ &\quad + (-1)^n \mathbb{P}[J = n \bmod 2, B(1) \in R_{n,0}(I)]. \end{aligned}$$

Note that

$$R_{2,0}(x) = \mathbf{r}_{a_1}(2a_0 - x) = 2a_1 - 2a_0 + x = 2\ell + x,$$

so that

$$R_{2k,0}(x) = 2k\ell + x, \quad R_{2k+1,0}(x) = 2a_0 - 2k\ell - x = 2a_1 - 2(k+1)\ell - x.$$

This shows that $\mathbb{P}[B(1) \in R_{n,0}(I)]$ goes to 0 exponentially fast as $n \rightarrow \infty$. We conclude that

$$p_0(a_0, a_1, I) = \sum_{k \geq 1} (-1)^{k+1} \mathbb{P}[B(1) \in R_{k,0}(I)]. \quad (3.4.15)$$

Define

$$R_{2k,1} := (r_{a_0} r_{a_1})^k, \quad R_{2k+1,1} := r_{a_1} R_{2k,1}.$$

Arguing as above we deduce

$$p_1(a_0, a_1, I) = \sum_{k \geq 1} (-1)^{k+1} \mathbb{P}[B(1) \in R_{k,1}(I)] \quad (3.4.16)$$

Note that

$$\begin{aligned} R_{2,1}(x) &= -2\ell + x, \quad R_{2k,1}(x) = -2k\ell x \\ R_{2k+1,1}(x) &= 2a_1 + 2k\ell - x. \end{aligned}$$

The equality (3.4.10) now follows from (3.4.14), (3.4.15), (3.4.16).

Let us prove the positivity claim (3.4.11). Set

$$\begin{aligned} \rho_{a_0, a_1}(x) &:= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left(e^{-\frac{(x-2k\ell)^2}{2}} - e^{-\frac{(x-2a_1+2k\ell)^2}{2}} \right) \\ &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left(e^{-\frac{(x-2k(a_1-a_0))^2}{2}} - e^{-\frac{(x-2a_0+(2k-1)(a_1-a_0))^2}{2}} \right). \end{aligned}$$

For fixed $a_0 < 0$ and $x \in (a_0, 0)$ we have a real analytic function of $a_1 > 0$

$$(0, \infty) \ni a_1 \mapsto f(a_1) = \rho_{a_0, a_1}(x).$$

Moreover

$$f(a_1) = \lim_{\varepsilon \searrow 0} \frac{1}{2\varepsilon} p(a_0, a_1, I_\varepsilon(x)), \quad I_\varepsilon(x) = [x - \varepsilon, x + \varepsilon].$$

Clearly the function $a_1 \mapsto p(a_0, a_1, I_\varepsilon(x))$ is nondecreasing for any $\varepsilon \geq 0$ due to the monotonicity of the probability measure. We deduce that the function $(0, \infty) \ni a_1 \mapsto f(a_1)$ is nondecreasing.

Note that for any $C > 0$ and $|\lambda| > C$, $y \in [-C, C]$ we have

$$(\lambda - y)^2 = |\lambda - y|^2 \geq d_C(\lambda)^2, \quad d_C(\lambda) := \text{dist}(\lambda, [-C, C])$$

Fix $C > 0$ such that $x, x - 2a_0 \in [-C, C]$. Then for $|k|$ sufficiently large

$$e^{-\frac{(x-2k(a_1-a_0))^2}{2}} \leq e^{-d_C(2k\ell)^2}, \quad e^{-\frac{(x-2a_0+(2k-1)(a_1-a_0))^2}{2}} \leq e^{-d_C((2k-1)\ell)^2}.$$

Finally, for $\ell = a_1 - a_0$ sufficiently large and $|k| > C$ we have

$$e^{-d_C(k\ell)^2} \leq e^{-d_C(k)^2}.$$

The series

$$\sum_{k \in \mathbb{Z}} e^{-d_C(k)^2}$$

converges since its terms decay very fast as $|k| \rightarrow \infty$. Using the Dominated Convergence Theorem we deduce that

$$\lim_{a_1 \nearrow \infty} f(a_1) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} > 0,$$

and

$$\lim_{a_1 \searrow 0} f(a_1) = 0.$$

Thus $(0, \infty) \ni a_1 \mapsto f(a_1)$ is a non-decreasing, nonnegative *real analytic* function, positive at ∞ .

Let us show that $f(a_1) > 0, \forall a_1 > 0$. We argue by contradiction. If f vanished at some point $a_1^* > 0$, then f would be identically zero on the open interval $(0, a_1^*)$. By unique continuation f would be identically zero on $(0, \infty)$, contradicting the fact that f is positive at infinity. Hence $\rho_{a_0, a_1}(x) > 0$, for $x \in (-a_0, 0)$ and $a_1 > 0$.

To deal with the case $x \in (0, a_1)$ observe that

$$\rho_{a_0, a_1}(x) = \rho_{-a_1, -a_0}(-x) > 0.$$

Thus $\rho_{a_1, a_1}(x) > 0$ for all $x \in (a_0, 0) \cup (0, a_1)$ and we conclude that

$$p(a_0, a_1, I) = \int_I \rho_{a_0, a_1}(x) dx > 0.$$

for any $a_0 < 0 < a_1$ and any nontrivial interval $I \subset [a_0, a_1]$. $\square$

An immediate consequence of the above results is Lévy's "*lois à trois variables*"⁷, [135, Thm. 42.6]. It describes the joint distribution of the random vector $(m(t), M(t), B(t))$,

$$m(t) = \min_{s \in [0, t]} B(s), \quad M(t) = \max_{s \in [0, t]} B(s).$$

Corollary 3.4.16. *Let $a_0 < 0 < a_1$, $I \subset [a_0, a_1]$ a nontrivial interval and $c > 0$. Set*

$$p(a_0, a_1, I | c) := \mathbb{P} \left[a_0 \leq \min_{s \in [0, c]} B(s) < \max_{s \in [0, c]} B(s) \leq a_1, \quad B(c) \in I \right].$$

Then

$$p(a_0, a_1, I | c) = p(c^{-1/2}a_0, c^{-1/2}a_1, c^{-1/2}I) > 0. \quad (3.4.17)$$

Proof. Let $B_c(t) = c^{-1/2}B(ct)$. Then $B_c(t)$ is a Brownian motion. Moreover

$$\begin{aligned} p(\varepsilon; c) &:= \sup_{t \in [0, c]} B(t) = c^{1/2} \sup_{t \in [0, 1]} B_c(t), \quad \inf_{t \in [0, c]} B(t) = c^{1/2} \inf_{t \in [0, 1]} B_c(t) \\ &\{ B(c) \in I \} = \{ B_c(1) \in c^{-1/2}I \}. \end{aligned}$$

$\square$

Corollary 3.4.17. *For any $\varepsilon, c > 0$ we have*

$$\begin{aligned} &\mathbb{P} \left[\sup_{t \in [0, c]} |B(t)| \leq \varepsilon \right] = p(-\varepsilon, \varepsilon, [-\varepsilon, \varepsilon] | c) \\ &= \mathbb{P} [|B(c)| \leq \varepsilon] + \sum_{n \geq 1} (-1)^n \mathbb{P} [(2n-1)\varepsilon \leq |B(c)| \leq 2(n+1)\varepsilon] > 0 \\ &= \frac{1}{\sqrt{2\pi c}} \int_{-\varepsilon}^{\varepsilon} e^{-\frac{x^2}{2c}} dx + \frac{1}{\sqrt{2\pi c}} \sum_{n \geq 1} \int_{(2n-1)\varepsilon}^{(2n+1)\varepsilon} e^{-\frac{x^2}{2c}} dx. \end{aligned}$$

$\square$

Corollary 3.4.18.

$$\mathbb{P} \left[\sup_{0 \leq t \leq 1} |B(t)| \leq a, \quad |B(1)| < \varepsilon \right] = \frac{1}{\sqrt{2\pi}} \int_{-\varepsilon}^{\varepsilon} \sum_{k \in \mathbb{Z}} (-1)^k e^{-2a^2(x-k)^2} dx$$

⁷the three-variable law

Proof. This is (3.4.10) where $a_0 = -1$, $a_1 = a$, $I = [-\varepsilon, \varepsilon]$. $\square$

Theorem 3.4.19 (Support theorem). *Let $f : [0, 1] \rightarrow \mathbb{R}$ be a continuous function such that $f(0) = 0$. Then, for any $\varepsilon > 0$,*

$$\mathbb{P} \left[\sup_{t \in [0, 1]} |B(t) - f(t)| \leq \varepsilon \right] > 0.$$

Proof. Fix $\varepsilon > 0$. The function f is uniformly continuous so there exists $n \in \mathbb{N}$ sufficiently large so that

$$\sup_{|s-t| \leq 1/n} |f(s) - f(t)| < \frac{\varepsilon}{3}.$$

Set $t_0 = 0$. For $k = 1, \dots, n$ set

$$t_k := \frac{k}{n}, \quad I_k = [t_{k-1}, t_k], \quad \varepsilon_n := \frac{\varepsilon}{3n}.$$

Let

$$J_{k,n} := [f(t_k) - f(t_{k-1}) - \varepsilon_n, f(t_k) - f(t_{k-1}) + \varepsilon_n],$$

denote by E_k the event

$$\left\{ (B(t_k) - B(t_{k-1})) \in J_{k,n}, \sup_{t \in I_k} |B(t) - B(t_{k-1})| \leq \frac{\varepsilon_n}{3} \right\},$$

and set $E := E_1 \cap \dots \cap E_n$.

Note that

$$\mathbb{P}[E_k] = p \left(-\varepsilon_n/3, \varepsilon_n/3, J_k | 1/n \right) \stackrel{(3.4.17)}{>} 0.$$

Since the Brownian motion has independent increments we deduce that

$$\mathbb{P}[E] = \prod_{k=1}^n \mathbb{P}[E_k] > 0.$$

If the event E occurs, then, $\forall t \in I_k$ we have

$$\begin{aligned} |B(t) - f(t)| &\leq |B(t) - B(t_{k-1})| \\ &+ |B(t_{k-1}) - f(t_{k-1})| + |f(t) - f(t_{k-1})| \\ &\leq \frac{2\varepsilon_n}{3} + |B(t_{k-1}) - f(t_{k-1})|. \end{aligned} \tag{3.4.18}$$

Set $\Delta_k := B(t_k) - f(t_k)$. Then,

$$|\Delta_k - \Delta_{k-1}| = |(B(t_k) - B(t_{k-1})) - (f(t_k) - f(t_{k-1}))| \leq \varepsilon_n$$

so that

$$|\Delta_k| = \left| \sum_{j=1}^k (\Delta_j - \Delta_{j-1}) \right| \leq \sum_{j=1}^k |\Delta_j - \Delta_{j-1}| \leq \sum_{j=1}^k \varepsilon_n = \frac{k\varepsilon}{3n} \leq \frac{\varepsilon}{3}.$$

Using this in (3.4.18) we deduce that, $\forall k = 1, \dots, n$,

$$\sup_{t \in I_k} |B(t) - f(t)| \leq \varepsilon.$$

Thus

$$\mathbb{P} \left[\sup_{t \in [0, 1]} |B(t) - f(t)| \leq \varepsilon \right] \geq \mathbb{P}[E] > 0.$$

$\square$

Remark 3.4.20. (a) Suppose that $B : (\Omega, \mathcal{S}, \mathbb{P}) \times [0, 1] \rightarrow \mathbb{R}$, $(\omega, t) \mapsto B_\omega(t)$ is a Brownian motion. Then the map

$$\Omega \ni \omega \mapsto B_\omega(-) \in C([0, 1])$$

is measurable with respect to the Borel sigma-algebra of the Banach space $C([0, 1])$. The push-forward of $\mathbb{P}$ is a Borel probability measure on $C([0, 1])$ called the Wiener measure; compare with Remark 2.5.16.

The support of Borel probability measure μ on $C([0, 1])$ is the smallest closed subset $S \subset C([0, 1])$ such that $\mu[S] = 1$. Theorem 3.4.19 shows that the support of the Wiener measure is the entire space $C([0, 1])$, whence the name support theorem. For a more conceptual proof of the support theorem I refer to [203, Cor. 4.1.2].

(b) Let me describe an amusing application of the support theorem. Suppose that $(B_t^i)_{t \geq 0}$, $i = 1, 2$, are two independent Brownian motions and $f^i : [0, 1] \rightarrow \mathbb{R}$, $i = 1, 2$ are two continuous functions such that $f^i(0) = 0$.

The support theorem implies immediately that for any $\varepsilon > 0$ we have

$$\begin{aligned} & \mathbb{P} \left[\max_i \sup_{t \in [0, 1]} |B_t^i - f^i(t)| \leq \varepsilon \right] \\ &= \mathbb{P} \left[\sup_{t \in [0, 1]} |B_t^1 - f^1(t)| \leq \varepsilon \right] \mathbb{P} \left[\sup_{t \in [0, 1]} |B_t^2 - f^2(t)| \leq \varepsilon \right] > 0. \end{aligned} \quad (3.4.19)$$

The pair of functions (f^1, f^2) defines a path

$$F : [0, 1] \rightarrow \mathbb{R}^2, \quad F(t) = (f^1(t), f^2(t)).$$

Think of $F(t)$ as tracing the motion of the tip of an infinitesimally fine pen as you sign a planar piece of paper, starting at the origin.

Any other path $G = (g^1, g^2) : [0, 1] \rightarrow \mathbb{R}^2$ satisfying

$$|g^i(t) - f^i(t)| < \varepsilon, \quad \forall t \in [0, 1], \quad i = 1, 2,$$

will follow closely the original motion of the fine pen, producing a curve essentially indistinguishable with the naked eye from the original signature. In fact, if $\varepsilon > 0$ is atomic scale small, one cannot distinguish the two curves, even using a magnifying glass.

The random path (B_t^1, B_t^2) is the so called *planar Brownian motion* started at the origin. The equality (3.4.19) shows that the probability p_0 that this random path follows very closely the motion of the tip of the fine pen is positive. For this reason the support theorem is sometimes referred to as *Lévy's forgery theorem*. $\square$

3.5. Continuous-time martingales

3.5.1. Definition and examples of continuous-time martingales. Fix a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$.

Definition 3.5.1. A random process $(X_t)_{t \geq 0}$ adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ such that $X_t \in L^1, \forall t$, is called a

- *martingale* if,

$$\mathbb{E}[X_t \mid \mathcal{F}_s] = X_s, \quad \forall 0 \leq s < t,$$

- *submartingale* if,

$$\mathbb{E}[X_t \mid \mathcal{F}_s] \geq X_s, \quad \forall 0 \leq s < t,$$

- *supermartingale* if,

$$\mathbb{E}[X_t \mid \mathcal{F}_s] \leq X_s, \quad \forall 0 \leq s < t.$$

□

Example 3.5.2 (Uniformly integrable martingales). To any integrable random variable X we can associate the martingale $X_t := \mathbb{E}[X \mid \mathcal{F}_t]$. □

Example 3.5.3 (Processes with independent increments). Suppose that the random process $(Z_t)_{t \geq 0}$ has *independent increments*, i.e., for any $n \in \mathbb{N}$ and any

$$0 \leq s_1 < t_1 \leq s_2 < t_2 \leq \dots \leq s_n < t_n,$$

the increments

$$Z_{t_1} - Z_{s_1}, Z_{t_2} - Z_{s_2}, \dots, Z_{t_n} - Z_{s_n}$$

are independent. The process (Z_t) is adapted to the natural filtration

$$(\mathcal{F}_t)_{t \geq 0}, \quad \mathcal{F}_t = \sigma(Z_s, s \leq t).$$

We deduce that, $\forall 0 \leq s < t$, the increment $Z_t - Z_s$ is independent of $\mathcal{F}_s$ so

$$\mathbb{E}[X_t \mid \mathcal{F}_s] - X_s = \mathbb{E}[(X_t - X_s) \mid \mathcal{F}_s] = \mathbb{E}[X_t - X_s].$$

Hence

$$\mathbb{E}[X_t - \mathbb{E}[X_t] \mid \mathcal{F}_s] = X_s - \mathbb{E}[X_s], \quad \forall 0 \leq s < t. \quad (3.5.1)$$

Then

- (i) if $Z_t \in L^1, \forall t \geq 0$, then $\tilde{Z}_t := Z_t - \mathbb{E}[Z_t]$ is a martingale;
- (ii) if $Z_t \in L^2, \forall t \geq 0$, then $\tilde{Z}_t^2 - \mathbb{E}[\tilde{Z}_t^2]$ is a martingale;
- (iii) if, for some $\theta \in \mathbb{R}$, we have $\mathbb{E}[e^{\theta Z_t}] < \infty, \forall t \geq 0$, then

$$X_t := \frac{e^{\theta Z_t}}{\mathbb{E}[e^{\theta Z_t}]}$$

is a martingale.

The case (i) follows from (3.5.1). The case (iii) is the continuous time analogue of Example 3.1.7 and the proof is similar. To prove (ii) note that

$$\begin{aligned} \mathbb{E}[\tilde{Z}_t^2 \mid \mathcal{F}_s] &= \mathbb{E}[(\tilde{Z}_s + \tilde{Z}_t - \tilde{Z}_s)^2 \mid \mathcal{F}_s] \\ &= \tilde{Z}_s^2 + 2\tilde{Z}_s \underbrace{\mathbb{E}[(\tilde{Z}_t - \tilde{Z}_s) \mid \mathcal{F}_s]}_{=0} + \mathbb{E}[(\tilde{Z}_t - \tilde{Z}_s)^2 \mid \mathcal{F}_s] = \tilde{Z}_s^2 + \mathbb{E}[(\tilde{Z}_t - \tilde{Z}_s)^2] \\ &= \tilde{Z}_s^2 + \mathbb{E}[\tilde{Z}_t^2] - 2\mathbb{E}[\tilde{Z}_s \tilde{Z}_t] + \mathbb{E}[\tilde{Z}_s^2] \\ &= \tilde{Z}_s^2 + \mathbb{E}[\tilde{Z}_t^2] - 2\mathbb{E}[\mathbb{E}[\tilde{Z}_s \tilde{Z}_t \mid \mathcal{F}_s]] + \mathbb{E}[\tilde{Z}_s^2] = \tilde{Z}_s^2 + \mathbb{E}[\tilde{Z}_t^2] - \mathbb{E}[\tilde{Z}_s^2]. \end{aligned}$$

Hence

$$\mathbb{E}[\tilde{Z}_t^2 - \mathbb{E}[\tilde{Z}_t^2] \mid \mathcal{F}_s] = \tilde{Z}_s^2 - \mathbb{E}[\tilde{Z}_s^2].$$

Classical examples of processes with independent increments are the Brownian motion, the Poisson process, or more generally the Lévy processes, [43, Chap. VII].

If B_t is a 1-dimensional Brownian motion started at 0, adapted to $\mathcal{F}_t$, then B_t is a normal random variable with mean 0 and variance t , for each $t > 0$. The moment generating function of B_t is

$$M_{B_t}(\theta) = \mathbb{E}[e^{\theta B_t}] = e^{\frac{\theta^2 t}{2}}.$$

We deduce from the above that

$$B_t, \quad B_t^2 - t, \quad e^{\theta B_t - \frac{\theta^2}{2}t}$$

are martingales, $\forall \theta \in \mathbb{R}$. The martingale

$$\left(e^{\theta B_t - \frac{\theta^2}{2}t} \right)_{t \geq 0}, \quad (3.5.2)$$

is called the *exponential martingale* of the Brownian motion.

Note that if we set $\lambda := \theta\sqrt{t}$, and $X = \frac{B_t}{\sqrt{t}}$, then

$$e^{\theta B_t - \frac{\theta^2}{2}t} = e^{\lambda X - \lambda^2/2} \stackrel{(1.6.6)}{=} \sum_{n \geq 0} H_n(X) \frac{\lambda^n}{n!}, \quad (3.5.3)$$

where $H_n(x)$ is the n -th Hermite polynomial (1.6.5). We can rewrite the above equality as

$$e^{\theta B_t - \frac{\theta^2}{2}t} = \sum_{n \geq 0} M_n(t) \frac{\theta^n}{n!}, \quad M_n(t) = t^{n/2} H_n(B_t/\sqrt{t}).$$

Each of the coefficients $M_n(t)$ is a continuous-time martingale. Note that

$$M_1(t) = B_t, \quad M_2(t) = B_t^2 - t.$$

□

Example 3.5.4 (New submartingales from old.). If $(X_t)_{t \geq 0}$ is a martingale and $f : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function such that $f(X_t) \in L^1, \forall t \geq 0$, then $(f(X_t))_{t \geq 0}$ is a submartingale. If $(X_t)_{t \geq 0}$ is only a submartingale and additionally, f is nondecreasing, then $(f(X_t))_{t \geq 0}$ is a submartingale. □

We conclude this subsection with a description of a large family of continuous martingales naturally associated to the Brownian motion. Suppose that

$$B_1(t), \dots, B_d(t) : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad \omega \mapsto B_k(\omega, t), \quad t \geq 0, \quad k = 1, \dots, d,$$

are independent Brownian motions defined on the same complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$. The d -dimensional Brownian motion is the stochastic process

$$\mathbf{B} : (\Omega, \mathcal{S}, \mathbb{P}) \times [0, \infty) \rightarrow \mathbb{R}^d, \quad (\omega, t) \mapsto \mathbf{B}(t) = (B_1(\omega, t), \dots, B_d(\omega, t))$$

Arguing as in Proposition 3.3.3 we deduce that $\mathbf{B}$ is progressively measurable, i.e., for any $t \geq 0$ the map $\mathbf{B} : \Omega \times [0, t] \rightarrow \mathbb{R}^d$ is measurable.

The distribution of $\mathbf{B}(t)$ is the Gaussian measure on $\mathbb{R}^d$

$$\Gamma_{t\mathbb{I}}[d\mathbf{x}] = \underbrace{(2\pi t)^{-d/2} e^{-\frac{|\mathbf{x}|^2}{2t}}}_{=: p_t(\mathbf{x})} \lambda[d\mathbf{x}].$$

The density $p_t(\mathbf{x})$ satisfies the heat equation

$$\partial_t p_t(\mathbf{x}) - \frac{1}{2} \Delta_{\mathbf{x}} p_t(\mathbf{x}) = 0, \quad \forall t > 0, \mathbf{x} \in \mathbb{R}^d \quad (3.5.4)$$

where Δ is the physicists'⁸ Laplacian

$$\Delta_{\mathbf{x}} = \partial_{x_1}^2 + \cdots + \partial_{x_d}^2.$$

Denote by L the partial differential operator

$$L = \partial_t + \frac{1}{2}\Delta_{\mathbf{x}},$$

and by $C_b^k(\mathbb{R}^d)$ the space of C^k -functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ whose derivatives up to order k are bounded. A function $f : [0, \infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is said to be *admissible* if

- it is C^1 in the t variable, C^2 in the $\mathbf{x}$ variables and
- there exists a locally bounded function $c : [0, \infty) \rightarrow (0, \infty)$ such that

$$\forall t > 0, \quad \|\partial_t f\|_{C_b^0(\mathbb{R}^d)} + \|f_t(\mathbf{x})\|_{C_b^2(\mathbb{R}^d)} \leq c(t). \quad (3.5.5)$$

If f is an admissible function, then an integration by parts shows coupled with the fact that $p_t(\mathbf{x})$ decays very fast as $\|\mathbf{x}\| \rightarrow \infty$ shows that

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}^d} (\Delta_{\mathbf{x}} f_t(\mathbf{x})) p_t(\mathbf{x}) d\mathbf{x} &= \frac{1}{2} \int_{\mathbb{R}^d} f_t(\mathbf{x}) (\Delta_{\mathbf{x}} p_t(\mathbf{x})) d\mathbf{x} \\ &= \int_{\mathbb{R}^d} f_t(\mathbf{x}) (\partial_t p_t(\mathbf{x})) d\mathbf{x}. \end{aligned} \quad (3.5.6)$$

Proposition 3.5.5. *Suppose that $f : [0, \infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is admissible. Define*

$$M_t = M_t^f := f(t, \mathbf{B}(t)) - f(0, \mathbf{B}(0)) - \int_0^t (Lf)(s, \mathbf{B}(s)) ds.$$

Then M_t is a continuous time martingale with continuous paths adapted to the natural filtration $\mathcal{F}_t = \sigma(\mathbf{B}(s))_{0 \leq s \leq t}$.

Proof. Observe first that since $\mathbf{B}(t)$ is progressively measurable and f is admissible the function

$$\Omega \times [0, t] \ni (\omega, s) \mapsto f(s, \mathbf{B}(s)) \in \mathbb{R}$$

is L^1 . Fubini-Tonelli's theorem 1.3.41 implies that M_t is $\mathcal{F}_t$ -measurable and L^1 . We first prove that for $t > s > 0$ we have

$$\mathbb{E}[M_t \mid \mathcal{F}_s] = M_s.$$

The increments $W_\tau = \mathbf{B}(s + \tau) - \mathbf{B}(s)$, $\tau \geq 0$ define a Brownian motion independent of $\mathcal{F}_s$.

For $z \in \mathbb{R}^d$ we set

$$\phi_z(\tau, u) := f(s + \tau, z + u).$$

and thus

$$\begin{aligned} \mathbb{E}[M_{s+\tau} - M_s \mid \mathcal{F}_s] &= \mathbb{E}[f(t, W_\tau + \mathbf{B}(s)) - f(s, \mathbf{B}(s)) \mid \mathcal{F}_s] \\ &\quad - \mathbb{E}\left[\int_0^\tau (Lf)(r, \mathbf{B}(s) + W(r)) dr \mid \mathcal{F}_s\right] \\ &= \mathbb{E}[\phi_z(\tau, W_\tau)] \big|_{z=\mathbf{B}(s)} - \mathbb{E}[\phi_z(0, 0)] \big|_{z=\mathbf{B}(s)} \\ &\quad - \mathbb{E}\left[\left(\int_0^\tau (L\phi_z)(r, W_r) dr\right)\right]_{z=\mathbf{B}(s)}. \end{aligned}$$

For simplicity we set $\phi := \phi_{z=\mathbf{B}(s)}$. We need the following technical fact.

⁸The geometers' Laplacian differs by sign $\Delta_{\text{geom}} = -\Delta_{\text{phys}} = -\sum_i \partial_{x_i}^2$.

Lemma 3.5.6. Let $g : [0, \infty) \times \mathbb{R}^d$ be a continuous function such that there exists a locally bounded function $c : [0, \infty) \rightarrow (0, \infty)$ so that

$$|g(r, u)| \leq c(r), \quad \forall (r, u) \in \mathbb{R}^d.$$

Then the function $[0, \infty) \ni r \mapsto \mathbb{E}[g(r, W_r)]$ is continuous. $\square$

Assuming the validity of this lemma we deduce

$$\mathbb{E}\left[\int_0^\tau (L\phi)(r, W_r) dr\right] = \lim_{\varepsilon \searrow 0} \mathbb{E}\left[\int_\varepsilon^\tau (L\phi)(r, W_r) dr\right].$$

On the other hand,

$$\begin{aligned} \mathbb{E}\left[\int_\varepsilon^\tau (L\phi)(r, W_r) dr\right] &= \mathbb{E}\left[\int_\varepsilon^\tau (\partial_r \phi)(r, W_r) + \frac{1}{2}(\Delta_u \phi)(r, W_r) dr\right] \\ &= \int_\varepsilon^\tau \int_{\mathbb{R}^d} \partial_r \phi(r, u) p_r(u) du + \frac{1}{2} \int_0^\tau \int_{\mathbb{R}^d} \Delta_u \phi(r, u) p_r(u) du \\ &\stackrel{(3.5.6)}{=} \int_\varepsilon^\tau \int_{\mathbb{R}^d} \partial_r (\phi(r, u) p_r(u)) du - \int_0^\tau \int_{\mathbb{R}^d} \phi(r, u) \partial_r p_r(u) du \\ &\quad + \frac{1}{2} \int_s^\tau \int_{\mathbb{R}^d} \phi(r, u) \Delta_u p_r(u) du \\ &\stackrel{(3.5.4)}{=} \int_{\mathbb{R}^d} \left(\int_\varepsilon^\tau \partial_r (\phi(r, u) p_r(u)) dr \right) du. \\ &= \int_{\mathbb{R}^d} f(\tau, \mathbf{x}) p_\tau(u) du - \int_{\mathbb{R}^d} \phi(\varepsilon, u) p_\varepsilon(u) du \\ &= \mathbb{E}[f(\tau, W_\tau)] - \mathbb{E}[f(\varepsilon, W_\varepsilon)]. \end{aligned}$$

Lemma 3.5.6 implies that

$$\lim_{\varepsilon \searrow 0} \left(\mathbb{E}[f(\tau, W_\tau)] - \mathbb{E}[f(\varepsilon, W_\varepsilon)] \right) = \mathbb{E}[\phi(\tau, W_\tau)] - \mathbb{E}[\phi(0, W_0)].$$

This proves that M_t^f is a martingale. Clearly its sample paths are continuous.

Proof of Lemma 3.5.6. Fix $\tau_0 \geq 0$, $T > r_0$ and set

$$C := \sup_{r \in [0, T]} c(r).$$

Then

$$|g(t, u)| \leq C, \quad \forall (t, u) \in [0, T] \times \mathbb{R}^d. \quad (3.5.7)$$

If $r_0 > 0$, then the equality

$$\lim_{r \rightarrow r_0} \mathbb{E}[g(r, W_r)] = \mathbb{E}[g(r_0, W_{r_0})]$$

follows from the Dominated Convergence Theorem aided by (3.5.7). Suppose that $r_0 = 0$. Then

$$\mathbb{E}[g(r, W_r)] = \int_{\mathbb{R}^d} g(r, u) p_r(u) du, \quad \mathbb{E}[g(r_0, W_{r_0})] = g(0, 0) = \int_{\mathbb{R}^d} g(0, 0) p_r(u) du$$

so that

$$|\mathbb{E}[g(r, W_r)] - \mathbb{E}[g(0, W_0)]| \leq \int_{\mathbb{R}^d} |g(r, u) - g(0, 0)| p_r(u) du$$

$$\begin{aligned}
&= \int_{|u| \leq \rho} |g(r, u) - g(0, 0)| p_r(u) du + \int_{|u| > \rho} |g(r, u) - g(0, 0)| p_r(u) du \\
&\leq \underbrace{\sup_{|u| < \rho} |g(r, u) - g(0, 0)|}_{=: \delta(r, \rho)} + 2 \int_{|u| > \rho} C p_r(u) du.
\end{aligned}$$

Note that $\delta(r, \rho)$ is non-decreasing in r and

$$\begin{aligned}
\int_{|u| > \rho} p_r(u) du &= \mathbb{P}[|W_r| > \rho] = \mathbb{P}[r^{1/2}|W_1| > \rho] \\
&= \mathbb{P}[|W_1| > r^{-1/2}\rho] \searrow 0 \text{ as } r \searrow 0,
\end{aligned}$$

so

$$0 \leq \limsup_{r \searrow 0} |\mathbb{E}[g(r, W_r)] - \mathbb{E}[g(0, W_0)]| \leq \underbrace{\lim_{r \searrow 0} \delta(r, \rho)}_{=: \delta(0, \rho)}, \quad \forall \rho > 0.$$

Since g is uniformly continuous on the compact set $[0, T] \times \{|u| \leq \rho\}$ we deduce that

$$\lim_{\rho \searrow 0} \delta_0(\rho) = \lim_{\rho \searrow 0} \sup_{|u| \leq \rho} |g(0, u) - g(0, 0)| = 0.$$

□

This completes the proof of Proposition 3.5.5. □

Remark 3.5.7. The growth condition (3.5.5) can be relaxed to

$$\exists C \geq 0, \quad \forall t > 0, \quad \|e^{-C|\mathbf{x}|} \partial_t f(\mathbf{x})\|_{C_b^0(\mathbb{R}^d)} + \|e^{-C|\mathbf{x}|} f_t(\mathbf{x})\|_{C_b^2(\mathbb{R}^d)} \leq c(t). \quad (3.5.8)$$

□

Corollary 3.5.8. For any $f \in C_b^2(\mathbb{R}^d)$ the stochastic process

$$M_t^f = f(\mathbf{B}(t)) - f(0) - \frac{1}{2} \int_0^t (\Delta f)(\mathbf{B}(s)) ds, \quad t \geq 0 \quad (3.5.9)$$

is a martingale with continuous paths. □

It turns out that the last corollary characterizes the Brownian motion.

Proposition 3.5.9. Suppose that $(X_t)_{t \geq 0}$ is a stochastic process with continuous paths such that, $X_0 = 0$ and for any $f \in C_b^2(\mathbb{R})$ the stochastic process

$$M_t^f = f(X_t) - f(0) - \frac{1}{2} \int_0^t f''(X_s) ds, \quad t \geq 0 \quad (3.5.10)$$

is a continuous martingale with respect to the filtration $\mathcal{F}_t = \sigma(X_s, 0 \leq s \leq t)$. Then $(X_t)_{t \geq 0}$ is a Brownian motion.

Proof. For ease of presentation we set $\mathbb{E}_s := \mathbb{E}[\cdot | \mathcal{F}_s]$. For any $f \in C_b^2(\mathbb{R})$, and any $0 \leq s \leq t$ we have

$$0 = \mathbb{E}_s[M_t^f - M_s^f] = \mathbb{E}_s\left[f(X_t) - f(X_s) - \frac{1}{2} \int_s^t f''(X_u) du\right].$$

Suppose now that $f(x) = e^{i\xi x} = \cos(\xi x) + i \sin(\xi x)$. Then

$$f(X_t) - f(X_s) = e^{i\xi(X_t - X_s)} e^{i\xi X_s}, \quad f''(x) = -\xi^2 f(x),$$

$$\mathbb{E}_s[f(X_t) - f(X_s)] = \mathbb{E}_s[e^{i\xi(X_t - X_s)} - 1] e^{i\xi X_s}$$

$$\frac{1}{2} \mathbb{E}_s \left[\int_s^t f''(X_u) du \right] = -\frac{\xi^2 e^{i\xi X_s}}{2} \int_s^{t+s} \mathbb{E}_s[e^{i\xi(X_u - X_s)}] du$$

Hence, if we set

$$g(t) = \mathbb{E}_s[e^{i\xi(X_t - X_s)}]$$

we deduce

$$g(t) = 1 - \frac{\xi^2}{2} \int_s^t g(u) du, \quad t \geq s.$$

Thus $g(t)$ is the solution of the initial value problem

$$g'(t) = -\frac{\xi^2}{2} g(t), \quad g(s) = 1,$$

so that $g(t) = e^{-\xi^2(t-s)/2}$. This is independent of ω and depends only on $\tau = (t - s)$.

We deduce from Corollary 2.2.30 that the increment $X_{s+\tau} - X_s$ is independent of $\mathcal{F}_s$ and its characteristic function is $e^{-\tau\xi^2/2}$. Hence $X_{s+\tau} - X_s$ is Gaussian with mean 0 and variance τ . This proves that X_t is a Brownian motion. $\square$

3.5.2. Limit theorems. Fix a filtered probability space $(\Omega, (\mathcal{F}_t)_{t \geq 0}, \mathcal{F}, \mathbb{P})$.

Definition 3.5.10. An R -(super)martingale is a (super)martingale $(X_t)_{t \geq 0}$ adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ such that the paths of X_t are a.s. R -functions. A continuous (sub/super)martingale is a (sub/super)martingale $(X_t)_{t \geq 0}$ adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ such that its paths are a.s. continuous. $\square$

Remark 3.5.11. Suppose that $(X_t)_{t \geq 0}$ is an R -supermartingale. Fix a negligible set $\mathcal{N} \subset \Omega$ such that $t \mapsto X_t(\omega)$ is an R -function for any $\omega \in \Omega \setminus \mathcal{N}$. Fix a dense countable subset D of $[0, \infty)$.

Note that for every open interval $I \subset [0, \infty)$ we have

$$\sup_{t \in D \cap I} X_t(\omega) = \sup_{t \in I} X_t(\omega), \quad \inf_{t \in D \cap I} X_t(\omega) = \inf_{t \in I} X_t(\omega), \quad \forall \omega \in \Omega \setminus \mathcal{N}. \quad (3.5.11)$$

This shows that $(X_t)_{t \geq 0}$ is a *separable process* in the sense of Doob, [61, II.2]. This means that there exist

- a countable dense subset $D \subset [0, \infty)$, and
- a negligible subset $\mathcal{N} \subset \Omega$,

such that, for any closed interval $I \subset \mathbb{R}$, and any open subset $\mathcal{O}$ of $[0, \infty)$, the sets

$$\{\omega; X_s(\omega) \in I, \forall s \in D \cap \mathcal{O}\} \text{ and } \{\omega; X_t(\omega) \in I, \forall t \in \mathcal{O}\}$$

differ by a subset of $\mathcal{N}$. A dense countable subset D with the above property is called a *separability set*. $\square$

Before we proceed investigating the properties of R -supermartingales it would help to understand how restrictive is the assumption that the paths are a.s. R -functions. The proof of Theorem 3.2.56 shows that if $(X_t)_{t \geq 0}$ is an R -submartingale, then, for any bounded set $S \subset [0, \infty)$ the family $(X_s)_{s \in S}$ is bounded in L^1 .

The Submartingale Convergence Theorem 3.2.4 and the Backwards Convergence Theorem 3.2.56 imply that for any $\tau > 0$ and any sequences of rational numbers $s_n \nearrow \tau$ and $t_n \searrow \tau$ the sequences (X_{s_n}) and (X_{t_n}) converge a.s. to L^1 random variables.

We have a more precise result, [43, Sec. V.7], [132, Sec. 3.3], [180, II.65-67].

Theorem 3.5.12 (Doob's regularization theorem). *If the filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfies the usual conditions, then a submartingale $(X_t)_{t \geq 0}$ adapted to this filtration admits an R -submartingale modification if and only if the function $t \mapsto \mathbb{E}[X_t]$ is right-continuous.* $\square$

☞ In the remainder of Section 3.5 I will assume that the filtrations satisfy the usual conditions.

Theorem 3.5.13 (Doob's maximal inequality). *Suppose that $(X_t)_{t \geq 0}$ is an R -submartingale adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$. Then, for any $a, t > 0$ we have*

$$a\mathbb{P}\left[\sup_{s \in [0, t]} |X_s| > a\right] \leq \mathbb{E}[|X_t^+|] \leq \mathbb{E}[|X_t|] + \mathbb{E}[|X_0|]. \quad (3.5.12)$$

Proof. For any $m \in \mathbb{N}$ we set

$$D_m := \left\{0, \frac{t}{m}, \dots, \frac{(m-1)t}{m}, t\right\}, \quad D := \bigcup_{m \in \mathbb{N}} D_m.$$

The discrete Doob maximal inequality (3.2.36) implies that

$$a\mathbb{P}\left[\sup_{s \in D_m} |X_s| > a\right] \leq \mathbb{E}[|X_t^+|] \quad \text{and} \quad a\mathbb{P}\left[\sup_{s \in D} |X_s| > a\right] \leq \mathbb{E}[|X_t^+|].$$

As observed in Remark 3.5.11 $(X_\bullet)$ is a separable process so (3.5.11)

$$\mathbb{P}\left[\sup_{s \in D} |X_s| > a\right] = \mathbb{P}\left[\sup_{s \in [0, t]} |X_s| > a\right].$$

$\square$

Theorem 3.5.14 (Doob's L^p -inequality). *Suppose that $(X_t)_{t \geq 0}$ is an R -martingale. Then, for any $t > 0$ and $p > 1$ we have*

$$\mathbb{E}\left[\sup_{s \in [0, t]} |X_s|^p\right]^{\frac{1}{p}} \leq q \|X_t\|_{L^p}, \quad \frac{1}{q} = 1 - \frac{1}{p}. \quad (3.5.13)$$

Proof. Argue as in the proof of Theorem 3.5.13 by relying on the separability of $(X_\bullet)$ and the discrete L^p -inequality (3.2.38). $\square$

Theorem 3.5.15. *Suppose that $(X_t)_{t \geq 0}$ is an R -submartingale and*

$$\sup_{t > 0} \mathbb{E}[|X_t|] < \infty. \quad (3.5.14)$$

Then there exists an integrable random variable X_∞ such that

$$\lim_{t \rightarrow \infty} X_t = X_\infty \quad \text{a.s..}$$

Proof. For any $m \in \mathbb{N}$ we set

$$D_m := \frac{1}{2^m} \mathbb{N}, \quad m \in \mathbb{N}, \quad D = \bigcup_{m \in \mathbb{N}} D_m.$$

For any function $f : [0, \infty) \rightarrow \mathbb{R}$, any rational numbers $a < b$ and any $S \subset [0, \infty)$ we denote by $N(f, S, [a, b])$ the supremum of the set of integers k such that there exist

$$s_1 < t_1 < \cdots < s_k < t_k$$

in S such that $f(s_i) \leq a$, $f(t_i) \geq b$, $\forall i = 1, \dots, k$.

For $m \in \mathbb{N}$ we set $N_m(f, [a, b]) := N(f, D_m, [a, b])$. Equivalently, $N_m(f, [a, b])$ is the number of upcrossings of the strip $[a, b]$ by the function $f|_{D_m}$. Note that

$$N_m(X, [a, b]) \leq N_{m+1}(X, [a, b]), \quad \forall m,$$

and

$$N(f, D, [a, b]) = \lim_{m \rightarrow \infty} N_m(X, [a, b]).$$

Doob's upcrossing inequality (3.2.2) implies

$$(b - a) \mathbb{E}[N_m(X, [a, b])] \leq \sup_{t > 0} \mathbb{E}[(X_t - a)^+] - \mathbb{E}[(X_0 - a)^+], \quad \forall m \in \mathbb{N}.$$

Letting $m \rightarrow \infty$ we deduce from the Monotone Convergence Theorem

$$(b - a) \mathbb{E}[N(X, D, [a, b])] \leq \sup_{t > 0} \mathbb{E}[(X_t - a)^+] - \mathbb{E}[(X_0 - a)^+] < \infty.$$

Thus $N(X, D, [a, b]) < \infty$ a.s. so the limit

$$X_\infty := \lim_{\substack{t \rightarrow \infty \\ t \in D}} X_t$$

exists a.s. We leave the reader convince her/himself that since the process $X_\bullet$ is *separable* (see Remark 3.5.11) the limit

$$X_\infty = \lim_{t \rightarrow \infty} X_t$$

exists a.s.. The boundedness assumption (3.5.14) coupled with Fatou's lemma implies that X_∞ is integrable. $\square$

The above theorem implies immediately the following continuous time counterpart of Theorem 3.2.23.

Theorem 3.5.16 (UI martingales). *Suppose that $(X_t)_{t \geq 0}$ is an UI R -martingale, i.e., the family $(X_t)_{t \geq 0}$ is uniformly integrable. Then*

$$X_\infty = \lim_{t \rightarrow \infty} X_t$$

exists a.s. and L^1 and

$$X_t = \mathbb{E}[X_\infty | \mathcal{F}_t], \quad \forall t > 0.$$

$\square$

3.5.3. Sampling and stopping. Suppose that $(X_t)_{t \geq 0}$ is an R -submartingale such that

$$X_\infty = \lim_{t \rightarrow \infty} X_t$$

exists a.s.. Let $T : \Omega \rightarrow [0, \infty]$ be a stopping time adapted to the filtration $(\mathcal{F}_t)$. The *optional sampling* of $X_\bullet$ at T is the random variable

$$X_T(\omega) = \mathbf{I}_{T < \infty} X_{T(\omega)}(\omega) + \mathbf{I}_{T = \infty} X_\infty(\omega).$$

Theorem 3.5.17 (Optional sampling). *Suppose that $(X_t)_{t \geq 0}$ is an UI R -martingale and S, T are stopping times such that $S \leq T$. Then the following hold.*

- (i) *The random variables X_S, X_T are integrable.*
- (ii) $X_S = \mathbb{E}[X_T \mid \mathcal{F}_S] = \mathbb{E}[X_\infty \mid \mathcal{F}_S]$.
- (iii) $\mathbb{E}[X_S] = \mathbb{E}[X_\infty] = \mathbb{E}[X_0]$.

Proof. We set

$$S_n = \sum_{k=0}^{\infty} \frac{k+1}{2^n} \mathbf{I}_{\{k2^{-n} < S \leq (k+1)2^{-n}\}} + \infty \mathbf{I}_{S=\infty},$$

$$T_n = \sum_{k=0}^{\infty} \frac{k+1}{2^n} \mathbf{I}_{\{k2^{-n} < T \leq (k+1)2^{-n}\}} + \infty \mathbf{I}_{T=\infty}$$

Observe that $S_n \geq S$, $T_n \geq T$ and $S_n \leq T_n$, $\forall n$.

Let us show that S_n is $\mathcal{F}_S$ measurable and T_n is T -measurable. In other words, we have to show that

$$\{S_n \leq c\} \cap \{S \leq s\} \in \mathcal{F}_s, \quad \forall c, s \geq 0.$$

Note that

$$\begin{aligned} \{S \leq s\} \cap \{S_n \leq c\} &= \{S \leq s\} \cap \left(\bigcup_{(k+1)2^{-n} \leq c} \left\{ k2^{-n} < S \leq (k+1)2^{-n} \right\} \right) \in \mathcal{F}_c. \\ &= \bigcup_{(k+1)2^{-n} \leq c} \left\{ k2^{-n} < S \leq \min(s, (k+1)2^{-n}) \right\} \in \mathcal{F}_s. \end{aligned}$$

Proposition 3.3.13(ii) now implies that S_n is a stopping time. A similar argument shows that T_n is a stopping time. Note that

$$S_n \searrow S \text{ and } T_n \searrow T \text{ as } n \rightarrow \infty.$$

For $n \in \mathbb{N}_0$ set $D_n = 2^{-n}\mathbb{N}_0$. For each $n \in \mathbb{N}_0$ the stochastic process

$$X^n := (X_t)_{t \in D_n},$$

is a UI *discrete* martingales with respect to the filtration $\mathcal{F}_\bullet^n := (\mathcal{F}_t)_{t \in D_n}$. The above arguments show that S_n and T_n are stopping times with respect to these filtrations. We deduce from the discrete Optional Sampling Theorems 3.2.28 that

$$X_{S_n} = X_{S_n}^n = \mathbb{E}[X_{T_n}^n \mid \mathcal{F}_{S_n}^n] = \mathbb{E}[X_{T_n} \mid \mathcal{F}_{S_n}],$$

and

$$X_{S_n} = \mathbb{E}[X_\infty \mid \mathcal{F}_{S_n}], \quad X_{T_n} = \mathbb{E}[X_\infty \mid \mathcal{F}_{T_n}].$$

Now observe that since (X_t) is a.s. right continuous we have

$$X_S = \lim_{n \rightarrow \infty} X_{S_n} \text{ and } X_T = \lim_{n \rightarrow \infty} X_{T_n} \text{ a.s..}$$

The families (X_{S_n}) and (X_{T_n}) are UI so the above convergences also hold in L^1 . Since $\mathcal{F}_S \subset \mathcal{F}_{S_n} \subset \mathcal{F}_{T_n}$ and the conditional expectation map

$$\mathbb{E}[\cdot | \mathcal{F}_S] : L^1(\Omega, \mathcal{F}, \mathbb{P}) \rightarrow L^1(\Omega, \mathcal{F}_S, \mathbb{P})$$

is a contraction we deduce

$$\begin{aligned} X_S &= \mathbb{E}[X_S | \mathcal{F}_S] = \lim_{n \rightarrow \infty} \mathbb{E}[X_{S_n} | \mathcal{F}_S] \\ &= \lim_{n \rightarrow \infty} \mathbb{E}[X_{T_n} | \mathcal{F}_S] = \mathbb{E}[X_T | \mathcal{F}_S], \end{aligned}$$

where the above converges are in L^1 . $\square$

Corollary 3.5.18. *Suppose that $(X_t)_{t \geq 0}$ is an R -martingale and S, T are bounded stopping times such that $S \leq T$ a.s.. Then the following hold.*

- (i) *The random variables X_S, X_T are integrable.*
- (ii) $X_S = \mathbb{E}[X_T | \mathcal{F}_S] = \mathbb{E}[X_\infty | \mathcal{F}_S]$.

Proof. Fix $t_0 > 0$ such $S, T \leq t_0$ a.s.. Then the stopped process $X_{t \wedge t_0}$ is an UI R -martingale. The conclusions now follow from Theorem 3.5.17 applied to this stopped martingale. $\square$

Corollary 3.5.19 (Optional stopping). *Suppose that $(X_t)_{t \geq 0}$ is an R -martingale adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ and T is a stopping time. Then the following hold.*

- (i) *The stopped process*

$$X_t^T := X_{t \wedge T}$$

is an R -martingale adapted to the same filtration $(\mathcal{F}_t)_{t \geq 0}$.

- (ii) *If additionally $(X_t)_{t \geq 0}$ is UI, then so is the stopped process X^T , and we have*

$$X_{T \wedge t} = \mathbb{E}[X_T | \mathcal{F}_t], \quad (3.5.15)$$

$$X_T = \lim_{t \rightarrow \infty} X_t \text{ a.s. and } L^1. \quad (3.5.16)$$

Proof. We begin by proving (ii). For $s < t$, the stopping times $s \wedge T$ and $t \wedge T$ are bounded and $s \wedge T \leq t \wedge T$. The random variables $X_{t \wedge T}$ are $\mathcal{F}_{t \wedge T}$ -measurable and thus $\mathcal{F}_t$ -measurable since $\mathcal{F}_{t \wedge T} \subset \mathcal{F}_t$. To prove (3.5.15) it suffices to check that for any $A \in \mathcal{F}_t$ we have

$$\mathbb{E}[X_T \mathbf{I}_A] = \mathbb{E}[X_{t \wedge T} \mathbf{I}_A].$$

Decompose $\mathbf{I}_A = \mathbf{I}_{A \cap \{T \leq t\}} + \mathbf{I}_{A \cap \{T > t\}}$. We have

$$X_T \mathbf{I}_{A \cap \{T \leq t\}} = X_{t \wedge T} \mathbf{I}_{A \cap \{T \leq t\}}$$

so that

$$\mathbb{E}[X_T \mathbf{I}_{A \cap \{T \leq t\}}] = \mathbb{E}[X_{t \wedge T} \mathbf{I}_{A \cap \{T \leq t\}}]. \quad (3.5.17)$$

On the other hand, we deduce from Theorem 3.5.17 that

$$X_{t \wedge T} = \mathbb{E}[X_T | \mathcal{F}_{t \wedge T}].$$

Now observe that

$$A \cap \{T > t\} \in \mathcal{F}_t \text{ and } A \cap \{T > t\} \in \mathcal{F}_T,$$

so $A \cap \{T > t\} \in \mathcal{F}_t \cap \mathcal{F}_T = \mathcal{F}_{t \wedge T}$. Hence

$$\begin{aligned} X_{t \wedge T} \mathbf{I}_{A \cap \{T > t\}} &= \mathbb{E}[X_T \mathbf{I}_{A \cap \{T > t\}} \mid \mathcal{F}_{t \wedge T}], \\ \mathbb{E}[X_{t \wedge T} \mathbf{I}_{A \cap \{T > t\}}] &= \mathbb{E}[X_T \mathbf{I}_{A \cap \{T > t\}}]. \end{aligned} \quad (3.5.18)$$

The desired conclusion follows by adding (3.5.17) and (3.5.18). The assertion (3.5.16) follows from the fact that the stopped martingale X^T is UI. Part (i) now follows from (ii) applied to the sequence of UI martingales

$$(X_t^n)_{t \geq 0} := (X_{n \wedge t})_{t \geq 0}, \quad n \in \mathbb{N},$$

Indeed, the martingales X^n are compatible with $\mathcal{F}_t$ and for $s < t$ we have

$$\mathbb{E}[X_{T \wedge t}^n \mid \mathcal{F}_s] = \mathbb{E}[\mathbb{E}[X_T^n \mid \mathcal{F}_t] \mid \mathcal{F}_s] = \mathbb{E}[X_T^n \mid \mathcal{F}_s] = X_{T \wedge s}^n.$$

Now let $n \rightarrow \infty$ and observe that for $n > t$ we have $X_{T \wedge t}^n = X_{T \wedge t}$. $\square$

Definition 3.5.20. Let $X = (X_t)_{t \geq 0}$ be a martingale adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ and T is a stopping time compatible with the same filtration. We say that T reduces X , or that T is *reducing* if the stopped martingale $X^T = (X_{T \wedge t})_{t \geq 0}$ is UI. $\square$

The excellent interaction between UI martingales and the operation of sampling essentially characterizes the UI martingales. We have the following result that is particularly useful in the theory of stochastic integration.

Theorem 3.5.21. Assume that the filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfies the usual conditions. Let

$$M = (M_t)_{1 \leq t \leq \infty}$$

be a progressive process such that, for any finite or infinite stopping time T we have $\mathbb{E}[|M_T|] < \infty$ and $\mathbb{E}[M_T] = 0$. Then M is a UI martingale.

Proof. I will follow the argument in the proof of [180, Thm.II.77.6]. The assumptions show that M_∞ is integrable and $\mathbb{E}[M_\infty] = 0$.

Let $t \geq 0$ and $F \in \mathcal{F}_t$. Consider the stopping time $T = t\mathbf{I}_F + \infty \cdot \mathbf{I}_{F^c}$. We have

$$\begin{aligned} 0 &= \mathbb{E}[M_T] = \mathbb{E}[M_t \mathbf{I}_F] + \mathbb{E}[M_\infty \mathbf{I}_{F^c}], \\ 0 &= \mathbb{E}[M_\infty] = \mathbb{E}[M_\infty \mathbf{I}_F] + \mathbb{E}[M_\infty \mathbf{I}_{F^c}]. \end{aligned}$$

We deduce that

$$\mathbb{E}[M_\infty \mathbf{I}_F] = \mathbb{E}[M_t \mathbf{I}_F], \quad \forall F \in \mathcal{F}_t,$$

i.e., $M_t = \mathbb{E}[M_\infty \mid \mathcal{F}_t]$. $\square$

Corollary 3.5.22. Assume that the filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfies the usual conditions. Let $M = (M_t)_{t \geq 0}$ be a UI R-martingale and $T_1 \leq T_2$ two (possibly infinite) stopping times. Let Z be a bounded $\mathcal{F}_{T_1}$ -measurable random variable. Then

$$X_t = Z \cdot (M_{t \wedge T_2} - M_{t \wedge T_1})$$

is a UI R-martingale.

Proof. According to the previous theorem it suffices to show that for any stopping time S we have

$$\mathbb{E}[|X_S|] < \infty, \quad (3.5.19a)$$

$$\mathbb{E}[X_S] = 0. \quad (3.5.19b)$$

Set $\|Z\| := \|Z\|_{L^\infty}$. We have

$$\mathbb{E}[|X_S|] \leq \|Z\| \cdot \mathbb{E}[|M_{T_2 \wedge S}| + |M_{T_1 \wedge S}|].$$

From the optional sampling theorem we deduce

$$M_{T_i \wedge S} = \mathbb{E}[M_\infty | \mathcal{F}_{T_i \wedge S}].$$

Since the conditional expectation is an L^1 -contraction we deduce that

$$\mathbb{E}[|M_{T_2 \wedge S}|] \leq \mathbb{E}[|M_\infty|] < \infty.$$

This proves (3.5.19a). The Monotone Class Theorem shows that it suffices to prove (3.5.19b) only in the special case when $A = I_A$, $A \in \mathcal{F}_{T_1}$. Set

$$T_i^A := I_A T_i + \infty \cdot I_{A^c}, \quad i = 1, 2.$$

In this case

$$\mathbb{E}[X_S] = \mathbb{E}[M_{S \wedge T_2^A} - M_{S \wedge T_1^A}] = 0,$$

where the last equality follows from the optional sampling theorem. $\square$

The stopping and sampling theorems are very versatile tools for computing expectations. The next example illustrates this aspect.

Example 3.5.23. Suppose that $(B_t)_{t \geq 0}$ is a Brownian motion started at 0 and $(\mathcal{F}_t)_{t \geq 0}$ is its canonical filtration. For any $a \in \mathbb{R}$ we set

$$T_a := \inf \{t \geq 0 : B_t = a\}.$$

According to Proposition 3.4.4(ii), $\mathbb{P}[T_a < \infty] = 1$.

(a) We want to show that if $u < 0 < v$, then

$$\mathbb{P}[T_u < T_v] = \frac{v}{v-u}, \quad \mathbb{P}[T_u > T_v] = \frac{-u}{v-u}. \quad (3.5.20)$$

Consider the stopping time $T = T_{u,v} = T_u \wedge T_v$. In other words,

$$T_{u,v} = \inf \{t \geq 0, B_t \in \{u, v\}\}.$$

The stopped martingale $M_t = B_{T \wedge t}$ is UI since $|M_t| \leq \max(|u|, |v|)$. We deduce

$$0 = \mathbb{E}[M_0] = \mathbb{E}[M_\infty] = \mathbb{E}[B_T] = u\mathbb{P}[T_u < T_v] + v\mathbb{P}[T_v < T_u].$$

The equalities (3.5.20) follow by observing that the probabilities $\mathbb{P}[T_u < T_v]$ and $\mathbb{P}[T_u > T_v]$ satisfy a second linear constraint

$$\mathbb{P}[T_u < T_v] + \mathbb{P}[T_u > T_v] = 1.$$

Next, we want to prove that

$$\mathbb{E}[T_{u,v}] = |uv| \quad (3.5.21)$$

To see this consider the martingale of Example 3.5.1(ii), $X_t = B_t^2 - t$. The stopped process $X_{t \wedge T_{u,v}}$ is still a UI martingale and we have

$$\mathbb{E}[X_{t \wedge T_{u,v}}] = \mathbb{E}[X_0] = 0 \quad \text{and} \quad \mathbb{E}[B_{t \wedge T_{u,v}}^2] = \mathbb{E}[t \wedge T_{u,v}].$$

The Monotone Convergence Theorem implies that

$$\lim_{t \rightarrow \infty} \mathbb{E}[t \wedge T_{u,v}] = \mathbb{E}[T_{u,v}].$$

We deduce that

$$\begin{aligned} \mathbb{E}[T_{u,v}] &= \lim_{t \rightarrow \infty} \mathbb{E}[t \wedge T_{u,v}] = \lim_{t \rightarrow \infty} \mathbb{E}[B_{t \wedge T_{u,v}}^2] = \mathbb{E}[B_{T_{u,v}}^2] \\ &= u^2 \mathbb{P}[T_u < T_v] + v^2 \mathbb{P}[T_v < T_u] \\ &\stackrel{(3.5.20)}{=} \frac{u^2 v}{v - u} - \frac{v^2 u}{v - u} = -uv = |uv|. \end{aligned}$$

Observe that the distribution of the random variable $B(T_{u,v})$ is

$$D_{u,v} = \frac{v}{v - u} \delta_u - \frac{u}{v - u} \delta_v.$$

This is the dipole discussed in Example 1.4.23.

(b) Fix $a > 0$. We want to compute the Laplace transform of T_a . To this aim, we consider for any $\lambda \in \mathbb{R}$ the martingale of Example 3.5.1(iii)

$$X_t^\lambda := \exp\left(\lambda B_t - \frac{\lambda^2 t}{2}\right). \quad (3.5.22)$$

For $\lambda > 0$ the stopped martingale $Y_y^\lambda = X_{t \wedge T_a}^\lambda$ is bounded thus UI and we deduce

$$1 = \mathbb{E}[Y_0^\lambda] = \mathbb{E}[Y_\infty^\lambda] = e^{\lambda a} \mathbb{E}[e^{-\frac{\lambda^2 T_a}{2}}]$$

Replacing λ with $\sqrt{2\lambda}$ we deduce

$$\mathbb{E}[e^{-\lambda T_a}] = e^{-a\sqrt{2\lambda}}.$$

This can be alternatively verified using the distribution of T_a computed in Corollary 3.4.13.

(d) We want to compute the Laplace transform of $T = T_{-a,a}$. Consider the stopped martingale $Z_t^\lambda := X_{t \wedge T}^\lambda$, where X_t^λ is defined as in (3.5.22). We deduce as above that

$$1 = \mathbb{E}[e^{\lambda B_T} e^{-\lambda^2 T/2}].$$

The computations in (a) show that

$$\mathbb{P}[B_T = a] = \mathbb{P}[B_T = -a] = \frac{1}{2}.$$

Note that

$$\mathbb{P}[T \leq t] = \mathbb{P}[B_T = a, T \leq t] + \mathbb{P}[B_T = -a, T \leq t].$$

Using the symmetry $B_t \mapsto -B_t$ we deduce

$$\begin{aligned} \mathbb{P}[B_T = a, T \leq t] &= \mathbb{P}[B_T = -a, T \leq t] = \frac{1}{2} \mathbb{P}[T \leq t] \\ &= \mathbb{P}[B_T = a] \mathbb{P}[T \leq t] = \mathbb{P}[B_T = -a] \mathbb{P}[T \leq t], \end{aligned}$$

proving that B_T and T are independent. Hence

$$1 = \mathbb{E}[e^{\lambda B_T}] \mathbb{E}[e^{-\lambda^2 T/2}] = \cosh(\lambda a) \mathbb{E}[e^{-\lambda^2 T/2}]. \quad \square$$

3.6. Some applications

Let us illustrate the strength of the techniques discussed above on some classical examples.

3.6.1. The invariance theorem and applications. Recall that we constructed the Brownian motion as a limit of suitably rescaled random walks. Given a sequence of independent Rademacher random variables (X_n) we constructed a sequence of zig-zags $W_\nu(t)$. They are continuous functions, linear on each of the intervals $[(k-1)/\nu, k/\nu]$ and such that

$$W_\nu(k/\nu) = \nu^{-1/2} S_k, \quad S_k = \sum_{j=1}^k X_j.$$

More explicitly, $\forall t \in [k/\nu, (k+1)/\nu)$,

$$\begin{aligned} W_\nu(t) &= \nu^{-1/2} (S_k + (\nu t - k) X_{k+1}) \\ &= \nu^{1/2} \left(\left(t - \frac{k}{\nu}\right) S_{k+1} + \left(\frac{k+1}{\nu} - t\right) S_k \right) \\ &= \nu^{-1/2} (S_{\lfloor \nu t \rfloor} + (\nu t - \lfloor \nu t \rfloor) X_{\lfloor \nu t \rfloor + 1}). \end{aligned} \tag{3.6.1}$$

that converge in distribution to the Brownian motion. The invariance theorem shows that this is the manifestation of a more general universality phenomenon, namely, in the above example we can assume only that $(X_n)_{n \geq 1}$ is a sequence of i.i.d random variables with mean zero and variance 1. However, in the proof I will rely essentially the fact that Brownian motion exists and has certain properties.

Theorem 3.6.1 (Skhorokhod Representation). *Let X be a real valued random variable with mean zero and variance one. Then there exist*

- a complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$,
- a Brownian motion B defined on this space,
- a B -admissible filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ of $\mathcal{S}$ (see Definition 3.4.6)
- and a sequence stopping times $(T_n)_{n \geq 1}$ adapted to $\mathcal{F}_\bullet$.

such that

- (i) $\mathbb{E}[T] < \infty$.
- (ii) $0 = T_0 = T_1 \leq T_2 \leq \dots \leq T_n \leq \dots$ and,
- (iii) the random variables $(T_n - T_{n-1})$ are i.i.d. with mean $\mathbb{E}[T_n - T_{n-1}] = 1$,
- (iv) the random variables $X_n = B_{T_n} - B_{T_{n-1}}$, $n \geq 1$ are independent have the the same distribution as X .

Proof. Clearly, given the strong Markov property of the Brownian motion (Theorem 3.4.8) it suffices only to construct a B -admissible filtration $\mathcal{F}_\bullet$ and a $\mathcal{F}_\bullet$ -stopping time T such that $\mathbb{E}[T] < \infty$ and B_T has the same distribution as X .

Let $\mu \in \text{Prob}(\mathbb{R})$ be the distribution of X . As discussed in Example 1.4.23 μ is a mixtures of dipoles. These are probability measures on $\mathbb{R}$ whose supports consist of at most two point and the origin is their center of mass.

Following Example 1.4.23, we denote by Q the quadrant

$$Q := (-\infty, 0] \times (0, \infty).$$

To μ we can associate a Borel probability measure m_μ on $\mathbb{R}^2$ defined by (1.4.31).

There exist a Brownian motion and random variables U_n, V_n defined on the same complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that B is independent of U_n, V_n and for any n the distribution of the random vector (U_n, V_n) is $\mathbf{m}_\mu$. We get B -admissible filtration $\mathcal{F}_\bullet$ defined by

$$\mathcal{F}_t = \sigma(B_s, s \leq t) \vee \sigma(\mathcal{N}) \vee \sigma(U_n, V_n, n \in \mathbb{N})$$

where $\mathcal{N} \subset \mathcal{S}$ denote the collection of $\mathbb{P}$ -negligible events. Set $(U, V) := (U_1, V_1)$.

As shown in Example 3.5.23 (a), for any $u \leq 0 < v$ we have a stopping time

$$T_{u,v} = \inf \{ t \geq 0, B_t \in \{u, v\} \}.$$

Moreover,

$$\mathbb{E}[T_{u,v}] = |uv|.$$

The distribution of the random variable $B_{T_{u,v}}$ is the dipole $\mathbf{D}_{u,v}$ supported on the set $\{u, v\}$.

The random variable $T = T_{U,V}$ is a.s. finite and it is adapted to $\mathcal{F}_\bullet$. Moreover

$$\mathbb{E}[T_{U,V}] = \int_{\mathbb{R}^2} |uv| \mathbf{m}_\mu[dudv] \stackrel{(1.4.32)}{=} 1.$$

The random variable B_T has the same distribution as X since, for any Borel set $S \subset \mathbb{R}$,

$$\begin{aligned} \mathbb{P}_{B_T}[S] &= \int_{\mathbb{R}^2} \mathbb{P}[B_{T_{U,V}} \in S | U = u, V = v] \mathbb{P}_{U,V}[dudv] \\ &= \int_{\mathbb{R}^2} \mathbb{P}[B_{T_{u,v}} \in S] \mathbf{m}_\mu[dudv] = \int_{\mathbb{R}^2} \mathbf{D}_{u,v}[S] \mathbf{m}_\mu[dudv] \stackrel{(1.4.34)}{=} \mu[S]. \end{aligned}$$

Define

$$T_n = T_{U_1, V_1} + \cdots + T_{U_n, V_n}.$$

□

We now have all the tools we need to prove the promised invariance theorem. First we need a brief foundational digression.

Denote by $\mathbf{F}$ the Banach space $C([0, 1])$ equipped with the sup-norm. For each $t \in \mathbb{R}$ we have an *evaluation map*

$$\mathbf{Ev}_t : \mathbf{F} \rightarrow \mathbb{R}, \quad \mathbf{Ev}_t[f] = f(t).$$

The Borel sigma-algebra of $\mathbf{F}$ is generated by these evaluation maps; see Exercise 1.4. Suppose that

$$X : [0, 1] \times \Omega \rightarrow \mathbb{R}, \quad (t, \omega) \mapsto X_t(\omega)$$

is a continuous stochastic process. It defines a tautological map

$$X_\square : \Omega \rightarrow \mathbf{F}, \quad \omega \mapsto X_\square(\omega) \in \mathbf{F}, \quad X_\square(\omega)(t) = X_t(\omega).$$

For each $t \in \Omega$ we have $\mathbf{Ev}_t \circ X_\square = X_t$. By definition X_t is measurable and we deduce that the map

$$X_\square : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbf{F}, \mathcal{B}_{\mathbf{F}})$$

is measurable. The pushforward of $\mathbb{P}$ via $X_\square$ is a Borel probability measure on $\mathbf{F}$ called the distribution of the process X and denoted by $\mathbb{P}_X$. The distribution of the Brownian motion B restricted to $[0, 1]$ is the Wiener measure on $\mathbf{F}$ and we denote it by $\mathbb{W}$.

Theorem 3.6.2 (Donsker's Invariance Theorem). *Let X be random variable with mean zero and variance 1. Fix a sequence of i.i.d. random variables with the same distribution as X defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. For $n \in \mathbb{N}$ we set*

$$S_n = \sum_{k=1}^n X_k$$

and we consider the continuous stochastic processes

$$W_\nu = W_\nu^X : [0, 1] \times \Omega \rightarrow \mathbb{R}$$

defined by (3.6.1). Let $\mathbb{P}_\nu = \mathbb{P}_{W_\nu} \in \text{Prob}(\mathbf{F})$ be the distribution of W_ν . Then $\mathbb{P}_\nu$ converge weakly to the Wiener measure $\mathbb{W}$ as $\nu \rightarrow \infty$.

Proof. I follow the elegant approach in [180, Sec.I.8]. We first invoke Skorokhod's Representation Theorem.

Choose a complete probability space, a Brownian motion $B(t)$ defined on it, a B -admissible filtration $(\mathcal{F}_t)_{t \geq 0}$ and $\mathcal{F}_\bullet$ -stopping times

$$0 = T_0 \leq T_1 \leq T_2 \leq \dots$$

such that, for any $n \in \mathbb{N}$, the random variable $B(T_n) - B(T_{n-1})$ has the same distribution as X . In particular, $B(T_n)$ has the same distribution as S_n .

Let $B_\nu(t) := \nu^{-1}B(\nu t)$. The processes $B_\nu(-)$ and $B(-)$ are different but they have the same distribution namely, Wiener's measure. Moreover the gaps

$$T_1 - T_0, T_2 - T_1, \dots,$$

form a sequence of i.i.d. random variables with mean 1.

The conclusion of the theorem is a consequence of the following key fact.

$$\forall \varepsilon > 0, \lim_{\nu \rightarrow \infty} \mathbb{P}[\|W_\nu - B_\nu\| > \varepsilon] = 0, \quad (3.6.2)$$

where $\|\cdot\|$ the sup-norm on $\mathbf{F}$. The above equality says that the random variable $\|W_\nu - B_\nu\|$ converges in probability to 0. Let us first prove that it implies that $\mathbb{P}_\nu$ converges weakly to the Wiener measure.

To do this we will rely on the Portmanteau Theorem 2.2.11, the implication (ii) $\Rightarrow$ (i). We have to show that, for any bounded uniformly continuous function $\Phi : \mathbf{F} \rightarrow \mathbb{R}$ we have

$$\lim_{\nu \rightarrow \infty} \int_{\mathbf{F}} \Phi(f) \mathbb{P}_\nu[df] = \int_{\mathbf{F}} \Phi(f) \mathbb{W}[df].$$

Note that for any $\nu \in \mathbb{N}$,

$$\int_{\mathbf{F}} \Phi(f) \mathbb{P}_\nu[df] = \mathbb{E}[\Phi(W_\nu)] \quad \text{and} \quad \int_{\mathbf{F}} \Phi(f) \mathbb{W}[df] = \mathbb{E}[\Phi(B_\nu)].$$

Since Φ is uniformly continuous, for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\|W_\nu - B_\nu\| \leq \delta \Rightarrow |\Phi(W_\nu) - \Phi(B_\nu)| < \varepsilon.$$

We have

$$\begin{aligned} |\mathbb{E}[\Phi(W_\nu)] - \mathbb{E}[\Phi(B_\nu)]| &= \mathbb{E}\left[\mathbf{I}_{\{\|W_\nu - B_\nu\| \leq \delta\}} |\Phi(W_\nu) - \Phi(B_\nu)|\right] \\ &\quad + \mathbb{E}\left[\mathbf{I}_{\{\|W_\nu - B_\nu\| > \delta\}} |\Phi(W_\nu) - \Phi(B_\nu)|\right] \end{aligned}$$

$$\leq \varepsilon + 2 \sup |\Phi| \cdot \mathbb{P}[\|W_\nu - B_\nu\| > \delta].$$

Letting $\nu \rightarrow \infty$ we deduce from (3.6.2) that

$$\forall \varepsilon > 0, \quad 0 \leq \limsup_{\nu \rightarrow \infty} |\mathbb{E}[\Phi(W_\nu)] - \mathbb{E}[\Phi(B_\nu)]| \leq \varepsilon.$$

This proves that $\mathbb{P}_\nu \Rightarrow \mathbb{W}$.

To prove (3.6.2) we need a technical fact, consequence of the continuity of the Brownian motion.

Lemma 3.6.3. *For any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that*

$$\mathbb{P}\left[\exists s, t \in [0, 2], \quad |s - t| \leq \delta, \quad |B(t) - B(s)| \geq \varepsilon\right] \leq \frac{\varepsilon}{2}.$$

Proof of Lemma 3.6.3. For $s, t, \delta \in (0, 2] \cap \mathbb{Q}$, we set

$$E_{s,t} = \{|B(s) - B(t)| \geq \varepsilon\}, \quad E_\delta = \bigcup_{|s-t| \leq \delta} E_{s,t}.$$

Note that if $\delta < \delta'$, then $E_\delta \subset E_{\delta'}$. Note that

$$\bigcap_{\delta > 0} E_\delta = \left\{ \omega; \quad \forall \delta, \quad \exists |s - t| \leq \delta, \quad |B(s, \omega) - B(t, \omega)| \geq \varepsilon \right\} = \emptyset$$

since the function $[0, 2] \ni t \mapsto B(t, \omega)$ is uniformly continuous for any ω . Hence, there exists $\delta \in \mathbb{Q} \cap (0, 2]$ such that $\mathbb{P}[E_\delta] < \frac{\varepsilon}{2}$. $\square$

Fix $\varepsilon > 0$ and $\delta(\varepsilon) > 0$ as in the above lemma. We can assume $\delta < 1$. Note that they depend only on the Wiener measure on $C([0, 2])$, i.e., the distribution of the Brownian motion restricted to $[0, 2]$.

The Strong Law of Large Numbers implies that

$$\frac{T_\nu}{\nu} \rightarrow 1, \quad \text{a.s..}$$

Set

$$Y_\nu^* := \sup_{k \leq n} |T_k - k|$$

Then

$$\lim_{\nu \rightarrow \infty} \frac{Y_\nu^*}{\nu} = 0, \quad \text{a.s..}$$

This is a consequence of the following elementary result whose proof we postpone.

Lemma 3.6.4. *Suppose that (a_n) is a sequence of real numbers such that $\frac{a_n}{n} \rightarrow 0$. Set*

$$b_n := \max_{1 \leq k \leq n} |a_k|.$$

Then $\frac{b_n}{n} \rightarrow 0$. $\square$

Hence $\frac{1}{\nu} Y_\nu^*$ also converges to 0 in probability. We deduce that there exists $N = N(\varepsilon) > 0$ such that

$$\forall \nu \geq N, \quad \mathbb{P}\left[\frac{1}{\nu} \sup_{k \leq \nu} |T_k - k| > \frac{\delta}{3}\right] \leq \frac{\varepsilon}{2}.$$

Let $\nu \geq N(\varepsilon)$. Note that $W_\nu(k/\nu) = \nu^{-1} S_k = \nu^{-1} B(T_k)$.

For any $t \in [(k-1)/\nu, k/\nu]$, the $W_\nu(t)$ is between $\nu^{-1/2}B(T_{k-1})$ and $\nu^{-1/2}B(T_k)$. The intermediate value property of continuous functions shows that there exists $\tau_k \in [T_{k-1}, T_k]$ such that

$$\nu^{-1/2}B(\tau_k) = W_\nu(t).$$

Set $u_{k,\nu} := \frac{\tau_k}{\nu} \in [T_{k-1}/\nu, T_k/\nu]$, so that $B_\nu(u_{k,\nu}) = W_\nu(t)$. Note that

$$\begin{aligned} \mathbb{P}[\|W_\nu - B_\nu\| > \varepsilon] &\leq \mathbb{P}[\|W_\nu - B_\nu\| > \varepsilon, \sup_{k \leq \nu} |T_k/\nu - k/\nu| \leq \frac{\delta}{3}] \\ &\quad + \mathbb{P}\left[\frac{1}{\nu} \sup_{k \leq \nu} |T_k - k| > \frac{\delta}{3}\right]. \end{aligned}$$

Now let us observe that if $\nu > \frac{3}{\delta}$, then

$$\begin{aligned} &\underbrace{\|W_\nu - B_\nu\| > \varepsilon, \frac{1}{\nu} \sup_{k \leq \nu} |T_k - k| \leq \frac{\delta}{3}}_{=: E_0(\varepsilon)} \\ \Rightarrow &\underbrace{\exists s, t \in [0, 2], |s - t| \leq \delta, |B_\nu(t) - B_\nu(s)| \geq \varepsilon}_{=: E_1(\varepsilon)}. \end{aligned}$$

Indeed, if $E_0(\varepsilon)$ holds, then there exists $k \in \llbracket 1, \nu \rrbracket$ and $t \in [(k-1)/\nu, k/\nu]$ such that $|W_\nu(t) - B_\nu(t)| > \varepsilon$.

On the other hand, there exists $u_{k,\nu} \in [T_{k-1}/\nu, T_k/\nu]$ such that

$$W_\nu(t) = B_\nu(u_{k,\nu}).$$

Note that

$$[T_{k-1}/\nu, T_k/\nu] \subset \left[\left((k-1)/\nu - \delta/3 \right)_+, k/\nu + \delta/3 \right] \subset [0, 1 + \delta/3].$$

Thus both t and $u_{k,\nu}$ are located in the same interval of length at most δ contained in $[0, 2]$. By construction

$$|B_\nu(u_{k,\nu}) - B_\nu(t)| = |W_\nu(t) - B_\nu(t)| > \varepsilon$$

so that $E_1(\varepsilon)$ holds.

We deduce that if $n \geq \max(N(\varepsilon), 3/\delta)$, then

$$\begin{aligned} \mathbb{P}[\|W_\nu - B_\nu\| > \varepsilon] &\leq \mathbb{P}[E_0(\varepsilon)] + \mathbb{P}\left[\frac{1}{\nu} \sup_{k \leq \nu} |T_k - k| > \frac{\delta}{3}\right] \\ &\leq \mathbb{P}[E_1(\varepsilon)] + \frac{\varepsilon}{2} < \varepsilon. \end{aligned}$$

□

Proof of Lemma 3.6.4. Clearly $\frac{|a_n|}{n} \rightarrow 0$. Hence, there exists $N_\varepsilon > 0$ such that, for any $n \geq N_\varepsilon$ we have $|a_n| \leq n\varepsilon$. We deduce that for $n \geq N_\varepsilon$ we have $|b_n| \leq \max(n\varepsilon, b_{N_\varepsilon})$ so that

$$\forall n \geq N_\varepsilon, \quad 0 \leq \frac{b_n}{n} \leq \max\left(\varepsilon, \frac{b_{N_\varepsilon}}{n}\right).$$

Letting $n \rightarrow \infty$ we deduce that

$$\forall \varepsilon > 0, \quad 0 \leq \liminf_{n \rightarrow \infty} \frac{b_n}{n} \leq \limsup_{n \rightarrow \infty} \frac{b_n}{n} \leq \varepsilon.$$

□

3.6.2. The Kolmogorov-Smirnov statistics. We conclude with an application of the invariance theorem to the topic of uniform laws of large numbers discussed in Section 2.4. We recall some of the concepts introduced there.

Suppose we are given a sequence of i.i.d. random variables with common cdf $F(x)$.

The empirical cdf-s F_n are the random variables $F_n(x)$ defined by

$$F_n(x) = \frac{1}{n} \sum_{k=1}^n \mathbf{I}_{(-\infty, x]}(X_k) = \frac{1}{n} \times \# \{ 1 \leq k \leq n; X_k \leq x \}.$$

We define the random variables

$$D_n^F := \sup_{x \in \mathbb{R}} |F_n(x) - F(x)|.$$

Let $U \in \text{Unif}([0, 1])$. In Lemma 2.4.1 we proved that $D_n^F \leq D_n := D_n^{F_U}$ with equality if the cdf F is continuous. The Glivenko-Cantelli Theorem 2.4.2 shows that $D_n^F \rightarrow 0$ a.s.

Let (U_n) be a sequence of i.i.d. random variables uniformly distributed on $[0, 1]$. Denote by $G_n(t)$ the associated sequence of empirical cdf-s

$$G_n(t) = \begin{cases} 0, & t < 0, \\ \frac{1}{n} \# \{ 1 \leq k \leq n; U_k \in [0, t] \}, & t \in [0, 1], \\ 1, & t > 1. \end{cases}$$

Note that for any $t \in [0, 1]$ and any $n \in \mathbb{N}$ the random variable $nG_n(t)$ is a sum of independent binomial random variables $\mathbf{I}_{[0, t]}(U) \in \text{Bin}(t)$ so that

$$\mathbb{E}[nG_n(t)] = nt, \quad \text{Var}[nG_n(t)] = nt(1-t).$$

For any $t \in [0, 1]$ we have $F_U(t) = t$ and the Strong Law of Large Numbers shows that

$$\lim_{n \rightarrow \infty} (G_n(t) - t), \quad \text{a.s.}$$

while the Central Limit Theorem implies that $\sqrt{n}(G_n(t) - t)$ converge in distribution to normal random variable with mean zero and variance $t(1-t)$.

Fix a Brownian motion $(B(t))_{t \geq 0}$ and denote by $B^*(t)$ the associated *Brownian bridge*

$$B^*(t) := B(t) - tB(1), \quad t \in [0, 1]$$

The Brownian bridge is a continuous Gaussian process. Note that $B^*(0) = B^*(1) = 0$ and, for any $t \in [0, 1]$, $B^*(t)$ is also a normal random variable with mean zero and variance $t(1-t)$. Hence the random variables $\sqrt{n}(G_n(t) - t)$ converge in distribution to $B^*(t)$.

Theorem 3.6.5. *If the cdf F is continuous, then the random variables $\sqrt{n}D_n^F$ converge in distribution to the Kolmogorov-Smirnov random variable*

$$M_\infty := \sup_{t \in [0, 1]} |B^*(t)|. \quad (3.6.3)$$

Proof. We follow the approach in [26, Sec. 13.6]. I plan to use Donsker's invariance theorem to reach the desired conclusion. To be able to use it we will make some very ingenious cosmetic changes to D_n^F .

Since F is continuous, for any $t \in [0, 1]$ we have

$$D_n^F = D_n := \sup_{t \in \mathbb{R}} |G_n(t) - F_U(t)| = \sup_{t \in [0, 1]} |G_n(t) - t|.$$

Let $(U_n)_{n \in \mathbb{N}}$ be a sequence of i.i.d. random variables uniformly distributed on $[0, 1]$. For any $n \in \mathbb{N}$ we denote by $(U_{(1)}, \dots, U_{(n)})$ the order statistics of $(U_1, \dots, U_n)$; see Exercise 1.57. Recall that for n real numbers $x_1, x_2, \dots, x_n$ we set

$$x_{(1)} = \min\{x_1, \dots, x_n\}, \quad x_{(i+1)} = \min(\{x_1, \dots, x_n\} \setminus \{x_{(1)}, \dots, x_{(i)}\}).$$

In other words, the random variables $U_{(1)}, \dots, U_{(n)}$ are obtained by rearranging $U_1, \dots, U_n$ in increasing order. Note that

$$G_n(t) = \frac{k(t)}{n}, \quad k(t) = \max\{1 \leq k \leq n; U_{(k)} \leq t\},$$

so

$$G_n(t) \in \{k/n; k \in \llbracket 0, n \rrbracket\}.$$

We have

$$\left| (G_n(t) - t) - \left(\frac{k(t)}{n} - U_{k(t)} \right) \right| \leq \frac{2}{n}.$$

Note that

$$\sup_{t \in [0, 1]} \left| \frac{k(t)}{n} - U_{k(t)} \right| = \sup_{1 \leq k \leq n} \left| \frac{k}{n} - U_{(k)} \right|.$$

We deduce that

$$\sup_{t \in (0, 1)} |G_n(t) - t| = \sup_{1 \leq k \leq n} \left| \frac{k}{n} - U_{(k)} \right| + O(1/n), \quad (3.6.4)$$

where the constant implied by the above O -symbol is *deterministic*.

Fix independent exponential variables $T_n \in \text{Exp}(1)$, $n \in \mathbb{N}$. Set

$$Z_n := T_1 + \dots + T_n \in \text{Gamma}(n),$$

$$X_n := T_n - 1, \quad S_n = X_1 + \dots + X_n = Z_n - n.$$

The variables X_n are i.i.d. with mean 0 and variance 1. At this point we want to invoke a fortunate accident.

Lemma 3.6.6. *The random vectors*

$$(U_{(1)}, \dots, U_{(n)})$$

and

$$(Z_1/Z_{n+1}, \dots, Z_n/Z_{n+1})$$

have the same distribution. □

Taking for granted the assertion of Lemma 3.6.6 we have

$$\begin{aligned} \sqrt{n}D_n &= \sqrt{n} \sup_{1 \leq k \leq n} \left| \frac{k}{n} - \frac{Z_k}{Z_{k+1}} \right| + O(n^{-1/2}) \\ &= \frac{n}{Z_{n+1}} \max_{1 \leq k \leq n} \left| \frac{Z_k}{\sqrt{n}} - \frac{k}{n} \frac{Z_{n+1}}{\sqrt{n}} \right| + O(n^{-1/2}) \\ &= \frac{n}{Z_{n+1}} \max_{1 \leq k \leq n} \left| \frac{Z_k - k}{\sqrt{n}} - \frac{k}{n} \frac{Z_{n+1} - n}{\sqrt{n}} \right| + O(n^{-1/2}) \end{aligned}$$

$$= \frac{n}{Z_{n+1}} \times \max_{1 \leq k \leq n} \underbrace{\left| \frac{S_k}{\sqrt{n}} - \frac{k}{n} \frac{S_{n+1} + 1}{\sqrt{n}} \right|}_{=: V_n} + O(n^{-1/2}).$$

Denote by $W_n(t)$ the continuous piecewise linear function defined as in (3.6.1) using the random variables (X_n) . Thus, $W_n(t)$ is linear on each of the intervals $[(k-1)/n, k/n]$ and

$$W_n(k/n) = n^{-1/2} S_k,$$

Observe that

$$V_n = \max_{t \in [0,1]} \left| W_n(t) - tW_n(1) - t \frac{X_{n+1} + 1}{\sqrt{n}} \right|.$$

Indeed, the continuous function

$$\begin{aligned} t \mapsto \Xi_n(t) &= W_n(t) - tW_n(1) - t \frac{X_{n+1} + 1}{\sqrt{n}} \\ &= W_n(t) - \frac{t}{\sqrt{n}} S_n - t \frac{X_{n+1} + 1}{\sqrt{n}} \end{aligned}$$

is linear on each of the intervals $[(k-1)/n, k/n]$ and thus

$$\sup_{t \in [0,1]} |\Xi_n(t)| = \sup_{1 \leq k \leq n} |\Xi_n(k/n)|$$

and

$$\Xi_n(k/n) = \frac{S_k}{\sqrt{n}} - \frac{k}{n} \frac{S_{n+1} + 1}{\sqrt{n}}.$$

Donsker's invariance theorem shows that the sequence $W_n(t) \in \mathbf{F}$ converges in distribution to the Brownian motion restricted to $[0, 1]$. Consider the continuous linear map

$$L : \mathbf{F} \rightarrow \mathbf{F}, \quad L[f](t) = f(t) - tf(1).$$

The Mapping Theorem 2.2.3 shows that

$$L[W_n] = W_n(t) - tW_n(1)$$

converges in distribution to $L[B] = B^*$ = the Brownian bridge.

The sequence of independent random variables

$$\sup_{t \in [0,1]} \left| t \frac{X_{n+1} + 1}{\sqrt{n}} \right|$$

converges in probability to 0. Slutsky's Theorem 2.2.13 shows that the sequences of $\mathbf{F}$ -valued random variables $\Xi_n(t)$ and $W_n(t) - tW_n(1)$ converge in distribution to the same random function, the Brownian bridge M_∞ .

Consider the continuous function $\Psi : \mathbf{F} \rightarrow \mathbb{R}$,

$$\Psi(f) = \|f\| = \sup_{t \in [0,1]} |f(t)|.$$

Then $V_n = \Psi(\Xi_n)$ so

$$V_n \Rightarrow \Psi(B^*) = M_\infty.$$

We have

$$\sqrt{n}D_n = \frac{n}{Z_{n+1}} V_n + O(n^{-1/2}).$$

The Strong Law of Large Numbers shows that $\frac{n}{Z_{n+1}}$ converge a.s. to 1. Hence $\frac{n}{Z_{n+1}} V_n$ converges in distribution to M_∞ . The constant implied by the above O -symbol is deterministic so $\sqrt{n}D_n - \frac{n}{Z_{n+1}} V_n$

converges a.s. to 0. Invoking Slutsky's Theorem one more time we deduce that $\sqrt{n}D_n$ converges in distribution to M_∞ . $\square$

Proof of Lemma 3.6.6. We first compute the distribution of the random vector

$$W = (Z_1/Z_{n+1}, \dots, Z_n/Z_{n+1}, Z_{n+1}).$$

Define $\mathcal{R} : \mathbb{R}_+^{n+1} \rightarrow \mathbb{R}_+^{n+1}$

$$\mathcal{R}(w_1, \dots, w_{n+1}) = (w_1 w_{n+1}, \dots, w_n w_{n+1}, w_{n+1}).$$

The Jacobian matrix of this map is

$$J_{\mathcal{R}}(w) = \begin{bmatrix} w_{n+1} & 0 & 0 & \cdots & 0 & w_1 \\ 0 & w_{n+1} & 0 & \cdots & 0 & w_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & w_{n+1} & w_n \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$$

Hence

$$\det J_{\mathcal{R}}(w) = w_{n+1}^n.$$

Note that

$$\mathcal{R}(W) = Z := (Z_1, \dots, Z_n, Z_{n+1})$$

We also have a random vector $T = (T_1, \dots, T_{n+1})$ and with i.i.d. components and

$$Z = L(T)$$

where L is the linear operator $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ described by the matrix

$$L = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & 1 \end{bmatrix}.$$

Thus

$$W = \Phi(T), \quad \Phi := \mathcal{R}^{-1} \circ L.$$

If we denote by p_W the probability density of W and by p_T the probability density of T we deduce from (1.2.28) that

$$p_W(w) = |\det J_{\Phi^{-1}}(w)| p_T(\Phi^{-1}w).$$

We have $\Phi^{-1} = L^{-1} \circ \mathcal{R}$, $\det L = 1$, and

$$\det J_{\Phi^{-1}}(w) = \det L^{-1} \det J_{\mathcal{R}}(w) = \det J_{\mathcal{R}}(w) = w_{n+1}^n.$$

Observing that

$$p_T(t_1, t_2, \dots, t_{n+1}) = e^{-(t_1 + t_2 + \cdots + t_{n+1})}$$

and

$$L^{-1} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ \vdots \\ z_{n+1} \end{bmatrix} = \begin{bmatrix} z_1 \\ -z_1 + z_2 \\ -z_2 + z_3 \\ \vdots \\ -z_n + z_{n+1} \end{bmatrix}$$

we deduce that

$$\begin{aligned} p_T(\Phi^{-1}(w)) &= w_{n+1}^n e^{-(w_1 w_{n+1} + w_2 w_{n+1} - w_1 w_{n+1} + \dots + w_{n+1} - w_n w_{n+1})} \\ &= w_{n+1}^n e^{-w_{n+1}}. \end{aligned}$$

Then

$$\begin{aligned} p_{W_1, \dots, W_n}(w_1, \dots, w_n) &= \mathbf{I}_{\{0 \leq w_1 \leq \dots \leq w_n \leq 1\}} \cdot \int_0^\infty w_{n+1}^n e^{-w_{n+1}} dw_{n+1} \\ &= \Gamma(n+1) \mathbf{I}_{\{0 \leq w_1 \leq \dots \leq w_n \leq 1\}} = n! \mathbf{I}_{\{0 \leq w_1 \leq \dots \leq w_n \leq 1\}}. \end{aligned}$$

This is precisely the probability density of the order statistics

$$(U_{(1)}, \dots, U_{(n)}).$$

This is what the elementary Exercise 1.57(i) asked you to do. $\square$

We conclude this subsection by finding an explicit description of the distribution of the Kolmogorov-Smirnov random variable M_∞ . This is based on an alternate description of the Brownian bridge.

Fix a Brownian motion $(B_t)_{t \geq 1}$ consider the associated Brownian bridge

$$(B_t^*)_{t \in [0,1]}, \quad B_t^* = B_t - tB_1.$$

For any $\varepsilon > 0$ consider the (conditioned) continuous stochastic process

$$B_t^\varepsilon = B_t \mid \{|B_1| \leq \varepsilon\}.$$

Its distribution is the unique Borel probability measure $\mathbb{P}^\varepsilon$ on $\mathbf{F} = C([0,1])$ uniquely determined by the equalities

$$\mathbb{P}[B_{t_k}^\varepsilon \leq b_k, \quad 1 \leq k \leq n] = \mathbb{P}[B_{t_k} \leq b_k, \quad 1 \leq k \mid |B_1| \leq \varepsilon],$$

$\forall t_1, \dots, t_n \in [0,1]$ and $\forall b_1, \dots, b_n \in \mathbb{R}$. Denote let $\mathbb{P}^* \in \text{Prob}(\mathbf{F})$ denote the distribution of the Brownian bridge.

Lemma 3.6.7. *The distribution $\mathbb{P}^\varepsilon$ converges weakly to $\mathbb{P}^*$ in $\text{Prob}(\mathbf{F})$.*

Proof. I follow the approach in the proof of [64, Prop.12.3.2]. Observe that

$$\forall t \in [0,1], \quad \mathbb{E}[B_t^* B_1] = 0.$$

For any $t_1, \dots, t_n \in [0,1]$, the random variables $B_{t_1}^*, \dots, B_{t_n}^*, B_1$ are jointly Gaussian and the components $(B_{t_1}^*, \dots, B_{t_n}^*)$ and B_1 are uncorrelated. Exercise 2.71(iii) implies that the Gaussian vector $(B_{t_1}^*, \dots, B_{t_n}^*)$ is independent of B_1 . In other words B_1 is independent of the Brownian bridge B^* .

Let $\Phi : \mathbf{F} \times \mathbb{R} \rightarrow \mathbf{F}$ denote the continuous linear map given

$$\Phi[f, b](t) = f(t) + bt, \quad t \in [0,1], \quad f \in \mathbf{F}, \quad b \in \mathbb{R}.$$

We have

$$B = \Phi[B^*, B_1].$$

Hence the Wiener measure $\mathbb{W}$, i.e., the distribution of B , can be described as a pushforward

$$\mathbb{W} = \Phi_\#(\mathbb{P}^* \otimes \mathbb{P}_{B_1}).$$

Let $\mathbb{Q}_\varepsilon$ be the conditional distribution of B_1 , given that $|B_1| \leq \varepsilon$. Let Z_ε be a random variable, independent of B^* , with distribution $\mathbb{Q}_\varepsilon$.

Lemma 3.6.8. *The processes B^ε and $X^\varepsilon = \Phi[B^*, Z_\varepsilon]$ have the same distributions.*

Proof. Let $n \in \mathbb{N}$, $t_1, \dots, t_n \in [0, 1]$

$$(B(t_1), \dots, B(t_n)) = \underbrace{(B^*(t_1), \dots, B(t_n)^*)}_U + \underbrace{B_1 \cdot (t_1, \dots, t_n)}_V.$$

The random vectors U and V are independent. Denote by $\mathbb{P}_U$ and respectively $\mathbb{P}_V$ their distributions. Then

$$\mathbb{P}_{B(t_1), \dots, B(t_n)} = \mathbb{P}_U * \mathbb{P}_V.$$

Set $\mathbb{P}_V^\varepsilon := \mathbb{P}_{V| \{|B_1| \leq \varepsilon\}}$. A direct computation shows that

$$\mathbb{P}_{B(t_1), \dots, B(t_n) | \{|B_1| \leq \varepsilon\}} = \mathbb{P}_{B^\varepsilon(t_1), \dots, B^\varepsilon(t_n)} = \mathbb{P}_U * \mathbb{P}_V^\varepsilon.$$

We have

$$(X^\varepsilon(t_1), \dots, X^\varepsilon(t_n)) = U + \underbrace{Z_\varepsilon \cdot (t_1, \dots, t_n)}_{W^\varepsilon}.$$

The random vectors U and W^ε are independent and we deduce that

$$\mathbb{P}_{X^\varepsilon(t_1), \dots, X^\varepsilon(t_n)} = \mathbb{P}_U * \mathbb{P}_{W^\varepsilon}.$$

Now observe that $\mathbb{P}_{W^\varepsilon} = \mathbb{P}_V^\varepsilon$ so that

$$\mathbb{P}_{X^\varepsilon(t_1), \dots, X^\varepsilon(t_n)} = \mathbb{P}_{B(t_1), \dots, B(t_n) | \{|B_1| \leq \varepsilon\}}.$$

□

For any bounded continuous function $\Xi : \mathbf{F} \rightarrow \mathbb{R}$ we have

$$\mathbb{E}[\Xi(B^\varepsilon)] = \mathbb{E}[\Xi \circ \Phi[B^*, Z_\varepsilon]].$$

On the other hand

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}[\Xi \circ \Phi[B^*, Z_\varepsilon]] = \mathbb{E}[\Xi \circ \Phi[B^*, 0]]$$

since $Z_\varepsilon \Rightarrow 0$ and $\Xi \circ \Phi$ is bounded and continuous. □

Remark 3.6.9. The above lemma implies that the stochastic process

$$t \mapsto B_t | \{B_1 = 0\}$$

has the same distribution as the Brownian bridge. More explicitly, for any $0 < t_1 < t_2 \cdots t_n \leq 1$ the Gaussian vectors $(B_{t_1}, \dots, B_{t_n}) | \{B_1 = 0\}$ and $(B_{t_1}^*, \dots, B_{t_n}^*)$ have the same distribution. □

Lemma 3.6.7 implies that for any $a > 0$ we have

$$\begin{aligned} \mathbb{P}[M_\infty \leq a] &\stackrel{(3.6.3)}{=} \mathbb{P}\left[\sup_{t \in [0,1]} |B_t^*| \leq a\right] \\ &= \lim_{\varepsilon \rightarrow 0} \mathbb{P}\left[\sup_{t \in [0,1]} |B_t| \leq a \mid |B_1| \leq \varepsilon\right]. \end{aligned}$$

Corollary 3.4.18 shows that

$$\mathbb{P}\left[\sup_{0 \leq t \leq 1} |B(t)| \leq a, \quad |B(1)| < \varepsilon\right] = \underbrace{\int_{-\varepsilon}^{\varepsilon} \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} (-1)^k e^{-2a^2(x-k)^2} dx}_{:= F_a(x)}.$$

On the other hand,

$$\mathbb{P}[|B_1| \leq \varepsilon] = \int_{-\varepsilon}^{\varepsilon} \gamma_1(x) dx, \quad \gamma_1(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}.$$

We conclude that

$$\mathbb{P}[M_\infty \leq a] = \frac{F_a(0)}{\gamma_1(0)} = \sum_{k \in \mathbb{Z}} (-1)^k e^{-2a^2 k^2} = 1 + 2 \sum_{k \geq 1} (-1)^k e^{-2a^2 k^2}.$$

Putting together the results obtained so far we obtain the following result.

Theorem 3.6.10 (Kolmogorov-Smirnov). *For any $a \geq 0$ we have*

$$\lim_{n \rightarrow \infty} \mathbb{P}[\sqrt{n} D_n \leq a] = 1 + 2 \sum_{k \geq 1} (-1)^k e^{-2a^2 k^2}.$$

□

Remark 3.6.11. The Kolmogorov-Smirnov theorem has important applications in statistics. It is used to test whether collection of data is distributed according to a given continuous cdf. One first application of was given by Kolmogorov in 1940.

Stalin supported Lysenko's school, results, allegedly refuted Mendel's laws of genetics. In a paper unpublished for political reasons Kolmogorov showed that Lysenko's conclusion was based on dubious data; see [5] for more details about this bit of history. □

3.7. Exercises

Exercise 3.1. Suppose that $(X_n)_{n \geq 0}$ is a sequence of integrable random variables and $(q_n)_{n \geq 1}$ is a sequence of nonzero real numbers such that, for any $n \in \mathbb{N}$

$$\mathbb{E}[X_n \mid \mathcal{F}_{n-1}] = q_n X_{n-1}, \quad \mathcal{F}_{n-1} := \sigma(X_0, \dots, X_{n-1}).$$

Define $Q_0 = 1$, $Q_n = q_1 \cdots q_n$, $\forall n \in \mathbb{N}$ and set $Y_n := \frac{1}{Q_n} X_n$. Prove that $(Y_n)_{n \geq 0}$ is a martingale with adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$. $\square$

Exercise 3.2. Suppose that $(X_n)_{n \geq 0}$ is a martingale with respect to a filtration $(\mathcal{F}_n)_{n \geq 0}$ such that $X_0 = 0$ and $\mathbb{E}[|X_n|^2] < \infty$, $\forall n$. Using the sequence of differences $D_n = X_n - X_{n-1}$, $n \geq 1$ we construct two new processes, the *optional quadratic variation*

$$Q_n = \sum_{k=1}^n D_k^2$$

and the *predictable quadratic variation*

$$V_n = \sum_{k=1}^n \mathbb{E}[D_k^2 \mid \mathcal{F}_{k-1}].$$

Prove that the processes

$$A_n = X_n^2 - Q_n \quad B_n = X_n^2 - V_n$$

are martingales with respect to the $(\mathcal{F}_n)_{n \geq 0}$. $\square$

Exercise 3.3 (S. Ulam). Let $x_1, \dots, x_r \in \mathbb{R}$. Fix a family $\{I_n, J_n; n \in \mathbb{N}\}$ of independent random variables such that I_n, J_n are uniformly distributed on $\{1, \dots, n-1\}$, $\forall n \geq 2$. Define inductively

$$X_n := \begin{cases} x_n, & n \leq r \\ X_{I_n} + X_{J_n}, & n > r, \end{cases}$$

and set

$$Y_n := \frac{1}{n(n+1)} \sum_{k=1}^n X_k.$$

Prove that the sequence $(Y_n)_{n \geq r}$ is a martingale with respect to the filtration

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n), \quad n \in \mathbb{N}. \quad \square$$

Exercise 3.4. Prove all the claims in Example 3.1.21. $\square$

Exercise 3.5 (Optional switching). Suppose that $\mathcal{F}_\bullet := (\mathcal{F}_n)_{n \geq 0}$ is a filtration of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and $(X_n)_{n \geq 0}$, $(Y_n)_{n \geq 0}$ are two $\mathcal{F}_\bullet$ -martingales. Let $T : \Omega \rightarrow \mathbb{N}_0 \cup \{\infty\}$ be a stopping time adapted to $\mathcal{F}_\bullet$. Suppose that $X_T = Y_T$. For $n \in \mathbb{N}_0$ define

$$Z_n : \Omega \rightarrow \mathbb{R}, \quad Z_n(\omega) = \begin{cases} X_n(\omega), & n \leq T(\omega), \\ Y_n(\omega), & n > T(\omega). \end{cases}$$

Prove that $(Z_n)_{n \geq 0}$ is a martingale adapted to $\mathcal{F}_\bullet$. $\square$

Exercise 3.6. Prove Lemma 3.1.34. $\square$

Exercise 3.7 (Dubins' inequality). Let $X_\bullet = (X_n)_{n \geq 0}$ be a nonnegative supermartingale adapted to the filtration $\mathcal{F}_\bullet$ of a probability space $(\Omega, \mathcal{S}, B)$. For $0 \leq a < b$ denote by $N_n([a, b], X)$ the number of upcrossings of $[a, b]$ by $X_\bullet$ up to time n ; see (3.2.1). Prove that for any $k = 1, 2, \dots, n$

$$\mathbb{P}[N_n(a, b, X) \geq k] \leq \left(\frac{a}{b}\right)^k \mathbb{E}[\min(1, X_0/a)].$$

□

Exercise 3.8. Prove Lemma 3.2.1 .

□

Exercise 3.9. Suppose that $X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P})$, $n \in \mathbb{N}$, is a uniformly integrable sequence of random variables that converges in law to the random variable X , $X_n \Rightarrow X$. Then $X \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{E}[X_n^\pm] &= \mathbb{E}[X^\pm], \quad \lim_{n \rightarrow \infty} \mathbb{E}[|X_n|] = \mathbb{E}[|X|], \\ \lim_{n \rightarrow \infty} \mathbb{E}[X_n] &= \mathbb{E}[X]. \end{aligned}$$

□

Exercise 3.10 (Pratt's Lemma). Let (X_n) , (Y_n) , (Z_n) be three sequences of integrable random variables with the following properties.

- (i) $X_n \leq Y_n \leq Z_n, \forall n$.
- (ii) $X_n \xrightarrow{p} X, Y_n \xrightarrow{p} Y, Z_n \xrightarrow{p} Z$.
- (iii) $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X], \mathbb{E}[Z_n] \rightarrow \mathbb{E}[Z]$.

Prove that $\mathbb{E}[Y_n] \rightarrow \mathbb{E}[Y]$.

□

Exercise 3.11. Suppose that $(X_n)_{n \geq 0}$ is a martingale defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that $\exists M > 0$,

$$\forall n \in \mathbb{N} \quad |X_n - X_{n-1}| \leq M, \quad \text{a.s..}$$

Define

$$\begin{aligned} A &:= \left\{ \omega \in \Omega; \lim_{n \rightarrow \infty} X_n(\omega) \text{ exists and is finite} \right\}, \\ B &:= \left\{ \omega \in \Omega; \liminf_{n \rightarrow \infty} X_n(\omega) = -\infty, \limsup_{n \rightarrow \infty} X_n(\omega) = \infty \right\}. \end{aligned}$$

Prove that $\mathbb{P}[A \cup B] = 1$. In other words, when a martingale (with bounded increments) does not have a limit, it oscillates wildly.

Hint. For $C > 0$ look at $T_C^\pm = \min \{n; \pm X_n > C\}$.

□

Exercise 3.12 (P. Lévy). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $(\mathcal{F}_n)_{n \geq 1}$ is a filtration of sigma-subalgebras. Let (F_n) be a sequence of events such that $F_n \in \mathcal{F}_n, \forall n$. We set

$$X_n = \sum_{k=1}^n (I_{F_k} - \mathbb{E}[I_{F_k} \mid \mathcal{F}_{k-1}]).$$

- (i) Prove that X_n is a martingale and $|X_n - X_{n-1}| \leq 1, \forall n$. **Hint.** Have a look at Example 3.1.14.
- (ii) Prove that

$$\{F_n \text{ i.o.}\} = \left\{ \sum_{n \geq 1} \mathbb{E}[I_{F_n} \mid \mathcal{F}_{n-1}] = \infty \right\}.$$

Hint. Use Exercise 3.11.

(iii) Deduce from (ii) the second Borel-Cantelli Lemma, Theorem 1.3.55(ii). □

Exercise 3.13. Consider the situation in Example 3.1.33. We have a finite set $\mathcal{A}$ called alphabet, a probability distribution π on $\mathcal{A}$ such that $\pi[a] \neq 0, \forall a \in \mathcal{A}$. Fix two words

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathcal{A}^k, \quad \mathbf{b} = (b_1, \dots, b_\ell) \in \mathcal{A}^\ell$$

and assume that $\mathbf{b}$ is not a subword of $\mathbf{a}$, i.e.,

$$(a_{i+1}, \dots, a_{i+\ell}) \neq (b_1, \dots, b_\ell), \quad \forall i = 0, \dots, k - \ell.$$

Let $(A_n)_{n \geq 1}$ be i.i.d. $\mathcal{A}$ valued random variable with common distribution π . As in Example 3.1.33 we denote by $T_{\mathbf{b}}$ the time to observe the pattern $\mathbf{b}$.

(i) Prove that

$$\mathbb{E}[T_{\mathbf{b}} | A_1 = a_1, \dots, A_k = a_k] - k = \Phi(\mathbf{b}, \mathbf{b}) - \Phi(\mathbf{a}, \mathbf{b})$$

where Φ is defined by (3.1.13).

(ii) Set $p_{\mathbf{a}} := \mathbb{P}[T_{\mathbf{a}} < T_{\mathbf{b}}]$, $p_{\mathbf{b}} := \mathbb{P}[T_{\mathbf{b}} < T_{\mathbf{a}}]$, $T = \min(T_{\mathbf{a}}, T_{\mathbf{b}})$. Prove that

$$p_{\mathbf{a}}\Phi(\mathbf{a}, \mathbf{a}) + p_{\mathbf{b}}\Phi(\mathbf{b}, \mathbf{a}) = \mathbb{E}[T] = p_{\mathbf{a}}\Phi(\mathbf{a}, \mathbf{b}) + p_{\mathbf{b}}\Phi(\mathbf{b}, \mathbf{b}).$$

(iii) Show that

$$\frac{p_{\mathbf{b}}}{p_{\mathbf{a}}} = \frac{\Phi(\mathbf{a}, \mathbf{a}) - \Phi(\mathbf{a}, \mathbf{b})}{\Phi(\mathbf{b}, \mathbf{b}) - \Phi(\mathbf{a}, \mathbf{b})}.$$

Hint. Consider the same martingale (X_n) as in Example 3.1.33. Observe that $X_k = \Phi(\mathbf{a}, \mathbf{b}) - k$ given that $A_j = a_j, j = 1, \dots, k$. (ii) Note that $\mathbb{E}[T_{\mathbf{b}}] = \mathbb{E}[T_{\mathbf{b}}] + \mathbb{E}[T_{\mathbf{b}} - T]$ and (i) gives a formula for $\mathbb{E}[T_{\mathbf{b}} - T | T = T_{\mathbf{a}}]$. □

Exercise 3.14. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability spaces and $\mathfrak{X} \subset \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$ a family of integrable random variables. Prove that the following are equivalent.

(i) The family $\mathfrak{X}$ is *UI*.

(ii) For any $\varepsilon > 0$ there exists $w_\varepsilon \in \mathcal{L}_+^1(\Omega, \mathcal{S}, \mathbb{P})$ such that

$$\sup_{X \in \mathfrak{X}} \mathbb{E}[|X| \mathbf{1}_{\{|X| > w_\varepsilon\}}] < \varepsilon.$$

□

Exercise 3.15. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a uniformly integrable sequence of random variables that converge in distribution to the random variable X . Prove that $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$.

Hint. Use Exercise 2.50. □

Exercise 3.16 (Kakutani). Let (X_n) be a sequence of independent positive random variables such that $\mathbb{E}[X_n] = 1$. Consider the product martingale

$$Y_n = \prod_{k=1}^n X_k.$$

Doob's convergence theorem shows that Y_n converges a.s. to a random variable Y_∞ satisfying $\mathbb{E}[Y_\infty] \leq 1$.

Set $a_n := \mathbb{E}[X_n^{1/2}]$. Prove that the following are equivalent.

(i) $\mathbb{E}[Y_\infty] = 1$.

- (ii) $Y_n \rightarrow Y_\infty$ in L^1 .
- (iii) The martingale $(Y_n)_{n \in \mathbb{N}}$ is UI.
- (iv) $\prod_n a_n > 0$.
- (v) $\sum_n (1 - a_n) > 0$.

Hint. The tricky implication is (iv) $\Rightarrow$ (iii). Define $Z_n = \prod_{i=1}^n (a_i^{-1} X_i^{1/2})$ and prove that it is an L^2 -bounded martingale. $\square$

Exercise 3.17. Consider the unbiased random walk in Example 3.1.5

$$S_0 = a \in \mathbb{Z}, \quad S_n = X_1 + \cdots + X_n, \quad n \geq 1,$$

where $(X_n)_{n \geq 1}$ are i.i.d. random variables such that $\mathbb{E}[X_n] = 0$, $\text{Var}[X_n] = 1$, $\forall n$. Set

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n), \quad n \in \mathbb{N}.$$

- (i) Show that the sequence $(S_n^2 - n)_{n \geq 0}$ is a martingale with respect to the filtration $\mathcal{F}_n$.
- (ii) Assume that $M(t) = \mathbb{E}[e^{tX_1}]$ exists for all $|t| < t_0$, $t_0 > 0$. For $|t| < t_0$ and $n \in \mathbb{N}$ we set

$$Z_n(t) := \frac{1}{M(t)^n} e^{tS_n}.$$

is a martingale with respect to the filtration $\mathcal{F}_n$.

- (iii) Set $D = \frac{d}{dx}$. We define $M(D) : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by the equality

$$\begin{aligned} M(D)[P](x) &= \sum_{k \geq 0} \frac{M^{(k)}(0)}{k!} D^k P(x) \\ &= \sum_{k \geq 0} \frac{\mu_k[X_1]}{k!} P^{(k)}(x), \quad \mu_k[X_1] = \mathbb{E}[X_1^k]. \end{aligned}$$

Prove that $M(D)$ is bijective and for any polynomial P the sequence

$$Y_n = M(D)^{-n} P(S_n), \quad n \geq 1,$$

is a martingale. Find Y_n when $P(x) = x$ and $P(x) = x^2$. **Hint.** Set $P_n := M(D)^{-n}[P]$ and express $\mathbb{E}[P_{n+1}(S_n + X_{n+1}) | X_1, \dots, X_n]$ using the operator $M(D)$.

- (iv) Compute $M(D)$ when (X_n) are independent Rademacher variables. Set $B_0(x) := 1$,

$$B_n(x) := \binom{x}{n} = \frac{x(x-1) \cdots (x-n+1)}{n!}, \quad n \in \mathbb{N}.$$

Express $M(D)^{-1}[B_n]$ in terms of the polynomials B_k , $k = 0, 1, \dots, n$. **Hint.** Consider the finite difference operator Δ that associates to any function $f : \mathbb{R} \rightarrow \mathbb{R}$ the new function Δf defined by $(\Delta f)(x) := f(x+1) - f(x)$. Observe that $\Delta B_n = B_{n-1}$, $\forall n \in \mathbb{N}$. $\square$

Exercise 3.18. Suppose that $(X_n)_{n \geq 0}$ is a martingale with respect to the filtration $\mathcal{F}_\bullet = (\mathcal{F}_n)_{n \geq 0}$ such that $\mathbb{E}[X_n^2] < \infty$, $\forall n$. The sequence $(X_n^2)_{n \geq 0}$ is a submartingale and thus, according to Proposition 3.1.13 it admits a Doob decomposition $X_n^2 = X_0 + M_n + C_n$, where $(M_n)_{n \geq 0}$ is a martingale and the compensator (C_n) is a predictable, nondecreasing process. Set

$$A_n = X_0 + C_n, \quad A_\infty = \lim_{n \rightarrow \infty} A_n = \sup_{n \in \mathbb{N}} A_n.$$

- (i) Prove that $\mathbb{E}[\sup_{n \geq 0} X_n] \leq 4\mathbb{E}[A_\infty]$. **Hint.** Use Doob's L^2 -maximal inequality.
- (ii) Prove that $\lim_{n \rightarrow \infty} X_n$ exists and is finite a.s. on the set $\{A_\infty < \infty\}$. **Hint.** For $a > 0$ we set $N_a = \min\{n; A_{n+1} > a^2\}$. Show that it is adapted to the filtration $\mathcal{F}_\bullet$. Apply (i) to the stopped martingale $X_{n \wedge N_a}$.
- (iii) Suppose that $f : [0, \infty) \rightarrow [1, \infty)$ is an increasing function such that

$$\int_0^\infty \frac{f(t)}{t^2} dt < \infty.$$

Prove that $\frac{X_n}{f(A_n)} \rightarrow 0$ a.s. on the set $\{A_\infty = \infty\}$. **Hint.** Set $H_n = \frac{1}{f(A_n)}$, $\forall n \in \mathbb{N}$. Let $Y_\bullet$ denote the martingale defined by the discrete stochastic integral $(H \cdot X)_\bullet$; see (3.1.3). Use the Doob decomposition of Y_n to prove that Y_n converges L^2 a.s. Conclude using Kronecker's lemma, Lemma 2.1.11.

□

Exercise 3.19 (Hunt). Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space. Suppose that $(\mathcal{F}_n)_{n \geq 0}$ a filtration of $\mathcal{S}$ and $(X_n \in L^1(\Omega, \mathcal{S}, \mathbb{P}))_{n \in \mathbb{N}}$ is a sequence of random variables dominated by an integrable random variable Y , i.e.,

$$\forall n, |X_n| \leq Y \text{ a.s.}$$

Prove that if $X_n \rightarrow X_\infty$ a.s., then

$$\mathbb{E}[X_n | \mathcal{F}_n] \rightarrow \mathbb{E}[X_\infty | \mathcal{F}_\infty] \text{ a.s.}$$

□

Exercise 3.20 (Dubins-Freedman). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $(\mathcal{F}_n)_{n \geq 1}$ is a filtration of sigma-subalgebras. Let (F_n) be a sequence of events such that $F_n \in \mathcal{F}_n$, $\forall n$. We set

$$X_n = \sum_{k=1}^n (I_{F_k} - f_k), \quad f_k := \mathbb{E}[I_{F_k} | \mathcal{F}_{k-1}].$$

- (i) Prove that $(X_n)_{n \geq 0}$ is a martingale and $\mathbb{E}[X_n^2] < \infty$, $\forall n \geq 0$.
- (ii) Define $S = \{\sum_n f_n = \infty\}$. Prove that

$$\frac{\sum_{k=0}^n I_{F_k}}{\sum_{k=0}^n f_k} \rightarrow 1, \text{ a.s. on } S. \quad (3.7.1)$$

- (iii) Deduce from (3.7.1) the conclusion of Exercise 3.12(ii). Thus (3.7.1) is a generalization of the second Borel-Cantelli lemma, Theorem 1.3.55(ii).

□

Exercise 3.21 (Conservation of fairness). A fair coin is flipped repeatedly and independently. A gambler starts with an initial fortune $f_0 > 0$. Before the n -th flip, his fortune is F_{n-1} . Based only on the information available to him at that moment, the gambler bets a sum $B_n \in (0, b)$, $0 \leq B_n \leq F_{n-1}$. If the n -th flip shows Heads he earns B_n dollars and if it shows Tails, he loses B_n dollars. The gambler stops gambling when he is broke or at the first moment when he reaches his goal, i.e., $F_n \geq g$ where $g > 0$ set in advance of his gambling.

- (i) Prove that the probability p_g that he reaches his goal is $\leq \frac{f_0}{g}$.
- (ii) Prove that if $B_n \leq \min(F_{n-1}, g - F_{n-1})$, $\forall n \geq 1$, then $p_g = \frac{f_0}{g}$.
- (iii) Find p_g if $B_n = \frac{1}{2}F_{n-1}$.

□

Remark 3.7.1. Note that if $f_0, g \in \mathbb{N}$ and the gambling strategy is $B_n = 1$ whenever his fortune is $< g$ the above problem reduces to the classical Gambler's ruin problem discussed in Example 3.2.37. The name “*conservation of fairness*” seems appropriate: whatever gambling strategy satisfying (ii) and based only on the information available at each moment, the probability of reaching the goal is the same, $\frac{f_0}{g}$. □

Exercise 3.22. Fix $a, g \in \mathbb{N}_0$, $a \leq g$. Consider the standard random walk $(S_n)_{n \geq 0}$ on $\mathbb{Z}$ started at a , i.e.,

$$S_0 = a, \quad S_n = X_1 + \cdots + X_n,$$

where $(X_n)_{n \geq 1}$ are i.i.d. with $\mathbb{P}[X_n = \pm 1] = \frac{1}{2}$. Set

$$T := \min \{ n \in \mathbb{N}_0; \ S_n = 0 \text{ or } S_n = g \}.$$

- (i) Show $\mathbb{P}[S_T = 0] = \frac{g-a}{g}$ and $\mathbb{P}[S_T = g] = \frac{a}{g}$. **Hint.** Use Theorem 3.2.35.
- (ii) Show that $\mathbb{E}[T] = a(g - a)$.
- (iii) Compute the pgf of T

$$f_a(s) := \mathbb{E}[s^T] = \sum_{n=0}^{\infty} \mathbb{P}[T = n] s^n.$$

Hint. Condition on X_1 . Alternatively, use the de Moivre martingale and the Optional Sampling Theorem 3.1.29.

□

Exercise 3.23. Suppose that $(S_n)_{n \geq 0}$ is the standard random walk on $\mathbb{Z}$. Thus

$$S_0 = 0, \quad S_n = X_1 + \cdots + X_n,$$

where $(X_n)_{n \in \mathbb{N}}$ are independent Rademacher random variable $\mathbb{P}[X_n = \pm 1] = \frac{1}{2}$, $\forall n$. Let $a \in \mathbb{Z}_{>0}$ and denote by T_a the first moment the random walk reaches the location a ,

$$T_a = \min \{ n \in \mathbb{N}; \ S_n = a \}.$$

- (i) Prove that

$$\mathbb{P}[T_0 > 2n] = \mathbb{P}[S_{2n} = 0] = \frac{1}{2^{2n}} \binom{2n}{n}$$

and deduce that $\mathbb{P}[T_0 = \infty] = 0$. **Hint.** Have a look at Example 1.2.38.

- (ii) Prove that $\mathbb{P}[T_a = \infty] = 0$.
- (iii) Compute the moment generating function of T_a .

□

Exercise 3.24. Suppose that $(X_n)_{n \geq 1}$ is a sequence of integrable random variables defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ such that for any $S \in \mathcal{S}$ the sequence $\mathbb{E}[X_N \mathbf{I}_S]$ has a finite limit.

- (i) Prove that

$$\sup_{S \in \mathcal{S}} \sup_{n \in \mathbb{N}} |\mathbb{E}[X_n \mathbf{I}_S]| < \infty.$$

and deduce that $\sup_{n \in \mathbb{N}} \mathbb{E}[|X_n|] < \infty$. **Hint.** Use the metric d in Exercise 2.13 and Baire's category theorem.

(ii) Prove that the sequence $(X_n)_{n \geq 0}$ is UI.

□

Exercise 3.25. Suppose that $(X_n)_{n \geq 0}$ adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ and T is a stopping time adapted to the same filtration such that $\mathbb{P}[T < \infty] = 1$ and $X_T \in L^1$. Prove that

$$\mathbb{E}[X_T \mid \mathcal{F}_n] = X_n \text{ on } \{T \geq n\}.$$

Hint. Have a look at the proof of Theorem 3.1.29.

Exercise 3.26. Suppose that $(X_n)_{n \geq 1}$ is a sequence of i.i.d., *nonnegative, integer valued* random variables with finite mean. Set

$$S_n := X_1 + \cdots + X_n.$$

For $k = 1 \dots, n$, set $\mathcal{F}_{-k} = \sigma(S_k, S_{k+1}, \dots, S_n)$, $Y_{-k} = S_k/k$.

(i) Prove that for $j \leq k$ we have

$$\mathbb{E}[X_j \mid \mathcal{F}_{-k}] = X_k.$$

(ii) Prove that $(Y_{-k})_{1 \leq k \leq n}$ is a martingale with respect to the filtration $(\mathcal{F}_{-k})_{1 \leq k \leq n}$. (Compare with Example 3.1.32.)

(iii) Show that

$$\mathbb{P}[S_k < k, \forall 1 \leq k \leq n \mid S_n] = (1 - S_n/n)^+.$$

Hint. (iii) Set $T = \inf \{-n \leq k \leq -1; Y_k \geq 1\}$, where we define $\inf \emptyset = -1$. Use Exercise 3.25.

□

Exercise 3.27. Suppose that $f : [0, 1] \rightarrow \mathbb{R}$ is a Lebesgue integrable function. For any $n \in \mathbb{N}_0$ we define the step function $f_n : [0, 1] \rightarrow \mathbb{R}$ by setting $f_n(0) = 0$ and

$$f_n(x) = \frac{1}{2^n} \int_{(k-1)/2^n}^{k/2^n} f(x) dx, \text{ if } 0 \leq \frac{(k-1)}{2^n} < x \leq \frac{k}{2^n} \leq 1.$$

Prove that f_n converges a.s. and L^1 to f as $n \rightarrow \infty$.

□

Exercise 3.28. Suppose that $(X_n)_{n \geq 0}$ is a supermartingale such that, there exist $f_0, g > 0$ with the property

$$X_0 = f_0 \text{ a.s., } 0 \leq X_n \leq g \text{ a.s., } \forall n \in \mathbb{N}.$$

Prove that for any stopping time T such that $\mathbb{P}[T < \infty] = 1$ we have $\mathbb{P}[X_T = g] \leq \frac{f_0}{g}$.

□

Exercise 3.29. Consider the branching process $(Z_n)_{n \geq 0}$ with initial condition $Z_0 = 1$ and reproduction law $\mu \in \text{Prob}(\mathbb{N}_0)$ such that

$$m := \mathbb{E}[\mu] = \sum_{n \geq 0} n \mu_n < \infty, \quad \mu_n := \mu[n].$$

Assume $\mu_0 > 0$. Denote by $f(s)$ the probability generating function (pgf) of μ

$$f(s) = \sum_{n \geq 0} \mu_n s^n = \mathbb{E}[s^{Z_1}].$$

We set

$$f_n(s) := \underbrace{f \circ \cdots \circ f}_n(s), \quad n \in \mathbb{N}.$$

- (i) Show that if $m > 1$ the equation $f(s) = s$ has a unique solution $r = r(\mu)$ in the interval $(0, 1)$. Compute $r(\mu)$ when

$$\mu_n = qp^n, \quad n \in \mathbb{N}_0,$$

where $p \in (1/2, 1)$, $q = 1 - p$.

- (ii) Prove that

$$\mathbb{E}[s^{Z_n}] = f_n(s), \quad \forall s \in [0, 1].$$

- (iii) Denote by E the extinction event

$$E = \bigcup_{n \geq 0} \{Z_n = 0\}.$$

Prove that

$$\mathbb{P}[E] = \lim_{n \rightarrow \infty} f_n(0) = \begin{cases} 1, & m \leq 1, \\ r(\mu), & m > 1. \end{cases}$$

- (iv) Assume $m > 1$. Prove that the sequence $(r^{Z_n})_{n \geq 0}$ is a martingale.

- (v) Set

$$W_n := \frac{1}{m^n} Z_n.$$

Assume

$$m > 1, \quad \mathbb{E}[Z_1^2] = \sum_{n \geq 1} n^2 \mu_n < \infty,$$

and set

$$W := \lim_{n \rightarrow \infty} W_n.$$

Denote by $\mathbb{P}_W$ the probability distribution of W and by $\varphi(\lambda)$ its Laplace transform

$$\varphi(\lambda) = \mathbb{E}[e^{-\lambda W}] = \int_{\mathbb{R}} e^{-\lambda w} \mathbb{P}_W(dw), \quad \lambda \in \mathbb{C}, \quad \operatorname{Re} \lambda \geq 0.$$

Prove that

$$\varphi'(0) = 1, \quad \varphi(\lambda) = f(\varphi(\lambda/m)) = \sum_{n=0}^{\infty} \mu_n \varphi(\lambda/m)^n, \quad \forall \operatorname{Re} \lambda \geq 0. \quad (3.7.2)$$

- (vi) Prove that there exists at most one probability measure $\nu \in \operatorname{Prob}([0, \infty))$ such that

$$\int_0^\infty t^2 \nu[dt] < \infty$$

and its Laplace transform

$$\varphi_\nu(\lambda) := \int_0^\infty e^{-\lambda t} \nu[dt], \quad \lambda \in \mathbb{C}, \quad \operatorname{Re} \lambda \geq 0,$$

satisfies (3.7.2).

Hint. Consider two such measures $\nu_k, k = 0, 1$, denote by $\Phi_k(t)$ their characteristic functions. Set $\Phi(t) = \Phi_1(t) - \Phi_0(t)$, $\gamma(t) = \Phi(t)/t, t \neq 0$. Prove that $|\gamma(mt)| \leq |\gamma(t)|$ and conclude that $\Phi \equiv 0$.

□

Exercise 3.30. Let $\mathfrak{S}_n$ denote the group of permutations of $\mathbb{I}_n := \{1, \dots, n\}$. We equip $\mathfrak{S}_n$ with the uniform probability measures. A *run* of a permutation π is a pair $(s, r) \in \mathbb{I}_n$, $s < r$ such that

$$\pi_{s-1} > \pi_s < \pi_{s+1} < \dots < \pi_r > \pi_{r+1},$$

where $\pi_0 := n+1$ and $\pi_{n+1} := 0$. We denote by $R_n(\pi)$ the number of runs of $\pi \in \mathfrak{S}_n$. Set

$$X_n := nR_n - \frac{1}{2}n(n+1).$$

(i) For $\pi \in \mathfrak{S}_{n+1}$ we set $k_\pi := \pi^{-1}(n+1)$ and denote by φ_π the unique increasing bijection

$$\varphi_\pi : \mathbb{I}_n \rightarrow \mathbb{I}_{n+1} \setminus \{k_\pi\}.$$

Set $\bar{\pi} := \pi \circ \varphi_\pi$. Show that the random maps

$$\mathfrak{S}_{n+1} \ni \pi \mapsto k_\pi \in \mathbb{I}_{n+1}, \quad \mathfrak{S}_{n+1} \ni \pi \mapsto \bar{\pi} \in \mathfrak{S}_n$$

are independent and uniformly distributed on their ranges.

(ii) Prove that (X_n) is a martingale.

(iii) Compute $\mathbb{E}[R_n]$ and $\mathbb{E}[R_n^2]$.

(iv) Show that

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\left(\frac{R_n}{n} - \frac{1}{2} \right)^2 \right] = 0.$$

□

Exercise 3.31. Suppose that $(X_n)_{n \geq 0}$ is an L^2 -martingale adapted to the filtration $(\mathcal{F}_n)_{n \geq 0}$ and $\langle X_\bullet \rangle$ is its quadratic variation; see Definition 3.1.15. Fix a bounded predictable process $(H_n)_{n \geq 0}$ and form the discrete stochastic integral $(H \bullet X)$; see Theorem 3.1.17.

(i) Show that

$$\mathbb{E}[X_n^2] - \mathbb{E}[X_0^2] = \mathbb{E}[\langle X \rangle_n].$$

(ii) Prove that the martingale $(H \bullet X)$ is an L^2 martingale.

(iii) Prove that

$$\langle H \bullet X \rangle_m = (H^2 \bullet \langle X \rangle)_n := \sum_{k=1}^n H_k^2 (\langle X \rangle_k - \langle X \rangle_{k-1}), \quad \forall n \geq 1.$$

(iv) Prove that

$$\mathbb{E}[(H \bullet X)_n^2] = \mathbb{E} \left[\sum_{k=1}^n H_k^2 (X_k - X_{k-1})^2 \right], \quad \forall n \geq 1.$$

Exercise 3.32. Suppose that $(X_n)_{n \in \mathbb{N}}$ is an exchangeable sequence of random variables and T is a stopping time adapted to the filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$. Prove that if $T < N$ a.s., then X_{T+1} has the same distribution as X_1 . □

Exercise 3.33. Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of random variables such that for any $n \in \mathbb{N}$ the distribution of the random vector $(X_1, \dots, X_n)$ is orthogonally invariant, i.e., for any $T \in O(n)$, $T_{\#} \mathbb{P}_{X_1, \dots, X_n} = \mathbb{P}_{X_1, \dots, X_n}$. Prove that $(X_n)_{n \in \mathbb{N}}$ are conditionally i.i.d. $N(0, \sigma^2)$ given a random variable $\sigma^2 \geq 0$. □

Exercise 3.34. Prove Lemma 3.3.8. □

Exercise 3.35. Finish the proof of Proposition 3.3.13. □

Exercise 3.36. Prove Proposition 3.3.15. □

Exercise 3.37 (Galmarino's test). Suppose that

$$X : (\Omega, \mathcal{S}, \mathbb{P}) \times [0, \infty) \rightarrow \mathbb{R}, \quad \omega, t \mapsto X_t(\omega)$$

is a continuous stochastic process. Set

$$\mathcal{F}_t = \sigma(\{X_s, 0 \leq s \leq t\}), \quad t \geq 0.$$

Let $T : \Omega \rightarrow \mathbb{R}$ be $\mathcal{S}$ -measurable. Prove that the following are equivalent.

- (i) T is a stopping time with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$.
- (ii) For any $\omega, \omega' \in \Omega$ such that $X_s(\omega) = X_s(\omega')$ for any $s \leq T(\omega)$ then $T(\omega') = T(\omega)$.

□

Exercise 3.38. Let $N(t)$ be a Poisson process with intensity λ as described in Example 1.3.7. Denote by $(\mathcal{F}_t)$ the natural filtration, $\mathcal{F}_t = \sigma(N(s), s \leq t)$.

- (i) Prove that $N(t)$ is an R -process.
- (ii) Prove that $\mathcal{F}_{t+} = \mathcal{F}_t, \forall t \geq 0$.
- (iii) Prove that $\mathbb{E}[N(t) \mid \mathcal{F}_s] = \mathbb{E}[N(t) \mid N(s)], \forall 0 \leq s < t$.
- (iv) Prove that the process $N(t) - t, t \geq 0$ is a martingale

□

Exercise 3.39. Suppose that $W : L^2([0, \infty)) \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$ is a Gaussian white noise; see Example 2.5.9. Fix $f \in L^2([0, \infty))$ and consider the Wiener integral (see Example 2.5.9 and Exercise 2.78)

$$X_t = \int_0^t f(s) dB(s) := W(I_{[0,t]} f), \quad t \geq 0.$$

- (i) Prove that (X_t) is an L^2 martingale adapted to the filtration $\mathcal{F}_t := \sigma(X_s, s \leq t)$.
- (ii) Use Kolmogorov's Continuity Theorem 2.5.12 to show that $(X_t)_{t \geq 0}$ admits a continuous modification.

□

Exercise 3.40. Let $B(t), t \geq 0$ be a one-dimensional Brownian motion started at 0. For each $n \in \mathbb{N}$ and each $t \geq 0$ we set

$$X_t^n := \sum_{k=1}^n B((k-1)t/n) \left(B(kt/n) - B((k-1)t/n) \right).$$

- (i) Prove that for any $n \in \mathbb{N}$ the stochastic process (X_t^n) is an L^2 -martingale.
- (ii) Prove that for each $t \geq 0$ X_t^n converges to $B(t)^2 - t$ in L^2 as $n \rightarrow \infty$.

□

Exercise 3.41. Suppose that $(W_t)_{t \geq 0}$ is a pre-Brownian motion defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$; see Definition 2.5.2. Let $t_0, \delta \geq 0$. Set

$$R(t_0, \delta) = \sup_{t \in \mathbb{Q} \cap [t_0, t_0 + \delta]} |B(t) - B(t_0)|.$$

(i) Prove that

$$\mathbb{P}[R(t_0, \delta) > \varepsilon] \leq \frac{3\delta^2}{\varepsilon^4}, \quad \forall \varepsilon, \delta > 0.$$

Hint. Use Doob's maximal inequalities.

(ii) Prove that W_t is a a.s. *uniformly* continuous on $\mathbb{Q}_{\geq 0}$ and conclude that (W_t) admits a modification continuous on $[0, \infty)$

□

Exercise 3.42. Let $d \in \mathbb{N}$ and suppose that $B_t^1, \dots, B_t^d$ are independent Brownian motions. Set

$$\mathbf{B}_t := (B_t^1, \dots, B_t^d), \quad t \geq 0.$$

Let U be a bounded open set containing the origin and set

$$T := \inf \{ t \geq 0; \mathbf{B}_t \in U^c \}.$$

Prove that $\mathbb{E}[T] < \infty$. **Hint.** Choose $a > 0$ such that $U \subset [-a, a]^d$ and use Example 3.5.23.

□

Exercise 3.43. Let $B(t)$ be a Brownian motion. For $u < 0 < v$ we set

$$T_{u,v} = \inf \{ t \geq 0; B(t) \in \mathbb{R} \setminus (u, v) \}.$$

Prove that for $\lambda > 0$ we have

$$\mathbb{E}[e^{-\lambda T_{u,v}}] = \frac{\sinh(\sqrt{2\lambda}|u|) + \sinh(\sqrt{2\lambda}v)}{\sinh(\sqrt{2\lambda}(|u| + v))}.$$

□

Markov chains

The Markov chains form a special but sufficiently general class of examples of stochastic processes. Their investigation requires a diverse arsenal of techniques, probabilistic and not only, and they reveal important patterns arising in many other instances.

The foundations of this theory were laid by the Russian mathematician A. A. Markov at the beginning of the twentieth century. By most accounts, Markov was a rather unconventional individual. He discovered what we now know as Markov chains in his attempts to contradict Pavel Nekrasov, a mathematician/theologian of that time who maintained on a theological basis that the Law of Large Numbers was specific to independent events/random variables and cannot be seen in other contexts. Markov succeeded in proving Nekrasov wrong and in the process laid the foundations of the theory of Markov chains. For more on this history of this concept we refer to the very readable article [103].

So what did Markov discovered? Think of a Markov chain as a random walk on a finite set $\mathfrak{X}$. From a given location x the walker can go to a location x' with probability $q_{x,x'}$. Suppose that at some location $x_0 \in \mathfrak{X}$ we placed a pile of sand consisting of giddy grains of sand: every second one of them starts this random walk and performs a billion steps (think of a fixed but very large number of steps). After all the grains of sand performed this ritual, the initial pile of sand is redistributed at various points of $\mathfrak{X}$. Denote by m_x^1 the mass of the pile of sand relocated at x . Next, collect the piles from their locations and move them back to the initial location x_0 .

Run the above experiment again we get a new distribution of piles of sand at the points of $\mathfrak{X}$. Denote the mass at x by m_x^2 . Markov observed that

$$\frac{m_x^1}{m_x^2} \approx 1, \quad \forall x.$$

Run the experiment a third time to obtain a third distribution of mass $(m_x^3)_{x \in \mathfrak{X}}$ and the conclusion is the same

$$\frac{m_x^1}{m_x^3} \approx 1, \quad \forall x.$$

To put it differently, if m is the mass of the pile of sand at x_0 , then, for any $x \in X$,

$$\frac{m_x^1}{m} \approx \frac{m_x^2}{m} \approx \frac{m_x^3}{m} \approx \dots$$

This phenomenon is one manifestation of the Law of Large Numbers for Markov chains.

During this more than a century since its creation, the theory of Markov chains has witnessed dramatic growth and generalizations, and has found applications in unexpected problems. For example, Google's PageRank algorithm is a special application of the Law of Large numbers for Markov chains.

The present chapter is an introduction to the theory of Markov chains. We present the classical results and spend some time on some more recent developments. As always, we try to illustrate the power of the theory on many concrete example. Needless to say, we barely scratch the surface of this subject.

4.1. Markov chains

In the sequel $\mathfrak{X}$ will denote a finite or countable set equipped with the discrete topology. We will refer to it as the *state space*. The Borel sigma-algebra of $\mathfrak{X}$ coincides with the sigma-algebra $2^{\mathfrak{X}}$ of all subsets of $\mathfrak{X}$.

4.1.1. Definition and basic concepts.

Definition 4.1.1. A *Markov chain with state space* $\mathfrak{X}$ is a sequence of random variables

$$X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathfrak{X}, 2^{\mathfrak{X}}), \quad n \in \mathbb{N}_0,$$

satisfying the *Markov property*

$$\mathbb{P}[X_{n+1} = x_{n+1} | X_n = x_n] = \mathbb{P}[X_{n+1} = x_{n+1} | X_n = x_n, \dots, X_0 = x_0], \quad (4.1.1)$$

$\forall n \in \mathbb{N}, x_0, x_1, \dots, x_n, x_{n+1} \in \mathfrak{X}$.

The *filtration associated to the Markov chain* is the sequence of sigma-subalgebras

$$\mathcal{F}_n := \sigma(X_0, \dots, X_n), \quad n \in \mathbb{N}_0.$$

The probability distribution of X_0 is called the *initial distribution* of the system.

The Markov chain is called *homogeneous* if, for any $x, x' \in \mathfrak{X}$, and any $n \in \mathbb{N}$ we have

$$\mathbb{P}[X_{n+1} = x' | X_n = x] = \mathbb{P}[X_1 = x' | X_0 = x]$$

In this case, the function

$$Q : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1], \quad Q(x_0, x_1) = Q_{x_0, x_1} = \mathbb{P}[X_1 = x_1 | X_0 = x_0]$$

is called the *transition matrix*¹ of the homogeneous Markov chain.

I will use the acronym HMC as a stand-in for Homogeneous Markov Chain. We denote by $\text{Markov}(\mathfrak{X}, \mu, Q)$ the collection of HMC-s with state space $\mathfrak{X}$, initial distribution μ and transition matrix Q . $\square$

Remark 4.1.2. (a) Let us observe that the Markov property can be written in the more compact form

$$\mathbb{P}[X_{n+1} = x | \mathcal{F}_n] = \mathbb{P}[X_{n+1} = x | \mathcal{F}_n], \quad \forall n \in \mathbb{N}, \quad x \in \mathfrak{X}. \quad (4.1.2)$$

In view of Proposition 1.4.18, the last property is equivalent to the conditional independence

$$X_{n+1} \perp\!\!\!\perp_{\mathcal{F}_n} X_n, \quad \forall n \in \mathbb{N}. \quad (4.1.3)$$

¹I made the decision to break with the tradition and use the letter Q to denote the transition matrix after teaching this topic and realizing that there were too many P 's on the blackboard and this sometimes confused the audience.

Exercise 1.64 shows that this is also equivalent to the condition

$$X_{n+1} \perp\!\!\!\perp_{X_n} \mathcal{F}_n, \quad \forall n \in \mathbb{N}. \quad (4.1.4)$$

One can show that this further equivalent to that

$$\sigma(X_{n+1}, X_{n+2}, \dots) \perp\!\!\!\perp_{X_n} \mathcal{F}_n. \quad (4.1.5)$$

This is colloquially expressed as saying that *the future is conditionally independent of the past given the present*.

(b) It is convenient to think of a Markov chain with state space $\mathfrak{X}$ as describing the random walk of a grasshopper hopscotching on the elements of $\mathfrak{X}$. The decision where to jump next is not influenced by the past, but only by the current location and the current time. For a homogeneous Markov chain the decision where to jump next depends only on the current location and not on the “time” n when the grasshopper reaches that state. Thus Q_{x_0, x_1} is the probability that the grasshopper, currently located at x_0 , will jump to x_1 .

We can represent an HMC with state space $\mathfrak{X}$ and transition matrix Q as a directed graph (loops allowed) with vertex set $\mathfrak{X}$ constructed as follows: there is a directed edge from x_0 to x_1 if and only if $Q_{x_0, x_1} > 0$. $\square$

If $(X_n)_{n \geq 0}$ is a homogeneous Markov chain then its transition matrix Q is *stochastic*, i.e.,

$$Q_{x_0, x_1} \geq 0, \quad \sum_{x \in \mathfrak{X}} Q_{x_0, x} = 1, \quad \forall x_0, x_1 \in \mathfrak{X}. \quad (4.1.6)$$

In other words, the entries of the matrix Q are nonnegative and the sum of the entries in each row is equal to 1.

If μ_n is the distribution of X_n , then, for any $x \in \mathfrak{X}$ we have

$$\mathbb{P}[X_{n+1} = x] = \sum_{x' \in \mathfrak{X}} \mathbb{P}[X_n = x'] Q_{x', x} = \sum_{x' \in \mathfrak{X}} \mu_n[x'] Q_{x', x}.$$

Think of μ_n and μ_{n+1} as matrices consisting of a single row. We can rewrite the above equality as an equality of matrices $\mu_{n+1} = \mu_n Q$. In particular,

$$\mu_n = \mu_0 Q^n, \quad (4.1.7)$$

where Q^n denotes the n -th power of the matrix Q , $Q^n = (Q_{x,y}^n)_{x,y \in \mathfrak{X}}$. From (4.1.7) we deduce that

$$\mathbb{P}[X_n = x_n | X_0 = x_0] = Q_{x_0, x_n}^n. \quad (4.1.8)$$

For this reason the matrix Q^n is also known as the n -th step transition matrix.

Let us show that given any matrix $Q : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1]$ satisfying (1.2.23) and any probability measure μ on $\mathfrak{X}$, there exists a homogeneous Markov chain, with state space $\mathfrak{X}$, initial distribution μ and transition matrix Q , i.e., $\text{Markov}(\mathfrak{X}, \mu, Q) \neq \emptyset$.

Observe that we can view Q as a kernel or random probability measure

$$\hat{Q} : \mathfrak{X} \times 2^{\mathfrak{X}} \rightarrow [0, 1], \quad (x, A) \mapsto \hat{Q}_x[A] = \sum_{a \in A} Q_{x,a}.$$

Note that $\hat{Q}_x[\cdot]$ is a probability measure on $\mathfrak{X}$. It is described by row x of the matrix Q .

Consider the set $\mathfrak{X}^{\mathbb{N}_0}$ equipped with the natural product sigma algebra $\mathcal{E}$; see Definition 1.5.3. In this case it coincides with the sigma algebra generated by π -system consisting of the *cylinders*

$$C_{s_0, s_1, \dots, s_k} := \{ \underline{x} = (x_n)_{n \in \mathbb{N}_0} \in \mathfrak{X}^{\mathbb{N}_0}; \ x_i = s_i, \ \forall i = 0, \dots, k \}.$$

Let us observe that there exists a probability measure $\mathbb{P}_\mu : \mathcal{E} \rightarrow [0, 1]$ uniquely determined by the conditions

$$\mathbb{P}_\mu [C_{s_0, s_1, \dots, s_k}] = \mu[s_0] \prod_{i=1}^k Q_{s_{i-1}, s_i}. \quad (4.1.9)$$

To prove that such a measure does indeed exist for any μ and Q we will rely on Kolmogorov's existence theorem, Theorem 1.5.5.

The equalities (4.1.9) define probability measures $\mathbb{P}_k = \mathbb{P}_k^{\mu, Q}$ on the product spaces $\mathfrak{X}^{\{0, 1, \dots, k\}}$ by setting

$$\mathbb{P}_k [(s_0, \dots, s_k)] := \mu[s_0] \prod_{i=1}^k Q_{s_{i-1}, s_i}. \quad (4.1.10)$$

Note that for $f : \mathfrak{X}^{\{0, 1, \dots, k\}} \rightarrow \mathbb{R}$ we have

$$\begin{aligned} & \int_{\mathfrak{X}^{\{0, 1, \dots, k\}}} f(x_0, \dots, x_k) \mathbb{P}_k [dx_0 \cdots dx_k] \\ &= \sum_{x_0 \in \mathfrak{X}} \sum_{x_1 \in \mathfrak{X}} \cdots \sum_{x_k \in \mathfrak{X}} \mu[x_0] Q_{x_0, x_1} \cdots Q_{x_{k-1}, x_k} f(x_0, \dots, x_k). \end{aligned} \quad (4.1.11)$$

The family of measures $(\mathbb{P}_k)_{k \geq 0}$ is projective since the transition matrix Q is stochastic. Indeed,

$$\begin{aligned} \mathbb{P}_{k+1} [(s_0, \dots, s_k) \times \mathfrak{X}] &= \sum_{x \in \mathfrak{X}} \mathbb{P}_{k+1} [(s_0, \dots, s_k, x)] \\ &= \left(\mu[s_0] \prod_{i=1}^k Q_{s_{i-1}, s_i} \right) \underbrace{\sum_x Q_{s_k, x}}_{=1} \\ &= \mu[s_0] \prod_{i=1}^k Q_{s_{i-1}, s_i} = \mathbb{P}_k [(s_0, \dots, s_k)]. \end{aligned} \quad (4.1.12)$$

Kolmogorov's existence theorem, then implies the existence of $\mathbb{P}_\mu \in \text{Prob}(\mathfrak{X}^{\mathbb{N}_0})$ satisfying (4.1.9). Moreover, the equality

$$\mu = \sum_{x \in \mathfrak{X}} \mu[x] \delta_x$$

implies that

$$\mathbb{P}_\mu = \sum_{x \in \mathfrak{X}} \mu[x] \mathbb{P}_x, \quad \mathbb{P}_x := \mathbb{P}_{\delta_x}. \quad (4.1.13)$$

For $n \in \mathbb{N}_0$ we denote by $\mathcal{E}_n$ the sub-sigma-algebra of $\mathcal{E}$ generated by $X_0, X_1, \dots, X_n$. Note that $\mathbb{P}_n$ can be identified with the restriction of $\mathbb{P}_\mu$ to $\mathcal{E}_n$.

For $\mu \in \text{Prob}(\mathfrak{X})$ we denote by $\mathbb{E}_\mu$ the expectation (integral) with respect to $\mathbb{P}_\mu$

$$\mathbb{E}_\mu : L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu) \rightarrow \mathbb{R}, \quad \mathbb{E}_\mu[F] = \int_{\mathfrak{X}^{\mathbb{N}_0}} F(\underline{x}) \mathbb{P}_\mu[d\underline{x}]. \quad (4.1.14)$$

For $x \in \mathfrak{X}$ we set

$$\mathbb{E}_x := \mathbb{E}_{\delta_x}. \quad (4.1.15)$$

We have a shift operator

$$\Theta : \mathfrak{X}^{\mathbb{N}_0} \rightarrow \mathfrak{X}^{\mathbb{N}_0}, \quad \Theta(x_0, x_1, x_2, \dots) = (x_1, x_2, \dots).$$

Note that $X_n = X_0 \circ \Theta^n$, $\Theta^n = \underbrace{\Theta \circ \dots \circ \Theta}_n$.

Theorem 4.1.3. *Consider the random variables*

$$X_n : \mathfrak{X}^{\mathbb{N}_0} \rightarrow \mathfrak{X}, \quad X_n(\underline{x}) = x_n, \quad n \in \mathbb{N}_0.$$

*Then the stochastic process $(X_n)_{n \in \mathbb{N}_0}$ is an HMC, defined on $(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu)$ with transition state space $\mathfrak{X}$ matrix Q and initial distribution μ . The probability space $(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu)$ is called the **path space** of this HMC.*

Moreover, if $F \in L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu)$, then

$$\mathbb{E}_\mu[F \circ \Theta^n \mid \mathcal{E}_n] = \mathbb{E}_\mu[F \mid X_n]. \quad (4.1.16)$$

Proof. For each x we have a probability measure Q_x on $\mathfrak{X}$ given by

$$Q_x[\{x'\}] = Q_{x,x'}, \quad \forall x' \in \mathfrak{X}.$$

We will show that for any $A \subset \mathfrak{X}$ we have the equality of random variables

$$\mathbb{P}[X_{n+1} \in A \mid \mathcal{E}_n] = Q_{X_n}[A] = \sum_{a \in A} Q_{X_n, a}. \quad (4.1.17)$$

Let $B \in \mathcal{E}_n$. It is a cylinder of the form

$$B = \{X_0 \in B_0, \dots, X_n \in B_n\}, \quad B_0, B_1, \dots, B_n \subset \mathfrak{X}.$$

Then

$$\begin{aligned} \mathbb{E}[I_A(X_{n+1})I_B] &= \mathbb{P}_\mu[\{X_{n+1} \in A\} \cap B] = \mathbb{P}_\mu[X_0 \in B_0, \dots, X_n \in B_n, X_{n+1} \in A] \\ &\stackrel{(4.1.11)}{=} \int_B Q_{X_n}[A] d\mathbb{P}_\mu. \end{aligned}$$

This proves (4.1.17).

The random measure Q_{X_n} is a regular version of the conditional probability $\mathbb{P}[X_{n+1} \in \cdot \mid X_n]$, i.e.,

$$Q_{X_n}[S] = \mathbb{P}[X_{n+1} \in S \mid X_n], \quad \forall S \subset \mathfrak{X}.$$

Using Proposition 1.4.25 we deduce that for every bounded function $f : \mathfrak{X} \rightarrow \mathbb{R}$ we have

$$\mathbb{E}[f(X_{n+1}) \mid \mathcal{E}_n] = \sum_{x \in \mathfrak{X}} Q_{X_n, x} f(x). \quad (4.1.18)$$

Let $\mathcal{M} \subset L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu)$ denote the collection of functions F satisfying (4.1.16). Clearly $\mathcal{M}$ is a vector space and if F_n is a sequence in $\mathcal{M}$ such that $F_n \nearrow F$, $F \in L^1$, then $F \in \mathcal{M}$. To show that $\mathcal{M} = L^1$ we use Monotone Class Theorem so it suffices to show that there exists a π -system $\mathcal{C} \subset \mathcal{E}$ that generates $\mathcal{E}$ such that $I_C \in \mathcal{M}$.

Denote by $\mathcal{C}$ the set of cylinders

$$C_{A_0, A_1, \dots, A_N} := \{\underline{x} \in \mathfrak{X}^{\mathbb{N}_0}; \quad x_i \in A_i, \quad i = 0, \dots, N\}.$$

Note that

$$\mathbf{I}_{C_{A_0, \dots, A_N}} = \prod_{k=0}^N \mathbf{I}_{\{X_k \in A_k\}},$$

and

$$\mathbf{I}_{C_{A_0, \dots, A_N}} \circ \Theta^n = \prod_{k=0}^N \mathbf{I}_{\{X_{n+k} \in A_k\}}$$

By definition $\mathcal{C}$ generates $\mathcal{E}$. Since $\mathcal{M}$ is a vector space it suffices to check that $\mathbf{I}_C \in \mathcal{M}$ for $C \in \mathcal{C}$ of the form

$$C = C_{A_0, \dots, A_N}, \quad A_k = \{x_k\}, \quad x_k \in \mathfrak{X}, \quad k = 0, 1, \dots, N.$$

To verify (4.1.16) for sets of this form and arbitrary n we argue by induction on N . For $N = 1$ this follows from (4.1.17). For the inductive step note that

$$\begin{aligned} \mathbb{E}[\mathbf{I}_{\{X_n=x_0, X_{n+1}=x_1, \dots, X_{n+N}=x_N\}} \mid \mathcal{E}_n] &= \mathbb{E}\left[\prod_{k=0}^N \mathbf{I}_{\{X_{n+k}=x_k\}} \mid \mathcal{E}_n\right] \\ &= \mathbb{E}\left[\mathbf{I}_{\{X_n=x_0\}} \mathbb{E}\left[\prod_{k=1}^N \mathbf{I}_{\{X_{n+k}=x_k\}} \mid \mathcal{E}_{n+1}\right] \mid \mathcal{E}_n\right] \\ &= \mathbb{E}\left[\mathbf{I}_{\{X_n=x_0\}} \underbrace{\mathbb{E}\left[\prod_{k=1}^N \mathbf{I}_{\{X_{n+k}=x_k\}} \mid X_{n+1}\right]}_{=: f(X_{n+1})} \mid \mathcal{E}_n\right] \end{aligned}$$

(use the inductive assumption)

$$= \mathbb{E}\left[\mathbf{I}_{\{X_n=x_0\}} f(X_{n+1}) \mid X_n\right] = \mathbb{E}\left[\mathbf{I}_{\{X_n=x_0\}} \underbrace{\mathbb{E}\left[\prod_{k=1}^N \mathbf{I}_{\{X_{n+k}=x_k\}} \mid \mathcal{E}_{n+1}\right]}_{=: f(X_{n+1})} \mid X_n\right]$$

($\sigma(X_n) \subset \mathcal{E}_{n+1}$)

$$= \mathbb{E}\left[\prod_{k=0}^N \mathbf{I}_{\{X_{n+k}=x_k\}} \mid X_n\right].$$

□

Remark 4.1.4. We have deduced (4.1.16) relying on the Markov property. The above proof shows that the Markov property (4.1.17) is a special case of (4.1.16). For this reason we can take (4.1.16) as definition of Markov's property. □

Given a homogeneous Markov chain $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}$, $n \geq 0$, with state space $\mathfrak{X}$, initial distribution μ and transition matrix Q , we obtain a measurable map

$$\vec{X} : (\Omega, \mathcal{S}) \rightarrow (\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}), \quad \omega \mapsto \vec{X}(\omega) = (X_n(\omega))_{n \geq 0}.$$

The *distribution of the Markov chain* is the pushforward measure

$$\mathbb{P}_{\vec{X}} := \vec{X}_\# \mathbb{P} \in \text{Prob}(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}).$$

It is uniquely determined by the equalities

$$\mathbb{P}_{\vec{\mathcal{X}}} [C_{s_0, s_1, \dots, s_k}] := \mathbb{P} [X_0 = s_0, \dots, X_k = s_k] = \mu_0[s_0] \prod_{i=1}^k Q_{s_{i-1}, s_i}. \quad (4.1.19)$$

We deduce

$$\mathbb{P}_{\vec{\mathcal{X}}} = \mathbb{P}_{\mu}.$$

For every, $F \in L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_{\mu})$, we have

$$\mathbb{E}_{\mathbb{P}} [F(X_0, X_1, \dots)] = \mathbb{E}_{\mu} [F] = \int_{\mathfrak{X}^{\mathbb{N}_0}} F(\underline{x}) \mathbb{P}_{\mu} [d\underline{x}].$$

This is a special case of the change in variables formula (1.2.25).

This shows that the distribution of the Markov chain is uniquely determined by the initial distribution $\mu \in \text{Prob}(\mathfrak{X})$ and the transition matrix Q .

Remark 4.1.5. One can define any HMC on probability spaces other than $\mathfrak{X}^{\mathbb{N}_0}$. Here is a such a construction corresponding to state space $\mathfrak{X}$ transition matrix Q and initial probability distribution μ . We set $\mu_x := \mu[x]$. Assume that $\mathfrak{X}$ is a subset of $\mathbb{N}$. As such it is equipped with a total order.

First, a little bit of terminology. We say that an interval is *convenient* if it is either empty or the form $[a, b)$, $a < b$. If $[a, b)$, $[c, d)$ are nonempty convenient intervals, then we say that $[a, b)$ precedes $[c, d)$ and we write $[a, b) \prec [c, d)$ if $b \leq c$. The empty set is allowed to precede or succeed any nonempty convenient interval.

The probability space is the unit interval $[0, 1)$ equipped with the Lebesgue measure. The random variables X_n , depend on the choice of initial distribution, and are defined inductively as follows.

- Partition $[0, 1)$ into convenient intervals $I_x = I_x^0$, $x \in \mathfrak{X}$ of Lebesgue measures $\mu_x = \lambda[I_x^0]$, such that

$$x < x' \Rightarrow I_x \prec I_{x'}.$$

- Partition each interval $I_{x_0}^0$ into convenient intervals I_{x_0, x_1}^1 of sizes $\mu_{x_0} Q_{x_0, x_1}$, $x_0, x_1 \in \mathfrak{X}$, such that

$$x < x' \Rightarrow I_{x_0, x}^1 \prec I_{x_0, x'}^1.$$

- Inductively, partition each interval $I_{x_0, x_1, \dots, x_n}^n$ into convenient intervals $I_{x_0, x_1, \dots, x_n, x_{n+1}}^{n+1}$ of sizes

$$\lambda[I_{x_0, x_1, \dots, x_n}^n] Q_{x_n, x_{n+1}} = \mu_{x_0} \prod_{j=0}^n Q_{x_j, x_{j+1}},$$

such that

$$x < x' \Rightarrow I_{x_0, \dots, x_n, x}^{n+1} \prec I_{x_0, x_1, \dots, x_n, x'}^{n+1}.$$

Now define $X_n : [0, 1) \rightarrow \mathfrak{X}$ by setting

$$X_n(t) = x_n \text{ if } t \in \bigcup_{x_0, x_1, \dots, x_{n-1} \in \mathfrak{X}} I_{x_0, x_1, \dots, x_n}^n$$

Note that these random variables are defined on the *same probability space*

$$([0, 1], \mathcal{B}_{[0, 1]}, \lambda),$$

but they depend on the choice of the initial distribution.

This is different from the construction based on path spaces. In that case we are given measurable maps defined on the same measurable space $(\Omega, \mathcal{S})$, but we obtain different HMC's by choosing *different probability measures* on $(\Omega, \mathcal{S})$. $\square$

4.1.2. Examples. The homogeneous Markov chains appear in many and diverse situations. According to the discussion in the previous subsection, to describe an HMC it suffices to describe the state space $\mathfrak{X}$ and the transition matrix Q . We will remain vague about the initial distribution μ .

Example 4.1.6 (Gambler's ruin). Consider the gambler's ruin problem discussed in Example 3.2.37. The state space is $\mathfrak{X} = \{0, 1, \dots, N\}$. Then X_n is the fortune of a gambler at time n . The gambler flips a fair coin with faces labeled ± 1 . If its fortune is strictly in between 0 and N , then its fortune changes by the amount shown on the face of the coin. The game stops when its fortune reaches either 0 or N . Concretely

$$\begin{aligned} Q_{N,k} &= 0, \quad \forall k < N, \quad Q_{N,N} = 1, \\ Q_{0,j} &= 0, \quad \forall j > 0, \quad Q_{0,0} = 1, \\ Q_{k,k+1} &= Q_{k,k-1} = \frac{1}{2}, \quad Q_{k,j} = 0, \quad \text{if } |k-j| > 1, \quad 0 < k, j < N. \end{aligned}$$

The directed graph describing this HMC is depicted in Figure 4.1 where, for clarity, we have omitted




Figure 4.1. The gambler's ruin chain

the loops at 0 and N $\square$

Example 4.1.7 (The Ehrenfest Urn). Consider the following situation. There are B balls in two urns. Equivalently, think of an urn with two chambers. Pick one of these B balls uniformly at random and move it in the other box/chamber. Denote by X_n the number of balls in the left box at time n . Then $(X_n)_{n \geq 0}$ is an HMC with transition probabilities

$$\begin{aligned} Q_{i,i+1} &= \frac{B-i}{B}, \quad i = 0, 1, \dots, B-1, \quad Q_{i,i-1} = \frac{i}{B}, \quad i = 1, \dots, B, \\ Q_{i,j} &= 0, \quad |i-j| > 1. \end{aligned}$$

This HMC is known as the *Ehrenfest urn*. Note that during this process it is more likely that a ball moves from the more crowded box to the less crowded one, similarly to what happens in diffusion processes. $\square$

Example 4.1.8 (Bernoulli-Laplace Urn). Consider again an urn with two chambers. There are N balls in each chamber and, of the total of $2N$ balls, half are black and half are white. We simultaneously draw one ball from each chamber, and then switch their chamber locations. After we perform n such experiments we will have W_n white balls in the first chamber. The sequence $(W_n)_{n \geq 0}$ is an HMC with state space $\{0, 1, \dots, N\}$ and transition probabilities

$$Q_{k,k-1} = \left(\frac{k}{N}\right)^2, \quad Q_{k,k+1} = \left(1 - \frac{k}{N}\right)^2, \quad Q_{k,k} = k(N-k)/N^2.$$

$\square$

Example 4.1.9 (Birth-and-death chains with two absorbing barriers). The state space is

$$\mathcal{J}_N = \{0, 1, \dots, N\}$$

and the transition matrix Q is tridiagonal, i.e., $Q_{ij} = 0$ if $|i - j| > 1$. For $i = 1, \dots, N - 1$ we set

$$p_i := Q_{i,i+1}, \quad q_i := Q_{i,i-1}, \quad r_i := Q_{i,i}.$$

We assume reflecting conditions at the the boundary $Q_{0,1} = 0 = Q_{N,N-1}$. $\square$

Example 4.1.10 (Random placement of balls). Consider a sequence of independent trials each consisting in randomly placing a ball in one of r given urns. We say that the system is in state k if exactly k urns are occupied.

We obtain an HMC with state space $\{0, 1, \dots, r\}$ with transition probabilities

$$Q_{j,j} = \frac{j}{r}, \quad Q_{j,j+1} = \frac{r-j}{r}, \quad 0 \leq j < r,$$

and of course $Q_{j,k} = 0$ for any other pairs (j, k) . If $X_0 = 0$, so initially all boxes are empty, then $X_n = r - N_{r,n}$, where $N_{r,n}$ is the number of empty boxes investigated in Exercise 2.31. $\square$

Example 4.1.11 (Random walk on $\mathbb{Z}^d$). Suppose that $(X_n)_{n \geq 1}$ are i.i.d. $\mathbb{Z}^d$ -valued random variables. Denote by π their common distribution. Set

$$S_0 = 0, \quad S_n = X_1 + \dots + X_n.$$

Then the random process $(S_n)_{n \in \mathbb{N}_0}$ is an HMC with transition matrix

$$Q_{\mathbf{m}, \mathbf{n}} = \mathbb{P}[X_1 = \mathbf{n} - \mathbf{m}] = \pi[\mathbf{n} - \mathbf{m}], \quad \mathbf{m}, \mathbf{n} \in \mathbb{Z}^d$$

One can imagine this process as a person starting at the origin of $\mathbb{Z}^d$ and walking with random step sizes, with X_n the size of the n -th step.

A standard random walk is obtained as follows. Denote by $e_1, \dots, e_d$ the canonical basis $\mathbb{Z}^d$ and choose π to be uniformly distributed on the set $\{\pm e_1, \dots, \pm e_d\}$, i.e.,

$$\pi[\pm e_k] = \frac{1}{2d}, \quad k = 1, \dots, d.$$

For example, when $d = 1$, this corresponds to a random walk on $\mathbb{Z}$ where, at each moment, going one step ahead or one step back is decided by flipping a fair coin. $\square$

Example 4.1.12 (Simple random walk on a graph). Consider an undirected graph $G = (V, E)$, where V is the set of vertices, and E denotes the set of edges. We do not allow for multiple edges connecting two vertices. For every vertex v we denotes by $\deg(v)$ is degree, i.e., the number of edges of E at v . We assume that the graph is locally finite, i.e., $\deg(v) < \infty, \forall v \in V$.

Suppose now that a grasshopper hopscoches on the set vertices V according to the following rule: if situated at a vertex v_0 , the grasshopper will jump to one of the neighbors of v_0 in V chosen uniformly randomly. Denote by X_n the location of the grasshopper at time n . then $(X_n)_{n \geq 0}$ is an HMC with state space V with transition matrix

$$Q_{v_0, v_1} = \begin{cases} \frac{1}{\deg(v_0)}, & \text{if } v_0 \text{ and } v_1 \text{ are neighbors,} \\ 0, & \text{otherwise.} \end{cases}$$

$\square$

Example 4.1.13 (The branching process). Consider again the branching process with reproduction law μ described in Example 3.1.8. Recall that it deals with the evolution of a population of individuals of a species with $\mu[j]$ denoting the probability that a given individual will have $j \in \mathbb{N}_0$ offsprings.

Denote by Z_n the size of the n -th generation population. We assume that $Z_0 = 1$. Then $(Z_n)_{n \geq 0}$ is an HMC with state space $\mathbb{N}_0$.

To see this, choose a sequence of i.i.d. random variables $(\xi_k)_{k \in \mathbb{N}}$ with common distribution μ . Then

$$\mathbb{P}[Z_{n+1} = j | Z_n = i] = \mathbb{P}[\xi_1 + \cdots + \xi_i = j].$$

The distribution of the random variable $\xi_1 + \cdots + \xi_i$ is the convolution of μ^{*i} , the convolution of i copies of μ . More precisely,

$$\mu^{*i}[j] = \sum_{k_1 + \cdots + k_i = j} \mu[k_1] \cdots \mu[k_i].$$

The transition matrix is then $Q_{i,j} = \mu^{*i}[j]$. □

Example 4.1.14 (Queing). Customers arrive for service and take their place in a waiting line. During each period of time one customer is served, if at least one customer is present. During a service period new customers may arrive. We assume that the number of customers that arrive during the n -th service period is a random variable ξ_n , and that the random variables $\xi_1, \xi_2, \dots$ are i.i.d. with common distribution $\mu \in \text{Prob}(\mathbb{N}_0)$. We set $\mu_i := \mu[i]$, $i \in \mathbb{N}_0$. For notation convenience we set $\mu_n = 0$ for $n < 0$.

We denote by X_n the number of customers in line at the end of the n -th period. Note that

$$X_{n+1} = (X_n - 1)^+ + \xi_n.$$

The sequence $(X_n)_{n \geq 0}$ is an HMC with state space $\mathbb{N}_0$ and transition matrix

$$Q_{i,j} = \begin{cases} \mu_j, & i = 0, \\ \mu_{j-i+1}, & i > 0. \end{cases}$$
□

Example 4.1.15 (Noisy dynamical systems). Suppose that $T : \mathfrak{X} \rightarrow \mathfrak{X}$ is a selfmap of an at most countable set $\mathfrak{X}$. This defines a dynamical system $(T^n)_{n \in \mathbb{N}}$ which can be viewed as a trivial Homogeneous Markov Chain with transition matrix

$$Q_{x,x'} = \begin{cases} 1, & x' = T(x), \\ 0, & x' \neq T(x). \end{cases}$$

We can obtain more general Markov chains if we work with “noisy” selfmaps.

More precisely, suppose that $(S, \mathcal{S})$ is a measurable space and

$$T : \mathfrak{X} \times S \rightarrow \mathfrak{X}, \quad (x, s) \mapsto T_s(x)$$

is a measurable map. In other words, (T_s) is a measurable family of maps $\mathfrak{X} \rightarrow \mathfrak{X}$.

Fix an S -valued “noise”, i.e., a sequence

$$Z_n : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow (S, \mathcal{S}), \quad n \in \mathbb{N}$$

of i.i.d. S -valued random variables and an $\mathfrak{X}$ -valued random variable X_0 independent of the Z 's we obtain a *noisy dynamical system*

$$X_{n+1} = T_{Z_{n+1}}(X_n), \quad \forall n \in \mathbb{N}.$$

Hence

$$X_n = T_{Z_n} \circ \cdots \circ T_{Z_1}(X_0).$$

The sequence $(X_n)_{n \geq 0}$ is an HMC. Indeed,

$$\begin{aligned} & \mathbb{P}[X_{n+1} = x_{n+1} \mid X_n = x_n, \dots, X_0 = x_0] \\ &= \mathbb{P}[T_{Z_{n+1}}(x_n) = x_{n+1} \mid X_n = x_n, \dots, X_0 = x_0] \\ &= \mathbb{P}[T_{Z_{n+1}}(x_n) = x_{n+1}] \end{aligned}$$

since the event $\{X_n = x_n, \dots, X_0 = x_0\}$ belongs to the sigma algebra generated by $X_0, Z_1, \dots, Z_n$ and thus is independent of Z_{n+1} . On the other hand, obviously

$$\mathbb{P}[T_{Z_{n+1}}(x_n) = x_{n+1}] = \mathbb{P}[X_{n+1} = x_{n+1} \mid X_n = x_n].$$

This Markov chain is *homogenous* since the random variables Z_n are identically distributed.

The standard random walk on $\mathbb{Z}$ is a Markov system generated in an obvious way by a random dynamical system defined by the map

$$T : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}, \quad (m, z) \mapsto T_z(m) = m + z$$

and the noise is described by a sequence of i.i.d. Rademacher random variables $(Z_n)_{n \in \mathbb{N}}$. Then $X_{n+1} = X_n + Z_{n+1}$. One can show that any Markov chain can be produced in this fashion, as iterates of random maps. $\square$

4.2. The dynamics of homogeneous Markov chains

In this section we will consistently adopt the dynamical point of view on Markov chains described in Remark 4.1.2 (b) and isolate some useful features.

4.2.1. Classification of states. Suppose that $(X_n)_{n \geq 0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q .

Definition 4.2.1. (a) A state $x_1 \in \mathfrak{X}$ is said to be *accessible* from a state $x_0 \in \mathfrak{X}$, and we denote this by $x_0 \rightarrow x_1$, if $Q_{x_0, x_1}^n > 0$ for some $n \in \mathbb{N}_0$.

(b) The states x_0 and x_1 *communicate* if $x_0 \rightarrow x_1$ and $x_1 \rightarrow x_0$. We indicate this using the notation $x_0 \leftrightarrow x_1$. $\square$

Recall that to an HMC with state space $\mathfrak{X}$ we can associate a directed graph with vertex set $\mathfrak{X}$; see Remark 4.1.2 (b). A *walk* from x to x' in this graph is a sequence of vertices

$$x = x_0, x_1, \dots, x_n = x'$$

such that, for any $i = 1, \dots, n$, there exists a directed edge from x_{i-1} to x_i . If $x \neq x'$, then x' is accessible from x if there is a walk from x to x' .

Proposition 4.2.2. *The communication relation “ $\leftrightarrow$ ” is an equivalence relation.*

Proof. Reflexivity. $x \leftrightarrow x$ since $Q_{x,x}^0 = 1$

Symmetry. The relation is symmetric by definition.

Transitivity. Suppose that $x_0 \leftrightarrow x_1$ and $x_1 \leftrightarrow x_2$. Then, there exist $m, n \in \mathbb{N}_0$ such that

$$Q_{x_0,x_1}^m > 0 \text{ and } Q_{x_1,x_2}^n > 0.$$

Observe that

$$Q_{x_0,x_2}^{m+n} = Q_{x_0,x_1}^m Q_{x_1,x_2}^n + \underbrace{\sum_{x \in \mathfrak{X} \setminus \{x_1\}} Q_{x_0,x}^m Q_{x,x_2}^n}_{\geq 0} > 0.$$

Hence $x_0 \rightarrow x_2$. The opposite relation $x_2 \rightarrow x_0$ is proved in identical fashion. $\square$

Definition 4.2.3. The equivalence classes of the relation $\leftrightarrow$ are called the *communication classes* of the given HMC. $\square$

Example 4.2.4. (a) Consider the HMC associated to the gamblers's ruin problem described in Example 4.1.6. The state space is $\{0, 1, \dots, N\}$ and there are three communication classes

$$C_0 = \{0\}, C = \{1, \dots, N-1\}, C_N = \{N\}.$$

Note that no state in C is accessible from C_0 or C_N .

(b) The HMC associated to the Ehrenfest urn model in Example 4.1.7 has state space $\{0, 1, \dots, N\}$ and any two states communicate so that there is only a single communication class.

(c) The HMC corresponding to the random placement of balls problem in Example 4.1.10 has state space $\{0, 1, \dots, r\}$ and communication classes

$$C_0 = \{0\}, C_1 = \{1\}, \dots, C_r = \{r\}.$$

Note that for $j > i$, the class C_j is accessible from the class C_i . $\square$

Definition 4.2.5. Let $(X_n)_{n \in \mathbb{N}_0}$ be an HMC with state space $\mathfrak{X}$ and transition matrix Q .

- (i) A subset $C \subset \mathfrak{X}$ is *closed* with respect to this HMC if no state outside C is accessible from a state in C .
- (ii) A subset of $\mathfrak{X}$ is called *irreducible* if it is closed and contains no proper closed subset.
- (iii) A state $x \in \mathfrak{X}$ is called *absorbing* if the set $\{x\}$ is closed.
- (iv) The HMC is called *irreducible* if its state space is irreducible.

$\square$

Example 4.2.6. For the HMC corresponding to the random placement of balls problem in Example 4.1.10 with state space $\{0, \dots, r\}$, all the subsets $\{k, k+1, \dots, r\}$ are closed and the state r is absorbing. This is not an irreducible Markov chain. $\square$

Note that a subset $C \subset \mathfrak{X}$ is closed if and only if for any $x \in C$ we have

$$\sum_{y \in C} Q_{x,y} = 1.$$

Equivalently, this means that $\mathbb{P}[X_n \in C \mid X_0 \in C] = 1$.

Using the intuition of the randomly hopping grasshopper, this says that, once the grasshopper steps in a closed set it will be trapped there. Thus, a closed set $C \subset \mathfrak{X}$ defines a Markov sub-chain with state space C and transition matrix $(Q_{x,y})_{x,y \in C}$. In particular, this proves the following result.

Proposition 4.2.7. *A closed subset of the state space of an HMC is a union of communication classes.* $\square$




Figure 4.2. An HMC with a single irreducible subset.




Figure 4.3. Another HMC with a single irreducible subset.

Example 4.2.8. Consider an HMC with associated digraph depicted in Figure 4.2. It consists of three communication classes

$$C_1 := \{1, 2, 3, 4\}, \quad C_2 := \{5, 7, 8\}, \quad C_3 := \{6\}.$$

The communication class C_3 is closed while C_1 and C_2 are not. The only irreducible set is C_3 . In particular the state 6 is absorbing.

Suppose now that we change the directions of the edges $4 \rightarrow 5$ and $4 \rightarrow 6$ as depicted in Figure 4.3. This HMC has the same communication classes C_1, C_2, C_3 , but this time only C_1 is closed. $\square$

Lemma 4.2.9. *Let $C \subset \mathfrak{X}$ be a closed subset. Then the following are equivalent.*

- (i) C is irreducible.
- (ii) C is a communication class.

In particular, an HMC is irreducible if and only if it consists of a single communication class.

Proof. The implication (ii) $\Rightarrow$ (i) follows from Proposition 4.2.7: a closed set is a union of communication classes.

(i) $\Rightarrow$ (ii) Suppose that C is an irreducible subset of $\mathfrak{X}$. In particular, C is a union of communication classes

$$C = \bigcup_{j=1}^N C_j, \quad N \in \mathbb{N} \cup \{\infty\}.$$

Suppose that $N \geq 2$. Set $j_0 := 1$. Since C_{j_0} is not closed, there exists $j_1 \neq j_0$, $x_{j_0} \in C_{j_0}$ and $x_{j_1} \in C_{j_1}$ such that $x_{j_0} \rightarrow x_{j_1}$. In fact, any class in C_{j_1} is accessible from any class in C_{j_0} . We write this $C_{j_0} \rightarrow C_{j_1}$.

Next, we can find $j_2 \notin \{j_0, j_1\}$ such that $C_{j_1} \rightarrow C_{j_2}$. Clearly no class in $C_{j_0} \cup C_{j_1}$ is accessible from C_{j_2} . Iterating, we obtain a (possible finite) subsequence in $\mathbb{N} \cap [1, N]$

$$j_0, j_1, \dots, j_k, \dots,$$

where the j_k 's are pairwise distinct, such that

$$C_{j_0} \rightarrow C_{j_1} \rightarrow C_{j_2} \rightarrow \dots$$

and no state in $C_{j_0} \cup \dots \cup C_{j_k}$ is accessible from $C_{j_{k+1}}$. Note that

$$C' = \bigcup_{k \geq 1} C_{j_k} \subset C \setminus C_{j_0}$$

is a proper closed subset of C , contradicting the fact that C is irreducible. $\square$

Definition 4.2.10. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q .

(i) The set of *periods* of a state $x \in \mathfrak{X}$ is

$$\mathcal{P}_x := \{n \in \mathbb{N}; \quad Q_{x,x}^n > 0\}.$$

(ii) The *period* of a state x is $d = d(x) := \gcd \mathcal{P}_x$, where “gcd” stands for greatest common divisor. When $\mathcal{P}_x = \emptyset$ we set $d(x) := \infty$.

(iii) A state x is called *aperiodic* if $d(x) = 1$. $\square$

Lemma 4.2.11. Let $(X_n)_{n \geq 0}$ be an HMC with state space $\mathfrak{X}$ and transition matrix Q . Suppose that $x \in \mathfrak{X}$ and $d(x) < \infty$. Then the following hold.

(i) The set $\mathcal{P}_x$ is a semigroup of the additive semigroup $(\mathbb{N}, +)$.

(ii) There exists $N \in \mathbb{N}$ such that $nd(x) \in \mathcal{P}_x$, $\forall n \geq N$.

(iii) If $x \leftrightarrow y$, then $d(x) = d(y)$.

Proof. (i) Follows from the fact that $Q_{x,x}^{m+n} \geq Q_{x,x}^m Q_{x,x}^n$.

(ii) Set $d := d(x)$. We claim that there exist $k \geq 2$ and $m_1, \dots, m_k \in \mathcal{P}_x$ such that

$$d = \gcd(m_1, m_2, \dots, m_k).$$

Set

$$m_1 := \min \{ \{m; \quad m \in \mathcal{P}_x\}.$$

Then $d|d_1$. If $d < m_1$ $m_2 \in \mathcal{P}_x$ such that

$$\gcd(m_1, m_2) = d_2 := \min \{ \gcd(m_1, m); m \in \mathcal{P}_x \}.$$

Then $d|d_2|d_1$, $d_2 < d_1$. If $d < d_2$ choose $m_3 \in \mathcal{P}_x$ such that

$$\gcd(m_1, m_2, m_3) = d_3 := \min \{ \gcd(m_1, m_2, m); m \in \mathcal{P}_x \} \geq d.$$

Then $d|d_3|d_2$, $d_3 < d_2$. If $d < d_3$ define

$$d_4 := \min \{ \gcd(m_1, m_2, m_3, m); m \in \mathcal{P}_x \}.$$

and choose $m_4 \in \mathcal{P}_x$ such that $d_4 = \gcd(m_1, m_2, m_3, m_4)$.

Then $d|d_4|d_3$, $d_4 < d_3$. If $d_4 = d$ we stop. If $d_4 > d$, we iterate the above procedure. Clearly this procedure will stop after finitely many iterations.

An old result of I. Schur [223, Thm.3.15.2] implies that the set

$$\{m_1 n_1 + \cdots + m_k n_k; n_1, \dots, n_k \in \mathbb{N}_0\}$$

contains all the sufficiently large multiples of d .

(iii) For $x, y \in \mathfrak{X}$ we set

$$\mathcal{P}_{x,y} := \{n \in \mathbb{N}; Q_{x,y}^n > 0\}.$$

Thus $\mathcal{P}_x = \mathcal{P}_{x,x}$. Suppose $x \leftrightarrow y$. Note that

$$\mathcal{P}_{x,y} + \mathcal{P}_{y,x} \subset \mathcal{P}_x.$$

Hence $d(x)|(\mathcal{P}_{x,y} + \mathcal{P}_{y,x})$. From the inclusion

$$\mathcal{P}_{x,y} + \mathcal{P}_y + \mathcal{P}_{y,x} \subset \mathcal{P}_x$$

we deduce that $d(x)|\mathcal{P}_y$ so $d(x)|d(y)$. Reversing the roles of x, y in the above argument we deduce $d(y)|d(x)$ so $d(x) = d(y)$. $\square$

According to the above result, all the states of an irreducible HMC have the same period so we can speak of the period of that HMC.

Definition 4.2.12. An irreducible HMC is called *aperiodic* if each of its states has period 1. $\square$

Example 4.2.13. (i) Each state in the standard random walk on $\mathbb{Z}$ locally finite graph has period 2.

More generally, given a vertex v in a locally finite, connected graph, its set of periods with respect to the standard random walk coincides with the set of lengths of paths in the graph that start and end at v . Since there is such a path of length 2 we deduce that the vertex is aperiodic if and only if there exists a path of odd length starting and ending at x .

(ii) The Ehrenfest urn in Example 4.1.7 is irreducible with period 2. $\square$

Proposition 4.2.14. Let $(X_n)_{n \geq 0}$ be an irreducible HMC with state space $\mathfrak{X}$, transition matrix Q , and period $d < \infty$. Fix $x_0 \in \mathfrak{X}$. Consider the HMC $(Y_n)_{n \geq 0}$ with state space $\mathfrak{X}$, initial state $Y_0 = x_0$ and transition matrix $T = Q^d$. Denote by $\mathcal{C}_T$ the set of communication classes of T . For each $x \in \mathfrak{X}$ we denote by $[x]_T$ the T -communication class of x . Then the following hold.

- (i) There exists a bijection $r = r_{x_0} : \mathcal{C}_T \rightarrow \mathbb{Z}/d\mathbb{Z}$ such that $r([x]_T) = k \bmod d$ iff there exists $n \in \mathbb{N}_0$ such that $Q_{x_0,x}^{nd+k} > 0$.
- (ii) If $Q_{x,y} > 0$, then $r(y) \equiv r(x) + 1 \bmod d$.

(iii) *Each T -communication class is T -closed.*

Proof. As in the proof of Lemma 4.2.11 we set

$$\mathcal{P}_{x,y} := \{n \in \mathbb{N}; Q_{x,y}^n > 0\}.$$

Let us observe that

$$\forall x, y \in \mathfrak{X}, \forall n, m \in \mathcal{P}_{x,y} : n \equiv m \pmod{d}. \quad (4.2.1)$$

The claim is obviously true if $n = m$. Suppose that $n > m$. Then $n - m \in \mathcal{P}_{y,y}$ so $d = d(y)$ divides $n - m$.

Thus, for any $x, y \in \mathfrak{X}$ there exists $r = r(x, y) \in \{0, 1, \dots, d-1\}$ such that

$$\mathcal{P}_{x,y} \subset r(x, y) + d\mathbb{N}_0.$$

For any $x \in \mathfrak{X}$ we set $r(x) := r(x_0, x)$. We want to prove that

$$[x]_T = [y]_T \iff r(x) = r(y). \quad (4.2.2)$$

Indeed, suppose that $[x]_T = [y]_T$. Then there exists n such that $T_{x,y}^n > 0$, i.e., $Q_{x,y}^{nd} > 0$. Fix $m \in \mathbb{N}_0$ such that $Q_{x_0,x}^m > 0$. Then $Q_{x_0,y}^{m+nd} > 0$. Clearly

$$r(y) \equiv m + nd \equiv m \equiv r(x) \pmod{d}.$$

Conversely, suppose that $r(x) = r(y)$. Fix $n_x, n_y \in \mathbb{N}_0$ such that $Q_{x_0,x}^{n_x}, Q_{x_0,y}^{n_y} > 0$. Choose N large enough such that $Nd > n_x$ and $Nd \in \mathcal{P}_{x_0}$. Then $Nd - n_x \in \mathcal{P}_{x,x_0}$ and $n_y \in \mathcal{P}_{x_0,x}$ so $Nd - n_x + n_y \in \mathcal{P}_{x,y}$. Hence

$$0 = r(y) - r(x) \equiv n_y - n_x \pmod{d}.$$

We deduce that there exists $m \in \mathbb{N}_0$ such that $md = Nd - n_x + n_y$. Hence $T_{x,y}^m > 0$. In other words y is T -accessible from x . A symmetric argument shows that x is T -accessible from y so that $[x]_T = [y]_T$. This proves (4.2.2) and (i).

The statements (ii) and (iii) follow immediately from the equality

$$r(x, z) \equiv r(x, y) + r(y, z) \pmod{d},$$

so $r(y) = r(x) + r(x, y)$. □

Remark 4.2.15. Suppose that $(X_n)_{n \geq 0}$ is an HMC as in the above proposition and

$$C_{\text{cpt}}, C_1, \dots, C_{d-1} \subset \mathfrak{X}$$

are the communication classes of Q^d . If $X_0 \in C_i$, then $X_n \in C_{i+n \bmod d}$, for any n . Thus a grasshopper hopscotching following the prescriptions of this Markov chain will jump from a region C_i to somewhere in the next region C_{i+1} and so on, returning after d jumps to the region where he started.

Observe also that the map $r : \mathcal{C}_T \rightarrow \mathbb{Z}/d\mathbb{Z}$ depends on the choice of x_0 . On the other hand, the action of $\mathbb{Z}/d\mathbb{Z}$ on $\mathcal{C}_T$ is independent of x_0 and it is free and transitive. Thus $\mathcal{C}_T$ is naturally a $\mathbb{Z}/d\mathbb{Z}$ -torsor. □

4.2.2. The strong Markov property. Suppose that $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}$, $n \geq 0$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q . As usual, we denote by $(\mathcal{F}_n)_{n \geq 0}$ the filtration of $\mathcal{S}$ determined by this random process, i.e., $\mathcal{F}_n = \sigma(X_0, \dots, X_n)$, $n \in \mathbb{N}_0$.

Suppose now that T is a stopping time adapted to this filtration. In (3.1.7) we defined the sigma-algebra $\mathcal{F}_T$ associated to T by the requirements

$$S \in \mathcal{F}_T \iff S \cap \{T \leq n\} \in \mathcal{F}_n, \quad \forall n \in \mathbb{N}_0.$$

Example 4.2.16 (Return times). Let $(X_n)_{n \in \mathbb{N}_0}$ be an HMC with state space $\mathfrak{X}$. For $A \subset \mathfrak{X}$ we define

$$T_A := \min \{n \geq 1; X_n \in A\}.$$

We will refer to T_A as the *return time* to A . This is a stopping time with respect to the canonical filtration $\mathcal{F}_n$. For $x \in \mathfrak{X}$ we set $T_x := T_{\{x\}}$.

Note that the event S belongs to $\mathcal{F}_{T_A}$ if at any moment n we can decide using the information collected up to that point in $\mathcal{F}_n$ whether S occurred and we have returned to A up to that moment. $\square$

Example 4.2.17 (Début times). Let $(X_n)_{n \in \mathbb{N}_0}$ be an HMC with state space $\mathfrak{X}$. For $A \subset \mathfrak{X}$ we define

$$D_A := \min \{n \geq 0; X_n \in A\}.$$

We will refer to D_A as the *début time* or *hitting time* of A . This is a stopping time with respect to the canonical filtration $\mathcal{F}_n$. For $x \in \mathfrak{X}$ we set $D_x := D_{\{x\}}$. $\square$

Theorem 4.2.18. Let $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}$, $n \in \mathbb{N}_0$, be an HMC with initial distribution μ and transition matrix Q . Suppose that T is a stopping time adapted to the canonical filtration $(\mathcal{F}_n)_{n \in \mathbb{N}_0}$. Conditional on $X_T = x \in \mathfrak{X}$ and $T < \infty$ the stochastic process

$$Y_n := X_{T+n}, \quad n \in \mathbb{N}_0,$$

is in $\text{Markov}(\mathfrak{X}, \delta_x, Q)$ and independent of $\mathcal{F}_T$. More explicitly, if Λ is the event

$$\Lambda = \{T < \infty, X_T = x\},$$

and $\mathbb{P}_\Lambda : \mathcal{S} \rightarrow [0, 1]$ is the probability measure $\mathbb{P}_\Lambda[S] = \mathbb{P}[S | \Lambda]$, then the stochastic process

$$Y_n : (\Omega, \mathcal{S}, \mathbb{P}_\Lambda) \rightarrow \mathfrak{X}$$

is $\text{Markov}(\mathfrak{X}, \delta_x, Q)$ and independent of $\mathcal{F}_T$.

Proof. For $n \in \mathbb{N}$ denote by T_n the stopping time $T_n := T + n$. Denote by S the event

$$\begin{aligned} S &:= \{T < \infty\} \cap \{X_T = x_0 = x, X_{T+1} = x_1, \dots, X_{T+n} = x_n\} \\ &= \Lambda \cap \{X_{T+1} = x_1, \dots, X_{T+n} = x_n\}. \end{aligned}$$

Note that $S \in \mathcal{F}_{T_n}$. We have to show that

$$\mathbb{P}_\Lambda[Y_{T+n+1} = x_{n+1} | S] = Q_{x_n, x_{n+1}},$$

i.e.,

$$\frac{\mathbb{P}_\Lambda[\{X_{T+n+1} = x_{n+1}\} \cap S \cap \Lambda]}{\mathbb{P}_\Lambda[S]} = Q_{x_n, x_{n+1}}.$$

or

$$\frac{\mathbb{P}[\{X_{T+n+1} = x_{n+1}\} \cap S \cap \Lambda]}{\mathbb{P}[S \cap \Lambda]} = \frac{\mathbb{P}[\{X_{T+n+1} = x_{n+1}\} \cap S]}{\mathbb{P}[S]} = Q_{x_n, x_{n+1}}.$$

We have

$$\mathbb{P}[\{X_{T+n+1} = x_{n+1}\} \cap S] = \sum_{k=0}^{\infty} \mathbb{P}[\{X_{k+n+1} = x_{n+1}\} \cap S \cap \{T = k\}]$$

(use the Markov property (4.1.2) and the definition of S)

$$\begin{aligned} &= \sum_{k=0}^{\infty} \mathbb{P}[\{T = k\} \cap S] Q_{x_n, x_{n+1}} \\ &= Q_{x_n, x_{n+1}} \sum_{k=0}^{\infty} \mathbb{P}[\{T = k\} \cap S] = Q_{x_n, x_{n+1}} \mathbb{P}[S]. \end{aligned}$$

To prove the independence of $\mathcal{F}_T$ given $\Lambda = \{X_T = x, T < \infty\}$ it suffices to show that each of the events

$$\Gamma_0 = \{X_T = x\}, \dots, \Gamma_n = \{X_T = x_0 = x, X_{T+1} = x_1, \dots, X_{T+n} = x_n\}, \dots$$

are independent of $\mathcal{F}_T$ given $\Lambda = \{X_T = x, T < \infty\}$. Let $S \in \mathcal{F}_T$. We have

$$\mathbb{P}[S \cap \Gamma_n \cap \Lambda] = \sum_{k=0}^{\infty} \mathbb{P}[S \cap \Gamma_n \cap \{T = k\}]$$

(use the Markov property repeatedly)

$$\begin{aligned} &= \sum_{k=0}^{\infty} \mathbb{P}[\{T = k\} \cap S \cap \Gamma_0] Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n} \\ &= \mathbb{P}[S \cap \Gamma_0 \cap \{T < \infty\}] Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n} = \mathbb{P}[S \cap \Lambda] Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n}, \end{aligned}$$

i.e.,

$$\mathbb{P}[S \cap \Gamma_n \cap \Lambda] = \mathbb{P}[S \cap \Lambda] Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n}.$$

Then

$$\mathbb{P}[S \cap \Gamma_n | \Lambda] = \frac{\mathbb{P}[S \cap \Lambda]}{\mathbb{P}[\Lambda]} \cdot Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n}, \quad x_0 = x.$$

Since the stochastic process $Y_n : (\Omega, \mathcal{S}, \mathbb{P}_\Lambda) \rightarrow \mathfrak{X}$ is Markov($\mathfrak{X}, \delta_x, Q$) we deduce

$$Q_{x_0, x_1} \cdots Q_{x_{n-1}, x_n} = \mathbb{P}[\Gamma_n | \Lambda].$$

Hence $\mathbb{P}[S \cap \Gamma_n | \Lambda] = \mathbb{P}[S | \Lambda] \cdot \mathbb{P}[\Gamma_n | \Lambda]$. □

In the following subsections we will have plenty of opportunities to see the strong Markov principle at work.

4.2.3. Transience and recurrence. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q . For any $x \in \mathfrak{X}$ we denote by T_x the return time to x , i.e.,

$$T_x := \min\{n \geq 1; X_n = x\}.$$

Moreover,

$$\mathbb{P}_x := \mathbb{P}[- | X_0 = x], \quad \mathbb{E}_x := \mathbb{E}[- | X_0 = x].$$

Definition 4.2.19. A state $x \in \mathfrak{X}$ is called *recurrent* or *persistent* if $\mathbb{P}_x[T_x < \infty] = 1$. Otherwise it is called *transient*. □

Example 4.2.20. If $\mathfrak{X}$ is *finite* and irreducible, then any state of $\mathfrak{X}$ is recurrent. Indeed the ‘sooner-rather-than-later’ Lemma 3.1.34 implies that $\mathbb{E}_x[T_x] < \infty, \forall x \in \mathfrak{X}$. $\square$

We set $T_x^1 := T_x$ and we define inductively

$$T_x^{k+1} := \min \{ n > T_x^k; X_n = x \},$$

$$N_x := \# \{ k \in \mathbb{N}; T_x^k < \infty \} = \# \{ n \in \mathbb{N}; X_n = x \} \in \mathbb{N} \cup \{\infty\}.$$

Thus T_x^k is the time of the k -th return to x . We will refer to N_x *the number of returns to x* .

We set

$$p = p_x := \mathbb{P}_x[T_x < \infty].$$

Lemma 4.2.21. For any $n \in \mathbb{N}_0$ we have $\mathbb{P}_x[N_x \geq n] = p^n$. In particular, if x is recurrent, i.e., $p = 1$, then $N_x = \infty$ a.s. and, if X is transient, then

$$\mathbb{E}_x[N_x] = \frac{p}{1-p}.$$

Proof. Set $p_n := \mathbb{P}_x[N_x \geq n]$. We will prove inductively that $p_n = p^n$.

Clearly $\mathbb{P}[N_x \geq 1] = \mathbb{P}[T_x^1 < \infty] = p$. Suppose that $p_n = p^n$. The post- T_x^n process $Y_n = X_{T_x^n+n}$ starts at x and the strong Markov property implies that it is a HMC with the same transition matrix. In particular, the probability that it returns to x is p . On the other hand, Y_n returns to x if and only if $N_x \geq n+1$. Since the post- T_x^n process is independent of $\mathcal{F}_{T_x^n}$ we deduce

$$\mathbb{P}[N_x \geq n+1] = p \mathbb{P}[N_x \geq n] = p^{n+1}.$$

$\square$

Corollary 4.2.22. Assume that $X_0 = x$ a.s.. Then the following hold

$$x \text{ is recurrent} \iff N_x = \infty \text{ a.s.} \iff \mathbb{E}_x[N_x] = \infty.$$

$$x \text{ is transient} \iff \mathbb{E}_x[N_x] < \infty.$$

$\square$

Clearly the recurrence/transience of a state depends only on the transition matrix. The next result characterizes these features in terms of the transition matrix

Theorem 4.2.23. Let $x \in \mathfrak{X}$. The following statements are equivalent.

- (i) The state x is recurrent.
- (ii)

$$\sum_{n \in \mathbb{N}} Q_{x,x}^n = \infty.$$

Proof. Observe that

$$N_x = \sum_{n \in \mathbb{N}} \mathbf{I}_{\{X_n = x\}}$$

and

$$\mathbb{E}_x[N_x] = \sum_{n \in \mathbb{N}} \mathbb{E}_x[\mathbf{I}_{\{X_n = x\}}] = \sum_{n \in \mathbb{N}} Q_{x,x}^n.$$

The result now follows from Corollary 4.2.22. $\square$

Corollary 4.2.24. *Let $x, y \in \mathfrak{X}$. If $x \rightarrow y$ and x is recurrent, then*

- (i) $x \leftrightarrow y$,
- (ii) $\mathbb{P}_y[T_x < \infty] = 1$,
- (iii) *the state y is recurrent.*

Proof. The state x is recurrent so $N_x = \infty$ a.s.. Since $x \rightarrow y$ we deduce that

$$\mathbb{P}_x[T_y < \infty] = \mathbb{P}[T_y < \infty | X_0 = x] > 0.$$

The post- T_y chain $Y_n = X_{n+T_y}$, $n \geq 0$, will almost surely reach x since $N_x = \infty$ a.s.. Using the strong Markov property at T_y we deduce that Y_n has the same transition matrix Q . Hence $y \rightarrow x$, i.e. $x \leftrightarrow y$. In particular, the original chain, started at y will almost surely reach x , i.e., $\mathbb{P}[T_x < \infty | X_0 = y] = 1$.

Since $x \leftrightarrow y$ there exist $j, k \in \mathbb{N}$ such that

$$c = \min\{Q_{x,y}^j, Q_{y,x}^k\} > 0.$$

We deduce

$$Q_{y,y}^{n+j+k} \geq Q_{y,x}^k Q_{x,x}^n Q_{x,y}^j \geq c^2 Q_{x,x}^n, \quad \forall n \in \mathbb{N}.$$

Hence

$$\sum_{m \geq 1} Q_{y,y}^m \geq \sum_{m > j+k} Q_{y,y}^m \geq c^2 \sum_{n \geq 1} Q_{x,x}^n = \infty.$$

□

The above result shows that if C is a communication class then, either all states in C are recurrent, or all states in C are transient. In the first case C is called a *recurrence class* and in the second case C is called a *transience class*.

An irreducible HMC consists of a single communication class. Accordingly *an irreducible HMC can be either transient, or recurrent.*

Proposition 4.2.25. *Suppose that $(X_n)_{n \geq 0}$ is an irreducible transient HMC with state space $\mathfrak{X}$, transition matrix Q and initial distribution μ . Then,*

$$\mathbb{E}_\mu[N_x] < \infty, \quad \forall x \in \mathfrak{X}.$$

Proof. We first prove that given $x_0 \in \mathfrak{X}$ there exists $C = C_{x_0} > 0$ such that

$$\mathbb{E}_y[N_{x_0}] \leq C, \quad \forall y \in \mathfrak{X}.$$

Indeed, using the strong Markov property as in the proof of Lemma 4.2.21 we deduce that for any $y \in \mathfrak{X}$ we have

$$\begin{aligned} \mathbb{E}_y[N_{x_0}] &= \sum_{n \geq 1} \mathbb{P}_y[N_{x_0} \geq n] = \sum_{n \geq 1} \mathbb{P}_{x_0}[N_{x_0} \geq n-1] \mathbb{P}_y[T_{x_0} < \infty] \\ &= \underbrace{\mathbb{P}_{x_0}[N_{x_0} \geq 0]}_{=1} \mathbb{P}_y[T_{x_0} < \infty] + \mathbb{P}_y[T_{x_0} < \infty] \underbrace{\sum_{m \geq 1} \mathbb{P}_{x_0}[N_{x_0} \geq m]}_{\mathbb{E}_{x_0}[N_{x_0}]} \\ &= \mathbb{P}_y[T_{x_0} < \infty] (1 + \mathbb{E}_{x_0}[N_{x_0}]) \leq \underbrace{1 + \mathbb{E}_{x_0}[N_{x_0}]}_{C_{x_0}}. \end{aligned}$$

Now observe that

$$\mathbb{E}_\mu[N_{x_0}] = \sum_{y \in \mathfrak{X}} \mu[y] \mathbb{E}_y[N_{x_0}] \leq C_{x_0}.$$

□

Theorem 4.2.26. *Suppose that C is a recurrence class and $X_0 = x \in C$ a.s.. Then,*

$$\mathbb{P}[N_y = \infty | X_0 = x] = 1, \quad \forall y \in C.$$

In particular, with probability one, the chain visits every state of C infinitely often, i.e.,

$$\forall y \in C, \quad \mathbb{P}[N_y = \infty, | X_0 \in C] = 1.$$

Proof. Let $x, y \in C$. We have

$$\mathbb{P}[N_x = \infty | X_0 = x] = 1, \quad \mathbb{P}[T_y < \infty | X_0 = x] = 1.$$

The strong Markov property shows that the post- T_y chain has the same distribution as the chain started at y . Since y is recurrent we have $\mathbb{P}[N_y = \infty | X_0 = x] = 1$ and we deduce that

$$\mathbb{P}[N_y < \infty | X_0 = x] = 0, \quad \forall x, y \in C,$$

$$\mathbb{P}[N_y < \infty | X_0 \in C] = 0, \quad \forall y \in C.$$

In particular,

$$\mathbb{P}[\exists y \in C, N_y < \infty | X_0 \in C] \leq \sum_{y \in C} \mathbb{P}[N_y < \infty | X_0 \in C] = 0.$$

Hence

$$\mathbb{P}[\forall y \in C, N_y = \infty, | X_0 \in C] = 1 - \mathbb{P}[\exists y \in C, N_y < \infty | X_0 \in C] = 1.$$

□

Proposition 4.2.27. *A recurrence communication class C is closed.*

Proof. Suppose that C is not closed. Then there exist states $c \in C$ and $x \in \mathfrak{X} \setminus C$ such that $c \rightarrow x$. Since C is a communication class x does not communicate with any $y \in C$. Fix $n_0 \in \mathbb{N}$ such that $p := \mathbb{P}[X_{n_0} = x | X_0 = c] > 0$. In particular, since x does not communicate with C we deduce that $\mathbb{P}[X_n \in \mathfrak{X} \setminus C, \forall n \geq n_0 | X_0 = c] \geq p$. In particular $\mathbb{P}[N_c < n_0 | X_0 = c] \geq p$. This contradicts the fact that $\mathbb{P}[N_c = \infty | X_0 = c] = 1$. □

The set of communication classes $\bar{\mathfrak{X}} := \mathfrak{X} / \leftrightarrow$ is itself the state space of an HMC with transition matrix

$$\bar{Q}_{C,C'} = \mathbb{P}[X_1 \in C' | X_0 \in C].$$

Each state of $\bar{\mathfrak{X}}$ is in itself a communication class of the new Markov chain. The state space $\bar{\mathfrak{X}}$ is partitioned into two types: transient states and recurrent states. The recurrent states are closed, i.e., they are absorbing as states in $\bar{\mathfrak{X}}$. Given a recurrent state $R \in \bar{\mathfrak{X}}$, no other communication class is accessible from R .

Example 4.2.28. Consider for example the gambler's ruin problem with total fortune $N \in \mathbb{N}$; see Example 4.1.6. This can be viewed as a Markov chain with state space $\{0, 1, \dots, N\}$ and transition probabilities

$$q_{i,i\pm 1} = \frac{1}{2}, \quad \forall 0 < i < N, \quad q_{0,0} = q_{N,N} = 1.$$

The communication classes of this Markov chain are

$$\{0\}, \{N\}, \{1, 2, \dots, N-1\}.$$

The first two are recurrent while the third is transient. $\square$

If $\mathcal{X}$ is finite, the argument in Example 4.2.20 shows that a communication class is closed iff it is recurrent.

Example 4.2.29 (G.Polya). (a) Consider the standard random walk on $\mathbb{Z}$. We denote by Q the transition matrix. This is an irreducible Markov chain and each state has period 2. To decide whether it is transient or recurrent it suffices to verify if the origin is such. Note that $Q_{0,0}^{2n-1} = 0, \forall n \in \mathbb{N}$. To compute $Q_{0,0}^{2n}$ we observe that a path of length $2n$ starts and ends at the origin if and only if it consists of exactly n steps to the right and n steps to the left. Since each such step occurs with probability $\frac{1}{2}$ we deduce

$$Q_{0,0}^{2n} = \frac{1}{2^{2n}} \binom{2n}{n} = \frac{(2n)!}{2^{2n}(n!)^2}.$$

Using Stirling's formula (A.1.7) we deduce that, as $n \rightarrow \infty$, we have

$$\frac{(2n)!}{2^{2n}(n!)^2} \sim \frac{\sqrt{4\pi n}}{2\pi n} \sim \frac{1}{\sqrt{\pi n}},$$

so

$$\sum_{n \in \mathbb{N}} Q_{0,0}^n = \infty.$$

Thus, the 1-dimensional standard random walk is recurrent.

(b) Consider the standard walk on $\mathbb{Z}^2$. It is irreducible. We want to decide if the origin is recurrent or transient. To compute $Q_{0,0}^{2n}$ we observe that a path of length $2n$ starts and ends at the origin if and only if the number of steps up is equal to the number of steps down and the number of steps to the right is equal to the number of steps to the left. We deduce that

$$Q_{0,0}^{2n} = \sum_{k=0}^n \frac{(2n)!}{4^{2n}(k!)^2((n-k)!)^2} = \frac{(2n)!}{4^{2n}(n!)^2} \sum_{k=0}^n \binom{n}{k}^2.$$

Using Newton's binomial formula in the equality

$$(x+y)^{2n} = (x+y)^n(x+y)^n$$

and identifying the coefficients of $x^n y^n$ on either side of the above equality we deduce

$$\binom{2n}{n} = \sum_{k=0}^n \binom{n}{k} \binom{n}{n-k} = \sum_{k=0}^n \binom{n}{k}^2,$$

so that

$$Q_{0,0}^{2n} = \left(\frac{1}{2^{2n}} \binom{2n}{n} \right)^2 \sim \frac{1}{\pi n} \text{ as } n \rightarrow \infty.$$

Hence, again

$$\sum_{n \in \mathbb{N}} Q_{0,0}^n = \infty,$$

so the standard 2-dimensional random walk is also recurrent.

(c) Consider the standard random walk on $\mathbb{Z}^3$. Arguing as in the 2-dimensional case we deduce

$$Q_{0,0}^{2n} = \frac{1}{6^{2n}} \sum_{j+k+\ell=n} \frac{(2n)!}{(j!)^2(k!)^2(\ell!)^2} = \frac{1}{2^{2n}} \binom{2n}{n} \sum_{j+k+\ell=n} \left(\frac{n!}{j!k!\ell!3^n} \right)^2.$$

Now observe that

$$\sum_{j+k+\ell=n} \underbrace{\frac{n!}{j!k!\ell!3^n}}_{=: p_{j,k,\ell}} = \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} \right)^n = 1.$$

Hence

$$\sum_{j,k,\ell} p_{j,k,\ell}^2 \leq \max_{j,k,\ell} p_{j,k,\ell} \sum_{j,k,\ell} p_{j,k,\ell} = \max_{j,k,\ell} p_{j,k,\ell},$$

so

$$Q_{0,0}^{2n} \leq \frac{1}{2^{2n}} \binom{2n}{n} \max_{j,k,\ell} p_{j,k,\ell}.$$

Let us observe that the maximum value of $p_{j,k,\ell}$ is achieved when j, k, ℓ are as close to $n/3$ as possible. Indeed, if $j \leq k \leq \ell$, $j < \ell$, then

$$(j+1)!(\ell-1)! = \frac{j+1}{\ell} j!\ell! \leq j!\ell!$$

so

$$p_{j+1,k,\ell-1} \geq p_{j,k,\ell}.$$

Assume now that $n = 3m$. We deduce

$$Q_{0,0}^{2n} \leq \frac{1}{2^{2n}} \binom{2n}{n} \frac{(3m)!}{(m!)^3 3^{3m}}.$$

Using again Stirling's formula we deduce that, as $m \rightarrow \infty$ we have

$$\frac{(3m)!}{(m!)^3 3^{3m}} \sim \frac{\sqrt{6\pi m}}{(2\pi m)^{3/2}} = \frac{\sqrt{3}}{2\pi m}.$$

On the other hand,

$$\frac{1}{2^{2n}} \binom{2n}{n} \sim \frac{1}{\sqrt{\pi n}} = \frac{1}{\sqrt{3\pi m}}.$$

We deduce that

$$Q_{0,0}^{6m} = O(m^{-3/2}) \text{ as } m \rightarrow \infty.$$

Arguing in a similar fashion we deduce

$$Q_{0,0}^{6m+2}, Q_{0,0}^{6m+4} = O(m^{-3/2}) \text{ as } m \rightarrow \infty.$$

We conclude that

$$\sum_{n \in \mathbb{N}} Q_{0,0}^n = \sum_{n \in \mathbb{N}} Q_{0,0}^{2n} < \infty,$$

so the standard 3-dimensional random walk is transient!

□

4.2.4. Invariant measures. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q . We will identify a sigma-finite measure λ on $\mathfrak{X}$ with a function

$$\lambda : \mathfrak{X} \rightarrow [0, \infty], \quad x \mapsto \lambda_x = \lambda[\{x\}].$$

Definition 4.2.30. An *invariant* or *stationary measure* for the HMC $(X_n)_{n \in \mathbb{N}_0}$ is a σ -finite measure λ on $\mathfrak{X}$ such that $\lambda = \lambda Q$, i.e.,

$$\lambda_x = \sum_{y \in \mathfrak{X}} \lambda_y Q_{y,x}, \quad \forall x \in \mathfrak{X}. \quad (4.2.3)$$

An *invariant* or *stationary distribution* is an invariant *probability* measure. $\square$

Example 4.2.31 (Time reversal). Suppose that π is an invariant *probability* distribution for $(X_n)_{n \geq 0}$ such that $\pi_x > 0, \forall x \in \mathfrak{X}$. Suppose additionally that

$$\mathbb{P}[X_0 = x] = \pi_x, \quad \forall x \in \mathfrak{X}.$$

Then

$$\mathbb{P}[X_1 = x] = \sum_{y \in \mathfrak{X}} \mathbb{P}[X_1 = x | X_0 = y] \pi_y = \sum_y \pi_y Q_{y,x} = \pi_x.$$

Iterating we deduce that the random variables $(X_n)_{n \in \mathbb{N}_0}$ are identically distributed. For $x, y \in \mathfrak{X}$ we set

$$R_{y,x} := \mathbb{P}[X_0 = x | X_1 = y] = \frac{\mathbb{P}[X_1 = y | X_0 = x] \mathbb{P}[X_0 = x]}{\mathbb{P}[X_1 = y]} = \frac{\pi_x}{\pi_y} Q_{x,y}.$$

Note that for every $x, y \in \mathfrak{X}$ we have

$$\sum_x R_{y,x} = \frac{1}{\pi_y} \sum_x \pi_x Q_{x,y} = 1,$$

so $(R_{y,x})_{x,y \in \mathfrak{X}}$ is a stochastic matrix describing the so called *time reversed chain*.

Suppose now that π is a probability distribution on $\mathfrak{X}$ such that $\pi_x > 0, \forall x \in \mathfrak{X}$ and satisfying

$$Q_{y,x} = R_{y,x} = \frac{\pi_x}{\pi_y} Q_{x,y}, \quad \forall x, y \in \mathfrak{X}. \quad (4.2.4)$$

From the equality

$$1 = \sum_x Q_{y,x} = \sum_x \frac{\pi_x}{\pi_y} Q_{x,y}$$

we deduce that π is a stationary distribution and the time reversed chain coincides with the initial chain. This is the reason why the chains satisfying (4.2.4) are called *reversible*. $\square$

Definition 4.2.32. An irreducible HMC with state space $\mathfrak{X}$ and transition matrix is called *reversible* if there exists a function $\lambda : \mathfrak{X} \rightarrow (0, \infty)$ satisfying satisfying the *detailed balance equations*

$$\lambda_y Q_{y,x} = \lambda_x Q_{x,y}, \quad \forall x, y \in \mathfrak{X}. \quad (4.2.5)$$

$\square$

Observe that if Q satisfies the detailed balance equation, then an argument as in Example 4.2.31 shows λ defines a Q -invariant measure

Example 4.2.33. (a) If $Q_{x,y} = Q_{y,x}$ for any $x, y \in \mathfrak{X}$, then the corresponding chain is reversible and any uniform measure on X is invariant. This happens for example if $(X_n)_{n \geq 0}$ describes the standard random walk on $\mathbb{Z}^d$.

(b) In the case the standard random walk on a locally finite connected graph we have

$$Q_{x,y} = \frac{1}{\deg x}, \quad \deg x \cdot Q_{x,y} = 1 = \deg y \cdot Q_{y,x}.$$

Hence Q is in detailed balance with invariant measure $x \mapsto \deg x$. If, additionally, $\mathfrak{X}$ is finite, then the probability measure

$$\pi_x = \frac{\deg x}{\sum_y \deg y}$$

is invariant. □

Example 4.2.34 (The Ehrenfest urn). Consider the Ehrenfest urn model detailed in Example 4.1.7. We recall that the state space is $\mathfrak{X} = \{0, 1, \dots, B\}$, $B \in \mathbb{N}$ and the only nontrivial transition probabilities are

$$Q_{k,k+1} = \frac{B-k}{B}, \quad Q_{k,k-1} = \frac{k}{B}.$$

Note that

$$\frac{Q_{k,k+1}}{Q_{k+1,k}} = \frac{B-k}{k+1} = \frac{\binom{B}{k+1}}{\binom{B}{k}}$$

Then the measure $k \rightarrow \lambda_k = \binom{B}{k}$ is invariant and

$$\pi_k = \frac{1}{2^B} \binom{B}{k}, \quad k = 0, 1, \dots, B,$$

so $\pi = \text{Bin}(n, 1/2)$ is the invariant probability distribution. □

Example 4.2.35 (Bernoulli-Laplace urn). Consider the Bernoulli-Laplace urn introduced in Example 4.1.8. Then

$$\frac{Q_{k,k+1}}{Q_{k+1,k}} = \left(\frac{N-k}{k+1} \right)^2 = \frac{\binom{N}{k+1}^2}{\binom{N}{k}^2},$$

and thus the function

$$\lambda : \{0, 1, \dots, B\} \rightarrow \mathbb{R}, \quad \lambda[k] = \binom{N}{k}^2$$

defines an invariant measure. Using the identity²

$$\sum_{j=0}^N \binom{N}{j}^2 = \binom{2N}{N}$$

we deduce that

$$k \rightarrow \pi_k = \frac{\binom{N}{k}^2}{\binom{2N}{N}}$$

is the invariant *probability* measure. Note that $\pi = \text{HGeom}(N, N; N)$; see Example 1.3.30. □

²To prove this identity look at the set of East-North zig-zags from $(0, 0)$ to (B, B) and partition it according to where a zig-zag intersects the line $x + y = B$.

Theorem 4.2.36. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an irreducible and recurrent HMC with state space $\mathfrak{X}$ and transition matrix Q . Assume $X_0 = x_0 \in \mathfrak{X}$, and denote by T_0 the time of first return to x_0 , i.e.,

$$T_0 := T_{x_0} = \min \{n \geq 1; X_n = x_0\}.$$

For any $x \in \mathfrak{X}$, define

$$N_x := \sum_{n \in \mathbb{N}} \mathbf{I}_{\{X_n = x\}} \mathbf{I}_{\{n \leq T_0\}} = \sum_{n=1}^{T_0} \mathbf{I}_{\{X_n = x\}}, \quad (4.2.6a)$$

$$\lambda_x = \lambda_{x, x_0} := \begin{cases} \mathbb{E}_{x_0} [N_x], & x \neq x_0, \\ 1, & x = x_0. \end{cases} \quad (4.2.6b)$$

In other words, λ_x is the expected number of visits to x before returning to x_0 when starting from x_0 . Then, the following hold.

- (i) $\lambda_x \in (0, \infty)$, $\forall x \in \mathfrak{X}$ and the associated measure λ on $\mathfrak{X}$, given by

$$\lambda[\{x\}] = \lambda_x, \quad \forall x \in \mathfrak{X}$$

is invariant.

- (ii) $\lambda[\mathfrak{X}] = \mathbb{E}_{x_0} [T_{x_0}]$.

- (iii) The measure λ is the unique invariant measure such that $\lambda_{x_0} = 1$.

Proof. (i) I follow the approach in [27, Thm. 3.2.1]. Clearly $\lambda_{x_0} = 1$. For $x \in \mathfrak{X} \setminus \{x_0\}$ and $n \in \mathbb{N}$ we set

$$p_x(n) := \mathbb{P}[X_n = x, n \leq T_0].$$

Thus, $p_x(n)$ is the probability of visiting state x at time n before returning to x_0 . The equality (4.2.6a) implies that

$$\lambda_x = \sum_{n \in \mathbb{N}} p_x(n), \quad \forall x \neq x_0. \quad (4.2.7)$$

Let us prove that λ is invariant, i.e., satisfies (4.2.3). Observe first that $p_x(1) = Q_{x_0, x}$. From the Markov property we deduce

$$p_x(n) = \sum_{y \neq x_0} p_y(n-1) Q_{y, x}. \quad (4.2.8)$$

Hence

$$\lambda_x \stackrel{(4.2.7)}{=} \sum_{n \in \mathbb{N}} p_x(n) = p_x(1) + \sum_{y \neq x_0} \underbrace{\left(\sum_{n \in \mathbb{N}} p_y(n) \right)}_{=\lambda_y} Q_{y, x}$$

$$(\lambda_{x_0} = 1, p_x(1) = Q_{x_0, x})$$

$$= \lambda_0 Q_{x_0, x} + \sum_{y \neq x_0} \lambda_y Q_{y, x} = \sum_y \lambda_y Q_{y, x}.$$

This proves (4.2.3).

Let us prove that $\lambda_x < \infty$, $\forall x$. Observe that

$$1 = \lambda_{x_0} = \sum_{x \in \mathfrak{X}} \lambda_x Q_{x, x_0}^n. \quad (4.2.9)$$

Suppose that $\lambda_x = \infty$ for some $x \neq x_0$. Since the chain is irreducible, the state x_0 communicates with x so there exists $n = n(x)$ such that $Q_{x_0,x}^n \neq 0$. The equality (4.2.9) implies $\lambda_x \leq \frac{1}{Q_{x,x_0}^n}$.

Finally, let us now show that the numbers λ_x defined by (4.2.6b) are *positive*. Suppose that $\lambda_x = 0$ for some $x \in \mathfrak{X}$. Obviously $x \neq x_0$. Moreover, from the equality $\lambda = \lambda Q^n$, $\forall n \in \mathbb{N}$ we deduce

$$0 = \lambda_x = Q_{x_0,x}^n + \sum_{y \neq x_0} \lambda_y Q_{y,x}^n.$$

Thus $Q_{x_0,x}^n = 0$, $\forall n \in \mathbb{N}$, which contradicts the fact that x_0 and x communicate.

(ii) We have

$$\sum_{x \in \mathfrak{X}} \sum_{n \geq 1} I_{\{X_n=x\}} I_{\{n \leq T_0\}} = \sum_{n \geq 1} \sum_{x \in \mathfrak{X}} I_{\{X_n=x\}} I_{\{n \leq T_0\}} = \sum_{x \in \mathfrak{X}} I_{\{n \leq T_0\}} = T_0.$$

Hence

$$\lambda[\mathfrak{X}] = \sum_{x \in \mathfrak{X}} \lambda_x = \sum_{x \in \mathfrak{X}} \sum_{n \geq 1} \mathbb{E}_{x_0} [I_{\{X_n=x\}} I_{\{n \leq T_0\}}] = \mathbb{E}_{x_0} [T_0].$$

(iii) We follow the approach in [2, 160]. Consider the matrix $K : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1]$

$$K_{x,y} = \begin{cases} Q_{x,y}, & y \neq x_0, \\ 0, & y = x_0. \end{cases}$$

We obtain the sequence $(\mu_n)_{n \geq 0}$ of measures on $\mathfrak{X}$ defined by

$$\mu_0 = \delta_{x_0}, \quad \mu_n[x] = p_x(n) = \mathbb{P}[X_n = x, \quad n < T_0], \quad x \in \mathfrak{X}.$$

Note that $\mu_n[x_0] = 0$, $\forall n$. The equality (4.2.8) implies that

$$\mu_n = \mu_{n-1}K, \quad \forall n \geq 1$$

so $\mu_n = \delta_{x_0}K^n$. Observe that

$$\lambda = \sum_{n \geq 0} \mu_n = \sum_{n \geq 0} \delta_{x_0}K^n.$$

Fix an invariant measure ν such that $\nu_{x_0} = 1$. The invariance condition reads $\nu = \delta_{x_0} + \nu K$. We deduce

$$\nu = \delta_{x_0} + (\delta_{x_0} + \nu K)K = \delta_{x_0} + \delta_{x_0}K + \nu K^2.$$

Arguing inductively we deduce

$$\nu = \sum_{m=0}^n \delta_{x_0}K^m + \nu K^{n+1} \geq \sum_{m=0}^n \delta_{x_0}K^m, \quad \forall n \in \mathbb{N}.$$

Letting $n \rightarrow \infty$ we deduce $\nu \geq \mu$. The difference $\sigma = \nu - \mu$ is also an invariant measure such that $\sigma[x_0] = 0$. Set $\sigma_x := \sigma[x]$.

Fix $x \in \mathfrak{X} \setminus \{x_0\}$. The Markov chain is irreducible so there exists $n \in \mathbb{N}$ such that $Q_{x,x_0}^n > 0$. From the equality $\sigma = \sigma Q^n$ we deduce

$$0 = \sigma_{x_0} = \sigma_x Q_{x,x_0}^n + \sum_{y \neq x} \sigma_y Q_{y,x_0}^n \geq \sigma_{x_1} Q_{x,x_0}^n.$$

Hence $\sigma_x = 0$, $\forall x \in \mathfrak{X}$ so that $\nu = \lambda$.

□

Remark 4.2.37. The example of the standard random walk on $\mathbb{Z}^3$ shows that even transient chains can admit invariant measures. $\square$

Suppose that $(X_n)_{n \geq 0}$ is irreducible and recurrent. For each $x \in \mathfrak{X}$ we denote by π^x the unique invariant measure on $\mathfrak{X}$ such that $\pi^x[x] = 1$. We know that for $x, y \in \mathfrak{X}$ the measure π^y is a positive multiple of π^x ,

$$\pi^y = c_{y,x} \pi^x.$$

From the equality $\pi^x[x] = 1$ we deduce $c_{y,x} = \pi^y[x]$ so that

$$\pi^y = \pi^y[x] \pi^x. \quad (4.2.10)$$

From Theorem 4.2.36(ii) we deduce that

$$\pi^x[\mathfrak{X}] = \mathbb{E}_x[T_x]. \quad (4.2.11)$$

In particular this shows that the following statements are equivalent

- (i) $\exists x \in \mathfrak{X}$ such that $\mathbb{E}_x[T_x] < \infty$.
- (ii) $\forall x \in \mathfrak{X}, \mathbb{E}_x[T_x] < \infty$.

Definition 4.2.38. Let $(X_n)_{n \geq 0}$ be an irreducible recurrent HMC.

- (i) The chain is called *positively recurrent* if $\mathbb{E}_x[T_x] < \infty$ for some $x \in \mathfrak{X}$.
- (ii) The chain is called *null recurrent* if $\mathbb{E}_x[T_x] = \infty$ for all $x \in \mathfrak{X}$.

$\square$

Corollary 4.2.39. An irreducible recurrent HMC is positively recurrent if for some (for all) $x \in \mathfrak{X}$ we have

$$\pi^x[\mathfrak{X}] < \infty. \quad (4.2.12)$$

In particular, a positively recurrent irreducible HMC admits a unique invariant probability measure π_∞ described by

$$\pi_\infty = \frac{1}{\mathbb{E}_x[T_x]} \pi^x, \quad \forall x \in \mathfrak{X}.$$

In other words,

$$\pi_\infty[x] = \frac{1}{\mathbb{E}_x[T_x]}, \quad \forall x \in \mathfrak{X}. \quad (4.2.13)$$

$\square$

Proof. The equality (4.2.13) follows from the equality $\pi^x[x] = 1$. $\square$

Corollary 4.2.40. Any irreducible HMC with finite state space $\mathfrak{X}$ admits a unique stationary probability measure. $\square$

Proof. As shown in Example 4.2.20 this chain is recurrent since the state space is finite. The finiteness of $\mathfrak{X}$ implies (4.2.12). $\square$

Example 4.2.41 (Random knight moves). Consider a regular 8×8 chess table and a knight that starts in the lower left-hand corner and then moves randomly along, making each permissible move with equal probability.

This is an example of random walk on a graph with vertex set $\mathfrak{X}$ consisting of the centers of the 64 squares of the board, where two vertices are connected by as many edges as possibilities for the knight to go from one square to the other in one move.

It is easily seen that this is a connected graph so the corresponding random walk is irreducible and has a unique invariant probability distribution given by

$$\pi[x] = \frac{\deg x}{Z}, \quad Z = \sum_{y \in \mathfrak{X}} \deg y.$$

Now observe that Z is twice the number of edges of this graph.

To count the number of edges observe that each 2×3 sub-rectangle of the chess table determines four edges in this graph, two for each diagonal. The same is true for the 3×2 rectangle. Moreover, any knight move corresponds to a such rectangle. If $N_{2 \times 3}$ and respectively $N_{3 \times 2}$ denote the number of 2×3 rectangles respectively 3×2 rectangles, then

$$N_{2 \times 3} = N_{3 \times 2} =: N.$$

There are two diagonals per rectangle, and $2N$ rectangles so there are $4N$ edges. Hence $Z = 8N$. Now observe that since a 3×2 rectangle is uniquely determined by the location of its lower-left corner we have $N = N_{3 \times 2} = 6 \times 7 = 42$ so $Z = 8 \cdot 42$.

If x_0 corresponds to the left-hand corner square of the chess table, then $\deg x_0 = 2$ so

$$\mathbb{E}_{x_0}[T_{x_0}] = \frac{1}{\pi[x_0]} = \frac{8 \cdot 42}{2} = 4 \cdot 42 = 168.$$

Thus, given that the knight starts at x_0 , the expected time to return to x_0 is 168. $\square$

Theorem 4.2.42. Suppose that (X_n) is an irreducible HMC with state space $\mathfrak{X}$ and transition matrix Q . Then the following are equivalent.

- (i) The chain is positively recurrent.
- (ii) There exists an invariant probability measure.

Proof. We have already shown that (i) $\Rightarrow$ (ii). To prove the implication (ii) $\Rightarrow$ (i) fix an invariant probability measure π . Thus $\pi = Q^n \pi, \forall n \in \mathbb{N}$, so that

$$\pi[y] = \sum_{x \in \mathfrak{X}} \pi[x] Q_{x,y}^n, \quad \forall y \in \mathfrak{X}, \quad n \in \mathbb{N} \quad (4.2.14)$$

Fix $y_0 \in \mathfrak{X}$ such that $\pi[y_0] \neq 0$. We prove first that if the chain is recurrent, then it has to be positively recurrent. Denote by λ_{y_0} the unique invariant measure such that $\lambda_{y_0}[y_0] = 1$. The measure λ_{y_0} is a constant multiple of π so it is finite. Hence

$$\lambda_{y_0}[\mathfrak{X}] = \mathbb{E}_{y_0}[T_{y_0}] = \frac{1}{\pi[y_0]} < \infty,$$

showing that the chain in fact positively recurrent.

We now argue by contradiction that the chain is indeed recurrent. Assume that our Markov chain is transient. Proposition 4.2.25 implies that for any $x \in \mathfrak{X}$ we have

$$\infty > \mathbb{E}_\pi[N_x] = \sum_n \mathbb{E}_\pi[\mathbf{I}_{N_x=n}] = \sum_n \sum_{x' \in \mathfrak{X}} Q_{x,x'}^n \pi[x'] \geq \pi[y_0] \sum_n Q_{x,y_0}^n.$$

Hence

$$\sum_n Q_{x,y_0}^n < \infty \text{ and } \lim_{n \rightarrow \infty} Q_{x,y_0}^n = 0, \quad \forall x \in \mathfrak{X}. \quad (4.2.15)$$

Set $q_n(x) = Q_{x,y_0}^n, \forall x \in \mathfrak{X}$. The equality (4.2.14) implies that

$$\int_{\mathfrak{X}} q_n(x) \pi[dx] = \pi[y_0] > 0, \quad \forall n \in \mathbb{N}$$

On the other hand, the equality (4.2.15) coupled with the Dominated Convergence theorem implies

$$\lim_{n \rightarrow \infty} \int_{\mathfrak{X}} q_n(x) \pi[dx] = 0.$$

This contradiction completes the proof. $\square$

Example 4.2.43. We have shown in Example 4.2.29 that the standard random walks on $\mathbb{Z}$ and $\mathbb{Z}^2$ are recurrent. Let us show that they are null recurrent.

Note that for $k = 1, 2$, the measure on $\mathbb{Z}^k$ defined by $\lambda[x] = 1, \forall x \in \mathbb{Z}^k$ is invariant. By Theorem 4.2.36, λ is the unique invariant measure such that $\lambda[0] = 1$. Since $\lambda[\mathbb{Z}^k] = \infty$ we deduce that there is no invariant finite measure. $\square$

Proposition 4.2.44. Suppose that $(X_n)_{n \geq 0}$ is an irreducible, positively recurrent HMC with state space $\mathfrak{X}$ and transition matrix Q . Then

$$\mathbb{E}_y[T_x] < \infty, \quad \forall x, y \in \mathfrak{X}.$$

Proof. Fix $x \in \mathfrak{X}$. Set

$$\mathcal{Y} := \{y \in \mathfrak{X}; \mathbb{E}_y[T_x] < \infty\}.$$

Note that $x \in \mathcal{Y}$.

For $y \in \mathcal{Y}$ we have

$$\mathbb{E}_y[T_x] = \mathbb{E}_y[T_x | X_1 = y] Q_{y,y} + \sum_{z \neq y} \mathbb{E}_y[T_x | X_1 = z] Q_{y,z}.$$

Hence

$$\forall z \in \mathfrak{X}, y \neq z, Q_{y,z} > 0 \Rightarrow \mathbb{E}_y[T_x | X_1 = z] < \infty$$

Now observe that for $z \neq y$

$$\mathbb{E}_y[T_x | X_1 = z] = \begin{cases} 1, & z = x, \\ 1 + \mathbb{E}_z[T_x], & z \neq x. \end{cases}$$

We deduce that $y \in \mathfrak{X}$ and $Q_{y,z} > 0$, then $z \in \mathcal{Y}$. We conclude iteratively that

$$\forall y \in \mathcal{Y}, \forall z \in \mathfrak{X}, y \rightarrow z \Rightarrow z \in \mathcal{Y}.$$

Since the chain is irreducible we deduce $\mathcal{Y} = \mathfrak{X}$. $\square$

Let $(X_n)_{n \geq 0}$ be an HMC with state space and transition matrix Q . Recall that for any set $A \subset \mathfrak{X}$ we denoted by T_A the time of first return to A

$$T_A := \inf \{ n \geq 1; X_n \in A \}.$$

Note that $T_A \leq T_a, \forall a \in A$, so

$$\mathbb{E}_x[T_A] \leq \mathbb{E}_x[T_a], \quad \forall x \in \mathfrak{X}, \quad \forall a \in A.$$

We deduce that if the chain is irreducible and positively recurrent then

$$\mathbb{E}_x[T_A] < \infty, \quad \forall x \in \mathfrak{X}, \quad \forall A \subset \mathfrak{X}.$$

We have a sort of converse.

Proposition 4.2.45. *Suppose that $(X_n)_{n \geq 0}$ is an irreducible HMC with state space $\mathfrak{X}$ and transition matrix Q . If there exists a finite subset $A \subset \mathfrak{X}$ such that*

$$\mathbb{E}_a[T_A] < \infty, \quad \forall a \in A,$$

then $(X_n)_{n \geq 0}$ is positively recurrent.

Proof. I follow the approach in [27, Chap.7, Sec. 1.1]. Set

$$M_A := \max_{a \in A} \mathbb{E}_a[T_A].$$

Consider the epochs of return to A ,

$$T^1 := T_A, \quad T^{k+1} := \min \{ n > T^k; X_n \in A \}.$$

Fix $a_0 \in A$ and suppose that $(X_n)_{n \geq 0}$ starts at $a_0, X_0 = a_0$ a.s.. We set

$$Y_0 := X_0, \quad Y_k := X_{T^k}, \quad k \in \mathbb{N}.$$

The strong Markov property shows that $(Y_k)_{k \geq 0}$ is an HMC with state space A . Since (X_n) is irreducible we deduce that $(Y_k)_{k \geq 0}$ is such. Since A is finite, the chain (Y_k) is positively recurrent. Denote by $\widehat{T}_0$ the time of first return to a_0 of the chain $(Y_k)_{k \geq 0}$. Set

$$S_0 = T^1, \quad S_k = T^{k+1} - T^k.$$

If T_{a_0} denotes the time of first return to a_0 of the original chain, then

$$T_{a_0} = \sum_{k=0}^{\infty} S_k \mathbf{I}_{\{k < \widehat{T}_0\}}, \quad \mathbb{E}_{a_0}[T_{a_0}] = \sum_{k=0}^{\infty} \mathbb{E}_{a_0}[S_k \mathbf{I}_{\{k < \widehat{T}_0\}}].$$

On the other hand,

$$\mathbb{E}_{a_0}[S_k \mathbf{I}_{\{k < \widehat{T}_0\}}] = \sum_{a \in A} \mathbb{E}_{a_0}[S_k \mathbf{I}_{\{k < \widehat{T}_0\}} \mathbf{I}_{\{X_{T^k} = a\}}].$$

Observe that the event $\{k < \widehat{T}_0\}$ belongs to $\mathcal{F}_{T^k}$. We deduce

$$\mathbb{E}_{a_0}[S_k \mathbf{I}_{\{k < \widehat{T}_0\}} \mathbf{I}_{\{X_{T^k} = a\}}] = \mathbb{E}_{a_0}[S_k | k < \widehat{T}_0, X_{T^k} = a] \mathbb{P}_{x_0}[k < \widehat{T}_0, X_{T^k} = a]$$

(use the strong Markov property for T^k)

$$\begin{aligned} &= \mathbb{E}_{a_0}[S_k | X_{T^k} = a] \mathbb{P}_{x_0}[k < \widehat{T}_0, X_{T^k} = a] = \mathbb{E}_a[T_A] \mathbb{P}_{x_0}[k < \widehat{T}_0, X_{T^k} = a] \\ &\leq M_A \mathbb{P}_{a_0}[k < \widehat{T}_0, X_{T^k} = a]. \end{aligned}$$

Hence

$$\begin{aligned}\mathbb{E}_{a_0}[T_{a_0}] &\leq M_A \sum_{k=0}^{\infty} \left(\sum_{a \in A} \mathbb{P}_{x_0}[\hat{T}_0 > k, X_{T^k} = a] \right) \\ &= M_A \sum_{k=0}^{\infty} \mathbb{P}_{a_0}[\hat{T}_0 > k] = M_A \mathbb{E}_{a_0}[\hat{T}_0] < \infty,\end{aligned}$$

since the chain (Y_k) is positively recurrent. $\square$

4.3. Martingale techniques

This section is a side trip in our journey through the fundamentals of the theory of Markov chains. It could be skipped at the first reading. However, the theory of martingales is affording us the opportunity to reveal surprising applications involving Markov chains and I could not help sharing some of them.

The first subsection describes several useful criteria for deciding the nature of an irreducible chain: transient, null or positively recurrent. The second subsection is mostly an invitation to a optimal stopping theory, a theory with many modern applications.

4.3.1. Recurrence, transience and Lyapunov functions. Suppose that $(X_n)_{n \geq 0}$ is an HMC with state space $\mathfrak{X}$, transition matrix Q and initial distribution π_0 . We assume that all the random variables X_n are defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$\pi_n := \pi_0 \cdot Q^n, \quad \forall n \in \mathbb{N}.$$

We denote by $\mathcal{F}_n$ the filtration

$$\mathcal{F}_n = \sigma(X_0, X_1, \dots, X_n) \subset \mathcal{S}, \quad n \geq 0.$$

We want to investigate the (sub/super)martingales with respect to this filtration and show some of their applications to the dynamics of the underlying HMC.

Note that any function $\mathfrak{X} \rightarrow (-\infty, \infty]$ is measurable with respect to the sigma-algebra $2^{\mathfrak{X}}$. For this reason we will denote by $\mathcal{L}^0(\mathfrak{X})$ the space of functions $\mathfrak{X} \rightarrow (-\infty, \infty]$. We set

$$\mathcal{L}_*^0(\mathfrak{X}) := \{ f \in \mathcal{L}^0(\mathfrak{X});; f(x) < \infty, \quad \forall x \in \mathfrak{X} \}.$$

We think of a function $f \in \mathcal{L}^0(\mathfrak{X})$ as a *column* vector $(f(x))_{x \in \mathfrak{X}}$ and we denote by Qh the function described by the multiplication of the matrix Q with the column vector f . More precisely,

$$(Qf)(x) = \sum_{y \in \mathfrak{X}} Q_{x,y} f(y) \in (-\infty, \infty], \quad \forall x \in \mathfrak{X}.$$

There is a small problem with this definition namely, if $\mathfrak{X}$ is infinite, then the above series may be divergent. Since the rows of Q define probability distributions on $\mathfrak{X}$ we see that the above sums are finite if f is bounded or the transition matrix is *locally finite*, i.e., each of its rows has only finitely many nonzero entries. Equivalently, at each state $x \in \mathfrak{X}$ the system can transition only to finitely many states.

☞ In the remainder of this subsection, in order to keep the technicalities at bay, we will assume that Q is locally finite.

In this case Q defines a map

$$Q : \mathcal{L}^0(\mathfrak{X}) \rightarrow \mathcal{L}^0(\mathfrak{X}).$$

It restricts to a linear map

$$Q : \mathcal{L}_*^0(\mathfrak{X}) \rightarrow \mathcal{L}_*^0(\mathfrak{X}), \quad f \rightarrow Qf.$$

Note that

$$Q\mathbf{I}_{\mathfrak{X}} = \mathbf{I}_{\mathfrak{X}}, \quad f \geq 0 \Rightarrow Qf \geq 0.$$

If we think of π_n as a row vector, then for any $g \in \mathcal{L}^1(\mathfrak{X}, \pi_n)$ we have

$$\int_{\mathfrak{X}} g d\pi_n = \pi_n \cdot g,$$

where the “ $\cdot$ ” denotes the multiplication of a one-row matrix π_n with a one-column matrix g . We deduce

$$\pi_n \cdot g = (\pi_0 \cdot Q^n) \cdot g = \pi_0 \cdot (Q^n g).$$

Thus

$$g \in L^1(\mathfrak{X}, \pi_n), \quad \forall n \geq 0 \iff Q^n g \in L^1(\mathfrak{X}, \pi_0), \quad \forall n \geq 0.$$

Observe also that if $g \in L^1(\mathfrak{X}, \pi_0)$, then $g(x) < \infty$ if $\pi_0[\{x\}] > 0$.

Definition 4.3.1. A function $f : \mathfrak{X} \rightarrow (-\infty, \infty]$ is called *excessive* or *Lyapunov* of the HMC $(X_n)_{n \geq 0}$ if $f \geq Qf$. $\square$

Let $f : \mathfrak{X} \rightarrow \mathbb{R}$. Since the distribution of X_n is π_n , we deduce that

$$f(X_n) \in L^1(\mathfrak{X}, \pi_0) \iff f \in L^1(\mathfrak{X}, \pi_n).$$

The Markov condition implies $\mathbb{E}[f(X_{n+1}) \mid \mathcal{F}_n] = \mathbb{E}[f(X_{n+1}) \mid X_n]$. Let us observe that

$$\mathbb{E}[f(X_{n+1}) \mid \mathcal{F}_n] = \mathbb{E}[f(X_{n+1}) \mid X_n] = Qf(X_n), \quad \forall n \geq 0. \quad (4.3.1)$$

Indeed,

$$\begin{aligned} \mathbb{E}[f(X_{n+1}) \mid X_n = x] &= \sum_{y \in \mathfrak{X}} f(y) \mathbb{P}[X_{n+1} = y \mid X_n = x] \\ &= \sum_{y \in \mathfrak{X}} Q_{x,y} f(y) = (Qf)(x). \end{aligned}$$

Proposition 4.3.2. Suppose that

$$f \in \bigcap_{n \geq 0} \mathcal{L}^1(\mathfrak{X}, \pi_n).$$

Then the sequence $(f(X_n))_{n \geq 0}$ is a martingale (resp. supermartingale) iff $Qf = f$ (resp. $Qf \leq f$). $\square$

Definition 4.3.3. (a) The operator $\Delta := \mathbb{I} - Q : \mathcal{L}_*^0(\mathfrak{X}) \rightarrow \mathcal{L}_*^0(\mathfrak{X})$ is called the Laplacian³ of the HMC. More explicitly that for any $f \in \mathcal{L}_*^0(\mathfrak{X})$ we have

$$(\Delta f)(x) = \sum_{y \in \mathfrak{X}} Q_{x,y} (f(x) - f(y)).$$

(b) A function $f : \mathfrak{X} \rightarrow \mathbb{R}$ is called (Q) -harmonic (resp. Q -superharmonic) with respect to Q if $\Delta f = 0$, resp. $\Delta f \geq 0$. Note that a function $f : \mathfrak{X} \rightarrow \mathbb{R}$ is excessive iff it is superharmonic. $\square$

³Here we are using the geometers' convention. As defined, the Laplacian is *nonnegative* definite.

Let $f : \mathfrak{X} \rightarrow \mathbb{R}$. The sequence $f((X_n))_{n \geq 0}$ is a martingale iff

- $Q^n f \in \mathcal{L}^1(\mathfrak{X}, \pi_0)$, $\forall n \in \mathbb{N}_0$, and
- f is *harmonic* with respect to this Laplacian.

This sequence is a supermartingale iff

- $Q^n f \in \mathcal{L}^1(\mathfrak{X}, \pi_0)$, $\forall n \in \mathbb{N}_0$, and
- $\Delta f \geq 0$, i.e., f is *superharmonic* with respect to this Laplacian.

A function $f : \mathfrak{X} \rightarrow \mathbb{R}$ is said to be *harmonic on a subset* $U \subset \mathfrak{X}$ if

$$\Delta f(u) = 0, \quad \forall u \in U \iff f(u) = \sum_{x \in \mathfrak{X}} Q_{u,x} f(x), \quad \forall u \in U.$$

Example 4.3.4. (a) Fix a nonempty subset $S \in \mathfrak{X}$ and a bounded function $\varphi : S \rightarrow \mathbb{R}$. Denote by H_S the hitting/début time of S

$$H_S := \min \{ n \geq 0; X_n \in S \}.$$

Suppose that $H_S < \infty$ a.s. and define

$$F = F_{S,\varphi} : \mathfrak{X} \rightarrow \mathbb{R}, \quad F(x) = \mathbb{E}_x[\varphi(X_{H_S})].$$

Note that $F(s) = \varphi(s)$, $\forall s \in S$. If $u \in U = \mathfrak{X} \setminus S$, then

$$F(u) = \sum_{v \in U} Q_{u,v} F(v) + \sum_{s \in S} Q_{u,s} \varphi(s)$$

Hence $F_{\varphi,S}$ is a solution of the Dirichlet problem, $\Delta F = 0$ harmonic on U and $F|_S = \varphi$.

(b) Consider a nonempty subset $Y \subset \mathfrak{X}$ and $x_0 \in \mathfrak{X} \setminus Y$. We denote by H_Y the hitting time of Y , and by H_{x_0} hitting time of $\{x_0\}$. Set

$$S = \{x_0\} \cup Y, \quad \varphi = \mathbf{I}_{\{x_0\}}.$$

Then $H_S = \min(H_{x_0}, H_Y)$. Assuming that H_S is a.s. finite we have

$$F_{x_0,Y}(x) = \mathbb{E}_x[\varphi(X_{H_S})] = \mathbb{P}_x[H_{x_0} < H_Y], \quad \forall x \in \mathfrak{X},$$

i.e., $F_{x_0,Y}(x)$ is the probability that the system started at x hits x_0 before it hits Y . Thus, $F_{x_0,Y}(x)$ is harmonic on $\mathfrak{X} \setminus (Y \cup \{x_0\})$ and satisfies the boundary conditions, $F(x_0) = 1$ and $F(y) = 0$, $\forall y \in Y$. $\square$

Proposition 4.3.5 (Lévy's martingale). *Suppose that $f \in \mathcal{L}^\infty(\mathfrak{X})$. For each $n \in \mathbb{N}_0$ we set*

$$\begin{aligned} M_n^f &:= f(X_n) - f(X_0) + \sum_{k=0}^{n-1} \Delta f(X_k) \\ &= f(X_n) - f(X_0) + \sum_{k=0}^{n-1} (X_k - \mathbb{E}[f(X_{k+1}) | \mathcal{F}_k]). \end{aligned}$$

Then the sequence $(M_n^f)_{n \geq 0}$ is a martingale.

Proof. Note that

$$M_{n+1}^f - M_n^f = f(X_{n+1}) - f(X_n) + \Delta f(X_n)$$

and

$$\begin{aligned} \mathbb{E}[M_{n+1}^f - M_n^f \mid \mathcal{F}_n] &= \mathbb{E}[f(X_{n+1}) \mid \mathcal{F}_n] - f(X_n) - \Delta f(X_n) \\ &\stackrel{(4.3.1)}{=} Qf(X_n) - f(X_n) + \Delta f(X_n) = 0. \end{aligned}$$

□

Let $\mathbf{I} : \mathfrak{X} \rightarrow \mathbb{R}$ denote the indicator of $\mathfrak{X}$, $\mathbf{I}(x) = 1, \forall x \in \mathfrak{X}$. Since Q is a stochastic matrix we have $Q\mathbf{I} = \mathbf{I}$, so that the constant functions are harmonic.

Theorem 4.3.6. *Suppose that the HMC $(X_n)_{n \geq 0}$ is irreducible and recurrent. Then any bounded Lyapunov function is constant.*

Proof. We argue by contradiction. Suppose that h is non-constant bounded Lyapunov function on $\mathfrak{X}$. There exist $x_0, x_1 \in \mathfrak{X}$ such that $h(x_0) \neq h(x_1)$.

Suppose that $\pi_0 = \delta_{x_0}$. The sequence $h(X_n)$ is a bounded supermartingale. The Submartingale Convergence Theorem implies that the sequence $h(X_n)$ converges a.s..

On the other hand, since (X_n) is recurrent we deduce

$$\mathbb{P}[X_n = x_0 \text{ i.o.}] = \mathbb{P}[X_n = x_1 \text{ i.o.}] = 1.$$

Thus, the sequence $h(X_n)$ has a.s. two different limit points $h(x_0)$ and $h(x_1)$ and thus $h(X_n)$ is a.s. divergent! □

Corollary 4.3.7. *If the irreducible HMC $(X_n)_{n \geq 0}$ admits a nonconstant, bounded Lyapunov function, then it must be transient.* □

Example 4.3.8. Suppose that $(X_n)_{n \geq 0}$ describes the simple random walk on a locally finite connected graph $G = (V, E)$ with vertex set V and edge set E . A function $f : V \rightarrow \mathbb{R}$ is then superharmonic with respect to this Markov chain if its value at each vertex is at least the average of the values at neighbors

$$f(v) \geq \frac{1}{\deg v} \sum_{u \sim v} f(u), \quad \forall v \in V,$$

where $u \sim v$ indicates that the vertices u and v are neighbors, i.e., connected by an edge.

Suppose that G is a rooted binary tree. This means that G is a tree, it has a unique vertex v_0 of degree 1, and every other vertex has degree 3. One can think that any vertex other than the root has a unique direct ancestor and two direct successors. The root has a unique successor

One can think of v_0 as the generation zero vertex. It has a unique successor. This is the generation 1 vertex. Its two successors form the second generation of vertices. Their 4 successors determine the third generation etc. Equivalently, a vertex belongs to the n -th generation, $n > 1$, if its predecessor is in the $(n - 1)$ -th generation. We obtain in this fashion a generation function

$$g : V \rightarrow \mathbb{N}_0, \quad g(v) := \text{the generation of the vertex } v.$$

Define

$$f : V \rightarrow [0, 1], \quad f(v) = 2^{-g(v)}.$$

Any vertex $v \neq v_0$ has two vertices of generation $g(v) + 1$ and one vertex of generation $g(v) - 1$. Hence

$$\sum_{u \sim v} f(u) = 2^{-g(v)+1} + 2 \cdot 2^{-f(v)-1} = 3 \cdot 2^{-g(v)} = 3f(v),$$

so that

$$f(v) = \frac{1}{3} \sum_{u \sim v} f(u), \forall v \in V \setminus \{v_0\}.$$

Obviously $f(v_0) > f(v)$, $\forall v \in V \setminus \{v_0\}$. This proves that f is superharmonic, nonconstant and bounded so the random walk on G is transient. $\square$

Example 4.3.9. Suppose that $(X_n)_{n \geq 0}$ describes the regular random walk on a d -regular tree $\mathcal{T}_d$. This is an connected graph $\mathcal{T}_d = \mathcal{T}_d(V, E)$ with no cycles such that each vertex has degree d . We will show that if $d \geq 3$, then the random walk is transient.

To prove this we will construct a nonconstant bounded super harmonic function on $\mathcal{T}_d$. In this case a function $f : V \rightarrow \mathbb{R}$ is harmonic if

$$f(v) = \frac{1}{d} \sum_{u \sim v} f(u).$$

Recall that a *path in a graph* is a sequence of distinct vertices $v_0, v_1, \dots, v_n$ such that $v_{i-1} \sim v_i$, $\forall i = 1, \dots, n$. The natural number n is called the *length of the path*.

Fix a vertex $v_0 \in V$. For each $v \in V$ there exists a unique path in $\mathcal{T}_d$ connecting v_0 to v . We will denote by $\ell(v)$ its length. We denote by S_n the sphere

$$S_\ell := \{v; \ell(v) = n\}.$$

We seek a bounded function $f : V \rightarrow \mathbb{R}$ that is of the form $f(v) = r^{\ell(v)}$, $r \in (-1, 1)$, that is is harmonic on $V \setminus \{v_0\}$. Since $f(v_0) = 1 > r$, this function is superharmonic on V .

A vertex $v \in S_n$, $n > 0$, has $(d-1)$ neighbors on S_{n+1} and a vertex on S_{n-1} . Note that $f = r^{\ell(v)}$ is harmonic on $V \setminus \{v_0\}$ iff for any $n \in \mathbb{N}$ we have

$$r^n = \frac{1}{d} (r^{n-1} + (d-1)r^{n+1}) \iff r^{n-1} ((d-1)r^2 - dr + 1) = 0.$$

This happens iff r is a root of the quadratic equation

$$(d-1)x^2 - dx + 1 = 0.$$

The discriminant of this equation is $\Delta = d^2 - 4(d-1) = (d-2)^2$, so its roots are

$$1 = \frac{d+d-2}{2d-2}, \quad \frac{1}{d-1} = \frac{d-(d-2)}{2(d-1)}.$$

For $d \geq 3$ we have $\frac{1}{(d-1)} \in (0, 1)$ and the function $f(v) = (d-1)^{-\ell(v)}$ is nonconstant, bounded and superharmonic.

For a different yet related approach we refer to Example 4.5.17. $\square$

Corollary 4.3.7 can be substantially generalized.

Proposition 4.3.10. Let $(X_n)_{n \geq 0}$ be an irreducible HMC with state space $\mathfrak{X}$ and transition matrix Q . The following statements are equivalent.

- (i) The Markov chain (X_n) is transient.
- (ii) There exists a function $f : \mathfrak{X} \rightarrow [0, \infty]$, a nonempty subset $A \subset \mathfrak{X}$ and a point $x_0 \in \mathfrak{X} \setminus A$ such that
- (a) $Qf(x) \leq f(x), \forall x \in A^c := \mathfrak{X} \setminus A$ and
 - (b)

$$f(x_0) < \inf_{a \in A} f(a) < \infty.$$

Proof. I follow [97].

(ii) $\Rightarrow$ (i) Denote by H_A the début/hitting time of A ,

$$H_A := \min \{ n \geq 0; X_n \in A \}.$$

By multiplying f with an appropriate positive constant we can assume that

$$\inf_{a \in A} f(a) = 1.$$

We have

$$\begin{aligned} f(x_0) &\geq \sum_{x_1 \in \mathfrak{X}} Q_{x_0, x_1} f(x_1) = \sum_{x_1 \in A} Q_{x_0, x_1} f(x_1) + \sum_{x_1 \in A^c} Q_{x_0, x_1} f(x_1) \\ &\geq Q_{x_0} [A] + \sum_{x_1 \in A^c} Q_{x_0, x_1} f(x_1) = \mathbb{P}[X_1 \in A | X_0 = x_0] + \sum_{x_1 \in A^c} Q_{x_0, x_1} f(x_1). \end{aligned}$$

For any $x_1 \in A^c$ we have similarly

$$f(x_1) \geq \mathbb{P}[X_2 \in A | X_1 = x_1] + \sum_{x_2 \in A^c} Q_{x_1, x_2} f(x_2).$$

Hence

$$\begin{aligned} 1 > f(x_0) &\geq \mathbb{P}[X_1 \in A | X_0 = x_0] + \sum_{x_1 \in A^c} \mathbb{P}[X_2 \in A | X_1 = x_1] Q_{x_1, x_0} \\ &\quad + \sum_{x_1 \in A^c} \sum_{x_2 \in A^c} Q_{x_0, x_1} Q_{x_1, x_2} f(x_2). \end{aligned}$$

Iterating the above procedure we deduce that for any $n \in \mathbb{N}, n \geq 2$, we have

$$\begin{aligned} 1 > f(x_0) &\geq \mathbb{P}[X_1 \in A | X_0 = x_0] \\ &+ \underbrace{\sum_{k=1}^{n-1} \sum_{(x_1, \dots, x_k) \in (A^c)^k} \mathbb{P}[X_{k+1} \in A | X_k = x_k] \prod_{j=1}^k Q_{x_{j-1}, x_j}}_{\mathbb{P}[X_{k+1} \in A, X_1, \dots, X_k \in A^c, X_0 = x_0]} \\ &\quad + \underbrace{\sum_{(x_1, \dots, x_n) \in (A^c)^n} \left(\prod_{j=1}^n Q_{x_{j-1}, x_j} \right) f(x_n)}_{\geq 0} \\ &\geq \sum_{k=1}^n \mathbb{P}_{x_0} [H_A = k]. \end{aligned}$$

Hence

$$\mathbb{P}_{x_0} [H_A < \infty] = \sum_{n \in \mathbb{N}} \mathbb{P}_{x_0} [H_A = n] \leq f(x_0) < 1,$$

proving that the chain is transient.

(i) $\Rightarrow$ (ii) Let A be a proper subset of X . For any $x \in \mathfrak{X}$ we set

$$f(x) = \mathbb{P}_x[H_A < \infty].$$

Clearly $f(a) = 1, \forall a \in A$ and $f(x) < 1$, for $x \in \mathfrak{X} \setminus A$ since the chain is transient. The Markov property implies that

$$f(x) = Qf(x), \quad \forall x \in \mathfrak{X} \setminus A.$$

On the other hand,

$$\forall x \in \mathfrak{X} \setminus A \quad f(x) < \inf_{a \in A} f(a) = 1.$$

Hence f satisfies (ii.a) and (ii.b). $\square$

Definition 4.3.11. A function $f \in \mathcal{L}^0(\mathfrak{X})$ is called *coercive* if, for any $C > 0$, the set $\{f \leq C\}$ is a finite subset of $\mathfrak{X}$. $\square$

Proposition 4.3.12. Let $(X_n)_{n \geq 0}$ be an irreducible HMC with infinite state space $\mathfrak{X}$ and transition matrix Q . Suppose that there exists a nonnegative coercive function $f : \mathfrak{X} \rightarrow [0, \infty)$ and a finite set $A \subset \mathfrak{X}$ such that

$$\sum_{y \in \mathfrak{X}} Q_{x,y} f(y) \leq f(x), \quad \forall x \in \mathfrak{X} \setminus A. \quad (4.3.2)$$

Then the Markov chain $(X_n)_{n \geq 0}$ is recurrent.

Proof. I follow [97]. For $N > \sup_{a \in A} f(a)$ we set

$$A_N := \{f \geq N\}, \quad f_N := \frac{1}{N} f.$$

Since $\mathfrak{X}$ is infinite we deduce $S_N \neq \emptyset, \forall N \in \mathbb{N}$. By construction, the sets A and S_N are disjoint and $f_N(x) \geq 1, \forall x \in S_N$. Denote by T_A the return time to A and by T_{S_N} the return time to S_N .

Set $A_N := A \cup S_N$. The sets A_N^c are finite and nonempty. Let $x_0 \in A_N^c$. Arguing as in the proof of Proposition 4.3.10 we deduce

$$\begin{aligned} f_N(x_0) &= \sum_{x_1 \in A} Q_{x_0, x_1} f(x_1) + \sum_{x_1 \in S_N} Q_{x_0, x_1} \underbrace{f_N(x_1)}_{\geq 1} + \sum_{x_1 \in A_N^c} Q_{x_0, x_1} f_N(x_1) \\ &\geq \mathbb{P}[X_1 \in S_N | X_0 = x_0] + \sum_{x_1 \in A_N^c} Q_{x_0, x_1} f_N(x_1) \\ &\geq \mathbb{P}[X_1 \in S_N | X_0 = x_0] + \sum_{x_1 \in A_N^c} \mathbb{P}[X_2 \in S_N | X_1 = x_1] Q_{x_0, x_1} \\ &\quad + \sum_{x_1, x_2 \in A_N^c} Q_{x_0, x_1} Q_{x_1, x_2} f_N(x_2) \\ &\quad \vdots \\ &\geq \mathbb{P}[X_1 \in S_N | X_0 = x_0] + \underbrace{\sum_{k=1}^{n-1} \sum_{(x_1, \dots, x_k) \in (A_N^c)^k} \mathbb{P}[X_{k+1} \in S_N | X_k = x_k] \prod_{j=1}^k Q_{x_{j-1}, x_j}}_{\mathbb{P}[X_{k+1} \in S_N, X_1, \dots, X_k \in A_N^c, X_0 = x_0]} \end{aligned}$$

$$+ \underbrace{\sum_{(x_1, \dots, x_n) \in (A_N^c)^n} \left(\prod_{j=1}^n Q_{x_{j-1}, x_j} \right) f_N(x_n)}_{\geq 0} \geq \sum_{k=1}^n \mathbb{P}_{x_0} [T_{S_N} = k < T_A].$$

Letting $n \rightarrow \infty$ we deduce that for any $x_0 \in A_N^c$ we have

$$f_N(x_0) \geq \mathbb{P}_{x_0} [T_{S_N} < T_A] \geq \mathbb{P}_{x_0} [T_A = \infty].$$

If we fix $x_0 \in A^c$, then $x_0 \in A_N^c$ for all $N \gg 1$. Letting $N \rightarrow \infty$ we deduce that $\mathbb{P}_{x_0} [T_A = \infty] = 0$, $\forall x_0 \in \mathfrak{X} \setminus A$. This implies that X_n is recurrent.

To see this fix $x_0 \in A^c$ and set

$$Y_n := X_{n+T_A}.$$

This is a Markov chain with state space $\mathfrak{X}$ and transition matrix Q and initial distribution $\mu := \mathbb{P}_{X_{T_A}}$. The chain X_n is recurrent iff Y_n is recurrent. To prove that Y_n is recurrent we argue by contradiction. Suppose that Y_n is transient. For $a \in A$ we denote by N_a the number of returns to a of the chain Y_n . We deduce from Proposition 4.2.25 that $\mathbb{E}_\mu [N_a] < \infty$.

Denote by N_A the number of returns to A of Y_n ,

$$N_A := \sum_{n \in \mathbb{N}} \mathbf{I}_A(Y_n).$$

Since

$$\mathbf{I}_A = \sum_{a \in A} \mathbf{I}_{\{a\}}$$

We deduce that

$$N_A = \sum_{a \in A} N_a.$$

Since A is finite we conclude that $\mathbb{E}_\mu [N_A] < \infty$.

On the other hand since $\mathbb{P}_{x_0} [T_A < \infty] = 1$, $\forall x_0 \in \mathfrak{X} \setminus A$, we deduce that $\mathbb{P}_\mu [N_A = \infty] = 1$. We have reached a contradiction. $\square$

Remark 4.3.13. The sufficient condition (4.3.2) can be rewritten as $\Delta f(x) \geq 0$, $\forall x \in \mathfrak{X} \setminus A$. It is also necessary for recurrence. For a proof we refer to [73, Sec. 2.2] or [97]. $\square$

Theorem 4.3.14 (Foster). *Let $(X_n)_{n \geq 0}$ be an irreducible, locally finite HMC with state space $\mathfrak{X}$ and transition matrix Q . Suppose that there exists a function $f : \mathfrak{X} \rightarrow [0, \infty)$, a finite set $A \subset \mathfrak{X}$ and $\varepsilon > 0$ such that*

$$\sum_{y \in \mathfrak{X}} Q_{x,y} f(y) \leq f(x) - \varepsilon, \quad \forall x \in \mathfrak{X} \setminus A. \quad (4.3.3)$$

Then $(X_n)_{n \geq 0}$ is positively recurrent.

Proof. I follow [73, Sec.2.2]. Suppose $X_0 = x \in \mathfrak{X} \setminus A$. Denote by T_A the time of first return to A and set $Y_n := X_{n \wedge T_A}$. Then (4.3.3) reads

$$\mathbb{E}_x [f(Y_{n+1}) - f(Y_n) | Y_n] = \mathbb{E}_x [f(Y_{n+1}) | Y_n] - Y_n \leq -\varepsilon \mathbf{I}_{\{T_A > n\}}.$$

Thus $f(Y_n)$ is a nonnegative supermartingale and

$$\mathbb{E}_x [f(Y_{n+1})] - \mathbb{E}_x [f(Y_n)] \leq -\varepsilon \mathbb{P}_x [T_A > n].$$

Hence

$$\mathbb{E}_x[f(Y_{n+1})] - f(x) = \mathbb{E}_x[f(Y_{n+1})] - \mathbb{E}_x[f(Y_0)] \leq -\varepsilon \sum_{k=0}^n \mathbb{P}_x[T_A > k]$$

so that

$$\sum_{k=0}^n \mathbb{P}_x[T_A > k] \leq \frac{1}{\varepsilon} f(x).$$

Letting $n \rightarrow \infty$ we deduce

$$\mathbb{E}_x[T_A] \leq \frac{1}{\varepsilon} f(x), \quad \forall x \in \mathfrak{X} \setminus A. \quad (4.3.4)$$

Now let $a \in A$. Then

$$\begin{aligned} \mathbb{E}_a[T_A] &= \sum_{b \in A} Q_{a,b} + \sum_{x \in \mathfrak{X} \setminus A} Q_{a,x} (1 + \mathbb{E}_x[T_A]) \\ &= 1 + \sum_{x \in \mathfrak{X} \setminus A} Q_{a,x} \mathbb{E}_x[T_A] \stackrel{(4.3.4)}{\leq} 1 + \frac{1}{\varepsilon} \sum_{x \in \mathfrak{X} \setminus A} Q_{a,x} f(x) < \infty. \end{aligned}$$

The last sum above is finite since the Markov chain is locally finite. Thus $\mathbb{E}_a[T_A] < \infty$, $\forall a \in A$ and Proposition 4.2.45 implies that $(X_n)_{n \geq 0}$ is positively recurrent $\square$

Remark 4.3.15. (a) Note that condition (4.3.3) reads

$$\Delta f(x) \geq \varepsilon, \quad \forall x \in \mathfrak{X} \setminus A.$$

(b) If $(X_n)_{n \geq 0}$ positively recurrent, $x_0 \in \mathfrak{X}$, then the function $f : \mathfrak{X} \rightarrow [0, \infty)$

$$f(x) = \begin{cases} \mathbb{E}_x[T_{x_0}], & x \neq x_0, \\ 0, & x = x_0 \end{cases}$$

satisfies the conditions of Theorem 4.3.14 with $A = \{x_0\}$, $\varepsilon = 1$.

(c) I refer to [14, Thm. 2.19] for generalizations and refinements of Proposition 4.3.12 and Theorem 4.3.14. $\square$

Example 4.3.16. Consider the biased random walk on $\mathbb{N}_0 = \{0, 1, \dots\}$ with transition probabilities

$$Q_{0,1} = 1, \quad Q_{n,n+1} = p_n, \quad Q_{n,n-1} = q_n := 1 - p_n, \quad \forall n \in \mathbb{N}.$$

Above $p_n, q_n > 0$, $\forall n \in \mathbb{N}$, so that the corresponding Markov chain is irreducible. Consider the coercive function

$$f : \mathbb{N}_0 \rightarrow [0, \infty), \quad f(n) = n.$$

Then, $\forall n \geq 1$ we have

$$\Delta f(n) = n - (p_n(n+1) + q_n(n-1)) = q_n - p_n.$$

Thus, if $q_n \geq p_n$, this random walk is recurrent. Moreover, if

$$\inf_{n \in \mathbb{N}} (q_n - p_n) > 0,$$

then this random walk is positively recurrent. $\square$

4.3.2. Optimal stopping times. We have encountered optimal stopping problems before in Example 1.4.14 and, as an application, we solved the secretary problem. I want to approach this subject again, in the context of Markov chains, because we have developed enough technology to be able to say something more substantial.

To motivate the general theory I will use a concrete example that seems to be part of the mathematical folklore. I found out from a graduate student that this question was asked at job interviews with some investment firms. Recall that $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{Z}$, $\forall a, b \in \mathbb{R}$.

Example 4.3.17. Bob plays the following game of chance. He rolls a fair die. If he rolls a 6, then the game is over, and he earns no money. If he rolls a number $k \in \llbracket 1, 5 \rrbracket$ he has two options: either he ends the game and pockets $\$k$, or he rolls the die again. And so on. Devise a stopping strategy that will maximize Bob's expected return.

Mathematically we are dealing with a Markov chain with state space $\mathfrak{X} = \llbracket 1, 6 \rrbracket$. The state 6 is absorbing, and from any state $k < 6$ we can equally likely transition to any state in $\mathfrak{X}$. In other words, the transition matrix is

$$Q = \begin{bmatrix} 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

We also have a *reward function* $R : \mathfrak{X} \rightarrow \mathbb{R}$

$$R(k) = \begin{cases} k, & k \neq 6, \\ 0, & k = 6. \end{cases}$$

We are supposed to find an a.s. finite stopping time T so $\mathbb{E}[R(X_T)]$ is as large as possible. Think of this stopping time as a stopping strategy. $\square$

We can now state the general *optimal stopping problem*.

A *game* is a triplet $(\mathfrak{X}, Q, R)$ where $\mathfrak{X}$ is the state space of a homogeneous Markov chain $(X_n)_{n \geq 0}$ with state transition matrix Q and $R : \mathfrak{X} \rightarrow \mathbb{R}$ is a reward function. Since this subsection is meant only to give a taste of the main ideas with fewer technicalities I will make many simplifying assumptions. In that spirit I will assume that $\mathfrak{X}$ is finite.

Denote by $\mathcal{T}$ the collection of a.s. finite stopping times adapted to this stochastic process. The *value* of the game is the function

$$v = v_R : \mathfrak{X} \rightarrow \mathbb{R}, \quad v(x) = \sup_{T \in \mathcal{T}} \mathbb{E}_x[R(X_T)]. \quad (4.3.5)$$

The problem asks to determine the value of the game and an optimal stopping time, i.e., a stopping time T such that

$$v(x) = \mathbb{E}_x[R(X_T)].$$

My presentation is a mix of the presentations in [42, Sec.7.3], [128, Sec.4.1] and [192, Sec.8.9].

First some general observations. Since R is bounded we have $v(x) < \infty$ for any x . Observe that if we modify the reward function R to $R + c$, $c \in \mathbb{R}$, then $v_{R+c} = v + c$. Hence, after possibly adding a constant to R we can assume that $R \geq 0$.

Clearly $v \leq \max R$. Let

$$\mathcal{M}_R := \left\{ y \in \mathfrak{X}; \ R(y) = \max_{x \in \mathfrak{X}} R(x) \right\}.$$

Clearly $v(x) = R(x)$, $\forall x \in \mathcal{M}_R$. If the chain is recurrent and D_R is the début time in $\mathcal{M}_R$,

$$D_R = \inf \{ n \geq 0; \ X_n \in \mathcal{M}_R \},$$

then D_R is an optimal stopping time and, for any $x \in \mathfrak{X}$,

$$v(x) = \max_{y \in \mathfrak{X}} R(y) = \max_{x \in \mathcal{M}_R} R(x).$$

Note also that if $x_0 \in \mathfrak{X}$ is an absorbing state, then $v(x_0) = R(x_0)$.

Let $\mathcal{E}_R$ denote the set of Q -excessive majorants of R , i.e., the set of functions $u : \mathfrak{X} \rightarrow \mathbb{R}$

$$u(x) \geq \max (R(x), Qu(x)), \quad \forall x \in \mathfrak{X}.$$

In our special case the Q -excessive functions (see Definition 4.3.1) are also the Q -subharmonic functions.

Theorem 4.3.18. *The value function $v = v_R$ is the minimal excessive majorant of R , i.e., $v \in \mathcal{E}_R$ and $v \leq u$, $\forall u \in \mathcal{E}_R$. Moreover, v satisfies the Wald-Bellman equation*

$$v(x) = \max (R(x), Qv(x)). \quad (4.3.6)$$

Proof. For any $\varepsilon > 0$, and any $x \in \mathfrak{X}$ there exists an a.s. stopping time $T_x = T_{x,\varepsilon}$ such that

$$v(x) \geq \mathbb{E}_x [R(X_{T_x})] > v(x) - \varepsilon.$$

For each x the stopping time is a function

$$T_x : \{x\} \times \mathfrak{X}^{\mathbb{N}} \rightarrow \mathbb{N}_0$$

such that the $\{T_x = n\} \in \sigma(x, X_1, \dots, X_n)$. We can combine these in a new stopping time /strategy T^+ . Take the first step and then, once at state X_1 , use the strategy T_{X_1} . Thus

$$T^+(x_0, x_1, x_2, \dots) = 1 + T_{x_1}(x_1, x_2, \dots).$$

We have

$$\begin{aligned} v(x) &\geq \mathbb{E}_x [R(X_{T^+})] = \sum_{y \in \mathfrak{X}} Q_{x,y} \mathbb{E}_y [R(X_{T_y})] \\ &> \sum_{y \in \mathfrak{X}} Q_{x,y} (v(y) - \varepsilon) = Qv(x) - \varepsilon. \end{aligned}$$

Thus

$$\forall \varepsilon > 0, \quad v(x) > Qv(x) - \varepsilon,$$

so that $v \geq Qv$.

Suppose that u is another excessive majorant of R . Then $u(X_n)$ is a (bounded) supermartingale and we deduce from the optional stopping theorem that for any $T \in \mathcal{T}$ we have

$$u(x) \geq \mathbb{E}_x (u(X_T)) \geq \mathbb{E}_x (R(X_T)).$$

Hence

$$u(x) \geq \sup_{T \in \mathcal{T}} \mathbb{E}_x (R(X_T)) = v(x).$$

This proves that v is indeed the minimal excessive majorant of R .

Finally, let us prove that v satisfies (4.3.6). Set

$$v_1 := \max (R(x), Qv(x)).$$

Then $R(x) \leq v_1(x) \leq v(x)$ so

$$QR(x) \leq Qv_1(x) \leq Qv(x) \leq \max (R(x), Qv(x)) = v_1(x)$$

so v_1 is an excessive majorant of R . Hence $v \leq v_1$, i.e.

$$\max (R(x), Qv(x)) \leq v(x) \leq v_1(x) = \max (R(x), Qv(x)).$$

□

Remark 4.3.19. (a) The minimal excessive majorant v of R is the solution of the linear programming problem

$$\min \sum_{x \in eX} u(x)$$

subject to

$$\begin{cases} u \geq R \geq 0, \\ u \geq Qu. \end{cases}$$

(b) Any solution of equation (4.3.6) is an excessive majorant of R . However, there are solutions that may not be the minimal Q -excessive majorant of R . For example, the constant the function $u^* : \mathfrak{X} \rightarrow \mathbb{R}$, $u^*(x) = \max R$ is always a of solution (4.3.6). □

Example 4.3.20. Consider the game in Example 4.3.17. Its value is a function $v : \llbracket 1, 6 \rrbracket \rightarrow \mathbb{R}$ such that

$$R(k) \leq v(k) \leq \max R, \quad \forall k \in \llbracket 1, 6 \rrbracket.$$

Moreover $v[5] = 5$, $v[6] = 0$. In this case, the fact that v is Q -excessive translates into the condition

$$v(k) \geq \mu[v] := \frac{1}{6} \sum_{j=1}^6 v(j), \quad \forall 1 \leq k \leq 5.$$

We are looking for the smallest v satisfying these conditions.

We seek a function v such that

$$v(k) = \begin{cases} x, & k = 1, 2, \\ R(k), & k \geq 3, \end{cases} \quad \mu[v] \leq x.$$

Note that

$$x \geq \mu[v] = \frac{1}{6}(2x + 12) = \frac{x}{3} + 2 \Rightarrow x \geq 3.$$

Thus the function v given by

$$v(k) = \begin{cases} 3, & k = 1, 2, \\ R(k), & k \geq 3, \end{cases}$$

is an excessive majorant of R . The above argument shows that it is indeed the minimal such majorant.

The gambling strategy is clear: roll the die until the first time you get $k \in \llbracket 3, 6 \rrbracket$. Note that

$$\llbracket 3, 6 \rrbracket = \{ k \in \llbracket 1, 6 \rrbracket; \ v(k) = R(k) \}.$$

□

Clearly the technique used in the above example is ad-hoc and does not generalize. We can certainly find the value of a game using the simplex method. The `boot` package that comes with R can be used for this purpose. We want to describe another iterative method for determining the value functions.

Define (nonlinear) transformations

$$\Phi_R, \Psi : \mathbb{R}^{\mathfrak{X}} \rightarrow \mathbb{R}^{\mathfrak{X}}, \quad \Phi_R[u] = \max(R, Qu), \quad \Psi[u] = \max(u, Qu).$$

Both transformations are monotone, i.e.,

$$u_0 \leq u_1 \Rightarrow \Phi[u_0] \leq \Phi[u_1], \quad \Psi[u_0] \leq \Psi[u_1].$$

The map Ψ is independent of R and its fixed points are precisely the excessive (superharmonic) functions. The fixed points of Φ_R are excessive majorants of R .

Theorem 4.3.21. *Suppose that $(\mathfrak{X}, Q, R)$ is a game. Assume for simplicity that $\mathfrak{X}$ is finite and $R \geq 0$. Define iteratively the functions $v_n : \mathfrak{X} \rightarrow \mathbb{R}$,*

$$v_0 = R, \quad v_{n+1} = \Phi_R[v_n], \quad \forall n \geq 0$$

Then the following hold.

- (i) *The sequence v_n is nondecreasing.*
- (ii) *The sequence v_n also satisfies the recurrence*

$$v_n = \Psi[v_{n-1}], \quad \forall n \in \mathbb{N}. \tag{4.3.7}$$

- (iii) *The sequence v_n converges to the value function v_R .*

Proof. (i) Note first that

$$v_1 = \Phi_R[v_0] \geq R = v_0.$$

The transformation Φ_R is monotone and we deduce iteratively that

$$0 \leq v_0 \leq v_1 \leq v_2 \leq \dots \leq \max R.$$

(ii) Let us prove that the sequence satisfies the recurrence (4.3.7). We argue by induction. Denote by w_n the sequence defined inductively by

$$w_0 = R, \quad w_{n+1} = \max(w_n, Qw_n).$$

We will prove inductively that $v_n = w_n$, for all n . Clearly $w_0 = v_0$, $w_1 = v_1$. Suppose that $v_k = w_k$, $\forall k \leq n$. Then,

$$w_{n+1} = \max(v_n, Qv_n) \geq \max(R, Qv_n) = v_{n+1}.$$

On the other hand, since $Qv_n \leq v_{n+1}$ and the sequence (v_n) is nondecreasing we have

$$w_{n+1} = \max(v_n, Qv_n) \leq \max(v_n, v_{n+1}) = v_{n+1}$$

(iii) The sequence of functions v_n is nondecreasing and bounded above so it converges pointwisely to a function $v_\infty : \mathfrak{X} \rightarrow [0, \max R]$. The limit v_∞ is a fixed point of both Φ_R and Ψ and thus it is an excessive majorant of R .

On the other hand, if v denotes the value function we deduce $v \geq R = v_0$ so

$$v = \Psi[v] \geq \Psi[R] = v_1.$$

We deduce inductively that $v \geq v_n$, $\forall n$, and thus $v \geq v_\infty$. Since v is the minimal excessive majorant we deduce $v_\infty = v$. $\square$

The iterates v_n described above have alternate, revealing interpretations. To describe them we need to introduce restricted classes of stopping times. For $n \leq N \in \mathbb{N}$ we set

$$\mathcal{T}_N := \{ T \in \mathcal{T}; T \leq N \text{ a.s.} \}, \quad \mathcal{T}_{N,n} := \{ T \in \mathcal{T}_N; T \geq n \}.$$

Theorem 4.3.22. *For each $N \in \mathbb{N}$ the function $v_N : \mathfrak{X} \rightarrow \mathbb{R}$ defined in Theorem 4.3.21, $v_N = \Phi_R^N[R]$, is the value of the restricted optimization problem*

$$v_N(x) = \sup_{T \in \mathcal{T}_N} \mathbb{E}_x[R(X_T)]. \quad (4.3.8)$$

Additionally, if we define

$$T_N^* := \min \{ n \in \llbracket 0, N \rrbracket; v_{N-n}(X_n) = R(X_n) \}, \quad (4.3.9)$$

then

$$v_N(x) = \mathbb{E}_x[R(X_{T_N^*})]. \quad (4.3.10)$$

Proof. I follow the approach in [190, Sec.2.2]. Consider the optimality conditions (1.4.17a), (1.4.17b), (1.4.18a) and (1.4.18b) in Example 1.4.14.

Define $Y_n = Y_n(x, X_1, \dots, X_n)$ by descending induction

$$Y_N = R(X_N), \quad Y_n = \max \left(R(X_{n-1}), \mathbb{E}_x[Y_n \mid X_0, \dots, X_{n-1}] \right).$$

The Markov condition implies

$$\mathbb{E}[Y_n \mid X_0, \dots, X_{n-1}] = \mathbb{E}[Y_n \mid X_{n-1}]$$

We have $Y_N = R(X_N)$,

$$\begin{aligned} Y_{N-1}(x) &= \max \left(R(X_{N-1}), \mathbb{E}_x[R(X_N) \mid X_{N-1}] \right) \\ &= \max \left(R(X_{N-1}), Q_R(X_{N-1}) \right) = \Phi_R[R](X_{N-1}) \end{aligned}$$

Iterating we deduce

$$Y_{N-n} = \Phi_R^n[R](X_{N-n})$$

In particular

$$Y_0(x) = \Phi_R^N[R](X_0 = x) = v_N(x).$$

According to the results Example 1.4.14, we have

$$Y_0(x) = \sup_{T \in \mathcal{T}_N} \mathbb{E}[R(X_T)].$$

An optimal stopping time prescribed by (1.4.17b) coincides with the optimal stopping time described in (4.3.9). $\square$

Example 4.3.23. I want to describe an R-implementation of the strategy in Theorem 4.3.21 for computing the value function in Example 4.3.17. Here's an R script that perform the Ψ -iterations.

```
options(rgl.useNULL = TRUE)
library(rgl)
#include the above two lines if you are using the 2025 MacOS update Tahoe

library(matlib)
p=1/6

#Define the transition matrix.
```

```

v<-c(rep(p,30))
v=c(v, 0,0,0,0,0,1)
Q<-matrix(v, ncol=6,byrow=TRUE)

#generate the LaTeX code describing the transition matrix Q
latexMatrix(Q, fractions=TRUE)

# Define the transformation Psi
game<-function(x) pmax(x,Q%%x )

#define the reward function
reward=matrix(c(1:5,0), ncol=1)

#Computing the N-th iterate of Psi
policy<-function(N, x){
  for( i in 1:N){
    x=game(x)
  }
  x
}
#Here's the fifth interate
v5=policy(5, reward)
v5

#generate the Latex code describing the function v5
latexMatrix(v5)

```

We deduce that

$$v_5 \approx \begin{bmatrix} 2.9938 \\ 2.9938 \\ 3.0000 \\ 4.0000 \\ 5.0000 \\ 0.0000 \end{bmatrix}$$

This is in perfect agreement with the result obtained by the ad-hoc arguments in Example 4.3.20. $\square$

Now that we have discussed ways of determining the value functions, I would like to conclude by describing an optimal gambling strategy that will maximize the expected payoff. This strategy assumes with have already determined the value function.

Theorem 4.3.24. *Consider a game $(\mathfrak{X}, Q, R)$ with $\mathfrak{X}$ finite. Denote by $v(x)$ the value of this game and set*

$$\mathfrak{X}_0 := \{ x \in \mathfrak{X}; v(x) = R(x) \}.$$

Denote by H_0 the début time in $\mathfrak{X}_0$,

$$H_0 := \{ n \geq 0; X_n \in \mathfrak{X}_0 \}.$$

Then the following hold

- (i) $H_0 < \infty$, a.s..
- (ii) $v(x) = \mathbb{E}_x[R(X_{H_0})]$.

Proof. (i) Note that

$$\{H_0 = \infty\} = E := \bigcap_{n \in \mathbb{N}} \{X_n \notin \mathfrak{X}_0\}.$$

We have to prove that $\mathbb{P}_x[E] = 0$ for all $x \in \mathfrak{X} \setminus \mathfrak{X}_0$. Set $\mathfrak{X}_* := \mathfrak{X} \setminus \mathfrak{X}_0$. Since $\mathfrak{X}$ is finite there exists $\varepsilon > 0$ such that

$$R(x) \leq v(x) - \varepsilon, \quad \forall x \in \mathfrak{X} \setminus \mathfrak{X}_0.$$

Then, for any a.s. finite stopping time T we have

$$\begin{aligned} \mathbb{E}_x[R(X_T)] &= \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}} \mathbb{P}_x[T = n, X_n = y] R(y) \\ &= \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}_*} \mathbb{P}_x[T = n, X_n = y] R(y) + \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}_0} \mathbb{P}_x[T = n, X_n = y] v(y) \\ &< \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}_*} \mathbb{P}_x[T = n, X_n = y] (v(y) - \varepsilon) + \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}_0} \mathbb{P}_x[T = n, X_n = y] v(y) \\ &= \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}} \mathbb{P}_x[T = n, X_n = y] v(y) - \underbrace{\varepsilon \sum_{n=0}^{\infty} \sum_{y \in \mathfrak{X}_*} \mathbb{P}_x[T = n, X_n = y]}_{\geq \mathbb{P}_x[E]} \\ &= \mathbb{E}_x[v(X_T)] - \varepsilon \mathbb{P}_x[E]. \end{aligned}$$

Since v is Q -excessive, and bounded the sequence $v(X_n)$ is a bounded supermartingale we deduce from the optimal stopping theorem that $\mathbb{E}_x[v(X_T)] \leq v(x)$. Hence

$$\forall \varepsilon > 0, \quad \forall T \in \mathcal{T}, \quad \mathbb{E}_x[g(X_T)] \leq v(x) - \varepsilon \mathbb{P}_x[E],$$

so that

$$\forall \varepsilon > 0, \quad v(x) = \sup_{T \in \mathcal{T}} \mathbb{E}_x[g(X_T)] \leq v(x) - \varepsilon \mathbb{P}_x[E].$$

Hence $\mathbb{P}_x[E] = 0$, i.e., H_0 is a.s. finite.

(ii) Set

$$v^*(x) := \mathbb{E}_x[R(X_{H_0})].$$

Note that $R(X_{H_0}) = v(X_{H_0})$ since $v = R$ on $\mathfrak{X}_0$. Since $v(X_n)$ is a bounded supermartingale we deduce

$$v^*(x) = \mathbb{E}_x[v(X_{H_0})] \leq v(x).$$

We will show that v^* is an excessive majorant of R .

Denote by T_0 the time of the first return to $\mathfrak{X}_0$, i.e.,

$$T_0 := \min \{ n \geq 1; X_n \in \mathfrak{X}_0 \}.$$

Clearly $H_0 \leq T_0$ and since $v(X_n)$ is a bounded submartingale we deduce from the Optional Sampling Theorem 3.2.44

$$v^*(x) = \mathbb{E}_x[v(X_{H_0})] \geq \mathbb{E}_x[v(X_{T_0})] = \sum_{y \in \mathfrak{X}} Q_{x,y} \mathbb{E}_y[v(X_{H_0})] = \sum_{y \in \mathfrak{X}} Q_{x,y} v^*(y).$$

This proves that v^* is Q -excessive. Finally, let us prove that $v^* \geq R$.

Choose $x_* \in \mathfrak{X}$ such that

$$v^*(x_*) - R(x_*) = c_* := \min_{x \in \mathfrak{X}} (v^*(x) - R(x)).$$

We will prove that $c_* \geq 0$. We argue by contradiction and assume $c_* < 0$. In particular $x_* \in \mathfrak{X}_*$ since, by definition, $v(x) = R(x)$ on $\mathfrak{X}_0$.

The function $w(x) = v^*(x) - c_*$ is excessive and

$$w(x) - R(x) = v^*(x) - R(x) - c_* \geq 0.$$

Hence w is an excessive majorant of R so that $w(x) \geq v(x)$, $\forall x \in \mathfrak{X}$. In particular

$$v(x_*) \leq w(x_*) = R(x_*),$$

Since v is a majorant of R we deduce $v(x_*) = R(x_*)$. In other words $x_* \in \mathfrak{X}_0$. This is impossible since $x_* \in \mathfrak{X}^*$. $\square$

Remark 4.3.25. As explained in Example 4.3.4 (a), the value function $v(x) = \mathbb{E}_x[R(X_{H_{\mathfrak{X}_0}})]$ is a solution of the Dirichlet boundary value problem

$$\Delta v = 0 \text{ on } \mathfrak{X} \setminus \mathfrak{X}_0 \text{ and } v = R \text{ on } \mathfrak{X}_0.$$

This is a more subtle boundary value problem since the “boundary” $\mathfrak{X}_0$ is not a priori known; it is determined by the sought after solution v . This is an example of a *free boundary problem*. This is a rather rich subject of investigation and we refer to [168] for more details. $\square$

I want to illustrate the effectiveness of the results established so far on another examples.

Example 4.3.26. Consider the Markov chain given by the standard random walk on $\llbracket 0, n \rrbracket$ with absorbing boundary, i.e., the walk ends when it reaches one of the endpoints 0 or n . Let us choose as reward the function

$$R : \llbracket 0, n \rrbracket \rightarrow \mathbb{R}, \quad R(k) = k^2.$$

To any function $v : \llbracket 0, n \rrbracket \rightarrow \mathbb{R}$ we canonically associate the unique continuous piecewise linear function $\hat{v} : [0, n] \rightarrow \mathbb{R}$ with nodes $(k, v(k))$, $k \in \llbracket 0, n \rrbracket$.

In this case, a function $v : \llbracket 0, n \rrbracket \rightarrow \mathbb{R}$ is excessive iff

$$\forall 0 < k < n, \quad v(k) \geq \frac{1}{2} (v(k-1)v(k+1)).$$

Equivalently, this means that $\hat{v}$ is concave. The smallest concave function $f : [0, n] \rightarrow \mathbb{R}$ such that

$$f(x) \geq x^2, \quad \forall x \in [0, n], \quad f(0) = R(0), \quad f(n) = R(n) = n^2$$

is the linear function $v_\infty : [0, n] \rightarrow \mathbb{R}$, $v_\infty(x) = nx$. This is the value of the game.

Let us specialize further and set $n = 6$. The value of the game in this case is $v_\infty(k) = 6k$. The transition matrix is

$$Q = \begin{bmatrix} 1.0 & 0.0 & 0.0 & 0.0 & 0.0 & 0.0 & 0.0 \\ 0.5 & 0.0 & 0.5 & 0.0 & 0.0 & 0.0 & 0.0 \\ 0.0 & 0.5 & 0.0 & 0.5 & 0.0 & 0.0 & 0.0 \\ 0.0 & 0.0 & 0.5 & 0.0 & 0.5 & 0.0 & 0.0 \\ 0.0 & 0.0 & 0.0 & 0.5 & 0.0 & 0.5 & 0.0 \\ 0.0 & 0.0 & 0.0 & 0.0 & 0.5 & 0.0 & 0.5 \\ 0.0 & 0.0 & 0.0 & 0.0 & 0.0 & 0.0 & 1.0 \end{bmatrix}$$

Using the function `policy(N, x)` described in Example 4.3.23 with $x(k) = k^2$ and the matrix Q above we the following approximations of the value function v_∞ .

$$v_3 = \begin{bmatrix} 0.00 \\ 3.00 \\ 6.75 \\ 12.00 \\ 18.75 \\ 27.00 \\ 36.00 \end{bmatrix}, \quad v_7 = \begin{bmatrix} 0.0000 \\ 4.3125 \\ 9.0469 \\ 14.6250 \\ 21.0469 \\ 28.3125 \\ 36.0000 \end{bmatrix}, \quad v_{20} = \begin{bmatrix} 0.0000 \\ 5.7372 \\ 11.5495 \\ 17.4744 \\ 23.5495 \\ 29.7372 \\ 36.0000 \end{bmatrix}, \quad v_{30} = \begin{bmatrix} 0.0000 \\ 5.9376 \\ 11.8931 \\ 17.8753 \\ 23.8931 \\ 29.9376 \\ 36.0000 \end{bmatrix}.$$




Figure 4.4. The piecewise-linear renditions of the functions $v_0 \leq v_3 \leq v_7 \leq v_\infty$.

Figure 4.4 depicts the approximations v_3, v_7 . At the bottom sits the reward and the value is the top function. I have generated this figure using the following R-code

```
options(rgl.useNULL = TRUE)
library(rgl)
library(matlib)
```

```
p1<-c(1,c(rep(0,6)))
```

```

p2=c(0.5,0,0.5, c(rep(0,4)))
p3=c(0,0.5,0,0.5,0,0,0)
p4=c(0,0,0.5,0,0.5,0,0)
p5=c(0,0,0,0.5,0,0.5,0)
p6=c(0,0,0,0,0.5,0,0.5)
p7=c(c(rep(0,6)),1 )

p=c(p1,p2,p3,p4,p5,p6,p7)

Q<-matrix(p, ncol=7,byrow=TRUE)

game<-function(x) pmax(x,Q%*%x )

#policy(N,x)  computes the N-th iterate of the function game
#with initial condition.

policy<-function(N, x){
  for( i in 1:N){
    x=game(x)
  }
  x
}
y=c(0:6)^2

reward=matrix(y, ncol=1,byrow=TRUE)

x=c(0:6)

v0=y
v3=t(policy(3,reward))
v7=t(policy(7,reward))
v=6*c(0:6)
plot(x,v0, col="red",pch=16,ylab="v")
lines(x,v0, col="red",pch=16)
points(x,v3, col="purple",pch=18)
lines(x,v3, col="purple",pch=18)
points(x,v7, col="blue",pch=20)
lines(x,v7, col="blue",pch=20)
points(x, v,col="red", pch=20)
lines(x, v,col="red", pch=20)
legend(x="topleft",c("Approximating the value of the game"),bg="grey")

```

□

Example 4.3.27 (The secretary problem revisited). I want to present another point of view on the secretary problem discussed in Example 1.4.16 and reveal a hidden Markov chain lurking in the background. My presentation is inspired from [70, §III.1].

Recall that we are given collection of N different real numbers

$$v_1 < v_2 < \cdots < v_N, \quad N \geq 3.$$

We do not know these numbers. They are sequentially and randomly revealed to us as a random sequence

$$V_1, \dots, V_N.$$

Formally, we are given a uniformly random permutation φ of $\llbracket 1, N \rrbracket$ such that $V_i = v_{\varphi(i)}$, $\forall i$. We are supposed to detect the index i such that $v_{\varphi(i)} = v_N$ using only information we gained from the knowledge of $v_{\varphi(1)}, \dots, v_{\varphi(i)}$

We are first shown $v_{\varphi(1)}$. We can accept it and stop, or we can reject it. If we reject it we no longer have to option to accept this number again, but we are shown for a new number $v_{\varphi(2)}$ and again we have to decide if this is the largest number. We can accept it and stop or reject it and continue etc.

As explained in Example 1.4.16, to a random permutation φ we can associate a sequence of independent random variables $X_j, j = 1 \dots, N$, where

$$\forall \varphi \in \mathfrak{S}_N, \quad X_j(\varphi) = \#\{i \leq j; \varphi(i) \leq \varphi(j)\}.$$

I will refer to these random variables as the ranks of a (random) permutation.

The random variables X_j is uniformly distributed in $\llbracket 1, j \rrbracket$. We say that an index or epoch j is leading if $X_j(\varphi) = j$. In other words, $v_{\varphi(j)}$ is the largest of the numbers $v_{\varphi(1)}, \dots, v_{\varphi(j)}$. Clearly the epoch i such that $\varphi(i) = N$ is among the leading epochs.

Let $J_1 < J_2 < \dots$ be the leading epochs, in the order they are revealed. The number of leading epochs is random. Note that $J_1 = 1, J_2 \in [2, N]$ etc. Thus, finding the largest number reduces to tracking the leading epochs and deciding when to stop. In fact, the last leading epoch is the winning index. The problem is that, once we reach a leading epoch we do not know if this is the last one.

For illustration, consider the permutation $\varphi \in \mathfrak{S}_{10}$

$$1, 6, 2, 5, 7, 10, 9, 4, 8, 3,$$

It has four leading epochs

$$J_1 = 1, \varphi(J_1) = 1, J_2 = 2, \varphi(J_2) = 6,$$

$$J_3 = 5, \varphi(J_3) = 7, J_4 = 6, \varphi(J_4) = 10.$$

Thus J_4 is the last leading epoch. Figure 4.5 depicts these facts.

Denote by J_L the last of the leading epochs. Define

$$J_{L+k} := \infty, \quad \forall k \in \mathbb{N},$$

and set

$$\mathfrak{X} := \llbracket 1, N \rrbracket \cup \{\infty\}.$$

We obtain a sequence of $\mathfrak{X}$ -valued random variables $J_k : \mathfrak{S}_N \rightarrow \mathfrak{X}$ with $J_1 = 1$. We will show that this sequence is an HMC with state space $\mathfrak{X}$ and determine its transition matrix.

Let us first compute $\mathbb{P}[J_k = m]$. Suppose $J_k(\varphi) = m$. Let us first show that for $m < n \leq N$

$$\mathbb{P}[J_{k+1} = n | J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1] = \mathbb{P}[J_{k+1} = n | J_k = m].$$

Set

$$\mathbf{L}_n := \{1, j_2, \dots, j_{k-1}, m, n\} \subset \llbracket 1, n \rrbracket, \quad \mathbf{L}_n^* := \llbracket 1, n \rrbracket \setminus \mathbf{L}_n,$$

$$\mathbf{L}_m := \{1, j_2, \dots, j_{k-1}, m\} \subset \llbracket 1, m \rrbracket, \quad \mathbf{L}_m^* := \llbracket 1, m \rrbracket \setminus \mathbf{L}_m.$$




Figure 4.5. A zig-zag depicting a permutation and its leading points, marked with a triangle. The zig-zag connects the points $(j, \varphi(j))$, $1 \leq j \leq n$, by line segments.

In other words, $\mathbf{L}_n$ is the collection of leading epochs $\leq n$ and $\mathbf{L}_m$ is the collection of leading epochs $\leq m < n$. Note that

$$\mathbf{L}_m \subset \mathbf{L}_n, \quad \mathbf{L}_n \setminus \mathbf{L}_m = \{n\},$$

$$\mathbf{L}_m^* \subset \mathbf{L}_n^*, \quad \mathbf{L}_n^* \setminus \mathbf{L}_m^* = \{m+1, \dots, n-1\}.$$

Since the rank variables X_i are independent, we deduce that

$$\begin{aligned} & \mathbb{P}[J_{k+1} = n, J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1] \\ &= \left(\prod_{\ell \in \mathbf{L}_n} \mathbb{P}[X_\ell = \ell] \right) \left(\prod_{\ell \in \mathbf{L}_n^*} \mathbb{P}[X_\ell < \ell] \right). \end{aligned}$$

On the other hand,

$$\begin{aligned} & \mathbb{P}[J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1] \\ &= \left(\prod_{\ell \in \mathbf{L}_m} \mathbb{P}[X_\ell = \ell] \right) \left(\prod_{\ell \in \mathbf{L}_m^*} \mathbb{P}[X_\ell < \ell] \right). \end{aligned}$$

Hence

$$\begin{aligned} & \mathbb{P}[J_{k+1} = n \mid J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1] \\ &= \left(\prod_{\ell \in \mathbf{L}_n \setminus \mathbf{L}_m} \mathbb{P}[X_\ell = \ell] \right) \left(\prod_{\ell \in \mathbf{L}_n^* \setminus \mathbf{L}_m^*} \mathbb{P}[X_\ell < \ell] \right) \\ &= \mathbb{P}[X_n = n] \prod_{m < \ell < n} \mathbb{P}[X_\ell < \ell] \end{aligned}$$

(X_j is uniformly distributed on $\llbracket 1, j \rrbracket$)

$$= \frac{m}{m+1} \cdots \frac{n-2}{n-1} \cdot \frac{1}{n} = \frac{m}{n(n-1)} =: Q_{m,n}.$$

Note that $\mathbb{P}[J_{k+1} = n | J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1]$ depends only on the J_k and J_{k+1} . We deduce from the conditional law of total probability (1.2.12) that

$$\begin{aligned} \mathbb{P}[J_{k+1} = n | J_k = m, J_{k-1} = j_{k-1}, \dots, J_1 = 1] \\ = \mathbb{P}[J_{k+1} = n | J_k = m] = Q_{m,n}. \end{aligned}$$

Define $Q_{\infty,\infty} = Q_{N,\infty} = 1$ and, for $m < N$,

$$\begin{aligned} Q_{m,\infty} &= 1 - \sum_{m < n \leq N} Q_{m,n} = 1 - m \sum_{n=m+1}^N \frac{1}{n(n-1)} \\ &= 1 - m \left(\frac{1}{m} - \frac{1}{N} \right) = \frac{m}{N}. \end{aligned}$$

The transition probability $Q_{m,\infty}$ is the probability that m is the last leading epoch, namely the winning epoch. You should compare this fact with (1.4.25).

We can outline the winning strategy. As mentioned earlier, we only have to keep track of the leading epochs $J_1, J_2, \dots$ and decide which epoch J_k is the last one, i.e., $\varphi(J_k) = N$.

Suppose we reached the leading epoch $J_k = j$. The probability that this is the winning one is $p(j) = \frac{j}{N}$.

Let us now compute the conditional probability that the next leading epoch is the winning epoch, i.e., the probability

$$q(j) = \mathbb{P}[J_{k+2} = \infty | J_{k+1} \neq \infty, J_k = j].$$

Using the Markov property we deduce

$$\begin{aligned} q(j) &= \sum_{n=j+1}^N Q_{j,n} Q_{n,\infty} = \sum_{n=j+1}^N \frac{j}{n(n-1)} \cdot \frac{n}{N} \\ &= p(j) \left(\frac{1}{j} + \dots + \frac{1}{N-1} \right). \end{aligned}$$

Note that the ratio

$$f_N(j) = \frac{q(j)}{p(j)} = \frac{1}{j} + \dots + \frac{1}{N-1}$$

decreases with j and

$$f_N(1) = 1 + \frac{1}{2} + \dots + \frac{1}{N-1}, \quad f_N(N-1) = \frac{1}{N-1}.$$

Set

$$j^* = j^*(N) = \min \{ j; f_N(j) \leq 1 \}.$$

Thus if we reached a leading epoch $j \geq j^*(N)$, the conditional probability that the next leading epoch is the winning one is not greater than the probability that the current leading epoch j is the winning one.

We can now describe an optimal stopping time. Consider the set

$$\mathcal{O}_N := \llbracket j^*(N), N \rrbracket \cup \{\infty\} \subset \mathfrak{X}.$$

Set

$$K = \min \{ k > 1; J_k \in \mathcal{O}_N \}, \quad T^* = \begin{cases} J_K, & J_K \leq N, \\ N, & J_K = \infty. \end{cases}$$

The choice V_{T^*} maximizes the probability of picking v_N . Informally we accept the number revealed to us at the leading epoch J_K , the first leading epoch $\geq j^*(N)$. In other words we accept all numbers up to $j^*(N) - 1$ and then we admit the first number bigger than all the previous ones.

When $J_K = \infty$, this means we have rejected all the numbers $V_1, \dots, V_N$, and all the leading epochs we encountered along occurred before the critical epoch $j^*(N)$.

Let us prove this claim and compute the value $v = v_N$ of the game. The reward of this game is $R(j) = \frac{j}{N}$. This is the probability that given that j is a leading epoch, it is the last leading epoch. We set $R(\infty) = 0$. The maximal probability is $v(1)$.

The value function $v(j)$ is the minimal solution of the system of linear inequalities

$$v(j) \geq \frac{j}{N}, \quad v(j) \geq \sum_{j < n \leq N} Q_{j,n} v(n) = j \sum_{j < n \leq N} \frac{v(n)}{n(n-1)}.$$

Hence, if $v(n)$ is known for all $n > j$ we have

$$v(j) = \max \left(\frac{j}{N}, j \sum_{j < n \leq N} \frac{v(n)}{n(n-1)} \right).$$

We will compute v by descending induction. Note that

$$\begin{aligned} v(N) = 1 = R(N), \quad v(N-1) &= \max \left(\frac{N-1}{N}, (N-1) \frac{1}{N(N-1)} \right) \\ &= \frac{N-1}{N} = R(N-1). \end{aligned}$$

Suppose that $v(n) = R(n)$, $\forall n > j$. Then

$$\begin{aligned} v(j) &= \max \left(\frac{j}{N}, \frac{j}{N} \sum_{j < n \leq N} \frac{1}{n-1} \right) \\ &= \max \left(\frac{j}{N}, \frac{j}{N} f_N(j) \right) = \frac{j}{N} \max(1, f_N(j)). \end{aligned}$$

Note that for $j \geq j^* = j^*(N)$ we have $f_N(j) \leq 1$ and we deduce inductively that

$$v(j) = \frac{j}{N}, \quad \forall j \in \llbracket j^*, N \rrbracket.$$

For $j < j^* - 1$ note that

$$j \sum_{j < n \leq N} \frac{v(n)}{n(n-1)} = \frac{v(j+1)}{(j+1)} + \frac{j}{j+1} v(j+1) = v(j+1).$$

Hence

$$v(j) = v(j^* - 1), \quad \forall j \leq (j^* - 1).$$

where

$$v(j^* - 1) = \frac{j^* - 1}{N} f_N(j^* - 1)$$

Thus the stopping strategy is as indicated and the maximum expected probability if

$$p_N = v_N(1) = \frac{j^* - 1}{N} f_N(j^* - 1).$$

This in agreement with the prediction (1.4.26) obtained in Example 1.4.16.

The value function v_N can be obtained using the R code


```

dynkin=function(N) {
  p=c(rep(1, N))
  for (i in 2:N) {
    p[i]=1/(i*(i-1))
  }
  v=c(1)
  for(i in 1:(N-1) ) {
    f= max( (N-i)/N, (N-i)*sum( p[(N-i+1):N]*v ))
    v=c(f, v)
  }
  v
}

```

Let us consider the case $N = 10$ and compare the vector y displayed in Table 1.4.27 with the value function $v = v_{10}$. In this case $j^*(10) = 4$.

n	1	2	3	4	5	6	7	8	9	10
y_n	0.3986	0.3986	0.3986	0.3986	0.3982	0.372	0.32	0.26	0.18	0.1
$v(n)$	0.3986	0.3986	0.3986	0.4	0.5	0.6	0.7	0.8	0.9	1

(4.3.11)

This shows clearly that the approach we have use here to compute p_N is different from the one used in Example 1.4.16. However, both approaches are based on the same remarkable fact: the rank variables X_j are independent. $\square$

4.4. Asymptotic behavior

Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an *irreducible, positively recurrent* HMC with state space $\mathfrak{X}$ and transition matrix Q . It thus has a unique stationary probability measure $\pi_\infty \in \text{Prob}(\mathfrak{X})$. In this section we will provide a dynamical description of π_∞ and prove a Law of Large Numbers that involves this measure.

4.4.1. The ergodic theorem. Fix an arbitrary state $x_0 \in \mathfrak{X}$, let $T_0 := T_{x_0}$ denote the time of first return to x_0 and denote by π^0 the unique invariant measure on $\mathfrak{X}$ such that $\pi^0[x_0] = 1$. The results in the previous section show that

$$\pi^0[x] = \mathbb{E}_{x_0} \left[\sum_{n \geq 1} \mathbf{I}_{\{X_n=x\}} \mathbf{I}_{\{n \leq T_0\}} \right] = \mathbb{E}_{x_0} \left[\sum_{n=1}^{T_0} \mathbf{I}_{\{X_n=x\}} \right], \quad \forall x \in \mathfrak{X}. \quad (4.4.1)$$

For $n \in \mathbb{N}$ we set

$$\nu(n) = \nu_{x_0}(n) := \sum_{k=1}^n \mathbf{I}_{\{X_k=x_0\}}.$$

In other words, the random variable $\nu_{x_0}(n)$ is the number of returns to x_0 during the interval $[1, n]$.

Proposition 4.4.1. *Suppose that $f \in L^1(\mathfrak{X}, \pi^0)$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{\nu_{x_0}(n)} \sum_{k=1}^n f(X_k) = \int_{\mathfrak{X}} f(x) \pi^0[dx] = \sum_{x \in \mathfrak{X}} f(x) \pi^0[x], \quad \mathbb{P}_{x_0} - \text{a.s.} \quad (4.4.2)$$

Proof. I will follow the proof in [27, Prop. 3.2.6]. Using the decomposition $f = f^+ - f^-$ we see that it suffices to consider only the case when f is nonnegative. Let $T_0 = \tau_1 \leq \tau_2 \leq \dots$ denote the successive times of return to x_0 . We set

$$U_p := \sum_{k=\tau_{p-1}+1}^{\tau_p} f(X_k).$$

The strong Markov property shows that the random variables $U_1, U_2, \dots$ are i.i.d.. We have

$$\begin{aligned} \mathbb{E}_{x_0}[U_1] &= \sum_{k=1}^{T_0} \mathbb{E}_{x_0}[f(X_k)] = \mathbb{E}_{x_0}\left[\sum_{k=1}^{T_0} \sum_{x \in \mathfrak{X}} f(x) \mathbf{I}_{\{X_k=x\}}\right] \\ &= \sum_{x \in \mathfrak{X}} f(x) \mathbb{E}_{x_0}\left[\sum_{k=1}^{T_0} \mathbf{I}_{\{X_k=x\}}\right] \stackrel{(4.4.1)}{=} \sum_{x \in \mathfrak{X}} f(x) \pi^0[x]. \end{aligned}$$

The Strong Law of Large Numbers implies

$$\lim_{p \rightarrow \infty} \frac{1}{p} \sum_{k=1}^p U_k = \sum_{x \in \mathfrak{X}} f(x) \pi^0[x], \quad \mathbb{P}_{x_0} - \text{a.s.}$$

In other words

$$\frac{1}{p} \sum_{k=1}^{\tau_p} f(X_k) \rightarrow \sum_{x \in \mathfrak{X}} f(x) \pi^0[x], \quad \mathbb{P}_{x_0} - \text{a.s.}$$

Observing that $\tau_{\nu(n)} \leq n < \tau_{\nu(n)+1}$, we deduce that for nonnegative f we have

$$\begin{aligned} \frac{1}{\nu(n)} \sum_{k=1}^{\tau_{\nu(n)}} f(X_k) &\leq \frac{1}{\nu(n)} \sum_{k=1}^n f(X_k) \leq \frac{1}{\nu(n)} \sum_{k=1}^{\tau_{\nu(n)+1}} f(X_k) \\ &= \frac{\nu(n)+1}{\nu(n)} \frac{1}{\nu(n)+1} \sum_{k=1}^{\tau_{\nu(n)+1}} f(X_k). \end{aligned}$$

Since the chain is recurrent we have $\nu(n) \rightarrow \infty$ so

$$\lim_{n \rightarrow \infty} \frac{\nu(n)+1}{\nu(n)} = 1.$$

Hence the extremes in the last inequality converge to $\sum_{x \in \mathfrak{X}} f(x) \pi^0[x]$, $\mathbb{P}_{x_0} - \text{a.s.}$ □

Corollary 4.4.2. *We have*

$$\frac{\nu(n)}{n} \rightarrow \frac{1}{\mathbb{E}_{x_0}[T_{x_0}]} = \pi_\infty[x_0], \quad \mathbb{P}_{x_0} - \text{a.s.} \quad (4.4.3)$$

Proof. Let $f = 1$ in Proposition 4.4.1. This f is integrable so

$$\frac{n}{\nu(n)} \rightarrow \pi^0[\mathfrak{X}] \stackrel{(4.2.11)}{=} \mathbb{E}_{x_0}[T_{x_0}].$$

□

Corollary 4.4.3 (Ergodic Theorem). *Suppose that $(X_n)_{n \geq 0}$ is a positively recurrent irreducible HMC with state space $\mathfrak{X}$, transition matrix Q and stationary distribution π_∞ . Let $f \in L^1(\mathfrak{X}, \pi_\infty)$. Then, for any $\mu \in \text{Prob}(\mathfrak{X})$ we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(X_k) = \int_{\mathfrak{X}} f(x) \pi_\infty[dx], \quad \mathbb{P}_\mu - \text{a.s.} \quad (4.4.4)$$

Proof. Assume first that (X_n) are defined on the path space $(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu)$.

Suppose $\mu = \delta_{x_0}$. If we divide both sides of (4.4.2) by $\mathbb{E}_{x_0}[T_{x_0}]$ we deduce

$$\lim_{n \rightarrow \infty} \frac{1}{\nu(n) \mathbb{E}_{x_0}[T_{x_0}]} \sum_{k=1}^n f(X_k) = \int_{\mathfrak{X}} f(x) \pi_\infty[dx].$$

Now observe that

$$\frac{1}{n} \sum_{k=1}^n f(X_k) = \frac{n}{\nu(n) \mathbb{E}_{x_0}[T_{x_0}]} \frac{1}{n} \sum_{k=1}^n f(X_k),$$

and (4.4.3) implies

$$\frac{n}{\nu(n) \mathbb{E}_{x_0}[T_{x_0}]} \rightarrow 1.$$

More generally, for any $\mu \in \text{Prob}(\mathfrak{X})$,

$$\mathbb{P}_\mu = \sum_{x \in \mathfrak{X}} \mu[x] \mathbb{P}_x \in \text{Prob}(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}).$$

we denote by $\mathcal{C}$ the subset of $\mathfrak{X}^{\mathbb{N}_0}$ consisting of sequences

$$\underline{x} = (x_0, x_1, \dots) \in \mathfrak{X}^{\mathbb{N}_0}$$

such that

$$\lim_{n \rightarrow \infty} \frac{1}{n+1} (f(x_0) + f(x_1) + \dots + f(x_n)) = \int_{\mathfrak{X}} f(x) \pi_\infty[dx].$$

From the above we deduce that $\mathbb{P}_x[\mathcal{C}] = 1, \forall x \in \mathfrak{X}$. Then

$$\mathbb{P}_\mu[\mathcal{C}] \stackrel{(4.1.13)}{=} \sum_{x \in \mathfrak{X}} \mu[x] \mathbb{P}_x[\mathcal{C}] = 1.$$

Suppose that random maps (X_n) are defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$, not necessarily the path space. Using the map $\vec{X} : \Omega \rightarrow \mathfrak{X}^{\mathbb{N}_0}$ we reduce this case to the situation we have discussed above. $\square$

The above Ergodic Theorem is a Law of Large Numbers for a sequence of *dependent* random variables!

4.4.2. Aperiodic chains. When $(X_n)_{n \geq 0}$ is an irreducible, *aperiodic*, positively recurrent HMC the Ergodic Theorem can be considerably strengthened. We need to introduce some terminology.

Let $\mathbb{X}$ be a Polish space with Borel sigma-algebra $\mathcal{B}$. The *variation distance* $d_v(\mu, \nu)$ between two Borel probability measures $\mu_0, \mu_1 \in \text{Prob}(\mathbb{X})$ is defined by

$$d_v(\mu_0, \mu_1) = \sup_{B \in \mathcal{B}} |\mu_0[B] - \mu_1[B]| = \sup_{B \in \mathcal{B}} (\mu_0[B] - \mu_1[B]) \quad (4.4.5)$$

The second equality follows from the elementary observation

$$\left| \mu_0[A] - \mu_1[A] \right| = \max \left\{ (\mu_0[B] - \mu_1[B]), (\mu_0[B^c] - \mu_1[B^c]) \right\}.$$

The variation distance defines a metric on the space $\text{Prob}(\mathbb{X})$ of Borel probability measures on $\mathbb{X}$. We will refer to it as the *variation metric*.

If X, Y are $\mathbb{X}$ -valued random variables, then the variation distance between them is defined to be the variation distance between their distributions $\mathbb{P}_X, \mathbb{P}_Y$,

$$d_v(X, Y) := d_v(\mathbb{P}_X, \mathbb{P}_Y).$$

Lemma 4.4.4. *Let $\mu_0, \mu_1 \in \text{Prob}(\mathbb{X})$. Suppose there exists a sigma-finite Borel measure ν on $\mathbb{X}$ such that both μ_0 and μ_1 are absolutely continuous with respect to ν . We denote by $p_i(x)$ the density of μ_i with respect to ν , i.e.*

$$\mu_i[dx] = p_i(x)\nu[dx], \quad i = 0, 1.$$

Then

$$d_v(\mu_0, \mu_1) := \frac{1}{2} \int_{\mathbb{X}} |p_0(x) - p_1(x)| \nu[dx]. \quad (4.4.6)$$

In particular, if $\mathbb{X}$ is finite or countable and ν is the standard counting measure, then

$$d_v(\mu_0, \mu_1) = \frac{1}{2} \sum_{x \in \mathbb{X}} |p_0(x) - p_1(x)|. \quad (4.4.7)$$

Proof. Define

$$D^\pm := \left\{ x \in \mathbb{X}; \pm(p_0(x) - p_1(x)) > 0 \right\}.$$

Note that

$$\begin{aligned} 0 &= \int_{\mathbb{X}} (p_0(x) - p_1(x)) \nu[dx] = \int_{D^+} (p_0(x) - p_1(x)) \nu[dx] \\ &\quad + \int_{D^-} (p_0(x) - p_1(x)) \nu[dx], \\ \int_{\mathbb{X}} |p_0(x) - p_1(x)| \nu[dx] &= \int_{D^+} (p_0(x) - p_1(x)) \nu[dx] \\ &\quad - \int_{D^-} (p_0(x) - p_1(x)) \nu[dx] \\ &= 2 \int_{D^+} (p_0(x) - p_1(x)) \nu[dx] = 2(\mu_0[D^+] - \mu_1[D^+]). \end{aligned}$$

Observe that for any $B \in \mathcal{B}$ we have

$$\begin{aligned} \mu_0[B] - \mu_1[B] &= \int_{B \cap D^+} (p_0(x) - p_1(x)) \nu[dx] + \int_{B \cap D^-} (p_0(x) - p_1(x)) \nu[dx] \\ &\leq \int_{B \cap D^+} (p_0(x) - p_1(x)) \nu[dx] \leq \int_{D^+} (p_0(x) - p_1(x)) \nu[dx]. \end{aligned}$$

In above inequality we have equality when $B = D^+$. Hence

$$\sup_{B \in \mathcal{B}} (\mu[B] - \nu[B]) = \mu_0[D^+] - \mu_1[D^+] = \frac{1}{2} \int_{\mathbb{X}} |p_0(x) - p_1(x)| \nu[dx].$$

□

Definition 4.4.5. The *coupling time* of two $\mathfrak{X}$ -valued stochastic processes $(X_n)_{n \in \mathbb{N}_0}$ and $(Y_n)_{n \in \mathbb{N}_0}$ is a stopping time T of the process (X_n, Y_n) such that

$$X_n = Y_n \quad \forall n \geq T.$$

The stochastic processes are said to *couple* if they admit an a.s. finite coupling time. $\square$

Lemma 4.4.6. Suppose that the $\mathfrak{X}$ -valued processes $(X_n)_{n \in \mathbb{N}_0}$ and $(Y_n)_{n \in \mathbb{N}_0}$ couple with coupling time T . Then

$$d_v(X_n, Y_n) \leq \mathbb{P}[T > n].$$

In particular, if T is a.s. finite,

$$\lim_{n \rightarrow \infty} d_v(X_n, Y_n) = 0.$$

Proof. For all $A \subset \mathfrak{X}$ we have

$$\begin{aligned} \mathbb{P}[X_n \in A] - \mathbb{P}[Y_n \in A] &= \boxed{\mathbb{P}[X_n \in A, T \leq n]} + \mathbb{P}[X_n \in A, T > n] \\ &\quad - \boxed{\mathbb{P}[Y_n \in A, T \leq n]} - \mathbb{P}[Y_n \in A, T > n] \\ (X_n = Y_n, \forall n \geq T) \\ &= \mathbb{P}[X_n \in A, T > n] - \mathbb{P}[Y_n \in A, T > n] \\ &\leq \mathbb{P}[X_n \in A, T > n] \leq \mathbb{P}[T > n]. \end{aligned}$$

$\square$

Theorem 4.4.7. Suppose that Q is a probability transition matrix on the state space $\mathfrak{X}$ such that the associated HMC's are irreducible, aperiodic and positively recurrent. Denote by π the unique invariant probability measure on $\mathfrak{X}$. Then, for any $\mu \in \text{Prob}(\mathfrak{X})$

$$\lim_{n \rightarrow \infty} d_v(\mu Q^n, \pi) = 0. \quad (4.4.8)$$

In particular, if $\mu = \delta_{x_0}$, we deduce that

$$\pi[x] = \lim_{n \rightarrow \infty} Q^n_{x_0, x}. \quad (4.4.9)$$

Proof. Consider two independent HMCs $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \geq 0}$ with state space $\mathfrak{X}$, transition matrix Q such that initial distribution of (X_n) is μ and the initial distribution of Y_n is π . Since π is stationary, the probability distribution of Y_n is π , $\forall n$.

We will construct a third HMC (Z_n) with state space $\mathfrak{X}$ and transition matrix Q such that Y_n and Z_n couple and $\mathbb{P}_{X_n} = \mathbb{P}_{Z_n}$, $\forall n$. Then $d_v(X_n, Y_n) = d_v(Z_n, Y_n)$ and Lemma 4.4.6 implies

$$\lim_{n \rightarrow \infty} d_v(Z_n, Y_n) = 0.$$

Consider the stochastic process (X_n, Y_n) . This is an HMC with state space $\mathfrak{X} \times \mathfrak{X}$ and transition matrix $\widehat{Q}$

$$\widehat{Q}_{(x_0, y_0), (x_1, y_1)} = Q_{x_0, x_1} \cdot Q_{y_0, y_1}.$$

Since Q is irreducible and aperiodic we deduce that for any $x_0, x_1, y_0, y_1 \in \mathfrak{X}$, $\exists N_0 > 0$ such that $Q^n_{x_1, x_1} \cdot Q^n_{y_1, y_1} > 0$, $\forall n \geq N_0$. We deduce that there exists $N \geq N_0$ such that

$$Q^n_{x_0, x_1} \cdot Q^n_{y_0, y_1} > 0, \quad \forall n \geq N.$$

This shows that $\widehat{Q}$ is irreducible and aperiodic.

The product measure $\pi \otimes \pi$ is an invariant probability measure of the chain (X_n, Y_n) . Theorem 4.2.42 implies that $\hat{Q}$ is positively recurrent. Proposition 4.2.44 shows that for any $x_{x_0} \in \mathfrak{X}$, the chain (X_n, Y_n) will almost surely return to (x_0, x_0) in finite time. In other words, for any $x_0 \in \mathfrak{X}$ the stopping time

$$T_{x_0} := \min \{ n \in \mathbb{N}; X_n = Y_n = x_0 \}$$

is a.s. finite. Fix $x_0 \in \mathfrak{X}$ set $T = T_{x_0}$ and define

$$Z_n = \begin{cases} X_n, & n < T, \\ Y_n, & n \geq T. \end{cases}$$

Clearly T is an a.s. finite coupling time for the processes (Y_n) and Z_n . It remains to prove that (X_n) and Z_n have the same distributions.

Indeed, the strong Markov property for the Markov chains with state space $\mathfrak{X} \times \mathfrak{X}$ and transition matrix $\hat{Q}$ shows that the process (X_{T+n}, Y_{T+n}) is a Markov chain on $\mathfrak{X} \times \mathfrak{X}$ has the same distribution as the Markov process (A_n, B_n) with the same transition matrix $\hat{Q}$ and initial condition $A_0 = B_0 = x_0 = X_T = Y_T$. In particular this shows that the processes X_{T+n} and Y_{T+n} have the same distributions. $\square$

Remark 4.4.8. Suppose that Q is the transition matrix of an irreducible, positively recurrent Markov chain with state space $\mathfrak{X}$ and invariant probability measure π . We form a new stochastic matrix

$$\tilde{Q} = \frac{1}{2}(1 + Q).$$

The chain with this transition matrix is called *the lazy version* of the original chain. It is irreducible and π is the invariant probability measure of the lazy version as well. However, the lazy chain is also *aperiodic*, even if the original chain is not. This follows from the equality

$$\tilde{Q}_{x,y}^n = \sum_{k=0}^n \frac{1}{2^n} \binom{n}{k} Q_{x,y}^k.$$

This shows that if $Q_{x,y}^k \neq 0$, then $\tilde{Q}_{x,y}^n > 0$, $\forall n \geq k$. Using the terminology of generalized convergence in [100], we can say that the *Euler means* of the sequence $(Q_{x,y}^n)_{n \geq 0}$ converge to the invariant measure. $\square$

4.4.3. The coupling technique. The technique behind the above proof of Theorem 4.4.7 was pioneered by W. Doeblin [59] in 1938. It took almost three decades for the mathematical community to appreciate the novelty of his ideas, distill the key principles, and organize them into what is now referred to as the *coupling technique*.

I want to describe a few coupling concepts that will put the results of the previous subsection in the proper context. For more details and applications I refer to [138, 209].

Definition 4.4.9. Let $\mathbb{X}$ be a Polish space with Borel algebra $\mathcal{B}$.

- (i) A *coupling* of a family $(\mu_i)_{i \in I}$ of Borel probability measures on $\mathbb{X}$ is a probability measure $\hat{\mu}$ on $(\mathbb{X}^I, \mathcal{B}^I)$ such that

$$\hat{\mu}_i = \mu_i, \quad \forall i \in I,$$

where $\hat{\mu}_i$ is the i -th marginal of $\hat{\mu}$, i.e., $\hat{\mu}_i$ is the pushforward of $\hat{\mu}$ under the natural projection

$$\pi_i : \mathbb{X}^I \rightarrow \mathbb{X}, \quad (x_i)_{i \in I} \mapsto x_i.$$

We will use the notation $\hat{\mu} \in \text{Couple}(\mu_i, i \in I)$ to indicate that $\hat{\mu}$ is a coupling of the family $(\mu_i)_{i \in I}$

- (ii) A *coupling* of a family $\mathbb{X}$ -valued random variables $(X_i)_{i \in I}$, defined on *possibly different probability spaces*, is a family of $\mathbb{X}$ -valued random variables $(\hat{X}_i)_{i \in I}$, on the same probability space,

$$\hat{X}_i : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{B}), \quad i \in I,$$

such that $\mathbb{P}_{\hat{X}_i} = \mathbb{P}_{X_i}, \forall i \in I$.

□

Let $\Delta \subset \mathbb{X}^2$ be the diagonal

$$\Delta = \{(x_0, x_1) \in \mathbb{X}^2; x_0 := x_1\}.$$

The set Δ is a closed subset of $\mathbb{X}^2$ and thus it is Borel measurable. Since $\mathbb{X}^2$ is a Polish space we have (see Exercise 1.5)

$$\mathcal{B}_{\mathbb{X}^2} = \mathcal{B}_{\mathbb{X}} \otimes \mathcal{B}_{\mathbb{X}}$$

so Δ is also $\mathcal{B}_{\mathbb{X}}^{\otimes 2}$ -measurable. In particular $\mathbb{X}_*^2 = \mathbb{X}^2 \setminus \Delta$ is also $\mathcal{B}_{\mathbb{X}}^{\otimes 2}$ -measurable.

The next result explains the relevance of couplings in estimating the variation distance between two measures.

Proposition 4.4.10. *Let $\mathbb{X}$ be a Polish and μ, ν two Borel probability measures on $\mathbb{X}$. Set*

$$\mathbb{X}_*^2 := \{(x_0, x_1) \in \mathbb{X}^2; x_0 \neq x_1\}.$$

Then,

$$d_v(\mu, \nu) \leq \lambda[\mathbb{X}_*^2], \quad \forall \lambda \in \text{Couple}(\mu, \nu). \quad (4.4.10)$$

Proof. For any $B \in \mathcal{B}$ we have

$$\mathbb{X}_*^2 \supset B \times B^c = B \times \mathbb{X} \setminus B \times B$$

so

$$\begin{aligned} \lambda[\mathbb{X}_*^2] &\geq \lambda[B \times \mathbb{X}] - \lambda[B \times B] \\ &\geq \lambda[B \times \mathbb{X}] - \lambda[\mathbb{X} \times B] = \mu[B] - \nu[B]. \end{aligned}$$

Hence

$$\lambda[\mathbb{X}_*^2] \geq \sup_{B \in \mathcal{B}} (\mu[B] - \nu[B]) = d_v(\mu, \nu).$$

□

Remark 4.4.11. The inequality (4.4.10) is optimal. When μ, ν are absolutely continuous with respect to a Borel measure β on $\mathbb{X}$ we can explicitly describe a coupling λ so that we have equality in (4.4.10). We call such a coupling *optimal*.

More precisely, if we set

$$\mu[dx] = p(x)\beta[dx], \quad \nu[dx] = q(x)\beta[dx].$$

we define

$$\lambda_{\mu, \nu, \beta}[dx_0 dx_1] = \rho(x_0, x_1)\beta^{\otimes 2}[dx_0 dx_1]$$

where

$$\rho(x_0, x_1) = \begin{cases} p(x_0) \wedge q(x_0), & x_0 = x_1, \\ \frac{(p(x_0) - p(x_0) \wedge q(x_0))(q(x_1) - p(x_1) \wedge q(x_1))}{d_v(\mu, \nu)}, & x_0 \neq x_1. \end{cases}$$

Above, d_v is defined by (4.4.6) and we use the convention $\frac{0}{0} := 0$. A simple computation shows that indeed $\lambda_{\mu, \nu, \beta}$ is an optimal coupling.

Let us observe any two Borel probability measures μ, ν on $\mathbb{X}$ are simultaneously absolutely continuous with respect to the probability measure $\beta = \frac{1}{2}(\mu + \nu)$. Hence for any two Borel probability measures on $\mathbb{X}$ there exists an optimal coupling. $\square$

Remark 4.4.12. Suppose that $\hat{Y}, \hat{Z} : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{X}$ define a coupling of the $\mathbb{X}$ -valued random maps Y, Z . A *coupling event* is an event whose occurrence guarantees that $\hat{Y} = \hat{Z}$.

Formally, a coupling event is a measurable set $S \subset \mathcal{S}$ such that $\hat{Y}(\omega) = \hat{Z}(\omega)$, $\forall \omega \in S$. If we denote by Φ the measurable map

$$\Phi : \Omega \rightarrow \mathbb{X}^2, \quad \Phi(\omega) = (\hat{Y}(\omega), \hat{Z}(\omega)),$$

then S is a coupling event iff $S \subset \Phi^{-1}(\Delta)$. This shows that if S is a coupling event, then $S^c \supset \Phi^{-1}(\mathbb{X}_*^2)$, so that

$$1 - \mathbb{P}[S] = \mathbb{P}[S^c] \geq \mathbb{P}_{(\hat{Y}, \hat{Z})}[\mathbb{X}_*^2] \geq d_v(Y, Z).$$

Suppose that the $\mathbb{X}$ -valued stochastic processes $(X_n)_{n \in \mathbb{N}_0}$ and $(Y_n)_{n \in \mathbb{N}_0}$ couple. Denote by T the coupling time T . Define a new process $(Z_n)_{n \in \mathbb{N}_0}$ by setting

$$Z_n = \begin{cases} Y_n, & T > n, \\ X_n, & T \leq n. \end{cases}$$

Note that for any n the pair (X_n, Z_n) is a coupling of (X_n, Y_n) . The event $\{T \leq n\}$ is a coupling event and we deduce

$$d_v(X_n, Y_n) \leq 1 - \mathbb{P}[T \leq n] = \mathbb{P}[T > n].$$

This is precisely the conclusion of Lemma 4.4.6. $\square$

4.5. Electric networks

In this section I will discuss a very versatile method, inspired from physics, of investigating the transience or the recurrence of reversible Markov chains.

4.5.1. Reversible Markov chains as electric networks. Suppose that $(X_n)_{n \geq 1}$ is an irreducible, reversible, *locally finite* HMC with state space $\mathfrak{X}$ and transition matrix Q . We recall that local finiteness means that

$$\forall x \in \mathfrak{X}, \quad \#\{y \in \mathfrak{X}; Q_{x,y} \neq 0\} < \infty.$$

The reversibility means that there exists a function $c : \mathfrak{X} \rightarrow (0, \infty)$ such that

$$c(y)Q_{y,x} = c(x)Q_{x,y} \quad \forall x, y \in \mathfrak{X}. \quad (4.5.1)$$

Observe that any positive multiple of c also satisfies (4.5.1). We set

$$c(x, y) := c(x)Q_{x,y}, \quad \forall x, y \in \mathfrak{X}.$$

The detail balance condition (4.5.1) shows that $c(x, y) = c(y, x)$ and $c(x, y) \neq 0$ iff $Q_{x,y} > 0$. Note that

$$Q_{x,y} = \frac{c(x, y)}{c(x)}, \quad c(x) = \sum_{y \sim x} c(x, y). \quad (4.5.2)$$

It is convenient to visualize this Markov chain as a random walk on an *undirected* graph with vertex set $\mathfrak{X}$ and *weighted* edges. Two vertices x, y are connected by an edge if and only if $Q_{x,y} > 0$. Since the Markov chain is irreducible, this graph is connected.

We use the notation $x \sim y$ to indicate that the vertices /nodes x, y are connected by an edge. We say that two vertices x, y are neighbors if $x \sim y$. For $x \in \mathfrak{X}$ we denote by $N(x)$ the set of neighbors of x . If $y \in N(x)$, then we weigh the connecting edge with the weight $c(x, y) = c(y, x)$.

We will assume $c(x, x) = 0, \forall x \in \mathfrak{X}$, i.e., the associated graph has no loops.

The Markov chain dynamics has the following equivalent description: if the system is at the state/vertex x it will transition to a neighbor y with probability $Q_{x,y}$ proportional to the weight $c(x, y)$. The weights $\{c(x)\}_{x \in \mathfrak{X}}$ define a Q -invariant measure μ_c on $\mathfrak{X}$

$$\mu_c[S] = \sum_{s \in S} c(s). \quad (4.5.3)$$

Formally, an *electric network* is a triplet $(\mathfrak{X}, E, c)$, where $(\mathfrak{X}, E)$ is a locally finite, connected, unoriented graph and $c : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, \infty)$ is a function. The set of vertices $\mathfrak{X}$ is assumed to be at most countable. We regard the set of edges E as a symmetric subset of $\mathfrak{X} \times \mathfrak{X}$, i.e., $(x, y) \in E \iff (y, x) \in E$. We assume there are no loops, i.e., $\forall (x, y) \in E, x \neq y$. We will frequently use the notation $x \sim y$ to indicate that $(x, y) \in E$.

The function $c : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, \infty)$ satisfies

- $c(x, y) > 0 \iff (x, y) \in E$.
- $c(x, y) = c(y, x), \forall (x, y) \in E$.

We have seen that a reversible Markov chain determines an electric network.

Conversely, an electric network $(\mathfrak{X}, E, c)$ determines a reversible Markov chain with state space $\mathfrak{X}$ and transition matrix $Q : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1]$

$$Q_{x,y} = \frac{c(x, y)}{c(x)}, \quad c(x) = \sum_{y \in N(x)} c(x, y)$$

An *electric network* corresponds to a real electric network in which an edge between two nodes x, y corresponds to a resistor between these two nodes with *resistance*

$$r(x, y) = \frac{1}{c(x, y)}.$$

The quantity $c(x, y)$ is called *conductance*.

4.5.2. Sources, currents and chains. The connection between electric networks and the dynamics associated Markov chain is through the classical physical laws of Kirchhoff and Ohm. As shown in the pioneering work of Nash-Williams [155], this point of view can shed remarkable insight into the behavior of the Markov chains. In the remainder of this section I will highlight some of this fruitful interplay between probability and physics. For more about this I refer to [62, 93, 141, 196] which served as my main sources of inspiration.

First, a matter of notation. For every pair of elements s, s' of a set S we denote by $\delta_{s,s'}$ the Kronecker symbol

$$\delta_{s,s'} = \begin{cases} 1, & s = s', \\ 0, & s \neq s'. \end{cases}$$

As observed by R. Bott and by H. Weyl, see e.g. [23], the physical laws of electric networks have simple geometric interpretations, best expressed in the language of Hodge theory.

The main objects in Hodge theory are the *chain/cochain complexes*. To define them we need to make some choices.

Consider a locally finite graph $(\mathfrak{X}, E)$. An *orientation* of the graph is a subset $E_+ \subset E$ such that for any edge $(x, y) \in E$ either $(x, y) \in E_+$ or $(y, x) \in E_+$, but not both.

One can obtain such an E_+ by assigning orientations (arrows) along the edges. Define E_+ as the collection of *positively oriented* edges. More precisely $(x, y) \in E_+$, if and only if the arrow of the oriented edge goes from x to y .

The vector space of 0-chains, denoted by C_0 , consists of formal sums of the type

$$j := \sum_{x \in \mathfrak{X}} j(x)[x], \quad j(x) \in \mathbb{R}, \quad \forall x \in \mathfrak{X}.$$

Equivalently, $C_0 = \mathbb{R}^{\mathfrak{X}}$. In physics a 0-chain is known as a *source* (of current) and $j(x) = 0$ for all but finitely many x .

The vector space C_1 of 1-chains consists of skew-symmetric functions

$$i : E \rightarrow \mathbb{R}, \quad (x, y) \mapsto i(x, y).$$

For any $(x, y) \in E$, define $[x] \otimes [y] : E \rightarrow \mathbb{R}$ by setting

$$[x] \otimes [y](e) = \begin{cases} 1, & e = (x, y) \\ -1, & e = (y, x), \\ 0 & \text{otherwise.} \end{cases}$$

If we fix an orientation E_+ , we will identify an oriented edge $e = (x, y) \in E_+$ with the current $[x] \otimes [y]$ and we write $i_e := [x] \otimes [y]$.

Once we fix an orientation E_+ we can describe each current as a formal sum of the type

$$i = \sum_{(x,y) \in E_+} i(x, y)[x] \otimes [y] = \frac{1}{2} \sum_{(x,y) \in E} i(x, y)[x] \otimes [y].$$

where E_+ is an orientation of the edges. In physics, 1-chains are known as *currents*.

One should think of $[x] \otimes [y]$ as representing the edge (x, y) oriented from x to y . A current can then visualized as an assignment of arrows and weights on edges with the understanding that we get the same current if we reverse any of the arrows and change its weight to the opposite.

A 0-chain j is called *compactly supported* if $j(x) = 0$ for all but finitely many x . Similarly, a 1-chain i is *compactly supported* if $i(x, y) = 0$ for all but finitely many edges $(x, y) \in E$. For $k = 0, 1$ we denote by C_k^{cpt} the space of compactly supported k -chains. There are *boundary operators*

$$\partial : C_1 \rightarrow C_0, \quad \partial : C_0^{\text{cpt}} \rightarrow \mathbb{R}$$

defined as follows.

- If $\mathbf{i} \in C_1$, then

$$\partial \mathbf{i} := \sum_{x \in \mathfrak{X}} w(x)[x], \quad w(x) = \sum_{y \in N(x)} i(x, y) = - \sum_{y \in N(x)} i(y, x), \quad \forall x \in \mathfrak{X}.$$

In particular, for $x_0, x_1 \in \mathfrak{X}$,

$$\partial[x_0] \oslash [x_1] = [x_1] - [x_0].$$

- If $\mathbf{j} \in C_0^{\text{cpt}}$, then

$$\partial \mathbf{j} = \sum_{x \in \mathfrak{X}} j(x) \in \mathbb{R}$$

Let us observe that for any *compactly supported* current $\mathbf{i}$ we have $\partial^2 \mathbf{i} = 0$. Indeed

$$\partial(\partial \mathbf{i}) = \sum_{x \in \mathfrak{X}} \sum_{y \in N(x)} i(x, y) = \sum_{(x, y) \in E} i(x, y) = 0$$

since $i(x, y) + i(y, x) = 0$ whenever $x \sim y$.

Remark 4.5.1. If $\mathfrak{X}$ is infinite, then there could exist 1-chains $\mathbf{i}$ such that $\partial \mathbf{i} \in C_0^0$, yet $\partial^2 \mathbf{i} \neq 0$. $\square$

The (finite) paths in the graph are special examples of compactly supported 1-chains. By a path of length n we understand a sequence of neighbors

$$x_0, x_1, \dots, x_n, \quad x_{k-1} \sim x_k, \quad x_{k-1} \neq x_k, \quad \forall k = 1, \dots, n.$$

The associated 1-chain $\mathbf{i} = \mathbf{i}_{x_0, x_1, \dots, x_n}$ is

$$\mathbf{i}_{x_0, x_1, \dots, x_n} = \sum_{k=1}^n [x_{k-1}] \oslash [x_k].$$

Note that

$$\partial \mathbf{i}_{x_0, x_1, \dots, x_n} = [x_n] - [x_0].$$

The path is closed if $x_0 = x_n$ or, equivalently, $\partial \mathbf{i}_{x_0, x_1, \dots, x_n} = 0$.

4.5.3. Kirchhoff's laws and Hodge theory. The actual sources and currents in real electric network are governed by Kirchhoff's laws. We refer to [11, Chap.12] for a more detailed description of the physical aspects. Fix an electrical network $(\mathfrak{X}, E, c)$.

Kirchhoff's first law states that the source of a (physical) current $\mathbf{i} \in C_1$ is the 0-chain $\mathbf{j} = -\partial \mathbf{i}$. More explicitly, this means that

$$j(x) + \sum_{y \in N(x)} i(x, y) = 0, \quad \forall x \in \mathfrak{X}. \quad (4.5.4)$$

This is a purely topological condition in the sense that it is independent of the choice of conductance function. The physics/geometry enters the scene through the conductance function.

The *energy* of a 0-chain $\mathbf{j}$ is the quantity

$$\mathcal{E}_r[\mathbf{j}] := \sum_{x \in \mathfrak{X}} \frac{1}{c(x)} j(x)^2.$$

We denote by C_0^∞ the space of finite energy 0-chains. It is a Hilbert space with respect to the inner product

$$\langle j_1, j_2 \rangle = \sum_{x \in \mathfrak{X}} \frac{1}{c(x)} j_1(x) j_2(x).$$

It contains C_0^{cpt} as a dense subspace.

The *energy*⁴ of a current i in an electric network is defined by

$$\mathcal{E}[i] := \frac{1}{2} \sum_{(x,y) \in E} r(x,y) i(x,y)^2. \quad (4.5.5)$$

If we fix an orientation E_+ of the edges we obtain the equivalent description

$$\mathcal{E}[i] = \sum_{(x,y) \in E_+} r(x,y) i(x,y)^2 = \sum_{(x,y) \in E_+} \frac{i(x,y)^2}{c(x,y)}. \quad (4.5.6)$$

From a geometric viewpoint we can interpret $r(x,y) i(x,y)^2$ as the square of the length of the edge (x,y) .

In physics currents are assumed to have finite energy. We denote by C_1^∞ the space of finite energy 1-chains. The space C_1^∞ is a Hilbert space with respect to the (resistor) inner product

$$\langle i_1, i_2 \rangle_r := \sum_{(x,y) \in E_+} r(x,y) i_1(x,y) i_2(x,y). \quad (4.5.7)$$

Thus, a physical current is an element of C_1^∞ .

To formulate Kirchhoff's second law we need to introduce the concept of *cochain*. The *cochains* are objects dual to chains.

The space of 0-cochains (resp. 1-cochains) is the dual vector space of C_0^{cpt} (resp. C_1^{cpt})

$$C^0 = \text{Hom}(C_0^{\text{cpt}}, \mathbb{R}), \quad C^1 := \text{Hom}(C_1^{\text{cpt}}, \mathbb{R}).$$

One we can think of a 0-cochain as a function $u : \mathfrak{X} \rightarrow \mathbb{R}$. Physicists call such functions *potentials*. For each $x \in \mathfrak{X}$ denote by $\delta^x \in C^0$ the elementary 0-cochain defined by

$$\delta^x([y]) = \delta_{x,y}, \quad \forall y \in \mathfrak{X}.$$

A 0-cochain is then a formal sum

$$u = \sum_{x \in \mathfrak{X}} u(x) \delta^x.$$

For each $(x,y) \in E$ denote by $dx \otimes dy : E \rightarrow \mathbb{R}$ the elementary 1-cochain defined by

$$dx \otimes dy(x', y') = \begin{cases} 1, & (x', y') = (x, y), \\ -1, & (x', y') = (y, x), \\ 0, & \text{otherwise.} \end{cases}$$

A 1-cochain should be viewed as a formal sum

$$v = \sum_{(x,y) \in E_+} v(x,y) dx \otimes dy = \frac{1}{2} \sum_{(x,y) \in E} v(x,y) dx \otimes dy, \quad v(x,y) = -v(y,x).$$

More concretely, we identify a 1-cochain with a skew-symmetric function on $v : E \rightarrow \mathbb{R}$. In physics such a function is called *voltage* and it is measured in Volts.

⁴The physical units of the expression in (4.5.6) are indeed the units for energy, Joules.

We define the “integral” of a 1-cochain v along a path

$$\gamma = x_0, x_1, \dots, x_n,$$

to be the real number

$$\int_{\gamma} v := \sum_{k=1}^n v(x_{k-1}, x_k).$$

There exists a *coboundary operator* $d : C^0 \rightarrow C^1$ that associates to each function $u : \mathfrak{X} \rightarrow \mathbb{R}$ its “differential”

$$du = \sum_{(x,y) \in E_+} (u(y) - u(x)) dx \otimes dy \quad (4.5.8)$$

A 1-cochain v is called *exact* if it is the differential of a 0-cochain. Table 4.1 summarizes the terminology introduced so far.

Dimension	Chain	Cochain
0	source	potential
1	current	voltage

Table 4.1. A geometry $\leftrightarrow$ physics dictionary

The following fact is left to the reader as an exercise.

Lemma 4.5.2. *A 1-cochain v is exact if and only if its integral along any closed path is 0. Equivalently, this means that the integral along a path depends only on the endpoints of the path.* $\square$

Denote by C_{∞}^0 space of finite energy 0-chains, i.e., 0-chains u satisfying

$$\sum_{x \in \mathfrak{X}} c(x) u(x)^2.$$

This defines an inner product on C_{∞}^0

$$\langle u_1, u_2 \rangle_c = \sum_{x \in \mathfrak{X}} c(x) u_1(x) u_2(x). \quad (4.5.9)$$

As such, C_{∞}^0 can be identified with the Hilbert space $L^2(\mathfrak{X}, \mu_c)$ where μ_c is the Q -invariant measure on $\mathfrak{X}$ determined by the detailed balance equations; see (4.5.3). The space C_{cpt}^0 the compactly supported functions $u : \mathfrak{X} \rightarrow \mathbb{R}$ is a dense subspace of C_{∞}^0 .

We have an isomorphism of Hilbert spaces

$$\mathcal{R}_0 : C_0^{\infty} \rightarrow C_{\infty}^0, \quad \mathcal{R}_0 \mathbf{j}(x) = \frac{1}{c(x)} j(x).$$

The *energy* of a 1-cochain

$$v = \sum_{(x,y) \in E_+} v(x,y) dx \otimes dy$$

is

$$\mathcal{E}[v] := \sum_{(x,y) \in E_+} c(x,y) v(x,y)^2 = \sum_{(x,y) \in E_+} \frac{1}{r(x,y)} v(x,y)^2 = \frac{1}{2} \sum_{(x,y) \in E} c(x,y) v(x,y)^2.$$

We denote by C_∞^1 the space of finite energy 1-cochains. It is a Hilbert space with (conductance) inner product

$$\langle v_1, v_2 \rangle := \sum_{(x,y) \in E_+} c(x,y) v_1(x,y) v_2(x,y). \quad (4.5.10)$$

Hence

$$\mathcal{E}[v] = \langle v, v \rangle.$$

We have a “resistor” duality map

$$\mathcal{R}_1 : C_1^\infty \rightarrow C_\infty^1, \quad C_1^\infty \ni \mathbf{i} \rightarrow \mathcal{R}_1 \mathbf{i} = \mathbf{i}^*,$$

$$\mathcal{R}_1 \left(\sum_{(x,y) \in E_+} i(x,y)[x] \otimes [y] \right) = \sum_{(x,y) \in E_+} r(x,y) i(x,y) dx \otimes dy.$$

Note that since $r(x,y) = \frac{1}{c(x,y)}$ we have

$$\mathcal{E}[\mathcal{R}_1 \mathbf{i}] = \sum_{(x,y) \in E_+} c(x,y) r(x,y)^2 i(x,y)^2 = \sum_{(x,y) \in E_+} r(x,y) i(x,y)^2 = \mathcal{E}[\mathbf{i}],$$

so that $\mathcal{R}_1$ induces a (bijective) isometry of Hilbert space $C_1^\infty \rightarrow C_\infty^1$. In fact C_∞^1 can be identified with the topological dual of C_1^∞ with the induced inner product and norm. For this reason, we will refer to $\mathcal{R}_1 \mathbf{i}$ as the dual of $\mathbf{i}$ and, when no confusion is possible, we will write $\mathbf{i}^* := \mathcal{R}_1(\mathbf{i})$.

Ohm's law states that for any finite energy current $\mathbf{i}$ in an electric network there is a difference of potential/voltage $u(x,y)$ between any two neighbors $x \sim y$ related to $i(x,y)$ via the equality

$$u(x,y) = r(x,y) i(x,y). \quad (4.5.11)$$

In other words, the voltage u associated to the current $\mathbf{i}$ is the dual 1-cochain, $u = \mathcal{R}_1 \mathbf{i}$

Kirchhoff's second law states that a finite energy current $\mathbf{i} \in C_1^\infty$ generated by a source $-\mathbf{j} = \partial \mathbf{i}$ must have a special property:

the dual 1-cochain of voltages is exact.

In other words, there exists a function $u : \mathfrak{X} \rightarrow \mathbb{R}$ such that $du = -\mathcal{R}_1 \mathbf{i}$, i.e.,

$$c(x,y)(u(y) - u(x)) = \frac{1}{r(x,y)}(u(y) - u(x)) = i(x,y), \quad \forall (x,y) \in E.$$

Note that

$$\mathcal{E}[du] := \langle du, du \rangle_c = \langle \mathbf{i}, \mathbf{i} \rangle_r = \mathcal{E}[\mathbf{i}] < \infty. \quad (4.5.12)$$

Definition 4.5.3. A *Kirchhoff current* is a finite energy current $\mathbf{i}$ such that its dual $\mathbf{i}^* = \mathcal{R}_1 \mathbf{i}$ is exact. A function $u \in C^0$ such that $\mathbf{i}^* = -du$ is called a *potential* of the Kirchhoff current. $\square$

Suppose that $\mathbf{i}$ is a Kirchhoff current and u is a potential of $\mathbf{i}$. If the graph is connected, then any other potential of $\mathbf{i}$ differs from u by an additive constant.

The source $\mathbf{j} = -\partial \mathbf{i}$ of the Kirchhoff current $\mathbf{i}$ can be described explicitly in terms of a potential u of $\mathbf{i}$. The equality $\partial \mathbf{i} = -\mathbf{j}$ reads

$$\sum_{y \in N(x)} c(x,y)(u(y) - u(x)) = \sum_{y \in N(x)} i(x,y) = -j(x), \quad \forall x \in \mathfrak{X}.$$

Since $c(x, y) = c(x)Q_{x,y}$ we deduce

$$\sum_{y \in N(x)} Q_{x,y} (u(y) - u(x)) = -\frac{1}{c(x)} j(x) = -\mathcal{R}_0 j(x), \quad \forall x \in \mathfrak{X}.$$

Equivalently, this means that any potential u of the Kirchhoff current i with source j satisfies

$$\Delta u(x) = \frac{1}{c(x)} j(x) = \mathcal{R}_0 j(x) \frac{1}{c(x)} = \sum_{y \in N(x)} i(y, x), \quad \forall x \in \mathfrak{X}, \quad (4.5.13)$$

where $\Delta = \mathbb{1} - Q$ is the Laplacian of the HMC with transition matrix Q ; see Definition 4.3.3.

Let us observe that if α_1, α_2 are k -cochains and at least one of them is compactly supported, then we can define $\langle \alpha_1, \alpha_2 \rangle_c$ using the same expressions (4.5.10), (4.5.9) as above.

Proposition 4.5.4 (Discrete integration by parts). *For any $u \in C^0$ and any $v \in C_{\text{cpt}}^0$ we have*

$$\langle \Delta u, v \rangle_c = \langle du, dv \rangle_c = \langle u, \Delta v \rangle_c. \quad (4.5.14)$$

Proof. We have

$$\begin{aligned} \langle \Delta u, v \rangle_c &= \sum_{x \in \mathfrak{X}} c(x) \Delta u(x) v(x) \\ &= \sum_{x \in \mathfrak{X}} \left(\sum_{y \in Y} c(x) Q_{x,y} (u(x) - u(y)) \right) v(x) = \sum_{x,y \in \mathfrak{X}} c(x, y) (u(x) - u(y)) v(x) \\ &= \sum_{(x,y) \in E_+} c(x, y) (u(x) - u(y)) v(x) + \sum_{(y,x) \in E_+} c(x, y) (u(x) - u(y)) v(x) \end{aligned}$$

(change variables $x \leftrightarrow y$ in the second sum)

$$\begin{aligned} &= \sum_{(x,y) \in E_+} c(x, y) (u(x) - u(y)) v(x) + \sum_{(x,y) \in E_+} c(x, y) (u(y) - u(x)) v(y) \\ &= \sum_{(x,y) \in E_+} c(x, y) (u(x) - u(y)) (v(x) - v(y)) = \langle du, dv \rangle_c. \end{aligned}$$

The same argument, with the roles of u and v reversed show that

$$\langle du, dv \rangle_c = \langle u, \Delta v \rangle_c.$$

The above expressions are well defined since both dv and Δv are compactly supported because the graph is locally finite. $\square$

Here is a simple consequence. We say that a set $S \subset \mathfrak{X}$ is *cofinite* if $\mathfrak{X} \setminus S$ is finite.

Corollary 4.5.5. *Suppose that S is a nonempty cofinite set and $u \in \mathfrak{X}$ is a solution of the boundary value problem*

$$\Delta u(x) = 0, \quad \forall x \in \mathfrak{X} \setminus S, \quad u(s) = 0, \quad \forall s \in S.$$

Then $u(x) = 0, \forall x \in \mathfrak{X}$.

Proof. We have $0 = \langle \Delta u, u \rangle_c = \langle du, du \rangle_c$. Hence $du = 0$ and since $\mathfrak{X}$ is connected, we deduce that u is constant. Since $S \neq \emptyset$, we deduce that u is identically zero. $\square$

4.5.4. A probabilistic perspective of Kirchoff's laws. Denote by $(X_n)_{n \geq 0}$ the random walk on the weighted graph defined by the electric network $(\mathfrak{X}, E, c)$

Let $S \subset \mathfrak{X}$ be a nonempty subset. Recall that we denote by H_S , respectively T_S , the hitting/début and respectively return time to S . Fix a bounded function $\varphi : S \rightarrow \mathbb{R}$ and define

$$u = u_\varphi : \mathfrak{X} \rightarrow \mathbb{R}, \quad u(x) = \mathbb{E}_x[\varphi(X_{H_S})].$$

Conditioning on neighbors we deduce from the Markov property that u is harmonic on $\mathfrak{X} \setminus S$ and $u = \varphi$ on S . Corollary 4.5.5 shows that if S is cofinite, then u is the unique function on $\mathfrak{X}$ that is harmonic on $\mathfrak{X} \setminus S$ and equal to φ on S .

Let us investigate a special case of this construction. Consider a cofinite set S_- and $x_+ \in \mathfrak{X} \setminus S_-$. Set $S := \{x_+\} \cup S_-$. If $\varphi = \mathbf{I}_{\{x_+\}} : S \rightarrow \mathbb{R}$, then the computation in Example 4.3.4 shows that u_φ is

$$u(x) = u_{x_+, S_-}(x) := \mathbb{P}_x[H_S = H_{x_+}] = \mathbb{P}_x[H_{x_+} < H_{S_-}]. \quad (4.5.15)$$

Thus $u(x)$ is the probability that the random walk started at x reaches x_+ before S_- . Clearly this function has finite energy since it has compact support. To this function we associate a current $\mathbf{i} = \mathbf{i}_{x_+, S_-}$ defined by $\mathcal{R}_1 \mathbf{i} = -du$. More precisely

$$\mathbf{i} = \sum_{(x,y)} c(x,y) (u(y) - u(x)) [x] \otimes [y],$$

and its source is

$$\mathbf{j} = \mathbf{j}_{x_+, S_-} : \mathfrak{X} \rightarrow \mathbb{R}, \quad \mathbf{j}(x) \stackrel{(4.5.13)}{=} c(x) \Delta u(x).$$

The current $\mathbf{i}$ has compact support contained in the finite set of edges with one end in the finite set $\mathfrak{X} \setminus S_-$. Hence $\partial^2 \mathbf{i} = 0$ so

$$0 = \sum_{x \in \mathfrak{X}} \mathbf{j}(x) = \sum_{x \in \mathfrak{X}} c(x) \Delta u(x). \quad (4.5.16)$$

The energy of u is

$$\begin{aligned} \langle du, du \rangle_c &= \langle u, \Delta u \rangle_c = \sum_{x \in \mathfrak{X}} c(x) u(x) \Delta u(x) \\ &= c(x_+) u(x_+) \Delta u(x_+) = u(x_+) \mathbf{j}(x_+). \end{aligned} \quad (4.5.17)$$

Now observe that $u(x_+) = 1$ so that

$$\begin{aligned} \Delta u(x_+) &= 1 - \sum_{x \in N(x_+)} Q_{x_+, x} u(x) \\ &= 1 - \sum_{x \in N(x_+)} Q_{x_+, x} \mathbb{P}_x[H_{S_-} > H_{x_+}] = \mathbb{P}_{x_+}[T_{x_+} > H_{S_-}]. \end{aligned}$$

Hence

$$\begin{aligned} \mathcal{E}[du] &\stackrel{(4.5.17)}{=} c(x_+) \Delta u(x_+) = \mathbf{j}_{x_+, S_-}(x_+) \\ &= c(x_+) \mathbb{P}_{x_+}[T_{x_+} > H_{S_-}] =: \kappa(x_+, S_-). \end{aligned} \quad (4.5.18)$$

The quantity $\kappa(x_+, S_-)$ is called the *effective conductance* from x_+ to S_- . Its inverse is called *effective resistance* between x_+ and S_- and it is denoted by $\mathcal{R}_{\text{eff}}(x_+, S_-)$. Thus

$$\mathcal{R}_{\text{eff}}(x_+, S_-) = \frac{1}{c(x_+) \mathbb{P}_{x_+}[T_{x_+} > H_{S_-}]}.$$

We set

$$\bar{u}(x) = \bar{u}_{x_+, S_-}(x) := \frac{1}{\kappa(x_+, S_-)} u_{x_+, S_-} = \frac{1}{\kappa(x_+, S_-)} \mathbb{P}_x [H_{x_+} < H_{S_-}].$$

This is the potential of the dual of a compactly supported Kirchhoff current

$$\bar{\mathbf{i}}_{x_+, S_-} = \frac{1}{\kappa(x_+, S_-)} \mathbf{i}_{x_+, S_-}, \quad \boxed{\mathcal{R}_1 \bar{\mathbf{i}}_{x_+, S_-} = d\bar{u}_{x_+, S_-}}, \quad (4.5.19)$$

with source

$$-\partial \bar{\mathbf{i}}_{x_+, S_-} = \bar{\mathbf{j}}_{x_+, S_-} = \frac{1}{\kappa(x_+, S_-)} \mathbf{j}_{x_+, S_-} = \frac{c(x)}{c(x_+) \mathbb{P}_{x_+} [T_{x_+} > H_{S_-}]} \Delta u(x), \quad (4.5.20)$$

where u is defined in (4.5.15), $u(x) = \mathbb{P}_x [H_{x_+} < H_{S_-}]$. Note that

$$\bar{\mathbf{j}}_{x_+, S_-}(x) = \begin{cases} 1, & x = x_+, \\ 0, & x \in \mathfrak{X} \setminus (\{x_+\} \cup S_-). \end{cases}$$

Its energy is

$$\mathbf{E}_{x_+, S_-} := \frac{1}{\kappa(x_+, S_-)^2} \mathcal{E}_c [du_{x_+, S_-}] = \frac{1}{\kappa(x_+, S_-)} = \bar{u}_{x_+, S_-}(x_+). \quad (4.5.21)$$

Since

$$u_{x_+, S_-}(x) = \mathbb{P}_x [H_{S_-} < T_{x_+}] \leq 1 = u_{x_+, S_-}(x_+), \quad \forall x \in \mathfrak{X},$$

we deduce

$$0 \leq \bar{u}_{x_+, S_-}(x) \leq \bar{u}_{x_+, S_-}(x_+) = \mathbf{E}_{x_+, S_-}, \quad \forall x \in \mathfrak{X}. \quad (4.5.22)$$

Let us observe that if $\mathfrak{X}$ is finite and $S_- = \{x_-\}$, then the equality (4.5.16) shows that

$$0 = \sum_{x \in \mathfrak{X}} \bar{\mathbf{j}}_{x_+, x_-}(x) = \bar{\mathbf{j}}_{x_+}(x_+) + \bar{\mathbf{j}}_{x_-}(x_-)$$

and thus

$$\bar{\mathbf{j}}_{x_+, x_-}(x) = \begin{cases} \pm 1, & x = x_{\pm}, \\ 0, & x \neq x_{\pm}. \end{cases}$$

Definition 4.5.6. A flow from x_+ to S_- on the electric network is any finite energy current $\mathbf{i} \in C_1^\infty$ such that $\partial \mathbf{i} = -\bar{\mathbf{j}}_{x_+, S_-}$. The source $\bar{\mathbf{j}}_{x_+, S_-}$, defined in (4.5.20), is called the *unit dipole* with source x_+ and sink S_- .

A flow from x_+ to S_- is called a *Kirchhoff flow* if it satisfies the second Kirchhoff law, i.e., the dual $\mathcal{R}_1 \mathbf{i} \in C_\infty^1$ is the differential of a function $u : \mathfrak{X} \rightarrow \mathbb{R}$. $\square$

Lemma 4.5.7. Suppose that $\mathbf{i}$ is a compactly supported current such that $\partial \mathbf{i} = 0$. Then for any $u : \mathfrak{X} \rightarrow \mathbb{R}$ we have $\langle \mathbf{i}^*, du \rangle_c = 0$.

Proof. We have

$$\sum_{y \in N(x)} i(x, y) = 0, \quad \forall x \in \mathfrak{X}.$$

We recall that $i(x, y) = -i(y, x)$, $\forall (x, y) \in E$. We have

$$\langle \mathbf{i}^*, du \rangle_c = \sum_{(x, y) \in E_+} r(x, y) c(x, y) i(x, y) (u(y) - u(x))$$

$$\begin{aligned}
&= \sum_{(x,y) \in E_+} i(x,y)(u(y) - u(x)) = 2 \sum_{(x,y) \in E} i(x,y)(u(y) - u(x)) \\
&= -2 \sum_{x \in \mathfrak{X}} u(x) \left(\sum_{y \in N(x)} i(x,y) \right) + 2 \sum_{y \in \mathfrak{X}} u(y) \left(\sum_{x \in N(y)} i(x,y) \right) = 0.
\end{aligned}$$

All the above sums involve only finitely many terms since i is compactly supported. $\square$

Theorem 4.5.8. *Suppose S_- is cofinite and $x_+ \in \mathfrak{X} \setminus S_-$. Then the following hold.*

- (i) *The current $\mathbf{i}_0 := \bar{\mathbf{i}}_{x_+, S_-}$ defined by (4.5.19) is the unique compactly supported Kirchhoff flow from x_+ to S_-*
- (ii) *The potential $\bar{u} = \bar{u}_{x_+, S_-}$ that determines $\mathbf{i}_0$ is the unique solution of the boundary value problem*

$$\begin{aligned}
\Delta v(x) &= 0, \quad \forall x \in \mathfrak{X} \setminus (S_- \cup \{x_+\}) \\
v(x) &= \begin{cases} \frac{1}{c(x_+)}, & x = x_+, \\ 0, & x \in S_-. \end{cases} \quad (4.5.23)
\end{aligned}$$

- (iii) *The energy of $\mathbf{i}_0$ is*

$$\mathcal{E}[\mathbf{i}_0] = \mathbf{E}_{x_+, S_-} = \bar{u}_{x_+, S_-}(x_+) = \frac{1}{c(x_+) \mathbb{P}_{x_+}[T_{x_+} > H_{S_-}]} = \mathcal{R}_{\text{eff}}(x_+, S_-).$$

- (iv) *If $\mathbf{i}_1$ is another compactly supported flow from x_+ to S_- , then*

$$\mathcal{E}[\mathbf{i}_1] \geq \mathcal{E}[\bar{\mathbf{i}}_{x_+, S_-}],$$

with equality iff $\mathbf{i}_1 = \mathbf{i}_0$.

Proof. (i) Set $u_0 := \bar{u}_{x_+, S_-}$. Recall that u_0 has compact support. Suppose that $\mathbf{i}_1$ is another compactly supported Kirchhoff flow from x_+ to S_- . Then there exists a function $u_1 : \mathfrak{X} \rightarrow \mathbb{R}$ such that $\mathbf{i}_1^* = du_1$. We deduce from (4.5.13) that the functions u_k , $k = 0, 1$ are solutions of the same equation

$$\Delta u_k(x) = \frac{1}{c(x)} j_0(x), \quad \forall x \in \mathfrak{X}.$$

If we write $u = u_1 - u_0$, then $\Delta u = 0$ on $\mathfrak{X}$. The function u may not have compact support, but du does since $du = \mathcal{R}_1(\mathbf{i}_1 - \mathbf{i}_0)$. We have

$$\begin{aligned}
\langle du, du \rangle_c &= \frac{1}{2} \sum_{(x,y) \in E} c(x,y)(u(x) - u(y))(u(x) - u(y)) \\
&= \frac{1}{2} \sum_{(x,y) \in E} c(x,y)(u(x) - u(y))u(x) - \frac{1}{2} \sum_{(x,y) \in E} c(x,y)(u(x) - u(y))u(y) \\
&= \frac{1}{2} \sum_{x \in \mathfrak{X}} u(x) \underbrace{\sum_{y \in N(x)} c(x,y)(u(x) - u(y))}_{=c(x)\Delta u(x)=0} + \frac{1}{2} \sum_{y \in \mathfrak{X}} u(y) \underbrace{\sum_{x \in N(y)} c(y,x)(u(y) - u(x))}_{=c(y)\Delta u(y)=0} \\
&= 0.
\end{aligned}$$

Hence $du = 0$ so that $\mathbf{i}_0 = \mathbf{i}_1$.

(ii) If v_1, v_2 are two compactly supported solutions of (4.5.23), then the argument above shows that $\langle dv, dv \rangle_c = 0$ and, since v is compactly supported, we deduce that $v = 0$. The equality (4.5.23) follows from (4.5.13). The equality (iii) follows from (4.5.21)

(iv) Set $\mathbf{i} = \mathbf{i}_1 - \mathbf{i}_0$. Then

$$\mathcal{E}[\mathbf{i}_1] = \mathcal{E}[\mathbf{i} + \mathbf{i}_0] = \langle \mathbf{i}_0^*, \mathbf{i}_0^2 \rangle_c + 2\langle \mathbf{i}_0^*, \mathbf{i}^* \rangle_c + \underbrace{\langle \mathbf{i}^*, \mathbf{i}^* \rangle_c}_{\geq 0}$$

($\mathbf{i}_0^* = du$)

$$\geq \mathcal{E}[\mathbf{i}_0] + 2\langle du, \mathbf{i}^* \rangle_c.$$

Lemma 4.5.7 shows that $\langle du, \mathbf{i}^* \rangle_c = 0$ since $\mathbf{i}$ has compact support and $\partial \mathbf{i} = 0$. $\square$

Remark 4.5.9. (a) Part (iv) of the theorem is known as the *Thompson or Dirichlet Principle*. It classically states that the Kirchhoff flow is *the* least energy compactly supported flow sourced by the dipole $\bar{\mathbf{j}}_{x_+, S_-}$. Observe that the energy of the Kirchhoff flow carries information about the dynamics of the Markov chain associated to the electric network.

(b) If S_- is cofinite, Kirchhoff flow from x_+ to S_- is the unique compactly supported current $\mathbf{i}$ such that

- $\partial \mathbf{i}(x_+) = -1$.
- There exists a function $u : \mathfrak{X} \rightarrow \mathbb{R}$, vanishing on S_- , such that $\mathbf{i}^* = du$.

$\square$

4.5.5. Degenerations. To proceed further we perform a reduction to a finite network. We set

$$S_+ := \mathfrak{X} \setminus S_-, \quad \partial S_+ := \{s_- \in S_-; N(s_-) \cap S_+ \neq \emptyset\}.$$

For simplicity we assume that x_+ does not have any neighbor in S_- . We obtain a new finite electric network $\mathfrak{X}/S_-$ described as follows.

- Its vertex set is $S_+ \cup \{x_-\}$. Think that we have identified all the vertices in S_- with a single point x_- .
- The conductances $c_*(x, y)$ of $\mathfrak{X}/S_-$ are defined according to the rule

$$c_*(x, y) = \begin{cases} c(x, y), & x, y \in S_+, \\ \sum_{s_+ \in \partial S_+} c(x, s_+), & y = x_- \\ \sum_{s_+ \in \partial S_+} c(s_+, y), & x = x_-. \end{cases}$$

Note that $c_*(x) = c(x)$, $\forall x \in S_+$. We denote by Δ_* the Laplacian determined by these conductances. The function $\bar{u} = \bar{u}_{x_0, S_-}$ is identically zero on S_- and thus descends to a function u_* on $\mathfrak{X}/S_-$ such that $u_*(x_-) = 0$. The set S_+ is also a subset of $\mathfrak{X}/S_-$.

$$\Delta_* u = \Delta \bar{u} \text{ on } S_+.$$

Moreover

$$c_*(x_{\pm}) \Delta_* u_*(x_{\pm}) = \pm 1.$$

Thus u_* is the potential of the Kirchhoff flow on $\mathfrak{X}/S_-$ from x_+ to x_- . We denote by E_{x_+, x_-} its energy.

Note that the induced Kirchhoff flow on $\mathfrak{X}/S_-$ has the same energy as the original Kirchhoff flow on $\mathfrak{X}$, i.e.,

$$\mathbf{E}_{x_+,x_-} = u_*(x_+) = u(x_+) = \mathbf{E}_{x_+,S_-}. \quad (4.5.24)$$

On the finite graph $\mathfrak{X}/S_-$ the energy of an edge (x, y) is $\frac{1}{c_*(x,y)}$. We define the energy of a path as the sum of energies of the edges along this paths. (An edge contributes to the energy of a path everytime it is encountered along the path.)

A flow from x_+ to x_- in the finite graph $\mathfrak{X}/S_-$ is the union of a path from x_+ and x_- and possibly a few other *closed* paths. They all have finite energy. The Kirchhoff flow is *the path* with minimal energy from x_+ to x_- .

In view of this reduction to finite graphs we concentrate on finite electric networks. Suppose $(\mathfrak{X}, E, c)$ is such a network and $x_+, x_- \in \mathfrak{X}$, $x_+ \neq x_-$. For finite graphs the finite energy condition is automatically satisfied and a flow from x_+ to x_- is simply a 1-chain $\mathbf{i}$ such that

$$\partial \mathbf{i} = [x_-] - [x_+].$$

The source $[x_+] - [x_-]$ is called a *dipole* with source x_+ and sink x_- .

☞ *On finite graphs the flow condition involves only the topology of graph and is independent of the physics of the network encoded by the conductance function. However, the Kirchhoff flow depends on the physics/geometry of the network.*

Denote by $\mathbf{i} = \mathbf{i}_{x_+,x_-}$ the Kirchhoff flow with source x_+ and sink x_- . Its *potential grounded at x_-* is the function $u = u_{x_+,x_-} : \mathfrak{X} \rightarrow \mathbb{R}$ uniquely determined by the equations

$$\Delta u = 0 \in \mathfrak{X} \setminus \{x_+, x_-\}, \quad u(x_-) = 0, \quad c(x_+) \Delta u(x_+) = 1. \quad (4.5.25)$$

Then $\mathbf{i}_{x_+,x_-} = \mathcal{R}^{-1} du_{x_+,x_-}$. The energy of this flow is

$$\mathbf{E}_{x_+,x_-} = u_{x_+,x_-}(x_+) = \frac{1}{c(x_+) \mathbb{P}_{x_+} [T_{x_+} > T_{x_-}]}. \quad (4.5.26)$$

This quantity is an invariant of the quadruplet $(\mathfrak{X}, c, x_+, x_-)$.

Clearly if we vary the conductance function the energy changes, and a flow that is minimal for a choice of conductance may fail to be so for another choice. In particular, a flow that has minimal energy with respect to a conductance function may not have this property if we change the conductance or, equivalently, the resistance function $r(x, y) = \frac{1}{c(x,y)} \in (0, \infty]$. We will indicate the dependence of $\mathbf{E}_{x_+,x_-}$ on r using the notation $\mathbf{E}_{x_+,x'}(r)$.

Suppose we change the conductance function to a new function c' that is bigger or, equivalently, such that $r'(x, y) \leq r(x, y)$. Then for any current $\mathbf{i}$ we have

$$\mathcal{E}_r[\mathbf{i}] = \frac{1}{2} \sum_{(x,y) \in E} r(x, y) i(x, y)^2 \geq \frac{1}{2} \sum_{(x,y) \in E} r'(x, y) i(x, y)^2 = \mathcal{E}_{r'}[\mathbf{i}].$$

This implies the following result known as the *Raleigh Principle*.

Theorem 4.5.10 (Raleigh). *The energy of the Kirchhoff flow with given source and sink increases with the increase of the resistance function or, equivalently, if the conductance function decreases.*

Proof. Suppose that we decrease the resistance of an edge from $r(x, y)$ to $r'(x, y)$. Denote by $\mathbf{i}(r)$ the Kirchhoff flow with source x_+ , sink x_- and choice of resistance r . Define $\mathbf{i}(r')$ in a similar fashion.

We have

$$\mathbf{E}_{x_+,x_-}(r) = \mathcal{E}_r[\mathbf{i}(r)] \geq \mathcal{E}_{r'}[\mathbf{i}(r)] \geq \mathcal{E}_{r'}[\mathbf{i}(r')] = \mathbf{E}_{x_+,x_-}(r')$$

□

We can use this principle to produce estimates for $\mathbf{E}_{x_+,x_-}(r)$ in terms $\mathbf{E}_{x_+,x_-}(r')$ if r' is chosen wisely making $\mathbf{E}_{x_+,x_-}(r')$ easier to compute. One way to simplify the computation of $\mathbf{E}_{x_+,x_-}$ is to modify the topology of the graph. We can achieve this by pushing r to extreme values. Let me describe two such degenerations.

Suppose $y_0, y_1 \in \mathfrak{X} \setminus \{x_+, x_-\}$ are two nodes connected by an edge. Upon rescaling c we can assume that $c(y_0, y_1) = 1 = r(y_0, y_1)$. We have a family of deformed resistances

$$r_t : E \rightarrow (0, \infty), \quad t > 0, \quad r_t(x, x') = \begin{cases} t, & (x, x') = (y_0, y_1) \text{ or } (y_1, y_0), \\ r(x, x'), & \text{otherwise.} \end{cases}$$

We denote by $\mathbf{i}^t$ the Kirchhoff flow with source x_+ , sink x_- and resistances r_t , by $\mathbf{E}^t$ its energy $\mathbf{E}^t = \mathbf{E}_{x_+,x_-}(r_t)$ and by u^t its potential grounded at x_- defined by (4.5.25).

The Raleigh Principle shows that $\mathbf{E}^t$ is an increasing function of t . We want to describe what happens with $\mathbf{E}^t$ and u^t as $t \rightarrow 0, \infty$.

Cutting. The behavior as $t \rightarrow \infty$ is described by the electric network $(\mathfrak{X}^\infty, c^\infty, E^\infty)$ obtained by *cutting* the edge (y_0, y_1) . Intuitively the resistance of the edge (y_0, y_1) becomes infinite, so there should not be any electric current passing through it. We can also see this probabilistically. The probability of transitioning from y_0 to y_1 is proportional to the conductance $c(y_0, y_1)$ that decays to 0.

More precisely

$$\begin{aligned} \mathfrak{X}^\infty &= \mathfrak{X}, \quad E^\infty = E \setminus \{(y_0, y_1), (y_1, y_0)\}, \\ c^\infty(x, x') &= \lim_{t \rightarrow \infty} c^t(x, x') = \begin{cases} 0, & (x, x') = (y_0, y_1) \text{ or } (y_1, y_0), \\ c(x, x'), & \text{otherwise.} \end{cases} \end{aligned}$$

Shorting. The behavior as $t \rightarrow 0$ is described by the network $(\mathfrak{X}^0, E^0, c^0)$ obtained by *shorting* the edge (y_0, y_1) . Intuitively, the shorted network is obtained by collapsing the vertices y_0, y_1 to single point $*$; see Figure 4.6. From a physical point of view the resistance along (y_0, y_1) and the probability of transitioning from y_0 to y_1 approaches 1.

More precisely

- $\mathfrak{X}^0 = (\mathfrak{X} \setminus \{y_0, y_1\}) \cup \{*\}$.
- If $x, x' \in \mathfrak{X} \setminus \{y_0, y_1\}$, then $c^0(x, x') = c(x, x')$ so that $(x, x') \in E \iff (x, x') \in E^0$.
- $x \in \mathfrak{X} \setminus \{y_0, y_1\}$, then $c(x, *) = c(x, y_0) + c(x, y_1)$ so that $(x, *) \in E^0$ if and only if $(x, y_0) \in E$ or $(x, y_1) \in E$.

Note that we have a natural projection $p : \mathfrak{X} \rightarrow \mathfrak{X}^0$

$$p(x) = \begin{cases} x, & x \neq y_0, y_1, \\ *, & x = y_0, y_1. \end{cases}$$




Figure 4.6. Shorting an electric network along the edge (y_0, y_1) .

For $\epsilon \in \{0, \infty\}$ denote by u^ϵ (resp. u^∞) the potential grounded at x_- of the Kirchhoff flow in $(\mathfrak{X}^\epsilon, E^\epsilon, c^\epsilon)$ with source x_+ and sink x_- . Denote by E^ϵ the energy of u^ϵ

$$E^\epsilon = \frac{1}{2} \sum_{(x,y) \in E^\epsilon} c^\epsilon(x,y) (u^\epsilon(x) - u^\epsilon(y))^2.$$

Theorem 4.5.11 (Maxwell-Raleigh). *Suppose that $y_0, y_1 \in \mathfrak{X} \setminus \{x_+, x_-\}$ have the property that the removal of the edge connecting them does not disconnect the graph $(\mathfrak{X}, E)$. Then*

$$\lim_{t \rightarrow \infty} u^t(x) = u^\infty(x), \quad \forall x \in \mathfrak{X},$$

$$\lim_{t \rightarrow 0} u^t(x) = u^0(p(x)), \quad \forall x \in \mathfrak{X},$$

and

$$\lim_{t \rightarrow \epsilon} E^t = E^\epsilon, \quad \epsilon = 0, \infty.$$

In particular, $E^0 \leq E^t \leq E^\infty$, $\forall t > 0$. Thus the energy of the Kirchhoff flow from x_+ to x_- is increased by cutting and decreased by shorting.

Proof. We will carry the proof in several steps. We set $r := r^1$.

1. Compactness. Fix a path γ in $\mathfrak{X}$ from x_+ to x_- that avoids the edge (y_0, y_1) ,

$$\gamma = x_-, x_0, x_1, \dots, x_n = x_-.$$

The r^t -energy of this path is

$$\mathcal{E}_{r^t}[\gamma] = \sum_{k=1}^n r(x_{k-1}, x_k) = \mathcal{E}_r[\gamma].$$

It is independent of t since the path avoids the only edge whose resistance depends on t . We deduce from Thompson's principle that

$$E^t \leq \mathcal{E}[\gamma], \quad \forall t > 0.$$

The local estimate (4.5.22) implies that

$$0 \leq u^t(x) \leq \mathbf{E}^t \leq \mathcal{E}_r[\gamma], \quad \forall t > 0.$$

This shows that the family of functions $u^t : \mathfrak{X} \rightarrow [0, \infty)$ is relatively compact with respect to the usual topology of the finite dimensional vector space $\mathbb{R}^{\mathfrak{X}}$.

2. $t \rightarrow \infty$. In this case observe that

$$\lim_{t \rightarrow \infty} c^t(x, y) = c^\infty(x, y), \quad \forall x, y \in \mathfrak{X}.$$

We will show that as $t \rightarrow \infty$ the family u^t has only one limit point. Suppose that for a sequence $t_n \rightarrow \infty$ the functions u^{t_n} converge to a function v . The function u^{t_n} satisfies the equation

$$\sum_{y \in \mathfrak{X}} c^{t_n}(x, y) (u^{t_n}(x) - u^{t_n}(y)) = \begin{cases} 0, & x \neq x_\pm, \\ \pm 1, & x = x_\pm, \quad u^{t_n}(x_-) = 0. \end{cases}$$

Letting $n \rightarrow \infty$ we deduce that v satisfies

$$\sum_{y \in \mathfrak{X}} c^\infty(x, y) (v(x) - v(y)) = \begin{cases} 0, & x \neq x_\pm, \\ \pm 1, & x = x_\pm, \quad v(x_-) = 0. \end{cases}$$

According to Theorem 4.5.8(ii) the above equation has a unique solution, the potential u^∞ of the Kirchhoff flow from x_+ to x_- grounded at x_- in $(\mathfrak{X}^\infty, c^\infty)$ proving that

$$\lim_{t \rightarrow \infty} u^t = u^\infty.$$

The equality

$$\lim_{t \rightarrow \infty} \mathbf{E}^t = \mathbf{E}^\infty$$

is obvious.

3. $t \rightarrow 0$. The above argument fails in this case because $c^t(y_0, y_1) = \frac{1}{t}$. Pick a sequence $t_n \searrow 0$ such that u^{t_n} has a limit u^0 as $t_n \searrow 0$. To simplify the presentation we will write u^t instead of u^{t_n} . We will show that

$$u^0(y_0) = u^0(y_1) \tag{4.5.27}$$

and the induced function $\bar{u}^0$ on $\mathfrak{X}^0$,

$$\bar{u}^0(x) = \begin{cases} u^0(x), & x \neq *, \\ u^0(y_0) = u^0(y_1), & x = *, \end{cases}$$

satisfies

$$\bar{u}^0(x_-) = 0, \tag{4.5.28a}$$

$$\sum_{y \in N^0(x)} c^0(x, y) (\bar{u}^0(x) - \bar{u}^0(y)) = \begin{cases} 0, & x \in \mathfrak{X}^0 \setminus \{x_+, x_-\}, \\ 1, & x = x_+, \end{cases} \tag{4.5.28b}$$

We set

$$\begin{aligned} N_*(y_0) &:= N(y_0) \setminus \{y_1\}, \quad N_*(y_1) := N(y_1) \setminus \{y_0\}, \\ c_*(y_i) &:= \sum_{y \in N_*(y_i)} c(y_i, y), \quad i = 0, 1. \end{aligned}$$

Denote by $N^0(*)$ the set of neighbors of $*$ in the graph $(\mathfrak{X}^0, E^0)$. Note that

$$N^0(*) = N_*(y_0) \cup N_*(y_1), \quad c^0(*) = c_*(y_0) + c_*(y_1).$$

Since $\Delta_{c^t} u^t(y_0) = 0$, $i = 0, 1$ we deduce

$$0 = \frac{1}{t} (u^t(y_0) - u^t(y_1)) + \sum_{y \in N_*(y_0)} c(y_0, y) (u^t(y_0) - u^t(y))$$

so that

$$(1 + tc_*(y_0))u^t(y_0) - u^t(y_1) = t \sum_{y \in N_*(y_0)} c(y_0, y)u^t(y).$$

A similar computation shows that

$$-u^t(y_0) + (1 + tc_*(y_1))u^t(y_1) = t \sum_{y \in N_*(y_1)} c(y_1, y)u^t(y).$$

Thus $(u^t(y_0), u^t(y_1))$ is the solution of the 2×2 non-homogeneous linear system

$$\underbrace{\begin{bmatrix} a_0(t) & -1 \\ -1 & a_1(t) \end{bmatrix}}_{=:A(t)} \cdot \begin{bmatrix} u^t(y_0) \\ u^t(y_1) \end{bmatrix} = t \cdot \begin{bmatrix} b_0(t) \\ b_1(t) \end{bmatrix},$$

where

$$a_i(t) = 1 + tc_*(y_i), \quad b_i(t) = \sum_{y \in N_*(y_i)} c(y_i, y)u^t(y), \quad i = 0, 1.$$

Note that

$$\det A(t) = a_0(t)a_1(t) - 1 = t(c_*(y_0) + c_*(y_1)) + O(t^2) = tc^0(*) + O(t^2).$$

Set

$$A_0(t) = \begin{bmatrix} b_0(t) & -1 \\ b_1(t) & a_1(t) \end{bmatrix}, \quad A_1(t) = \begin{bmatrix} a_0(t) & b_0(t) \\ -1 & b_1(t) \end{bmatrix}.$$

Using Cramer's rule we deduce

$$u^t(y_0) = \frac{t \det A_0(t)}{\det A(t)} = \frac{a_1(t)b_0(t) + b_1(t)}{c^0(*) + O(t)},$$

$$u^t(y_1) = \frac{a_0(t)b_1(t) + b_0(t)}{c^0(*) + O(t)}.$$

Now observe that

$$\lim_{t \rightarrow 0} a_i(t) = 1$$

and, since $N^0(*) = N_*(y_0) \cup N_*(y_1)$

$$\lim_{t \rightarrow 0} (b_0(t) + b_1(t)) = \sum_{y \in N^0(*)} c^0(*, y)u^0(y).$$

Hence

$$u^0(y_0) = u^0(y_1) = \bar{u}^0(*) := \frac{\sum_{y \in N^0(*)} c^0(*, y)u^0(y)}{c^0(*)}.$$

This proves (4.5.27). The equality (4.5.28a) is obvious. Observe that

$$\bar{u}^0(*) \sum_{y \in N^0(*)} c^0(*, y) = \sum_{y \in N^0(*)} c^0(*, y)\bar{u}^0(y),$$

i.e. ,

$$\sum_{y \in N^0(*)} c^0(*, y)(\bar{u}^0(*) - \bar{u}^0(y)) = 0.$$

This proves (4.5.28b) for $x = *$.

If $x \in \mathfrak{X} \setminus \{*, x_-\}$, then

$$\sum_{y \in N(x)} c^t(x, y) (u^t(x) - u^t(y)) = \begin{cases} 1, & x = x_+, \\ 0, & x \neq x_+. \end{cases}$$

The equality (4.5.28b) for $x \neq *, x_-$ follows by letting $t \rightarrow 0$ above and observing that

$$\lim_{t \rightarrow 0} u^0(y_i) = \bar{u}^0(*), \quad i = 0, 1, \quad \lim_{t \rightarrow 0} (c^t(x, y_0) + c^t(x, y_1)) = c^0(x, *)$$

and

$$N^0(x, *) = (N(x) \setminus \{y_0, y_1\}) \cup \{*\}.$$

This proves the equality (4.5.28b). This determines $\bar{u}^0$ uniquely and shows that

$$\lim_{t \rightarrow 0} u^t = \bar{u}^0(p(x)).$$

It remains to verify only the claim

$$\lim_{t \rightarrow 0} \mathbf{E}^t = \mathbf{E}^0.$$

Note that

$$\mathbf{E}^t = \frac{1}{2} \sum_{(x, y) \in E} c^t(x, y) (u^t(x) - u^t(y))^2.$$

There are two problematic terms in the above sum corresponding to $(x, y) = (y_0, y_1)$ or (y_1, y_0) and their contribution to the energy is

$$\frac{1}{t} (u^t(y_0) - u^t(y_1))^2.$$

Now observe that

$$u^t(y_0) - u^t(y_1) = \frac{b_0(t)(a_1(t) - 1) - b_1(t)(a_0(t) - 1)}{c^0(*) + O(t)} = t \frac{c_*(y_1)b_0(t) - c_*(y_0)b_1(t)}{c^0(*) + O(t)}.$$

Hence

$$\frac{1}{t} (u^t(y_0) - u^t(y_1))^2 = O(t) \quad \text{as } t \rightarrow 0,$$

so

$$\begin{aligned} \lim_{t \rightarrow 0} \mathbf{E}^t &= \frac{1}{2} \lim_{t \rightarrow 0} \sum_{(x, y) \in E \setminus \{(y_0, y_1), (y_1, y_0)\}} c^t(x, y) (u^t(x) - u^t(y))^2 \\ &= \frac{1}{2} \sum_{(x, y) \in E^0} c^0(x, y) (u^0(x) - u^0(y))^2 = \mathbf{E}^0. \end{aligned}$$

□

Remark 4.5.12. (i) Let us explain what happens if cut an edge (y_0, y_1) that disconnects the graph, but x_+, x_- lie in the same connected component of the resulting graph.

Denote by $(\mathfrak{X}_0, E_0)$ the connected component containing x_+, x_- and by $(\mathfrak{X}_*, E_*)$ the union of the other components. The compactness part of the argument still works since the energy of u^t is bounded by the energy of a path in $(\mathfrak{X}_0, E_0)$ connecting x_+ to x_- .

Denote by u_0^t the restriction of u to $\mathfrak{X}_0$ and by u_*^t its restriction of u to $\mathfrak{X}^*$. Then

$$\begin{aligned} \mathcal{E}[du^t] &= \underbrace{\frac{1}{2} \sum_{(x,y) \in E_0} c(x,y) (u^t(x) - u^t(y))^2}_{=: \mathcal{E}_0^t} + \frac{1}{t} (u^t(y_0) - u^t(y_1))^2 \\ &\quad + \underbrace{\frac{1}{2} \sum_{(x,y) \in E_*} c(x,y) (u^t(x) - u^t(y))^2}_{=: \mathcal{E}_*^t}. \end{aligned}$$

Note that

$$\lim_{t \rightarrow \infty} \frac{1}{t} (u^t(y_0) - u^t(y_1))^2 = 0.$$

Arguing exactly as in Step 2 of the proof of Theorem 4.5.11 one can show that u_0^t converges to u_{x_+, x_-}^0 the potential grounded at 0 of the Kirchhoff flow in $\mathfrak{X}^0$ from x_+ to x_- . If u_* is any limit point of u_*^t , then u_* satisfies $\Delta_* u_* = 0$ so

$$\langle du_*, du_* \rangle_* = \langle \Delta_* u_*, u_* \rangle_* = 0$$

so 0 is the only limit point of $\mathcal{E}_*^t$ as $t \rightarrow \infty$.

$$\mathcal{E}_c[du]_c \leq \lim_{t \rightarrow 0} \mathcal{E}_{c^t}[du^t] = \mathcal{E}_{c^0}[du^0],$$

so the energy of the Kirchhoff flow from x_+, x_- in $\mathfrak{X}$ is not greater than the energy of the similar flow in $\mathfrak{X}_0$.

(ii) To understand why shorting is tricky recall that $\mathfrak{X}$ is finite so the Markov chain defined by the conductance c_t has an invariant probability measure is given by

$$\pi_t(x) = \pi(x) = \frac{c_t(x)}{Z_t}, \quad Z_t = \sum_{x \in \mathfrak{X}} c_t(x).$$

If we let $t = c_t(y_0, y_1) \rightarrow \infty$ and leave the other conductances unchanged, then, as $t \searrow 0$

$$\pi_t(x) \rightarrow 0, \quad \forall x \neq y_0, y_1, \quad \pi_t(x) \rightarrow \frac{1}{2}, \quad x = y_0, y_1.$$

In other words, $\pi_0 = \frac{1}{2}(\delta_{x_0} + \delta_{x_1})$.

(iii) In view of the conservation of energy equality (4.5.24), the cutting and shorting procedures can be used in infinite graphs to estimate the energy E_{x_+, S_-} by reducing, them to cutting/shorting procedure on the collapsed graph X/S_- . Cutting has to be performed with care so that while cutting edges we do not disconnect x_+ from S_- . $\square$

4.5.6. Applications. I want to illustrate the usefulness of the above results on some concrete examples.

When the graph $(\mathfrak{X}, E)$ is finite and all the edges have *the same conductances*, the Kirchhoff flow from x_+ to x_- can be described explicitly in terms certain counts of spanning trees, [93, Thm. 1.16]. In particular, its energy $\mathcal{K}(x_+, x_-)$ is a topological invariant of the quadruplet $(\mathfrak{X}, E, x_+, x_-)$ described explicitly in terms of spanning trees.

If we now assign conductances c to the edges, the energy $\mathbf{E}_{x_+, x_-}(c)$ of the Kirchhoff flow from x_-, x_+ satisfies

$$\frac{1}{\sup c(x, y)} \mathcal{K}(x_+, x_-) \leq \mathbf{E}_{x_+, x_-}(c) \leq \frac{1}{\inf c(x, y)} \mathcal{K}(x_+, x_-).$$

The computation of $\mathcal{K}(x_+, x_-)$ is impractical for complicated graphs, but the above rather rough estimate expresses in a simple fashion the fact that $\mathbf{E}_{x_+, x_-}(c)$ depends on both the topology and the geometry of the electrical network.

Example 4.5.13. Suppose that $(\mathfrak{X}, E, c)$ is a finite electric network such that the underlying graph is a tree. Then for any pair of points x_+, x_- there exists a unique 1-chain $\mathbf{i}$ such that

$$\partial \mathbf{i} = [x_-] - [x_+].$$

It is described by a minimal path

$$x_+ = x_0, x_1, \dots, x_n = x_-.$$

This is the Kirchhoff flow from x_+ to x_- and its energy is

$$\mathbf{E}_{x_+, x_-} = \sum_{i=1}^n r(x_{i-1}, x_i) = \sum_{i=1}^n \frac{1}{c(x_{i-1}, x_i)}.$$

As a special case of this consider the Ehrenfest urn model. Recall that the state space is the set $\mathfrak{X} := \{0, 1, \dots, B\}$, $B \in \mathbb{N}$ and transition matrix Q given by

$$Q_{k, k-1} = \frac{k}{B}, \quad \forall k \geq 1, \quad Q_{j, j+1} = \frac{B-j}{B}, \quad \forall j < B.$$

As explained in Example 4.2.34, this can be described as an electric network whose underlining graph is a path

$$0 \rightarrow 1 \rightarrow \dots \rightarrow B,$$

and conductances

$$c(j, j+1) = \binom{B}{j} \frac{B-j}{B} = \binom{B-1}{j}.$$

In particular,

$$c(j) = \binom{B-1}{j} + \binom{B-1}{j-1} = \binom{B}{j}.$$

If B is even, $B = 2N$, then

$$\mathbf{E}_{0, N} = \mathbf{E}_{N, 0} = \sum_{j=0}^{N-1} \frac{1}{\binom{2N-1}{j}}.$$

Thus

$$\mathbb{P}_N[T_N > T_0] = \frac{1}{c(N)\mathbf{E}_{N, 0}}, \quad \mathbb{P}_0[T_0 > T_N] = \frac{1}{c(0)\mathbf{E}_{N, 0}}$$

Hence

$$\frac{\mathbb{P}_0[T_0 > T_N]}{\mathbb{P}_N[T_N > T_0]} = \frac{c(N)}{c(0)} = \binom{2N}{N} \sim \frac{4^N}{\sqrt{\pi N}}.$$

In particular, this shows that $\mathbb{P}_N[T_N > T_0]$ is extremely small for large N . Thus, if initially in the two chambers there were equal numbers of balls, the probability that during the random transfers of balls between them, one of the chambers will empty before it will have again half the number of balls

is extremely small. In fact, the expected time of emptying the left chamber while starting with equal numbers of balls in both is (see [113, Sec. VII.3, p.175] with $s = 2N$)

$$\mathbb{E}_N[T_0] \sim 4^N (1 + A/N) \text{ as } N \rightarrow \infty, \quad 1 \leq A \leq 2. \quad (4.5.29)$$

This example is historically important because it was used to explain an apparent contradiction between Boltzmann's kinetic theory of gases and classical thermodynamics. We refer to [19, 108] for more details. $\square$

Remark 4.5.14. There is a discrepancy between the estimate (4.5.29) proved in [113] and the estimate for $\mathbb{E}_N[T_0]$ proved in [15, Sec. III.5] which states that

$$\mathbb{E}_N[T_0] = \frac{4^N}{N} (1 + O(1/N)) \text{ as } N \rightarrow \infty. \quad (4.5.30)$$

The estimate (4.5.30) also contradicts the estimates [114, Eq. (4.27)] and [165, Eq.(7)]. $\square$

Example 4.5.15 (Random walks on infinite graphs). Let us investigate the standard random walk on an infinite, locally finite graph $(\mathfrak{X}, E, c)$. Thus we think of an electric network in which all edges have the same conductance 1. For $x, y \in \mathfrak{X}$ define $\text{dist}(x, y)$ to be the minimal length of a path joining x and y . Fix $x_+ \in \mathfrak{X}$ and set

$$B_n := \{x \in \mathfrak{X}; \text{dist}(x_+, x) \leq n\},$$

$$\Sigma_n = \{x \in \mathfrak{X}; \text{dist}(x_+, x) = n\} = B_n \setminus B_{n-1}, \quad S_n^- = \mathfrak{X} \setminus B_n.$$

Note that the balls B_n are finite. For $n \in \mathbb{N}$ we denote by $C(n)$ the total number of edges connecting a point in Σ_{n-1} to a point in Σ_n .




Figure 4.7. Shorting an infinite electric network inside spheres.

Form the collapsed electric network $(\mathfrak{X}^n, E^n, c^n)$, $\mathfrak{X}^n := X/S_n^-$. The set S_n^- corresponds to a unique vertex x_n^- in $\mathfrak{X}^n$; see the top of Figure 4.7. Denote by E_{x_+, x_n^-} the energy of the Kirchhoff flow in $\mathfrak{X}^n$ from x_+ to x_n^- .

As we have seen

$$\frac{1}{c(x_+) \mathbb{P}_{x_+} [T_{x_+} > H_{S_n^-}]} = E_{x_+, x_n^-}.$$

Observe that the collapsed network $\mathfrak{X}/S_n^-$ is obtained from the collapsed network $\mathfrak{X}/S_{n+1}^-$ by first shorting the edges in $\Sigma_n \subset \mathfrak{X}/S_{n+1}^-$ and then shorting the edge (x_n^-, x_{n+1}^-) . Hence

$$E_{x_+, x_n^-} \leq E_{x_+, x_{n+1}^-}.$$

We set

$$E_{x_+, \infty} := \lim_{n \rightarrow \infty} E_{x_+, x_n} = \lim_{n \rightarrow \infty} \frac{1}{c(x_+) \mathbb{P}_{x_+} [T_{x_+} > H_{S_n^-}]}$$

Thus

$$\lim_{n \rightarrow \infty} \mathbb{P}_{x_+} [T_{x_+} > H_{S_n^-}] = \frac{1}{c(x_+) E_{x_+, \infty}},$$

and

$$\lim_{n \rightarrow \infty} \mathbb{P}_{x_+} [T_{x_+} < \infty] = \lim_{n \rightarrow \infty} \mathbb{P}_{x_+} [T_{x_+} < H_{S_n^-}] = 1 - \frac{1}{c(x_+) E_{x_+, \infty}}.$$

We deduce that the associated Markov chain is recurrent if and only if $E_{x_+, \infty} = \infty$ and transient otherwise.

To estimate E_{x_+, x_n^-} from below we short edges in $\mathfrak{X}/S_n^-$. First we short the edges between points in Σ_k , $k = 1, \dots, n-1$. We obtain the electric network $\mathfrak{X}_*^n$ at the bottom of Figure 4.7. As explained in Example 4.5.13, energy of the Kirchhoff flow in $\mathfrak{X}_*^n$ from x_+ to x_n^- is

$$E^n = \sum_{k=1}^n \frac{1}{C(k)} \leq E_{x_+, x_n^-}.$$

Hence

$$E_{x_+, \infty} \geq \sum_{k=1}^{\infty} \frac{1}{C(k)}.$$

We deduce that if

$$\sum_{k=1}^{\infty} \frac{1}{C(k)} = \infty,$$

then the corresponding Markov chain is recurrent.

To estimate $E_{x_+, \infty}$ from above we use the cutting trick. We gradually remove edges such that the component containing x_+ has infinitely many vertices. Restricting to the component containing x_+ we obtain a electric network with bigger $E_{x_+, \infty}$ according to Theorem 4.5.11 and Remark 4.5.12(iii).

Thus if the graph $(\mathfrak{X}, E)$ contains a connected subgraph $(\mathfrak{X}_0, E_0)$ such that the random walk on $\mathfrak{X}_0$ is transient, then the random walk on $(\mathfrak{X}, E)$ is also transient. $\square$

Example 4.5.16 (Random walk on $\mathbb{Z}^2$). Suppose that $(\mathfrak{X}, E, c)$ corresponds to the standard random walk on $\mathbb{Z}^2$. Observe that the sphere Σ_{n-1} , $n-1 > 0$, is the square

$$\Sigma_{n-1} = \{ (x, y) \in \mathbb{Z}^2; |x| + |y| = n-1 \}.$$

Each of the four vertices of this square is connected to Σ_n through 3 edges. The interior of each of the four edges contains $(n - 2)$ lattice points and each of them is connected to Σ_n through 2-edges. Thus

$$C(n) = 12 + 8(n - 2) = 8n - 4, \quad \forall n \in \mathbb{N}.$$

Since

$$\sum_{n \geq 1} \frac{1}{8n - 4} = \infty$$

We deduce again that the random walk on $\mathbb{Z}^2$ is recurrent. □

Example 4.5.17 (Random walks on symmetric trees). Consider the unbiased random walk on an infinite locally finite tree $(\mathfrak{X}, E)$. Fix $x_+ \in \mathfrak{X}$ and think of x_+ as the root of the tree. Looking at the paths started at x_+ we see that every vertex has a unique predecessor and a number $s(x)$ of successors so the degree is

$$d(x) = \begin{cases} s(x) + 1, & x \neq x_+, \\ s(x_+), & x = x_+. \end{cases}$$

Define B_n, Σ_n, S_n^- as in the previous example. We assume that the tree is *radially symmetric* about the root i.e., for any $n \in \mathbb{N}$ the vertices on the sphere Σ_n have the same number s_n of successors. Set

$$\sigma_k := |\Sigma_k|.$$

Note that for any $k \geq 0$ we have

$$\sigma_{k+1} = s_0 s_1 \cdots s_k.$$

One can think of σ_k as the “volume” of the sphere Σ_k . For example if the tree is d regular, then $s_0 = d$ and $s_k = d - 1, \forall k \geq 1$, so $\sigma_k = d(d - 1)^{k-1}$.

We want to investigate the unbiased random walk on this tree. Equivalently, this means assigning conductance 1 to every edge. We want to solve the equation

$$\Delta u(x) = \begin{cases} 0, & x \in B_n \setminus \{x_+\}, \\ \frac{1}{d(x_+)}, & x = x_+, \end{cases}$$

subject to the boundary condition

$$u(x) = 0, \quad \forall x \in S_n^- := \mathfrak{X} \setminus B_n.$$

We know that this equation has a unique solution. We can invoke the symmetry of the graph and show that this solution must be constant along the spheres Σ_n but we do not really need to do this. If we can find a solution with this property then it has to be *it*. So make use of this Ansatz and seek a solution that is constant on the spheres.

Denote by u_k the value of u on Σ_k . We set $u_0 := u(x_+)$

$$\Delta_k = u_k - u_{k+1}, \quad \forall k \geq 0.$$

Note that $\Delta_n = u_n$. For $k \in \{1, n\}$ we have

$$u_k = \frac{s_k u_{k+1} + u_{k-1}}{s_k + 1}$$

so that

$$(s_k + 1)u_k = s_k u_{k+1} + u_{k-1} \iff \Delta_{k-1} = s_k \Delta_k.$$

Iterating we deduce

$$\Delta_{k-1} = s_{k+1} \cdots s_n \Delta_n = \frac{s_0 s_1 \cdots s_n}{s_0 \cdots s_k} \Delta_n = \frac{\sigma_n}{\sigma_k} \Delta_n = \frac{\sigma_n}{\sigma_k} u_n.$$

Hence

$$u_0 = u_0 - u_{n+1} = \sum_{k=0}^n \Delta_k = \sigma_n \left(\sum_{k=0}^n \frac{1}{\sigma_k} \right) u_n.$$

The equation

$$\Delta u(x_+) = \frac{1}{s_0}$$

is equivalent to $\Delta_0 = \frac{1}{s_0}$ so that

$$\frac{1}{s_0} = \frac{\sigma_n}{s_0} u_n, \quad u_n = \frac{1}{\sigma_n}, \quad E_{x_+, S_n^-} = u_0 = \sum_{k=0}^n \frac{1}{\sigma_k}.$$

Hence,

$$E_{x_+, \infty} = \sum_{k=0}^{\infty} \frac{1}{\sigma_k}.$$

This shows that if the number of vertices on Σ_n growth fast the random walk is transient and if it growth slow, the walk is recurrent. Intuitively, the more vertices far away, more opportunities to get lost. As an example fix $d \in \mathbb{N}$, $d \geq 2$. We denote by $\mathcal{T}_d$ the rooted radially symmetric tree with successor sequence (s_n) given by

$$s_n = \begin{cases} d, & n = 2^k - 1, \quad k \geq 0, \\ 1, & \text{otherwise.} \end{cases}$$

Thus

$$\sigma_n = d^{k+1} \quad 2^k \leq n < 2^{k+1}$$

and

$$\sum_{n=0}^{\infty} \frac{1}{\sigma_n} = \frac{1}{d} + \sum_{n=2}^3 \frac{1}{d^2} + \sum_{n=4}^7 \frac{1}{d^3} + \cdots = \frac{1}{d} \sum_{k=0}^{\infty} \left(\frac{2}{d} \right)^k = \begin{cases} \frac{1}{d-2}, & d \geq 3, \\ \infty, & d = 2. \end{cases}$$

Thus, the random walk on $\mathcal{T}_d$ is transient if $d \geq 3$ and recurrent if $d = 2$.

We can obtain a more striking example of recurrent random walk by choosing the successor sequence to be

$$s_n = \begin{cases} k, & n = k!, \quad k \geq 2 \\ 1, & \text{otherwise} \end{cases}$$

For more information about random walks on trees we refer to the very comprehensive monograph [143]. $\square$

4.6. Finite Markov chains

For HMC-s with finite state space the theory simplifies somewhat and new techniques are available.

4.6.1. The Perron-Frobenius theory. Consider a homogenous Markov chain with finite state space

$$\mathfrak{X} = \mathbb{I}_m := \{1, 2, \dots, m\}.$$

In this case the transition matrix Q is an $m \times m$ *stochastic matrix*, i.e., a matrix with nonnegative entries such that the sum of the entries on each row is 1. If we set

$$\mathbf{e} := \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} \in \mathbb{R}^m$$

then we see that an $m \times m$ matrix Q with nonnegative entries is stochastic iff

$$Q\mathbf{e} = \mathbf{e}$$

We view measures on $\mathfrak{X}$ as *row* vectors $\mu = [\mu_1, \dots, \mu_m]$.

For convenience I will denote by $\mathcal{R}_m$ the space of *row* vectors and by $\mathcal{C}_m$ the space of *column* vectors. I will denote the row vectors using greek letters and we will think of them as *signed* measures on $\mathfrak{X}$. The matrix Q acts on row vectors by right multiplication $\mu \rightarrow \mu \cdot Q$, and on column vectors by left multiplication, $v \mapsto Q \cdot v$.

A signed measure $\mu \in \mathcal{R}_m$ is a probability measure if

$$\mu_k \geq 0, \quad \forall k \in \mathbb{I}_m, \quad \mu \cdot \mathbf{e} = 1.$$

Let $\text{Prob}_m \subset \mathcal{R}_m$ denote the space of probability measures on $\mathbb{I}_m$. We equip $\mathcal{R}_m$ with the variation norm

$$\|\alpha\|_v := \sum_{k=1}^m |\alpha_k|.$$

Observe that if $\mu, \nu \in \text{Prob}_m$, then

$$d_v(\mu, \nu) = \frac{1}{2} \|\mu - \nu\|_v.$$

Note that a *column* vector

$$\mathbf{z} = \begin{bmatrix} z_1 \\ \vdots \\ z_m \end{bmatrix} \in \mathbb{C}^m$$

is a (right) eigenvector of Q^T corresponding to an eigenvalue $\lambda \in \mathbb{C}$ if and only if the row vector $\mathbf{z}^\top$ is a (left) eigenvector of Q since

$$\mathbf{z}^\top \cdot Q = \lambda \mathbf{z}^\top.$$

The matrix Q and its transpose $Q^\top$ have the same eigenvalues,⁵ the vector $\mathbf{e}$ is a (right) eigenvector of Q corresponding to the eigenvalue 1 we deduce that there exists a row vector $\alpha \in \mathcal{R}_m$ such that

$$\alpha \cdot Q = \alpha$$

If α had nonnegative entries, then it would be an invariant measure for the HMC defined by Q . The classical Perron-Frobenius theory explains when this is the case and much more.

Observe that the HMC defined by Q is irreducible if and only if

$$\forall i, j \in \mathfrak{X}, \quad \exists n > 0 \text{ such that } Q_{i,j}^n > 0.$$

⁵ $\det(\lambda \mathbf{1} - Q) = \det(\lambda \mathbf{1} - Q)^\top$.

Additionally, it is aperiodic if and only if Q is *primitive*, i.e., there exists $n_0 \in \mathbb{N}$ such that

$$\forall n > n_0, \forall i, j \in \mathfrak{X} \text{ such that } Q_{i,j}^n > 0, \forall 1 \leq i, j \leq m.$$

For a proof of the following result we refer to [82, Chap. XIII] or [188, Chap. 8].

Theorem 4.6.1 (Perron-Frobenius). *Suppose that Q is a stochastic $m \times m$ matrix. Then the following hold.*

- (i) *All the eigenvalues of $Q^\top$ are contained in the unit disk.*
- (ii) *If Q is irreducible, then there exists $p \in \mathbb{N}$ such that*

$$\lambda \in \text{Spec}(Q) \text{ and } |\lambda| = 1 \iff \lambda^p = 1.$$

Moreover, every eigenvalue on the unit circle has algebraic multiplicity 1, i.e., it is a simple root of the characteristic polynomial of Q

- (iii) *The eigenspace $\ker(\mathbb{1} - Q^\top)$ is spanned by a positive vector.*
- (iv) *The matrix Q is primitive if and only if $p = 1$. In this case*

$$\rho := \max \{ |\lambda|; \lambda \in \text{Spec}(Q), \lambda \neq 1 \} < 1.$$

□

Suppose that Q is primitive and denote by π the unique invariant probability distribution of Q , i.e., the unique row vector $\pi = (\pi_1, \dots, \pi_m)$ such that

$$\pi_k > 0, \forall k, \pi_1 + \dots + \pi_m = 1.$$

Denote by $\Delta(\lambda)$ the characteristic polynomial of Q , $\Delta(\lambda) = \det(\lambda \mathbb{1} - Q)$. Set $B(\lambda) = \frac{1}{\lambda-1} \Delta(\lambda)$.

Since 1 is a simple eigenvalue of Q the polynomials $\lambda - 1$ and $B(\lambda)$ have no common divisor and thus we have a decomposition of the row space $\mathcal{R}_m$ (see [127, Thm. XI. 4.1]) as a direct sum of (right) Q -invariant subspaces

$$\mathcal{R}_m = \ker_r(\mathbb{1} - Q) \oplus \ker_r B(Q),$$

where

$$\begin{aligned} \ker_r(\mathbb{1} - Q) &:= \{ \alpha \in \mathcal{R}_m; \alpha \cdot (\mathbb{1} - Q) = 0 \} = \text{span}(\pi), \\ \ker_r B(Q) &:= \{ \alpha \in \mathcal{R}_m; \alpha \cdot B(Q) = 0 \}. \end{aligned}$$

Thus any $\alpha \in \mathcal{R}_m$ admits a unique decomposition

$$\alpha = \alpha^0 + \alpha^\perp, \alpha^0 \in \ker_r(\mathbb{1} - Q), \alpha^\perp \in \ker_r B(Q).$$

More explicitly, choose polynomials $u(\lambda), v(\lambda)$ such that

$$u(\lambda)(\lambda - 1) + v(\lambda)B(\lambda) = 1.$$

Then

$$\alpha^\perp = \alpha \cdot u(Q)(Q - 1) \in \ker B(Q) \quad \alpha^0 = \alpha \cdot v(Q)B(Q).$$

Note that

$$\alpha^\perp \cdot e = \alpha \cdot \alpha(Q)(Q - 1) \cdot e = 0.$$

If $\mu \in \mathcal{R}_m$ is a probability measure, then it has a canonical decomposition

$$\mu = c\pi + \mu^\perp, \mu^\perp \in \ker_r B(Q).$$

Since $\mu \cdot e = 1$ and $\mu^\perp \cdot e = 0$ we deduce $c = 1$ so $\mu = \pi + \mu^\perp$ and thus

$$\mu \cdot Q^n = \pi + \mu^\perp \cdot Q^n,$$

i.e.,

$$\mu \cdot Q^n - \pi = \mu^\perp \cdot Q^n.$$

Since $\ker_r B(Q)$ is Q -invariant we deduce from Theorem 4.6.1 that for any $\gamma \in (\rho, 1)$ there exists $C > 0$ such that

$$\forall \alpha \in \ker_r B(Q), \quad \forall n \in \mathbb{N}, \quad \|\alpha \cdot Q^n\|_v \leq C\gamma^n \|\alpha\|_v.$$

Hence

$$\forall n \in \mathbb{N}, \quad \forall \mu \in \text{Prob}_m : \quad \|\mu \cdot Q^n - \pi\|_1 = \|\mu^\perp \cdot Q^n\|_v \leq C\gamma^n.$$

In particular, if we choose μ to be the Dirac measure concentrated at $k \in \mathbb{I}_m$, then $\delta^k \cdot Q^n$ is the k -th row of the matrix Q^n and we deduce

$$\sum_{\ell=1}^m |Q_{k,\ell}^n - \pi_\ell| \leq C\gamma^n, \quad \forall k \in \mathbb{N}.$$

Theorem 4.6.1 allows us sharpen the above estimate. If

$$\Delta(\lambda) = \det(\lambda - Q) = \lambda^m + \sum_{j=0}^{m-1} a_j \lambda^j$$

denotes the characteristic polynomial of Q , then Cayley-Hamilton theorem implies that the sequence of matrices $(Q^n)_{n \in \mathbb{N}_0}$ satisfies the linear recurrence relation

$$Q^{n+m} + \sum_{j=0}^{m-1} a_j Q^{n+j} = 0, \quad \forall n \in \mathbb{N}_0.$$

Let $1, \lambda_2, \dots, \lambda_s$ be the eigenvalues of Q ,

$$1 > \rho = |\lambda_2| \geq \dots \geq |\lambda_s|,$$

The eigenvalue λ_2 is usually referred to as the *second largest eigenvalue (or SLE)* of the transition matrix.

Denote by m_2 is the largest Jordan cell size of the eigenvalues of norm ρ . The above recurrence relation shows that, for any $1 \leq i, j \leq m$, the sequence $(Q_{i,j}^n)_{n \geq 0}$ admits a description of the form

$$Q_{i,j}^n = c_{ij} + \sum_{k=2}^r C_{i,j}^k(n) \lambda_k^n$$

where $C_{i,j}^k(z)$ is a complex polynomial of degree $\leq m_k - 1$. We deduce that

$$|Q_{ij}^n - c_{ij}| = O(n^{m_2-1} \rho^n).$$

We conclude that $c_{i,j} = \pi_j$ and thus

$$|Q_{ij}^n - \pi_j| = O(n^{m_2-1} \rho^n). \quad (4.6.1)$$

If the Markov chain is reversible, i.e.,

$$\pi_i Q_{ij} = \pi_j Q_{ji}, \quad \forall i, j \in \mathbb{I}_m,$$

then the operator $Q : \mathcal{C}_m \rightarrow \mathcal{C}_m$ is symmetric with respect to the $L^2(\pi)$ -inner product $\langle -, - \rangle_\pi$ on $\mathcal{C}_m = \mathbb{R}^m$

$$\langle x, y \rangle_\pi = \sum_{i=1}^n x_i y_i \pi_i, \quad \forall x, y \in \mathcal{C}_m.$$

Indeed

$$\begin{aligned}\langle Qx, y \rangle_\pi &= \sum_i \sum_j Q_{ij} x_j y_i \pi_i = \sum_j \sum_i \pi_j Q_{ji} x_j y_i \\ &= \sum_j \left(\sum_i Q_{ji} y_i \right) \pi_j x_j = \langle x, Qy \rangle_\pi.\end{aligned}$$

In this case all the eigenvalues are real and the operator Q is diagonalizable and (4.6.1) improves to

$$|Q_{ij}^n - \pi_j| = O(\rho^n). \quad (4.6.2)$$

In general finding or estimating the SLE can be a daunting task. If some symmetry is present this is sometimes manageable.

Example 4.6.2 (Random walks on groups). Suppose that G is a finite group and $H \subset G$ is a set of generators. The set H determines a random walk on G . From g one can transition to $h \cdot g$, $h \in H$, with probability $\frac{1}{|H|}$.

A frequently encountered case is when H is symmetric, i.e.,

$$x \in H \iff x^{-1} \in H.$$

The directed graph corresponding to this random walk is symmetric, i.e., there is a directed edge from g to g' if and only if there is a directed edge from g' to g . The resulting undirected graph is called the *Cayley graph* determined by the symmetric set of generators. The random walk on the groups is then the standard walk on the Cayley graph. The group structure behind the Cayley graph adds a lot of symmetry that we can use to our advantage. For a detailed presentation of this technique and many interesting applications we refer to the beautiful monograph [52].

We want illustrate this principle on a simpler situation. Suppose that G is the discrete torus

$$G := (\mathbb{Z}/n\mathbb{Z})^d$$

We will denote by $x = (x_1, \dots, x_d)$ the elements of G , $x_k \in \mathbb{Z}/n\mathbb{Z}$. As generators we choose

$$\pm e_k \bmod n\mathbb{Z}, \quad k \in \llbracket 1, d \rrbracket,$$

where

$$e_1 = (1, 0, \dots, 0), \dots, e_d = (0, \dots, 0, 1).$$

For $d = 2$ this random walk can be visualised as a random walk on the vertices of the square grid $S_n = [0, n]^2 \cap \mathbb{Z}^2$ where the opposite edges are identified. Thus from $(0, y)$ we can transition $(0, y \pm 1 \bmod n)$ or $(\pm 1 \bmod n, y)$ with equal probabilities. Note that when n is odd, the random walk is irreducible and aperiodic.

When $n = 2$ this becomes a random walk on the set of vertices of the hypercube $[0, 1]^d$ or, equivalently, on the set of subsets of $\llbracket 1, d \rrbracket$.

The invariant probability measure π is, up to a multiplicative constant, the uniform counting measure. We write

$$L^2(G) = L^2(G, \pi), \quad \|f\| = \|f\|_{L^2} = \frac{1}{|G|^{1/2}} \left(\sum_{x \in \mathbb{T}_n^d} |f(x)|^2 \right)^{1/2}.$$

Here we work with complex valued functions so the inner product is

$$\langle f, g \rangle = \frac{1}{|G|} \sum_{x \in G} f(x) \overline{g(x)}.$$

If Q denotes the transition matrix of this Markov chain, then for any $f \in L^2(G)$ we have

$$Qf(x) = \sum_{x' \in G} Q_{x,x'} f(x') = \frac{1}{d} \sum_{k=1}^d \frac{f(x + e_k) + f(x - e_k)}{2}, \quad (4.6.3a)$$

$$\Delta f(x) = f(x) - Qf(x) = -\frac{1}{d} \sum_{k=1}^d \frac{f(x + e_k) - 2f(x) + f(x - e_k)}{2}. \quad (4.6.3b)$$

One can verify that the induced operator $Q : L^2(G) \rightarrow L^2(G)$ is symmetric since Q is reversible but we will not rely on this fact in this example.

To compute the eigenvalues of $Q : L^2(G) \rightarrow L^2(G)$ we use Fourier analysis. This requires a little bit of representation theory and we will refer to [208] for the proofs of all the claims below.

A *character* of G is a group morphism

$$\chi : G \rightarrow S^1 := \{ z \in \mathbb{C}; |z| = 1 \}.$$

The set $\widehat{G}$ of characters is a group itself with respect to the pointwise multiplication of characters. It is called the *dual group*.

Denote by $\mathbf{R}_n$ the group of n -th roots of unity

$$\mathbf{R}_n := \{ z \in \mathbb{C}^*; z^n = 1 \}.$$

Observe that for any character χ , the complex numbers $\chi(e_k)$ are n -th roots of 1. In fact, the map

$$\rho : \widehat{G} \rightarrow \mathbf{R}_n^d, \quad \widehat{G} \ni \chi \mapsto (\rho_1, \dots, \rho_d) = (\chi(e_1), \dots, \chi(e_d)) \in \mathbf{R}_n^d$$

is a group isomorphism. The collection of functions

$$\chi : G \rightarrow \mathbb{C}, \quad \chi \in \widehat{G}$$

is an *orthonormal basis* of $L^2(G)$ and thus, for any $f \in L^2$ we have an orthogonal decomposition

$$f = \sum_{\chi \in \widehat{G}} \langle f, \chi \rangle \chi. \quad (4.6.4)$$

The function

$$\widehat{G} \ni \chi \mapsto \widehat{f}(\chi) := \langle f, \chi \rangle \in \mathbb{C}$$

is called the *Fourier transform* of f . More explicitly,

$$\widehat{f}(\chi) = \frac{1}{|G|} \sum_{x \in G} f(x) \overline{\chi(x)}.$$

The equality (4.6.4) can be rewritten

$$f(x) = \sum_{\chi \in \widehat{G}} \widehat{f}(\chi) \chi(x), \quad \forall x \in G, \quad (4.6.5)$$

and, as such, it is known as the *Fourier inversion formula*

If we identify $\chi \in \widehat{G}$ with $\rho(\chi) = (\rho_1, \dots, \rho_d) \in \mathcal{R}_n^d$, then we can view the Fourier transform $\widehat{f}$ as a function on $\mathcal{R}_n^d$

$$\widehat{f}(\rho_1, \dots, \rho_d) = \frac{1}{|G|} \sum_{x \in G} f(x) \rho_1^{-x_1} \cdots \rho_d^{-x_d},$$

and Fourier inversion formula reads

$$f(x) = \sum_{\substack{\rho_k^n = 1, \\ k=1, \dots, d}} \widehat{f}(\rho_1, \dots, \rho_d) \rho_1^{x_1} \cdots \rho_d^{x_d}.$$

Using (4.6.3a) and (4.6.5) we deduce

$$Qf(x) = \sum_{\chi} \widehat{f}(\chi) \cdot \underbrace{\frac{1}{2d} \left(\sum_{k=1}^d (\chi(e_k) + \chi(-e_k)) \right)}_{=: m(\chi)} \cdot \chi(x).$$

Thus

$$Qf = Q \left(\sum_{\chi} \widehat{f}(\chi) \chi \right) = \sum_{\chi} m(\chi) \widehat{f}(\chi) \chi.$$

In other words, the orthonormal basis $\{\chi; \chi \in \widehat{G}\}$ diagonalizes Q and

$$\text{Spec } Q = \{m(\chi), \chi \in \widehat{G}\}.$$

If we write

$$\chi(e_k) = \rho_k = \cos \theta_k + i \sin \theta_k \in \mathcal{R}_n,$$

then $\chi(e_k) + \chi(-e_k) = \rho_k + \bar{\rho}_k = 2 \cos \theta_k$ and

$$m(\chi) = \frac{1}{d} \sum_{k=1}^d \cos \theta_k, \quad \theta_k \in \left\{ 0, \frac{2\pi}{n}, \dots, \frac{2\pi(n-1)}{n} \right\}.$$

Thus $\text{Spec } Q \subset [-1, 1]$ and $1 \in \text{Spec } Q$. The SLE is

$$\lambda_2 = \lambda_2(d, n) = \frac{d-1 + \cos 2\pi/n}{d} = 1 - \frac{2 \sin^2 \pi/n}{d}.$$

Note that

$$\lambda_2(d, n) \sim 1 - \frac{2\pi^2}{dn^2} \text{ as } n \rightarrow \infty. \quad (4.6.6)$$

If $n = 2$, then the random walk on $(\mathbb{Z}/2)^d$ can be alternatively described as follows.

The states of the systems consists of strings of ± 1 -bits of length d ,

$$\vec{\epsilon} = (\epsilon_1, \dots, \epsilon_d), \quad \epsilon_j = \pm 1.$$

The transition corresponds to picking uniformly at random one bit in this string and then flipping it.

In this case all the characters/eigenfunctions are real valued. More precisely, for every

$$\vec{\epsilon} = (\epsilon_1, \dots, \epsilon_d) \in \{-1, 1\}^d$$

we have an eigenfunction $\chi_{\vec{\epsilon}}$ given by

$$\chi_{\vec{\epsilon}}(x) = \prod_{k=1}^d \epsilon_k^{x_k}, \quad \forall x = (x_1, \dots, x_d) \in \{0, 1\}^d. \quad (4.6.7)$$

The corresponding eigenvalue is

$$\lambda_\epsilon = \frac{1}{d}(\epsilon_1 + \cdots + \epsilon_d), \quad \epsilon_k = \pm 1.$$

Hence

$$\text{Spec}(Q) = \left\{ -1 + \frac{2k}{d}, \quad k = 0, 1, \dots, d \right\}.$$

In this case the SLE is

$$\lambda_2(d, 2) = 1 - \frac{2}{d}. \quad (4.6.8)$$

A probability measure μ on G can be identified with a continuous linear functional on $L^2(G, \pi)$ and, as such, can be identified with a function $\mu^* \in L^2(G, \pi)$

$$\mu^* = \sum_{\chi} \mu[\chi] \chi, \quad \mu[\chi] = \sum_{x \in \mathbb{T}_n^d} \mu[x] \chi(x) = \int_{\mathbb{T}_n^d} \chi d\mu.$$

Then

$$\mu \cdot Q^N = \sum_{\chi} m(\chi)^N \chi.$$

□

Example 4.6.3 (The Ehrenfest urn revisited). The random walk $(X_n)_{n \geq 0}$ on

$$\mathcal{V}_d := \{0, 1\}^d,$$

the set of vertices of the hypercube $[0, 1]^d$, is intimately related to Ehrenfest urn; see Example 4.1.7.




Figure 4.8. The random walk along the edges of the 3-dimensional cube $\mathcal{V}_3$.

To see this, consider the *states*

$$s_k := \{x = (x_1, \dots, x_d) \in \mathcal{V}_d; \quad |x| := x_1 + \cdots + x_d = k\}, \quad k \in \llbracket 0, d \rrbracket.$$

Geometrically, the state s_k consists of all vertices of the cube that lie in the hyperplane $x_1 + \cdots + x_d = k$.

If we think of the vertices $x \in \mathcal{V}_d$ as vectors of 0/1 bits, then the random walk has a simple description: if located at $x \in \mathcal{V}_d$, pick a random component of x and flip it to the opposite bit. Note that

$$\mathbb{P}[X_{n+1} \in s_{k+1} | X_n \in s_k] = \frac{d-k}{d}, \quad \mathbb{P}[X_{n+1} \in s_{k-1} | X_n \in s_k] = \frac{k}{d}.$$

We recognize here the transition rules for the Ehrenfest urn model with d particles/balls. Thus, if on our walk along the vertices of the hypercube, we only keep track of the state we are in, we obtain the Markov chain defined by Ehrenfest's urn model. For concrete computations it is convenient to have an alternate description of this phenomenon.

Denote by $\mathfrak{S}_d$ the group of permutations of $\{1, \dots, d\}$. There is an obvious *left* action of $\mathfrak{S}_d$ on $\mathcal{V}_d$,

$$\varphi \cdot (x_1, \dots, x_d) = (x_{\varphi(1)}, \dots, x_{\varphi(d)}), \quad \forall \varphi \in \mathfrak{S}_d, \quad (x_1, \dots, x_d) \in \{0, 1\}^d.$$

On the other hand, $\mathcal{V}_d$ is equipped with a metric, the so called *Hamming distance*,

$$\delta(x, y) = \sum_{i=1}^d |x_i - y_i|, \quad x, y \in \mathcal{V}_d.$$

Two vertices $x, y \in \mathcal{V}_d$ are neighbors (connected by an edge of the cube) iff $\delta(x, y) = 1$. Since the above action of $\mathfrak{S}_d$ preserves the Hamming distance we deduce that $\mathfrak{S}_d$ is a group of graph isomorphisms, i.e.,

$$\forall x, y \in \mathcal{V}_d, \quad \varphi \in \mathfrak{S}_d: \quad x \sim y \iff \varphi \cdot x \sim \varphi \cdot y.$$

Observe also that the states s_k , $k = 0, 1, \dots, d$, are the orbits of the above action of $\mathfrak{S}_d$. Thus, the state space of the Ehrenfest urn model can be identified with the quotient $\bar{\mathcal{V}}_d := \mathfrak{S}_d \backslash \mathcal{V}_d$, the space of orbits of the above left action. Denote by π the invariant probability measure of the random walk on $\mathcal{V}_d$ and $\bar{\pi}$ the invariant measure of the Ehrenfest urn model

$$\bar{\pi}[k] = \frac{1}{2^d} \binom{d}{k}.$$

If $\text{Proj} : \mathcal{V}_d \rightarrow \mathfrak{S}_d \backslash \mathcal{V}_d$ is the natural projection, then

$$\text{Proj}_\# \pi = \bar{\pi}.$$

The left action of $\mathfrak{S}_d$ on $\mathcal{V}_d$ induces a *right* action on the space $L^2(\mathcal{V}_d, \pi)$

$$(f \cdot \varphi)(x) = f(\varphi \cdot x), \quad \forall f : \mathcal{V}_d \rightarrow \mathbb{R}, \quad x \in \mathcal{V}_d, \quad \varphi \in \mathfrak{S}_d.$$

We denote by $L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}$ the subspace consisting of invariant functions, i.e., functions constant along the orbits of $\mathfrak{S}_d$. The pullback

$$\text{Proj}^* : L^2(\mathfrak{S}_d \backslash \mathcal{V}_d, \bar{\pi}) \rightarrow L^2(\mathcal{V}_d, \pi), \quad f \mapsto f \circ \text{Proj}$$

is an isometry onto $L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}$.

Let us observe that the induced linear operator

$$Q : L^2(\mathcal{V}_d, \pi) \rightarrow L^2(\mathcal{V}_d, \pi)$$

is $\mathfrak{S}_d$ -equivariant, i.e., for any $f \in L^2(\mathcal{V}_d, \pi)$, $\sigma \in \mathfrak{S}_d$,

$$Q(f \cdot \sigma) = (Qf) \cdot \sigma \tag{4.6.9}$$

In particular, this shows that

$$Q(L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}) \subset L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}.$$

If $\bar{Q}$ denotes the transition matrix of the Ehrenfest model, then

$$\begin{array}{ccc} L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d} & \xrightarrow{Q} & L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d} \\ \text{Proj}^* \downarrow & & \downarrow \text{Proj}^* \\ L^2(\bar{\mathcal{V}}_d, \bar{\pi}) & \xrightarrow{\bar{Q}} & L^2(\bar{\mathcal{V}}_d, \bar{\pi}) \end{array}$$

If $\lambda \in \text{Spec } Q$ and $\chi \in \ker(\lambda - Q)$ is an eigenfunction of Q , then (4.6.9) implies that $\chi \cdot \varphi \in \ker(\lambda - Q)$, $\forall \varphi \in \mathfrak{S}_d$.

For every $\epsilon \in \{-1, 1\}^d$ we set

$$w(\vec{\epsilon}) = \#\{j; \epsilon_j = -1\}.$$

Note that

$$\sum_j \epsilon_j = d - 2w(\vec{\epsilon}), \quad \lambda(\vec{\epsilon}) = 1 - \frac{2w(\vec{\epsilon})}{d}.$$

If $\lambda_j = 1 - \frac{2j}{d}$, then

$$\ker(\lambda_j - Q) = \text{span}\{\chi_{\vec{\epsilon}}; w(\vec{\epsilon}) = j\}.$$

The orthogonal projection Π onto $L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}$ is the symmetrization operator

$$L^2(\mathcal{V}_d) \ni f \mapsto \Pi f = \frac{1}{d!} \sum_{\varphi \in \mathfrak{S}_d} f \cdot \varphi \in L^2(\mathcal{V}_d, \pi)^{\mathfrak{S}_d}.$$

The above description shows that

$$\Pi \ker(\lambda - Q) \subset \ker(\lambda - \bar{Q}), \quad \forall \lambda \in \text{Spec } Q,$$

so that

$$\text{Spec } \bar{Q} \subset \text{Spec } Q.$$

Since

$$\chi_{\varphi \cdot \vec{\epsilon}}(x) = \chi_{\vec{\epsilon}}(\varphi^{-1} \cdot x), \quad \forall \varphi \in \mathfrak{S}_d, \quad x \in \{0, 1\}^d,$$

we deduce

$$\Pi \chi_{\vec{\epsilon}} = \Pi \chi_{\varphi \cdot \vec{\epsilon}}, \quad \forall \varphi \in \mathfrak{S}_d.$$

Thus $\Pi \chi_{\vec{\epsilon}}$ depends only on $w(\vec{\epsilon})$. We set

$$\Psi_j := \Pi \chi_{\vec{\epsilon}}, \quad w(\vec{\epsilon}) = j.$$

Note that

$$\Psi_j = \frac{1}{\binom{d}{j}} \sum_{w(\vec{\epsilon})=j} \chi_{\vec{\epsilon}}. \quad (4.6.10)$$

Since the eigenfunctions $\chi_{\vec{\epsilon}}$ with fixed weight $w(\vec{\epsilon}) = j$ span the eigenspace of Q corresponding to the eigenvalue λ_j we deduce that

$$\ker(\lambda_j - \bar{Q}) = \text{span}(\Psi_j)$$

so $\dim \ker(\lambda - \bar{Q}) \leq 1, \forall \lambda \in \text{Spec } \bar{Q} \subset \text{Spec } Q$. Hence

$$\# \text{Spec } \bar{Q} = \dim \bar{\mathcal{V}}_d = d + 1 = \# \text{Spec } Q$$

and thus

$$\text{Spec } Q = \text{Spec } \bar{Q} \quad \text{and} \quad \dim \ker(\lambda - \bar{Q}) = 1, \quad \forall \lambda \in \text{Spec } \bar{Q}.$$

Define $K : \mathcal{V}_d \times \mathbb{C} \rightarrow \mathbb{C}$,

$$K(x, z) := \prod_{i=1}^d (1 + (-1)^{x_i} z) = (1 - z)^{|x|} (1 + z)^{d-|x|}, \quad (4.6.11)$$

$$|x| = \sum_i x_i = \#\{i; x_i = 1\}.$$

Observe that

$$K(x, z) = \sum_{j=0}^d \left(\sum_{w(\vec{\epsilon})=j} \chi_{\vec{\epsilon}}(x) \right) z^j \stackrel{(4.6.10)}{=} \sum_{j=0}^d \binom{d}{j} \Psi_j(x) z^j.$$

Thus

$$(1 - z)^{|x|} (1 + z)^{d-|x|} = \sum_j \binom{d}{j} \Psi_j(x) z^j. \quad (4.6.12)$$

Integrating the equality $K(x, z)^2 = (1 - z)^{2|x|} (1 + z)^{2(d-|x|)}$ over $\mathcal{V}_d$ with the uniform probability measure π we deduce

$$\begin{aligned} \int_{\mathcal{V}_d} K(x, z)^2 \pi[dx] &= \frac{1}{2^n} \sum_{k=1}^d \binom{d}{k} (1 - z)^{2k} (1 + z)^{2(d-k)} \\ &= \frac{1}{2^d} ((1 - z)^2 + (1 + z)^2)^d = (1 + z^2)^d. \end{aligned}$$

This shows that

$$\|\Psi_j\|_{L^2(\pi)}^2 = \frac{1}{\binom{d}{j}}.$$

Identify $L^2(\bar{\mathcal{V}}_d, \bar{\pi})$ with the space $\mathbb{R}^{d+1}$

$$L^2(\bar{\mathcal{V}}_d, \bar{\pi}) \ni f \mapsto \begin{bmatrix} f(0) \\ \vdots \\ f(d) \end{bmatrix} \in \mathbb{R}^{1+d}$$

with the inner product

$$\langle u, v \rangle_\pi := \frac{1}{2^d} (Bu, v),$$

where $(-, -)$ denotes the canonical inner product on $\mathbb{R}^{d+1}$,

$$(u, v) = \sum_{i=0}^d u_i v_i,$$

and B is the diagonal matrix

$$B = \text{Diag} \left(\binom{d}{0}, \dots, \binom{d}{d} \right).$$

We denote by c_{kj} the coefficient of z^j in $(1-z)^k(1+z)^{d-k}$. If we think of the invariant eigenfunction Ψ_j as a function on $\bar{\mathcal{V}}_d$, $\Psi_j(k) := \Psi_j(x)$, $|x| = k$, then we have

$$\binom{d}{j} \Psi_j(k) \stackrel{(4.6.12)}{=} c_{kj}, \quad \binom{d}{j} \Psi_j = \underbrace{\begin{bmatrix} c_{0j} \\ \vdots \\ c_{dj} \end{bmatrix}}_{=: C_j}.$$

Denote by C the $(d+1) \times (d+1)$ matrix with columns C_j and by Λ the diagonal matrix

$$\Lambda = \text{Diag}(\lambda_0, \lambda_1, \dots, \lambda_d).$$

If we regard the columns C_j as functions in $L^2(\bar{\mathcal{V}}_d, \bar{\pi})$, then each is a multiple of an eigenfunction Ψ_j of $\bar{Q}$ so that

$$\bar{Q}C_j = \lambda_j C_j, \quad \forall j = 0, 1, \dots, d.$$

Hence $\bar{Q}C = C\Lambda$ so that C diagonalizes $\bar{Q}$,

$$C^{-1}QC = \Lambda, \quad \text{i.e., } Q = C\Lambda C^{-1}.$$

Remarkably, the inverse of C can be described explicitly.

From the equalities

$$\binom{d}{j} \Psi_j = C_j, \quad \|\Psi_j\|_{L^2(\pi)}^2 = \frac{1}{\binom{d}{j}}$$

we deduce

$$\langle C_i, C_j \rangle_\pi = \binom{d}{i} \delta_{ij}, \quad \forall i = 0, 1, \dots, d.$$

In other words ,

$$\frac{1}{2^d} (BCx, Cy) = (Bx, y), \quad \forall x, y \in \mathbb{R}^{d+1}.$$

Hence

$$C^\top BC = 2^d B. \tag{4.6.13}$$

The matrix C has another miraculous symmetry. To prove it we need to get back to the definition of the entries c_{kj} ,

$$(1-z)^k(1+z)^{d-k} = \sum_j c_{kj} z^j.$$

Consider the function

$$F(u, z) = \sum_k \binom{d}{k} u^k (1-z)^k (1+z)^{d-k} = ((1+u) + (1-u)z)^d.$$

On one hand, we have

$$\begin{aligned} F(u, z) &= \sum_k \binom{d}{k} u^k \boxed{(1-z)^k (1+z)^{d-k}} \\ &= \sum_k \binom{d}{k} u^k \boxed{\sum_j c_{kj} z^j} = \sum_{k,j} \binom{d}{k} c_{kj} z^j u^k. \end{aligned}$$

On the other hand, the binomial formula yields

$$\begin{aligned} F(u, z) &= ((1+u) + (1-u)z)^n = \sum_j \binom{d}{j} z^j \boxed{(1-u)^j u^{d-j}} \\ &= \sum_j \binom{d}{j} z^j \boxed{\sum_k c_{jk} u^k} = \sum_{k,j} \binom{d}{j} c_{jk} z^j u^k. \end{aligned}$$

Hence

$$\binom{d}{k} c_{kj} = \binom{d}{j} c_{jk}, \quad \forall j, k.$$

This can be written in more compact form as

$$(BC)_{kj} = (BC)_{jk} \iff BC = (BC)^\top = C^\top B.$$

Using this in (4.6.13) we deduce $BC^2 = 2^d B$ so that

$$C^{-1} = \frac{1}{2^d} C.$$

Hence

$$Q^n = C \Lambda^n C^{-1} = \frac{1}{2^d} C \Lambda^n C, \quad \forall n \geq 0. \quad (4.6.14)$$

The above formula was first obtained by M. Kac [108]. Since then, many different proofs were offered [113, 114, 205].

For more about the rich history and the ubiquity of the Ehrenfest urn we refer to [19, 205]. As a curiosity, we want to mention that the spectrum of $\bar{Q}$ was known to J. J. Sylvester in the 19th century.

One can use (4.6.14) to obtain important information about the dynamics of the Ehrenfest urn such that the return or first passage times $T_i, i = 0, 1, \dots, d$. We refer to [19, 108, 113, 114] for more details.

The above “miraculous” properties of the matrix C are manifestations of the remarkable symmetries of the Krawtchouk polynomials. We refer to [54, 55] for more about these polynomials and their applications in probability. $\square$

4.6.2. Variational methods. Consider a reversible, irreducible, *aperiodic* Markov chain with finite state space $\mathfrak{X}$ and transition matrix Q . Set $N := |\mathfrak{X}|$. Denote by π the invariant probability distribution. We have seen that Q is symmetric as a linear operator

$$L^2(\mathfrak{X}, \pi) \rightarrow L^2(\mathfrak{X}, \pi).$$

We denote by $\langle -, - \rangle_\pi$ the inner product in $L^2(\mathfrak{X}, \pi)$ and by $\| - \|_\pi$ the associated norm. We identify $L^2(\mathfrak{X}, \pi)$ with $\mathbb{R}^N$ equipped with the inner product

$$\langle u, v \rangle_\pi = \sum_{i=1}^N u_i v_i \pi_i.$$

The eigenvalues have variational characterizations. We order the eigenvalues of Q decreasingly

$$1 = \lambda_1 > \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_N \geq -1.$$

Above, each eigenvalue of Q appears as often as its multiplicity. The eigenspace corresponding to the eigenvalue 1 is spanned by the constant function $e = 1$ or, equivalently the column vector $e \in \mathbb{R}^N$ with all the coordinates equal to 1.

As we have seen, the second largest eigenvalue (or SLE) λ_2 controls the rate of convergence of the Markov chain. It has the variational description

$$\lambda_2 = \sup_{\substack{u \in \mathbb{R}^N \setminus \{0\}, \\ \langle u, e \rangle_\pi = 0}} \frac{\langle Qu, u \rangle}{\|u\|_\pi^2}.$$

Let us point out that $\langle u, e \rangle_\pi$ is the mean of u with respect to the probability measure π . We will use this variational characterization to provide upper estimates for λ_2 .

It is more convenient to work with the Laplacian $\Delta := \mathbb{1} - Q$. Note that $\ker \Delta = \text{span}\{e\}$. Its eigenvalues are $\mu_k = 1 - \lambda_k$,

$$0 = \mu_1 < \mu_2 \leq \mu_3 \leq \dots \leq \lambda_N \leq 2.$$

Note that lower estimates for μ_2 are equivalent with upper estimates for λ_2 .

The first positive eigenvalue μ_2 has a variational characterization in terms of the *Dirichlet form*

$$\mathcal{E}(-, -) : L^2(\mathfrak{X}, \pi) \times L^2(\mathfrak{X}, \pi) \rightarrow \mathbb{R}, \quad \mathcal{E}(u, v) = \langle \Delta u, v \rangle_\pi.$$

Lemma 4.6.4.

$$\mathcal{E}(u, v) = \frac{1}{2} \sum_{x, y \in \mathfrak{X}} \pi_x Q_{x, y} (u(x) - u(y)) (v(x) - v(y)).$$

Proof.

$$\begin{aligned} & \sum_{x, y \in \mathfrak{X}} \pi_x Q_{x, y} (u(x) - u(y)) (v(x) - v(y)) \\ &= \underbrace{\sum_{x, y \in \mathfrak{X}} Q_{x, y} (u(x) - u(y)) v(x) \pi_x}_{=: A} - \underbrace{\sum_{x, y \in \mathfrak{X}} \pi_x Q_{x, y} (u(x) - u(y)) v(y)}_{=: B}. \end{aligned}$$

Note that

$$\begin{aligned} A &= \sum_{x \in \mathfrak{X}} \sum_{y \in Y} Q_{x, y} (u(x) - u(y)) v(x) \pi_x \\ &= \sum_{x \in \mathfrak{X}} (u(x) - (Qu)(x)) v(x) \pi_x = \langle \Delta u, v \rangle_\pi. \end{aligned}$$

Using the detailed balance equations $\pi_x Q_{x, y} = \pi_y Q_{y, x}$ we deduce

$$\begin{aligned} B &= \sum_{y \in X} \left(\sum_{x \in \mathfrak{X}} Q_{y, x} (u(x) - u(y)) \right) v(y) \pi_y \\ &= \sum_{y \in \mathfrak{X}} ((Qu)(y) - u(y)) v(y) \pi_y = -\langle \Delta u, v \rangle_\pi. \end{aligned}$$

□

Let us observe that the reversible Markov chain is defined by an electric network with conductances $c(x, y)$, where

$$c(x, y) := \pi_x Q_{x, y}.$$

Then $\forall u, v \in L^2(\mathfrak{X}, \pi)$

$$\mathcal{E}(u, v) = \frac{1}{2} \sum_{x, y \in \mathfrak{X}} c(x, y) (u(x) - u(y)) (v(x) - v(y)) = \langle du, dv \rangle_c,$$

where $\langle -, - \rangle_c$ is the inner product (4.5.10) on 1-cochains and d is the coboundary operator (4.5.8).

The classical Ritz-Raleigh description of eigenvalues of a symmetric operator shows that

$$\mu_2 := \inf \{ \mathcal{E}(u, u); \|u\|_\pi = 1, \langle u, e \rangle_\pi = 0, \}.$$

Note that for any λ in $\mathbb{R}$ we have

$$\mathcal{E}(u + \lambda, u + \lambda) = \mathcal{E}(u, u).$$

If we think of $u \in L^2(\mathfrak{X}, \pi)$ as a random variable defined on the probability space $(\mathfrak{X}, \pi)$, then the above characterization of μ_2 can be rewritten as

$$\mu_2 := \inf_{\substack{\mathbb{E}_\pi[u] = 0 \\ u \neq 0}} \frac{\mathcal{E}(u, u)}{\text{Var}[u]}.$$

Lower bounds of μ_2 are classically known as *Poincaré inequalities*. Thus, a lower bound $\mu_2 > m > 0$ is equivalent to a statement of the form

$$\begin{aligned} \frac{1}{2} \sum_{x,y} \pi_x Q_{x,y} (u(x) - u(y))^2 &\geq m \sum_{x \in \mathfrak{X}} \pi_x u(x)^2 \text{ if } \sum_x u(x) \pi_x = 0 \\ \iff \frac{1}{2} \sum_{x,y} c(x,y) (u(x) - u(y))^2 &\geq m \sum_{x \in \mathfrak{X}} c(x) u(x)^2 \text{ if } \sum_x u(x) c(x) = 0. \end{aligned}$$

To state our first Poincaré type inequality we need a few geometric preliminaries.

To our reversible Markov chain we associate a graph G with vertex set $\mathfrak{X}$. Two vertices x, y are connected by an edge iff $Q(x, y) \neq 0$. We write $x \sim y$ if x and y are connected by an edge in G . This graph could have loops. It is connected since the Markov chain is irreducible. We set

$$\widehat{E} := \{ (x, y) \in \mathfrak{X} \times \mathfrak{X}; x \sim y \}. \quad (4.6.15)$$

We think of the elements of $\widehat{E}$ as edges of G equipped with an orientation. For any $u : \mathfrak{X} \rightarrow \mathbb{R}$ and $e = (x', x'') \in \widehat{E}$ we set

$$\delta_e u := u(x'') - u(x').$$

We can speak of the conductance $c(e)$ of any oriented edge $e = (x, y)$,

$$c(e) := c(x, y) = \pi_x Q_{x,y}.$$

Note that

$$\mathcal{E}(u, u) = \frac{1}{2} \sum_{e \in \widehat{E}} c(e) (\delta_e u)^2. \quad (4.6.16)$$

A *trail* in G between two vertices x, y is a succession of vertices

$$\gamma : x = x_0 \sim x_1 \sim \cdots \sim x_{\ell-1} \sim x_\ell = y,$$

where we do not allow repeated edges. The number ℓ is called the *length* of γ and it is denoted by $\ell(\gamma)$. The trail γ determines a collection of oriented edges

$$e_i = (x_{i-1}, x_i), \quad i = 1, \dots, \ell.$$

We will use the notation $e \in \gamma$ to indicate that e is one of the *oriented* edges determined by γ .

We denote by Γ the collection of trails in G . It comes with an obvious equivalence relation: two trails are equivalent if they have the same initial and final points.

Fix a collection $\mathcal{C}$ of representatives of this equivalence relation. Thus, $\mathcal{C}$ contains exactly one trail $\gamma_{x,y}$ for every pair (x, y) of vertices and this trail connects x to y . Following [57] we set

$$K(\mathcal{C}) := \sup_{e \in \hat{E}} K(\mathcal{C}, e), \quad K(\mathcal{C}, e) := \frac{1}{c(e)} \sum_{\mathcal{C} \ni \gamma_{x,y} \ni e} \ell(\gamma_{x,y}) \pi_x \pi_y. \quad (4.6.17)$$

If an oriented edge e is not contained in any trail $\gamma \in \mathcal{C}$ we set $K(\mathcal{C}, e) = 0$.

Theorem 4.6.5 (Diaconis-Stroock). *For any $u \in L^2(\mathfrak{X}, \pi)$ we have*

$$\text{Var} [u] \leq K(\mathcal{C}) \mathcal{E}(u, u). \quad (4.6.18)$$

Thus $\mu_2 \geq \frac{1}{K(\mathcal{C})}$ so that

$$\lambda_2(Q) \leq 1 - \frac{1}{K(\mathcal{C})}.$$

Proof. We follow the approach in the proof of [57, Proposition 1]. Set $K = K(\mathcal{C})$. Let $u \in L^2(\mathfrak{X}, \pi)$.

For any $x, y \in \mathfrak{X}$ we have the telescoping equality

$$u(y) - u(x) = \sum_{e \in \gamma_{x,y}} \delta_e u.$$

Using the Cauchy Schwartz inequality we deduce

$$(u(y) - u(x))^2 = \left(\sum_{e \in \gamma_{x,y}} \delta_e u \right)^2 \leq \ell(\gamma_{x,y}) \sum_{e \in \gamma_{x,y}} (\delta_e u)^2.$$

Now observe that

$$\begin{aligned} \text{Var} [u] &= \frac{1}{2} \sum_{x,y} (u(y) - u(x))^2 \pi_x \pi_y \leq \frac{1}{2} \sum_{x,y} \ell(\gamma_{x,y}) \sum_{e \in \gamma_{x,y}} (\delta_e u)^2 \\ &= \frac{1}{2} \sum_{e \in \hat{E}} (\delta_e u)^2 \sum_{\gamma_{x,y} \ni e} \gamma_{x,y} = \frac{1}{2} \sum_{e \in E} c(e) (\delta_e u)^2 \underbrace{\frac{1}{c(e)} \sum_{\gamma_{x,y} \ni e} \gamma_{x,y}}_{\leq K} \\ &\leq \frac{K}{2} \sum_{e \in E} c(e) (\delta_e u)^2 \stackrel{(4.6.18)}{=} K \mathcal{E}(u, u). \end{aligned}$$

□

Example 4.6.6. Suppose that our Markov chain corresponds to the random walk on the Cayley graph of the cyclic group $\mathbb{Z}/n\mathbb{Z}$, n odd; see Example 4.6.2. Equivalently, it is the random walk on the set

$$\mathfrak{X} = \{x_i\}_{i \in \mathbb{Z}/n\mathbb{Z}}$$

of vertices of a regular n -gon, where at each vertex we are equally likely to move to one of its two neighbors. In this case we have

$$\begin{aligned} \pi_x &= \frac{1}{n}, \quad Q_{x_i, x_{i+1}} = Q_{x_i, x_{i-1}} = \frac{1}{2}, \quad \forall i \in \mathbb{Z}/n\mathbb{Z}, \\ c(x_i, x_j) &= \frac{1}{2n} \times \begin{cases} 1, & i = j \pm 1, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

As collection $\mathcal{C}$, we choose geodesics (shortest paths) connecting the pair of vertices. Since n is odd, for every $x, y \in \mathfrak{X}$ there exists a unique such geodesic $\gamma_{x,y}$ and it has length $< \frac{n}{2}$. Due to the symmetry of the graph the quantity

$$K(e) := \frac{1}{c(e)} \sum_{\gamma_{x,y} \ni e} \ell(\gamma_{x,y}) \pi_x \pi_y = \frac{2}{n} \sum_{\gamma_{x,y} \ni e} \ell(\gamma_{x,y})$$

is independent of e so

$$K(\mathcal{C}) = K(e), \quad \forall e \in E.$$

Averaging over the n edges of the graph we deduce

$$\begin{aligned} K(\mathcal{C}) &= \frac{1}{n} \sum_e K(e) = \frac{2}{n^2} \sum_e \sum_{\gamma_{x,y} \ni e} \ell(\gamma_{x,y}) \\ &= \frac{2}{n^2} \sum_{x,y} \sum_{e \in \gamma_{x,y}} \ell(\gamma_{x,y}) = \frac{2}{n^2} \sum_{x,y} \ell(\gamma_{x,y})^2 \end{aligned}$$

$$(n = 2m + 1)$$

$$= \frac{2}{n} \sum_{i=1}^n \ell(\gamma_{x_1, x_i})^2 = \frac{4}{n} \sum_{i=1}^m i^2 = \frac{n^2}{6} + O(n), \quad \text{as } n \rightarrow \infty.$$

Hence

$$\lambda_2 \leq 1 - \frac{6}{n^2} + O(n^{-3}), \quad \text{as } n \rightarrow \infty.$$

Thus, for large n this lower estimate is of the same order, as the precise estimate (4.6.6) with $d = 1$. $\square$

We want to describe another geometric estimate for μ_2 of the type first described in Riemannian geometry by J. Cheeger [35].

The volume of a set $S \subset \mathfrak{X}$ is computed using the stationary measure π ,

$$V(S, Q) := \pi[S] = \sum_{s \in S} \pi_s.$$

The “boundary” of the set S is the collection of oriented edges

$$\partial S := \{ (s, s') \in \widehat{E}; \quad s \in S, \quad s' \in S^c \}.$$

The “area” of the boundary of $S \subset \mathfrak{X}$ is

$$A(\partial S, Q) := \sum_{e \in \partial S} c(e) = \sum_{(s, s') \in S \times S^c} \pi_s Q_{s, s'}, \quad S^c = \mathfrak{X} \setminus S.$$

Note that $A(\partial S, Q) = A(\partial S^c, Q)$. The ratio

$$h(S, Q) = \frac{A(\partial S, Q)}{V(S, Q)}$$

is the conditional probability that the Markov chain will transition from a state in S to a state in S^c given that initial distribution is the equilibrium distribution.

Remark 4.6.7. If Q is associated to an electric network with arbitrary conductances $\tilde{c}(x, y)$, then there exists $Z > 0$ such that

$$\tilde{c}(x) = \sum_y \tilde{c}(x, y) = Z \pi_x, \quad \forall x \in \mathfrak{X}.$$

Note that if we define

$$\tilde{V}(S, Q) := \sum_{s \in S} \tilde{c}(s), \quad \tilde{A}(\partial S, Q) := \sum_{e \in \partial S} \tilde{c}(e),$$

then

$$\frac{\tilde{A}(\partial S, Q)}{\tilde{V}(S, Q)} = \frac{A(\partial S, Q)}{V(S, Q)}. \quad \square$$

Now define the *Cheeger isoperimetric constant* or the *conductance* of $(\mathfrak{X}, Q)$ to be

$$\begin{aligned} h(Q) &:= \inf \left\{ h(S, Q); \quad 0 < \mu[S] < \frac{1}{2} \right\} \\ &= \inf \left\{ \max(h(S, Q), h(S^c, Q)); \quad \emptyset \neq S \subsetneq \mathfrak{X} \right\}. \end{aligned} \quad (4.6.19)$$

To get a feeling of the meaning of $h(Q)$ suppose that Q corresponds to the unbiased random walk on a connected graph G with vertex set $\mathfrak{X}$. For any $S \subset \mathfrak{X}$, the area $A(\partial S)$ is, up to a multiplicative constant, the number of edges connecting a vertex in S with a vertex outside S . The volume $V(S)$ is, up to a multiplicative constant the sum of degrees of vertices in S , or equivalently, $V(S) - A(\partial S)$ is twice the number of edges with both endpoints in S . Thus, a “large” $h(Q)$ signifies that, for any subset S of $\mathfrak{X}$, a large fraction of the edges with at least one endpoint in S have the other endpoint outside S .

As an example of graph with small h think of a “bottleneck”, i.e., a graph obtained by connecting with a single edge two disjoint copies of a complete graph.

Various versions of Cheeger’s isoperimetric constant of a (connected) graph play a key role in the definition of *expander families* of graphs, [121, 140]. It was in that context that the connection with random walks on graphs was discovered. For general reversible Markov chains we have the following result due to Jerrum and Sinclair [107].

Theorem 4.6.8. *Let Q denote the transition matrix of a reversible Markov chain with finite state space $\mathfrak{X}$. Then*

$$\mu_2 \geq \frac{h(Q)^2}{2}.$$

In particular,

$$\lambda_2 \leq 1 - \frac{h(Q)^2}{2}.$$

Proof. We follow the presentation in [57]. Let $u \in L^2(\mathfrak{X}, \pi)$. Set $u_+ := \max(u, 0)$,

$$S_u := \{u > 0\} \subset \mathfrak{X}, \quad h(u) = \inf_{S \subset S_u} \frac{A(\partial S, Q)}{V(S)}.$$

Lemma 4.6.9. *If $u \in L^2(\mathfrak{X}, \pi)$ and $u_+ \neq 0$, then*

$$\mathcal{E}(u_+, u_+) \geq \frac{h(u)^2}{2} \|u_+\|_\pi^2, \quad (4.6.20)$$

Proof. We can assume without any loss of generality that $u = u_+$. Then

$$2 \sum_{u(x) < u(y)} (u(y)^2 - u(x)^2) c(x, y) = \sum_{x, y} |u(x)^2 - u(y)^2| c(x, y) \leq$$

$$\begin{aligned}
&\leq \left(\underbrace{\sum_{x,y} (u(x) - u(y))^2 c(x,y)}_{=2\mathcal{E}(u,u)} \right)^{\frac{1}{2}} \left(\sum_{x,y} \underbrace{(u(x) + u(y))^2 c(x,y)}_{\leq 2(u(x)^2 + u(y)^2)} \right)^{\frac{1}{2}} \\
&\leq 2\mathcal{E}(u,u)^{1/2} \left(\underbrace{\sum_{x,y} (u(x)^2 + u(y)^2) c(x,y)}_{=2\|u\|_\pi^2} \right)^{\frac{1}{2}} = 2^{3/2} \mathcal{E}(u,u)^{1/2} \|u\|_\pi.
\end{aligned}$$

We deduce

$$\begin{aligned}
2^{3/2} \mathcal{E}(u,u)^{1/2} \|u\|_\pi &\geq 2 \sum_{u(x) < u(y)} (u(x)^2 - u(y)^2) c(x,y) \\
&= 4 \sum_{x,y} \left(\int_{u(x)}^{u(y)} t dt \right) c(x,y) = 4 \int_0^\infty t \left(\sum_{u(x) \leq t < u(y)} c(x,y) \right) dt.
\end{aligned}$$

If we write $S_t := \{u > t\} \subset S_u$ and observe that

$$\sum_{u(x) \leq t < u(y)} c(x,y) = A(\partial S_t, Q) \geq h(u) \pi[S_t] = h(u) \pi[u > t].$$

We deduce

$$\int_0^\infty t \left(\sum_{u(x) \leq t < u(y)} c(x,y) \right) dt \geq h(u) \int_0^\infty t \pi[u > t] dt \stackrel{(1.3.48)}{=} \frac{h(u)}{2} \|u\|_\pi^2.$$

□

Observe now that for any $x, y \in \mathfrak{X}$ we have

$$(u_+(x) - u_+(y))(u(x) - u(y)) \geq (u_+(x) - u_+(y))^2.$$

To see this, note first that above we have equality if both $u(x)$ and $u(y)$ are nonnegative or both nonpositive. We have strict inequality if one is positive and the other negative, say $u(x) > 0 > u(y)$. Indeed,

$$\begin{aligned}
(u_+(x) - u_+(y))(u(x) - u(y)) &= u(x)(u(x) - u(y)) \\
&> u(x)^2 = (u_+(x) - u_+(y))^2.
\end{aligned}$$

In particular, we deduce that

$$\mathcal{E}(u_+, u) \geq \mathcal{E}(u_+, u_+),$$

and thus,

$$\mu > 0, \quad \Delta u \leq \mu u \text{ on } \{u > 0\} \Rightarrow \mu \|u_+\|_\pi^2 \geq \mathcal{E}(u_+, u_+). \quad (4.6.21)$$

Indeed,

$$\mu \|u_+\|_\pi^2 \geq \lambda \langle u_+, \Delta u \rangle_\pi = \mathcal{E}(u_+, u) \geq \mathcal{E}(u_+, u_+).$$

Combining (4.6.20) and (4.6.21) we deduce that

$$\Delta u \leq \mu u \text{ on } \{u > 0\} \neq \emptyset \Rightarrow \mu \geq \frac{h(u)^2}{2}.$$

Suppose now that u is a nontrivial eigenfunction corresponding to the eigenvalue μ_2 of Δ . Since

$$\sum_x u(x)\pi_x = 0$$

we deduce that $\{u > 0\} \neq \emptyset$ and we conclude that

$$\mu_2 \geq \frac{h(u)^2}{2} \geq \frac{h(Q)^2}{2}$$

as claimed. $\square$

The quantity $h(Q)$ is rather difficult to compute, but lower estimates are easier to obtain. Consider a collection $\mathcal{C}$ of trails in G as in the definition (4.6.17). We set

$$\kappa(\mathcal{C}, e) = \frac{1}{c(e)} \sum_{\mathcal{C} \ni \gamma_{x,y} \ni e} \pi_x \pi_y, \quad \kappa(\mathcal{C}) = \sup_{e \in E} \kappa(\mathcal{C}, e).$$

If an oriented edge e is not contained in any trail $\gamma \in \mathcal{C}$ we set $\kappa(e) = 0$. We have the following result, [57, 195].

Proposition 4.6.10. *We have*

$$h(Q) \geq \frac{1}{2\kappa(\mathcal{C})},$$

so that

$$\lambda_2 \leq 1 - \frac{1}{8\kappa(\mathcal{C})^2}, \quad \forall \mathcal{C}.$$

Proof. Let $S \subset \mathcal{X}$ be a set of vertices with $V(S) = \pi[S] \leq \frac{1}{2}$. We set

$$W(S) = \sum_{\substack{\gamma_{x,y} \in \mathcal{C} \\ x \in S, y \in S^c}} \pi_x \pi_y.$$

Clearly

$$W(S) = \pi[S]\pi[S^c] \geq \frac{1}{2}\pi[S] = \frac{1}{2}V(S).$$

On the other hand

$$W(S) \leq \sum_{e \in \partial S} \sum_{\gamma_{x,y} \ni e} \pi_x \pi_y = \sum_{e \in \partial S} c(e)\kappa(e) \leq \kappa \sum_{e \in \partial S} c(e) = \kappa A(\partial S),$$

and we deduce

$$\kappa A(\partial S) \geq \frac{1}{2}V(S).$$

$\square$

4.6.3. Markov Chain Monte Carlo. Since this is only an invitation to this subject I will not attempt to formulate the most general situation or technique.

Suppose that we want to sample a very large but finite set $\mathfrak{X}$ according to a probability measure on it. The information we have about the set and the given distribution is not complete, but "obtainable".

The probability measure π is known only up to a multiplicative constant. More precisely, we know only a weight $w : \mathfrak{X} \rightarrow (0, \infty)$ that is proportional to π , i.e.,

$$\pi[x] = \frac{w(x)}{Z}, \quad Z = \sum_{x \in \mathfrak{X}} w(x).$$

For all intents and purposes, the normalizing constant Z is not effectively available to us. Still, we would like to produce an $\mathfrak{X}$ -valued random variable with distribution π . E.g., think that we would like to sample uniformly the set Internet sites.

We seek an irreducible, aperiodic, reversible Markov chain $(X_n)_{n \geq 0}$ with state space $\mathfrak{X}$ and invariant distribution π . Preferably, the transition probabilities $Q_{x,x'}$ of such chain would be functions of the ratios $\frac{\pi(x)}{\pi(x')}$, $x, x' \in \mathfrak{X}$. If we can find such a chain, then the distribution of X_n will be close in total variation distance to π . This is in essence Markov Chain Monte Carlo (MCMC) method.

The preferred Ansatz for the transition matrix Q of such a chain is

$$Q_{x,x'} = A_{x,x'} H_{x,x'}, \quad \forall x \neq x',$$

where A is the transition matrix of an irreducible aperiodic chain with state space $\mathfrak{X}$, and the numbers $H_{x,x'} \in (0, 1]$, $x \neq x'$, are such that Q satisfies the detailed balance equation (4.2.5)

$$\pi(x)Q_{x,y} = \pi(y)Q_{y,x}, \quad \forall x, y \in \mathfrak{X}.$$

Often the matrix A describes the random walk on a graph naturally related to the concrete problem at hand. The matrix A is known as the *candidate matrix*.

The idea of this choice is simple. We set

$$N_A(x) := \{y \in \mathfrak{X}; A_{x,y} > 0\}, \quad N_A^*(x) := N_A(x) \setminus \{x\}.$$

The transitions from a state x to a state y according to Q have an intuitive interpretation.

- (i) Pick $y \in N_A(x)$ of x with probability $A_{x,y}$. This is what we would do if we were to follow the Markov chain governed by A . This the *acceptance* part.
- (ii) If $y \neq x$, then we transition to y with probability $H_{x,y}$ and we remain at x with probability $1 - H_{x,y}$.

There are many solutions of the above Ansatz. One of them is provided by the *Metropolis algorithm*. The input of this algorithm is a triplet (A, w, h) , where A is the transition matrix of an irreducible chain with state space $\mathfrak{X}$ such that

$$A_{x,y} > 0 \iff A_{y,x} > 0,$$

$w : \mathfrak{X} \rightarrow (0, \infty)$ is a weight, and h is a function and $h : (0, \infty) \rightarrow (0, 1]$ subject to the constraint

$$h(u) = uh(1/u), \quad \forall u > 0. \quad (4.6.22)$$

The transition matrix A is called the *candidate matrix* and h is called the *activation function*. Often the acceptance matrix is suggested by the problem at hand. Frequently used activation functions are

$$h(u) = \min(1, u), \quad h(u) = \frac{u}{1+u}.$$

We then define $H_{x,y} = h\left(\frac{w(y)A_{y,x}}{w(x)A_{x,y}}\right)$ so the transition matrix Q is given by

$$Q_{x,y} = \begin{cases} 0, & y \notin N_A(x), \\ Q_{x,y} = A_{x,y}h\left(\frac{w(y)A_{y,x}}{w(x)A_{x,y}}\right), & y \in N_A^*(x), \\ 1 - \sum_{x' \in N_A^*(x)} A_{x,y'}h\left(\frac{w(y')A_{y',x}}{w(x)A_{x,y'}}\right), & y = x. \end{cases}$$

Let us show that the Metropolis chain satisfied the detailed balance equation (4.2.5)

$$w(x)Q_{x,y} = w(y)Q_{y,x}, \quad \forall x, y \in \mathfrak{X},$$

so that Q is reversible and its equilibrium distribution is proportional to w .

Indeed, for $x \neq y$, we have

$$\begin{aligned} w(x)Q_{x,y} &= w(x)A_{x,y}h\left(\frac{w(y)A_{y,x}}{w(x)A_{x,y}}\right) \\ &\stackrel{(4.6.22)}{=} w(y)A_{y,x}h\left(\frac{w(x)A_{x,y}}{w(y)A_{y,x}}\right) = w(y)Q_{y,x}. \end{aligned}$$

If the Markov chain A is primitive, or $Q_{x,x} > 0$ for any $x \in \mathfrak{X}$, then so is the Metropolis chain. If not, we replace the Metropolis chain with its lazy version; see Remark 4.4.8. Note that if $h(u) < 1$, $\forall u > 0$, then the Metropolis chain is irreducible and primitive.

Often the candidate matrix A describes the unbiased random walk on a connected unoriented graph G with vertex set $\mathfrak{X}$. In this case G is called the *candidate graph*, $G_A = G$ and

$$A_{x,y} = \frac{1}{d(x)}, \quad \forall x \in \mathfrak{X}$$

where $d(x)$ is the degree of x .

If the graph G is regular, i.e., $d(x) = d(y)$, $\forall x, y$, then the Metropolis chain has a simple interpretation. If situated at a vertex x , pick a neighbor y uniformly randomly. If $h\left(\frac{w(y)}{w(x)}\right) = 1$, then move to y . Otherwise, if $h\left(\frac{w(y)}{w(x)}\right) < 1$, move to y with probability $h\left(\frac{w(y)}{w(x)}\right) < 1$.

We refer to [96, 195] for applications of this algorithm to combinatorics. In general, it is difficult to estimate the SLE or the rate of converges of the Metropolis chain, but in practice it works well.

Example 4.6.11. A few years ago (2016-17) I asked *Mike McCaffrey*, at that time a student writing his senior thesis under my supervision, to read Diaconis' excellent survey [53] and then to try to implement numerically the decryption strategy described in that paper, based on the Metropolis algorithm. I want to report some of McCaffrey's nice findings. For more details I refer to his senior thesis [149].

Let me first outline the encryption problem and the decryption strategy proposed in [53]. The encryption method is a simple substitution cipher. Scramble the 26 letters of the English alphabet $\mathcal{E}$. The encryption is captured by a permutation φ of the set $\mathcal{E}$, or equivalently, an element of φ the symmetric group $\mathfrak{S}_{26}$.

The decryption problem asks to determine the decoding permutation φ^{-1} , given a text encoded by the (unknown) permutation φ . Thus, we need to find one element in a set of $26!$ elements. To appreciate how large $26!$ is, it helps to have in mind that a pile of $26!$ grains of sand will cover the

continental United States with a layer of sand 0.6 miles (approx. one kilometer) thick. We are supposed to find a single grain of sand in this huge pile. Needle in a haystack sounds optimistic!

The strategy outlined in [53] goes as follows. There are 26^2 pairs of letters in the English alphabet $\mathcal{E}$, and they appear as adjacent letters in English texts with a certain frequency. E.g., one would encounter quite frequently the pair “*th*”, less so pairs such as “*tt*” or “*tw*”. We denote by $f(s_1, s_2)$ the frequency of the pair of letters (s_1, s_2) . More precisely $f(s_1, s_2)$ is the conditional probability that in an English text the letter s_1 is followed by s_2 .

To any text of length n , viewed as string of n letters, $\underline{x} = x_1 \dots, x_n$ we associate the weight

$$w(\underline{x}) := \prod_{i=2}^n f(x_{i-1}, x_i).$$

We can use a given encrypted text $\underline{x}$ to define a weight on $\mathfrak{S}_{26}$

$$w(\varphi) := w(\varphi(x_1) \dots \varphi(x_n)).$$

If $\underline{x}$ is obtained from a genuine English text $a_1, \dots, a_n$ via a permutation φ_0 , $x_i = \varphi_0^{-1}(a_i)$, then φ_0 is the decoder $x_i \mapsto \varphi_0(x_i) = a_i$ and

$$w(\varphi_0) := w(a_1 \dots a_n).$$

Intuitively, the permutations with higher weight are closer to the decoding permutation since an incorrect decoder will assign lower frequencies to frequently encountered pairs of adjacent letters. In other words it is reasonable to expect that the decoder satisfies

$$w(\varphi_0) \approx \max_{\sigma \in \mathfrak{S}_{26}} w(\sigma),$$

so we can find it by determining $\operatorname{argmax}_{\sigma \in \mathfrak{S}_{26}} w(\sigma)$.

The weight function w defines a probability measure on $\mathfrak{S}_{26}$. It is reasonable to assume that this measure is highly concentrated around the decoding permutation. If we sample this probability measure there is a high probability that we will land near the decoding permutation.

To sample this probability measure he relied on the Metropolis algorithm. The symmetric group is generated by its $\binom{26}{2} = 325$ transpositions and as candidate graph we take the associated Cayley graph defined by this set of generators. This is a regular graph and the degree of each vertex is 325. As initial state we take the identity permutation. As activation function he used $h(u) = \min(1, u)$.

The question is how well does this work in practice. First, one needs to find the relative frequencies of adjacent pairs in English texts. One can do this by analyzing a large text. In [53] Diaconis suggested using “*War and Peace*”.

Mike McCaffrey used “*Moby Dick*” for this purpose. The table in Figure 4.9 (borrowed from [149]) depicts these relative frequencies.⁶

He then proceeded to test⁷ this method using first a shorter text

THE PROBABILITY THAT WE MAY FAIL IN THE STRUGGLE OUGHT
NOT TO DETER US FROM THE SUPPORT OF A CAUSE WE BELIEVE TO
BE JUST

The scrambled version looked like

⁶He actually used an alphabet consisting of 27 symbols, the 26 letters of the alphabet and a 27-th representing any symbol that is not a letter.

⁷An R-code implementing this algorithm was and is publicly available.




Figure 4.9. Moby Dick Transition Matrix

OVBJCTEADKDOM OVJO SB RJMHJDK DN OVB WOTYXXKB EYXVO
 NEO OE ZBOBT YW HTER OVB WYCCETO EH J QJYWB SB ABKDBLB
 OE AB UYWO

We expect the weight of the decoded text w_{true} to be a lot higher than the weight of the encoded text. In the above example, the weight of the original text is 2.6×10^{115} times higher than that of the cyphered text!!!

After 3,000 steps in the random walk governed by the above Metropolis algorithm, the output was reasonably close to the original text:

THE PROLALINITY THAT WE MAY FAIN ID THE STRUGGNE OUGHT
 DOT TO KETER US FROM THE SUPPORT OF A JAUSE WE LENIEVE TO
 LE BUST

Mike then tested this algorithm on a bigger text. He chose the easily recognizable Gettysburg address by Abraham Lincoln.

The most vivid confirmation of the power of this method came when he presented his results to a mixed group of students in the College of Science of the University of Notre Dame. He began his presentation by projecting the ciphered Gettysburg address, but the audience was left in the dark about the nature of original text. While Mike was describing the problem and the decoding strategy, his laptop was running the algorithm in the background, every few seconds it performed a step on the Metropolis random walk, and then the text on the screen would scramble revealing a new text resembling more and more an English text. Ten minutes or so into his presentation the audience was able to recognize without difficulty the Gettysburg address. It took about 120 steps in the Metropolis random walk to reach an easily recognizable, albeit misspelled text! $\square$

Example 4.6.12 (Gibbs sampler). Statistical physics problems lead to considering probability measures on finite sets $\mathcal{X}$ of the form $\mathcal{X} = \mathbb{B}^S$, where $\mathbb{B} = \{b_0, b_1\}$ is the set of bits and S is a finite set, typically very large, often the set of vertices of a graph. The points in S are called *sites*. The functions $x \in \mathbb{B}^S$ are known as *spin configurations*. The value $x(s)$ is the spin of the site s . Note that $\mathbb{B}$ is

equipped with a nontrivial involution

$$b_i \mapsto \bar{b}_i := b_{1-i}.$$

The physics appears through a given a potential energy function $V : \mathfrak{X} \rightarrow \mathbb{R}$. For any probability distribution $\pi \in \text{Prob}(\mathfrak{X})$ we have an expected potential $\mathbb{E}_\pi[V]$.

A *Gibbs distribution* determined by V and $E \in (\min V, \max V)$ is a distribution $\pi \in \text{Prob}(\mathfrak{X})$ with largest entropy, given that $\mathbb{E}_\pi[V] = E$.

We write $\pi \in \text{Prob}(\mathfrak{X})$ as $\pi[\mathbf{x}] = e^{-f(\mathbf{x})}$, $\mathbf{x} \in \mathfrak{X}$, where $f : \mathfrak{X} \rightarrow (0, \infty)$ satisfies

$$\sum_{\mathbf{x} \in \mathfrak{X}} e^{-f(\mathbf{x})} = 1.$$

Then

$$\text{Ent}[\pi] = \sum_{\mathbf{x}} f(\mathbf{x}) e^{-f(\mathbf{x})}, \quad \mathbb{E}_\pi[V] = \sum_{\mathbf{x} \in \mathfrak{X}} V(\mathbf{x}) e^{-f(\mathbf{x})} = E.$$

Suppose that π is a Gibbs distribution. The Lagrange multipliers theorem shows that there exist $\lambda, \mu \in \mathbb{R}$ such that

$$e^{-f(\mathbf{x})} - f(\mathbf{x}) e^{-f(\mathbf{x})} = -\lambda e^{-f(\mathbf{x})} - \mu V(\mathbf{x}) e^{-f(\mathbf{x})}, \quad \forall \mathbf{x} \in \mathfrak{X}.$$

We deduce that $f(\mathbf{x}) = (\lambda + 1) + \mu V(\mathbf{x})$. The constants μ and λ are determined from the equalities

$$\sum_{\mathbf{x} \in \mathfrak{X}} e^{-\mu V(\mathbf{x})} = e^{\lambda+1}, \quad \sum_{\mathbf{x} \in \mathfrak{X}} V(\mathbf{x}) e^{-\mu V(\mathbf{x})} = e^{\lambda+1} E.$$

More concretely, μ is determined from the equality

$$R(\mu) := \frac{\sum_{\mathbf{x} \in \mathfrak{X}} V(\mathbf{x}) e^{-\mu V(\mathbf{x})}}{\sum_{\mathbf{x} \in \mathfrak{X}} e^{-\mu V(\mathbf{x})}} = E \in (\min V, \max V).$$

Such a constant μ exists since

$$\lim_{\mu \rightarrow \infty} R(\mu) = \min V \quad \text{and} \quad \lim_{\mu \rightarrow -\infty} R(\mu) = \max V.$$

Thus the Gibbs distributions have the form

$$\pi[\mathbf{x}] = \pi_\mu[\mathbf{x}] = \frac{e^{-\mu V(\mathbf{x})}}{Z_\mu}, \quad Z_\mu = \sum_{\mathbf{x} \in \mathfrak{X}} e^{-\mu V(\mathbf{x})}.$$

In statistical physics Z_μ is called the *partition function*. Note that

$$\lambda = \log Z_\mu - 1, \quad E = \frac{d}{d\mu} \log Z_\mu.$$

We can use the Metropolis algorithm to sample a Gibbs distribution. Note that $\mathfrak{X} = \mathbb{B}^S$ can be identified with the set of vertices of a hypercube $\{0, 1\}^N$, $N = \#S$. As candidate matrix we choose the transition matrix for the random walk on the hypercube. In this case $d(\mathbf{x}) = N$, $\forall \mathbf{x} \in \mathfrak{X}$. Recall that two spin configuration $\mathbf{x}, \mathbf{y}$ are connected by an edge in the hypercube if they differ only in one bit, i.e., the Hamming distance $\text{dist}_H(\mathbf{x}, \mathbf{y})$ between $\mathbf{x}$ and $\mathbf{y}$ is 1

$$\sum_{s \in S} \delta_{\mathbf{x}(s), \mathbf{y}(s)} = 1.$$

Fixing an activation function h we obtain a Metropolis chain with transition matrix uniquely determined by

$$Q_{\mathbf{x}, \mathbf{y}} = \frac{1}{N} h(e^{\mu\{V(\mathbf{x}) - V(\mathbf{y})\}}), \quad \text{dist}_H(\mathbf{x}, \mathbf{y}) = 1. \quad (4.6.23)$$

In other words, if we are at a spin configuration $\mathbf{x}$, we first choose a site $s \in S$ uniformly randomly. We then switch the spin $x(s)$ to the opposite $\bar{x}(s)$ with probability $h(e^{\mu(V(\mathbf{x})-V(\mathbf{y}))})$.

There is another Markov chain $(X_n)_{n \geq 0}$ with invariant distribution μ . Suppose $X_n = \mathbf{x} \in \mathfrak{X}$. Then X_{n+1} will differ from $\mathbf{x}$ at most one site, i.e., the Hamming distance between X_n and X_{n+1} is at most 1. Here is how the transition from X_n to X_{n+1} occurs. Choose a site $s \in S$ uniformly randomly. Then

$$X_{n+1}(s') = X_n(s'), \quad \forall s' \neq s,$$

and, for $b \in \mathbb{B}$,

$$\mathbb{P}[X_{n+1}(s) = b \mid X_n = \mathbf{x}] = \mathbb{P}[y(s) = \sigma \mid \mathbf{y}|_{S \setminus \{s\}} = \mathbf{x}|_{S \setminus \{s\}}] =: p_{\mathbf{x},s,b}.$$

Let us observe that the computation of the above conditional probability does not require the knowledge of the partition function Z_μ . In other words it suffices to know the invariant measure only up to a multiplicative constant. For $\mathbf{x} \in \mathfrak{X}$, $s \in S$ and $b \in \mathbb{B}$ denote by $\mathbf{y}_{\mathbf{x},s,b}$ the function $\mathbf{y} \in \mathfrak{X}$ defined by

$$\mathbf{y}(s') = \begin{cases} b & s' = s, \\ \mathbf{x}(s'), & s' \neq s. \end{cases}$$

Thus

$$\mathbb{P}[X_{n+1} = \mathbf{y}_{\mathbf{x},s,b} \mid X_n = \mathbf{x}] = \frac{1}{N} p_{\mathbf{x},s,b}. \quad (4.6.24)$$

This prescription defines a reversible, irreducible, aperiodic Markov chain with invariant distribution $\propto e^{-\mu V(\mathbf{x})}$; see [27, Sec.11.3.1]. This Markov chain is known as the *Gibbs sampler*.

The numerical implementations of these chains work reasonably well if N is not too big. However if N is large, as needed in statistical physics, computing the transitions probabilities can be challenging because it requires computing the potential of V which may involve all the values of $\mathbf{x}$. Given that $\#\mathfrak{X} = 2^N$, it is not practical to compute $V(\mathbf{x})$ for all $\mathbf{x} \in \mathfrak{X}$.

However, they work rather well for potentials V that are relevant in statistical physics. Consider for exemplification the case of the *Ising potential* that arises in magnetism. In this case $\mathbb{B} = \{\pm 1\}$, S is the set of vertices of a simple unoriented graph G . Given $s_0 \neq s_1 \in S$ will use the notation $s_0 \sim s_1$ to indicate that s_0, s_1 are neighbors in G . Given nonzero constants J, H we define

$$V = V_{H,J} : \mathfrak{X} \rightarrow \mathbb{R}, \quad V(\mathbf{x}) = -J \sum_{s \in S} \mathbf{x}(s) - H \sum_{s \sim s'} \mathbf{x}(s) \mathbf{x}(s').$$

In this case

$$p_{\mathbf{x},s,b} = \frac{\exp \{ (-H - J \sum_{s' \sim s} \mathbf{x}(s')) (1+b) \}}{1 + \exp \{ -2(H + J \sum_{s' \sim s} \mathbf{x}(s')) \}}$$

The computation of the transition probability $p_{\mathbf{x},s,b}$ involves only the neighbors of s in the graph G which are typically few in numbers.

For the Ising potential the transition probabilities (4.6.23) are also computable. Indeed, in this case

$$V(\mathbf{x}) - V(\mathbf{y}) = J \sum_s (\mathbf{y}(s) - \mathbf{x}(s)) + H \sum_{s \sim s'} (\mathbf{y}(s) \mathbf{y}(s') - \mathbf{x}(s) \mathbf{x}(s')).$$

If $\text{dist}_H(\mathbf{x}, \mathbf{y}) = 1$, then there exists $s_0 \in S$ such that $\mathbf{y}(s) = \mathbf{x}(s)$, $\forall s \neq s_0$, and $\mathbf{y}(s_0) = \overline{\mathbf{x}(s_0)} = -\mathbf{x}(s_0)$. We deduce

$$V(\mathbf{x}) - V(\mathbf{y}) = -2J\mathbf{x}(s_0) - 4 \sum_{s \sim s_0} \mathbf{x}(s_0) \mathbf{x}(s)$$

$$= -2\mathbf{x}(s_0) \left(J + 2H \sum_{s \sim s_0} \mathbf{x}(s) \right).$$

The *Boltzmann machine* that is used in AI is the Markov chain defined by (4.6.23), where V is an Ising-like potential with $\mathbb{B} = \{0, 1\}$. More precisely

$$V(\mathbf{x}) = - \sum_{s \in S} w(s) \mathbf{x}(s) - \frac{1}{2} \sum_{s \sim s'} w(s, s') \mathbf{x}(s) \mathbf{x}(s'),$$

where

$$w(s) \in \mathbb{R}, \quad w(s, s') = w(s', s) \in \mathbb{R}.$$

In this case $\mathbf{y}(s_0) = \overline{\mathbf{x}(s_0)} = 1 - \mathbf{x}(s_0)$

$$\begin{aligned} V(\mathbf{x}) - V(\mathbf{y}) &= -w(s_0) (\mathbf{y}(s_0) - \mathbf{x}(s_0)) \\ &\quad - \sum_{s \sim s_0} w(s_0, s) (\mathbf{y}(s_0) \mathbf{x}(s) - \mathbf{x}(s_0) \mathbf{x}(s)) \\ &= (\mathbf{y}(s_0) - \mathbf{x}(s_0)) \left(w(s_0) + \sum_{s \sim s_0} w(s_0, s) \mathbf{x}(s) \right). \end{aligned}$$

Its stationary measure is proportional to $e^{-V(\mathbf{x})}$. □

4.7. Exercises

Exercise 4.1. Consider the construction in Remark 4.1.5 of an HMC with initial distribution μ and transition matrix Q as a sequence of random variables defined on $[0, 1)$ equipped with the Lebesgue measure λ . For every $t \in [0, 1)$ there exists

$$\underline{x} = \underline{x}(t) \in \mathfrak{X}^{\mathbb{N}_0}$$

uniquely determined by

$$t \in \bigcap_{n \geq 0} I_{x_0, \dots, x_n}^n$$

- (i) Prove that the resulting map $\Psi : [0, 1) \rightarrow \mathfrak{X}^{\mathbb{N}_0}$ given by $t \mapsto \underline{x}(t)$ is measurable and $\Psi_{\#}\lambda = \mathbb{P}_{\mu}$. **Hint.** Use the π - λ theorem.
- (ii) Prove that the map Ψ is injective and its image is shift-invariant and has $\mathbb{P}_{\mu}$ -negligible complement.
- (iii) Describe the map $t \mapsto \underline{x}(t)$ when $\mathfrak{X} = \{0, 1\}$, $\mu[0] = \mu[1] = \frac{1}{2}$ and

$$Q = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}.$$

Describe explicitly the random variables

$$X_n : [0, 1) \rightarrow \mathbb{R}, \quad X_n(t) = x_n(t),$$

where

$$\underline{x}(t) = (x_0(t), x_1(t), \dots) \in \{0, 1\}^{\mathbb{N}_0}.$$

□

Exercise 4.2. Two people A, B play the following game. Two dice are tossed. If the sum of the numbers showing is less than 7, A collects a dollar from B . If the total is greater than 7, then B collects a dollar from A . If a 7 appears, then the person with the fewest dollars collects a dollar from the other. If the persons have the same amount, then no dollars are exchanged. The game continues until one person runs out of dollars. Let A 's number of dollars represent the states. We know that each person starts with 3 dollars.

- (i) Show that the evolution of A is governed by a Markov chain. Describe its transition matrix.
- (ii) If A reaches 0 or 6, then he stays there with probability 1. What is the probability that B loses in 3 tosses of the dice?
- (iii) What is the probability that A loses in 5 or fewer tosses?

□

Exercise 4.3. Prove that (4.1.4) is equivalent to (4.1.5).

□

Exercise 4.4. Suppose that $(X_n)_{n \geq 0}$ is a homogeneous Markov chain with state space $\mathfrak{X}$ and transition matrix Q . Let $n \in \mathbb{N}$. Prove that X_n is independent of $\{X_k; k \neq n-1, n\}$ given X_{n-1}, X_{n+1} . □

Exercise 4.5. Let $\mathfrak{X}$ be a finite or countable subset. Construct a Markov chain with state space $\mathfrak{X}$ such that any subset of $\mathfrak{X}$ is a closed set of this Markov chain. □

Exercise 4.6. Suppose that $(Y_n)_{n \geq 0}$ is a sequence of i.i.d., $\mathbb{N}_0$ -valued random variables with common probability generating function

$$G(s) = \sum_{k \geq 0} p_k s^k, \quad p_k := \mathbb{P}[Y_n = k], \forall k, n \in \mathbb{N}_0.$$

Let X_n the amount of water in a reservoir at noon on day n . During the 24 hour period beginning at this hour a quantity Y_n flows into reservoir, and just before noon a quantity of one unit of water is removed, if this amount can be found. The maximum capacity of the reservoir is K , excessive inflows are spilled and lost. Show that $(X_n)_{n \geq 0}$ is an HMC, and describe the transition matrix and its stationary distribution in terms of G . $\square$

Exercise 4.7. Denote by X_n the capital of gambler at the end of the n -th game. He relies on the following gambling strategy. If his fortune is $\geq \$4$ he gambles \$2 expecting to win \$4, \$3, \$2 with respective probabilities 0.25, 0.30, 0.45. If his capital is 1, 2 or 3 dollars he bets \$1 expecting him to earn \$2 and \$0 with probabilities 0.45 and respectively 0.55. When his fortune is 0 he stops gambling.

(i) Show that $(X_n)_{n \geq 0}$ is a homogeneous Markov chain, compute its transition probabilities and classify its states.

(ii) Set

$$T := \inf \{ n \in \mathbb{N}; X_n = 0 \}.$$

Show that $\mathbb{P}[T < \infty] = 1$.

(iii) Compute $\mathbb{E}[T]$. $\square$

Exercise 4.8. Suppose that $(X_n)_{n \geq 1}$ is a sequence of nonnegative i.i.d., continuously distributed random variables. Consider the sequence of *records* $(R_n)_{n \in \mathbb{N}}$ defined inductively by the rule

$$R_1 = 1, \quad R_n = \inf \{ n > 1; X_n > \max(X_1, \dots, X_{n-1}) \}.$$

Show that the sequence (R_n) is an Markov chain with state space $\mathbb{N}$ and then compute its transition probabilities. Is this a homogeneous chain? $\square$

Exercise 4.9. At an office served by a single clerk arrives a Poisson stream of clients. More precisely, the n -th client arrives at time $T_n = S_1 + \dots + S_n$ where $(S_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables, $S_n \sim \text{Exp}(\lambda)$. The time to process the n -th client is Z_n , where $(Z_n)_{n \geq 1}$ is a sequence of i.i.d. nonnegative random variables with common distribution $\mathbb{P}_Z$. We assume that the random variables Z_n are independent of the arrival times T_m . For $n \geq 0$ we denote by X_n the number of customers waiting in line immediately after the n -th arrived customer was served.

(i) Show that $(X_n)_{n \geq 0}$ is a homogeneous Markov chain with transition probabilities

$$\mathbb{P}[X_{n+1} = k | X_n = j] = \begin{cases} q_{k-j}, & k \geq j, \\ 0, & k < j, \end{cases}$$

where

$$q_j = \int_0^\infty e^{-\lambda z} \frac{(\lambda z)^j}{j!} \mathbb{P}_Z[dz].$$

Hint. Use Exercise 1.60.

(ii) Set $\mu = \mathbb{E}[Z]$, $r := \lambda\mu$. Prove that the above chain is positively recurrent if and only if $r < 1$.

(iii) Assume that $r < 1$ and $c^2 := \mathbb{E}[Z^2] < \infty$. Prove that

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n] = r + \frac{\lambda^2 c^2}{2(1-r)}.$$

□

Exercise 4.10. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an irreducible HMC with state space $\mathfrak{X}$ and transition matrix Q . Prove that the following statements are equivalent.

(i) The chain is recurrent.

(ii) There exist $x, y \in \mathfrak{X}$ such that

$$\sum_{n \in \mathbb{N}} Q_{x,y}^n = \infty.$$

(iii) For any $x, y \in \mathfrak{X}$ we have

$$\sum_{n \in \mathbb{N}} Q_{x,y}^n = \infty.$$

□

Exercise 4.11. Suppose that $(X_n)_{n \geq 0}$ is an irreducible Markov chain with state space $\mathfrak{X}$ transition probability matrix Q and $x_0 \in \mathfrak{X}$.

(i) For $n \in \mathbb{N}$ set

$$\tau_x(n) = \mathbb{P}_x[T_{x_0} > n], \quad \tau_x = \lim_{n \rightarrow \infty} \tau_x(n)$$

Prove that

$$\tau_x = \sum_{y \neq x_0} Q_{x,y} \tau_y, \quad \forall x \in \mathfrak{X} \setminus \{x_0\}. \quad (4.7.1)$$

(ii) Show that if x_0 is transient, then there exists $x \in \mathfrak{X} \setminus \{x_0\}$ such that $\tau_x \neq 0$

(iii) Suppose there exists a function $\alpha : \mathfrak{X} \setminus \{x_0\} \rightarrow [-1, 1]$, not identically zero, satisfying (4.7.1). Prove that x_0 is transient.

Exercise 4.12. Suppose that $(X_n)_{n \geq 0}$ is a transient irreducible Markov chain with state space $\mathfrak{X}$. Prove that, with probability 1, the chain will exit any finite subset $F \subset \mathfrak{X}$, never to return, i.e.,

$$\mathbb{P} \left[\lim_{n \rightarrow \infty} \mathbf{I}_F(X_n) = 0 \right] = 1.$$

□

Exercise 4.13. Bobby's business fluctuates in successive years between three states between three states: 0 = bankruptcy, 1 = verge of bankruptcy, 2 = solvency. The transition matrix giving the probability of evolving from state to state is

$$Q = \begin{bmatrix} 1 & 0 & 0 \\ 0.5 & 0.25 & 0.25 \\ 0.5 & 0.25 & 0.25 \end{bmatrix}$$

(i) What is the expected number of years until Bobby's business goes bankrupt, assuming it starts in solvency.

- (ii) Bobby's rich father, decides that it is bad for the family name if his son goes bankrupt. Thus, when state 0 is entered, his father infuses Bobby's business with cash returning him to solvency with probability 1. Thus the transition matrix for this Markov chain is

$$P = \begin{bmatrix} 0 & 0 & 1 \\ 0.5 & 0.25 & 0.25 \\ 0.5 & 0.25 & 0.25 \end{bmatrix}$$

Show that this Markov chain is irreducible aperiodic and find the expected number of years between cash infusions from his father.

□

Exercise 4.14. Let Q be a stochastic $n \times n$ matrix and denote by C the $n \times n$ matrix such that $C_{i,j} = \frac{1}{n}, \forall i, j$.

- (i) Prove that for any $r \in (0, 1)$ the Markov chain defined by the stochastic matrix

$$Q(r) = (1 - r)Q + rC$$

is irreducible and aperiodic. Denote by π_r the unique stationary probability measure.

- (ii) Prove that π_r converges as $r \rightarrow 0$ to a stationary probability measure π_0 of the HMC defined by Q .
- (iii) Describe π_0 in the special case when the HMC determined by Q consists of exactly two communication classes C_1 and C_2 and there exist $x_i \in C_i, i = 1, 2$ such that $Q_{x_1, x_2} > 0$.

□

Exercise 4.15. The random walk of a chess piece on a chess table is governed by the rule: the feasible moves are equally likely. Suppose that a rook and a bishop start at the same corner of a 4×4 chess table and perform these random walks. Denote by T the time they meet again at the same corner. Find $\mathbb{E}[T]$.

□

Exercise 4.16. Consider the HMC with state space $\mathfrak{X} = \{0, 1, 2, \dots\}$ and transition matrix Q defined by

$$Q_{n, n+k} = \frac{1}{2^{n+1}} \binom{n}{k}, \quad \forall 0 \leq k \leq n, \quad n \geq 1,$$

$$Q_{0,1} = 1, \quad Q_{n,0} = \frac{1}{2}, \quad \forall n \geq 1.$$

Prove that the chain is irreducible, positively recurrent and aperiodic and find $\mathbb{E}_0[T_0]$.

□

Exercise 4.17. Let K_{n+1} denote the complete graph with $n + 1$ vertices $v_0, v_1, \dots, v_n$. Denote by $(X_n)_{n \geq 0}$ the random walk on K_{n+1} transition rules

$$Q_{v_i, v_j} = \frac{1}{n}, \quad \forall i > 0, \quad j \geq 0, \quad Q_{v_0, v_i} = 0, \quad Q_{v_0, v_0} = 1.$$

Thus the vertex v_0 is absorbent. For $i > 0$ we denote that the time to reach the vertex v_0 starting at v_i ,

$$H_i := \min \{ j \geq 0 : X_0 = i, \quad X_j = 0 \}.$$

Prove that $\mathbb{E}[H_i] = n, \forall i > 0$.

□

Exercise 4.18. Let C_N be the cyclic graph with $N \geq 3$ vertices. We identify the vertex set of C_N with the cyclic group of N -th roots of unity

$$\mathcal{R}_N = \{z \in \mathbb{C}; z^N = 1\}.$$

Denote by $(X_n)_{n \geq 0}$ the standard random walk on C_N . For $z \in \mathcal{R}_N$ denote by H_z the hitting time of z

$$H_z = \inf \{n \geq 0; X_n = z\}.$$

(i) Let $z_1, z_2 \in \mathcal{R}_N, z_1 \neq z_2$. Compute the function

$$\mathcal{R}_N \ni z \mapsto \mathbb{P}_z[H_{z_1} < H_{z_2}].$$

(ii) Let V_n denote the number of vertices visited during the random walk up to epoch n , i.e., the number of pairwise distinct vertices among $X_0, \dots, X_n$. Note that $V_n \leq N$. Set

$$W := \inf \{n \geq 0; V_n = N\}.$$

The stopping time W is called the *covering time* of the cyclic graph. Prove that for any $z \in \mathcal{R}_N^* := \mathcal{R}_N \setminus \{1\}$ we have

$$\mathbb{P}_1[X_W = z] = \frac{1}{N-1}.$$

In other words the random variable $X_W | (X_0 = 1)$ is uniformly distributed⁸ on $\mathcal{R}_N^*$

(iii) Compute $\mathbb{E}_1[W]$.

□

Exercise 4.19. Let K_N be the complete graph with vertex set $\llbracket 1, N \rrbracket$, $N \geq 3$. Denote by $(X_n)_{n \geq 0}$ the standard random walk on C_N . For $i \in \llbracket 1, N \rrbracket$ denote by H_i the hitting time of i

$$H_i = \inf \{n \geq 0; X_n = i\}.$$

(i) Let $i, j \in \mathcal{R}_N, i \neq j$. Compute the function

$$\llbracket 1, N \rrbracket \ni k \mapsto \mathbb{P}_k[H_i < H_j].$$

(ii) Let V_n denote the number of vertices visited during the random walk up to epoch n , i.e., the number of pairwise distinct vertices among $X_0, \dots, X_n$. Note that $V_n \leq N$. Set

$$W := \inf \{n \geq 0; V_n = N\}.$$

Prove that for any $i \in \llbracket 2, N \rrbracket$ we have

$$\mathbb{P}_1[X_W = i] = \frac{1}{N-1}.$$

In other words the random variable $X_W | (X_0 = 1)$ is uniformly distributed on $\llbracket 2, N \rrbracket$

(iii) Compute $\mathbb{E}_1[W]$.

□

⁸This shows that we can sample the uniform distribution on a discrete interval $\llbracket 1, N-1 \rrbracket$ using only a fair coin. Indeed, if $(R_n)_{n \geq 1}$ is a sequence of independent Rademacher random variables and $\chi := e^{\frac{2\pi i}{N}}$, then $\chi^{R_1 + \dots + R_W}$ is uniformly distributed on $\mathcal{R}_N^*$.

Exercise 4.20. A particle performs a random walk on the *nonnegative* integers with transition probabilities

$$p_{0,0} = q, \quad p_{i,i+1} = p, \quad p_{j,j-1} = q, \quad i \geq 0, \quad j > 0,$$

where $p \in (0, 1)$ and $q = 1 - p$.

Prove that the random walk is transient if $p > q$, null recurrent if $p = q$, and positively recurrent if $p < q$. In the last case determine the unique invariant probability distribution. $\square$

Exercise 4.21. We generate a sequence B_n of bits, i.e., 0's and 1's, as follows. The first two bits are chosen randomly and independently with equal probabilities. (Flip a fair 0/1 coin twice and record the results). If $B_1, \dots, B_n$ are generated, then we generate B_{n+1} according to the rules

$$\mathbb{P}[B_{n+1} = 0 \mid B_n = B_{n-1} = 0] = \frac{1}{2} = \mathbb{P}[B_{n+1} = 0 \mid B_n = 0, B_{n-1} = 1]$$

$$\mathbb{P}[B_{n+1} = 0 \mid B_n = 1, B_{n-1} = 0] = \frac{1}{4} = \mathbb{P}[B_{n+1} = 0 \mid B_n = B_{n-1} = 1]$$

What is the proportion of 0's in the long run? $\square$

Exercise 4.22. Consider the Markov chain with state space $\mathfrak{X} = \mathbb{N}_0$ and transition probabilities

$$Q_{n,n-1} = 1, \quad \forall n \in \mathbb{N},$$

$$Q_{0,n} = p_n, \quad \forall n \in \mathbb{N}_0, \quad \sum_{n \geq 0} p_n = 1.$$

Find a necessary and sufficient condition on the distribution $(p_n)_{n \geq 0}$ guaranteeing that the above HMC is positively recurrent. $\square$

Exercise 4.23. Suppose that a gambler plays a fair game with winning probability $p = \frac{1}{2}$. He starts with an initial fortune $X_0 = 1$ dollar. His goal is to reach a fortune of g dollars, $g \in \mathbb{N}$. He stops if he reaches this fortune or he is broke and he is employing a bold strategy: at every game he stakes the largest of money that will get him closest to but not above g . He cannot bet a sum greater than his fortune at that moment. Denote by X_n his fortune after the n -th game.

- (i) Prove that $(X_n)_{n \geq 0}$ is an HMC. Describe its state space and its transition matrix.
- (ii) Prove that the player reaches his goal with probability $\frac{1}{g}$ and goes broke with probability $\frac{g-1}{g}$.

$\square$

Exercise 4.24. Consider the standard random walk in $\mathbb{Z}^2$ started at the origin. For each $m \in \mathbb{Z}$ we denote by T_m the first moment the random walk reaches the line $x + y = m$ and we denote by (U_m, V_m) the point where this walk intersects the above line. Find the probability distributions of T_m, U_m and V_m . $\square$

Exercise 4.25. Suppose that $\mathfrak{X}$ is an at most countable set equipped with the discrete topology $\mu \in \text{Prob}(\mathfrak{X})$ and $Q : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1]$ is a stochastic matrix. Let $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}$ be a sequence of measurable maps.

- (i) Prove that $(X_n)_{n \geq 0} \in \text{Markov}(\mu, Q)$ if and only if

$$\mathbb{E}[f(X_{n+1}) \mid X_n, \dots, X_0] = Q^* f(X_n)$$

for any bounded function $f : \mathfrak{X} \rightarrow \mathbb{R}$.

- (ii) (Lévy) Prove that $(X_n)_{n \geq 0} \in \text{Markov}(\mu, Q)$ if and only if, for any $f \in L^\infty(\mathfrak{X}, \mu)$ the sequence

$$\begin{aligned} Y_0 &= X_0, \quad Y_n = f(X_n) - \sum_{k=0}^{n-1} (Qf(X_k) - f(X_k)) \\ &= f(X_n) + \sum_{k=0}^{n-1} \Delta f(X_k), \quad \Delta = \mathbb{1} - Q. \end{aligned}$$

is a martingale with respect to the filtration $\mathcal{F}_n = \sigma(X_0, X_1, \dots, X_n)$.

□

Exercise 4.26. Consider an irreducible HMC with finite state space $\mathfrak{X}$ and transition matrix Q . We denote by π the invariant probability distribution. For every $x \in \mathfrak{X}$ we denote by H_x the hitting time of x ,

$$H_x := \min \{ n \geq 0; \quad X_n = x \}$$

- (i) Show that

$$\tau(x) := \sum_{y \in \mathfrak{X}} \mathbb{E}_x[H_y] \pi[y]$$

is independent of x .

- (ii) Prove that

$$\tau(x) = \sum_y \mathbb{E}_\pi[H_y] \pi[y].$$

□

Exercise 4.27. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q . Suppose that $B \subset \mathfrak{X}$ and H_B is the *hitting* time of B

$$H_B := \min \{ n \geq 0, \quad X_n \in B \}.$$

We define

$$h_B : \mathfrak{X} \rightarrow [0, 1], \quad h_B(x) = \mathbb{P}_x[H_B < \infty] = \mathbb{P}[H_B < \infty \mid X_0 = x],$$

$$k_B : \mathfrak{X} \rightarrow [0, \infty], \quad k_B(x) = \mathbb{E}_x[H_B].$$

- (i) Show that h_B satisfies the linear system

$$\begin{aligned} h_B(x) &= 1, \quad \forall x \in B, \\ h_B(x) &= \sum_{y \in \mathfrak{X}} Q_{x,y} h_B(y), \quad x \in \mathfrak{X} \setminus B. \end{aligned} \tag{4.7.2}$$

- (ii) Show that if $h : \mathfrak{X} \rightarrow [0, \infty)$ is a solution of (4.7.2), then

$$h_B(x) \leq h(x), \quad \forall x \in \mathfrak{X}.$$

(iii) Show that k_B satisfies the linear system

$$\begin{aligned} k_B(x) &= 0, \quad \forall x \in B, \\ k_B(x) &= 1 + \sum_{y \in \mathfrak{X}} Q_{x,y} k_B(y), \quad x \in \mathfrak{X} \setminus B. \end{aligned} \quad (4.7.3)$$

(iv) Show that if $k : \mathfrak{X} \rightarrow [0, \infty]$ satisfies (4.7.3), then $k_B(x) \leq k(x)$, $\forall x \in \mathfrak{X}$

□

Exercise 4.28. Suppose that $(X_n)_{n \in \mathbb{N}_0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q . For $x \in \mathfrak{X}$ we denote by T_x the return time to x

$$T_x := \min \{ n \geq 1; X_n = x \}.$$

We set

$$\begin{aligned} f_{x,y}(n) &:= \mathbb{P}_x [T_y = n], \\ F_{x,y}(s) &:= \sum_{n \geq 0} f_{x,y}(n) s^n, \quad P_{x,y}(s) := \sum_{n \geq 0} Q_{x,y}^n s^n, \\ f_{x,y} &:= F_{x,y}(1) = \sum_{n \geq 0} f_{x,y}(n) = \mathbb{P}_x [T_y < \infty]. \end{aligned}$$

(i) Prove that

$$P_{x,y}(s) = \delta_{x,y} + F_{x,y}(s) P_{y,y}(s), \quad \forall x, y \in \mathfrak{X},$$

where

$$\delta_{x,y} = \begin{cases} 1, & x = y, \\ 0, & x \neq y. \end{cases}$$

(ii) Deduce from (i) that

$$\sum_{n \geq 0} Q_{x,x}^n < \infty \iff \mathbb{P}_x [T_x < \infty] < 1.$$

(iii) Set $T_x^{(1)} := T_x$ and define inductively $T_x^{(k)} := \min \{ n > T_x^{(k-1)}; X_n = x \}$, $k > 1$. Prove that

$$\mathbb{P} [T_x^{(k-1)} < \infty] = f_{x,y} f_{yy}^{(k-1)}.$$

□

Exercise 4.29. Suppose that we are given an HMC with state space $\mathfrak{X}$ and transition matrix $Q : \mathfrak{X} \times \mathfrak{X} \rightarrow [0, 1]$. For any subset $S \subset \mathfrak{X}$ denote by Q_S the restriction of Q to $S \times S$. For $n \in \mathbb{N}$ define $f_n = f_{S_n} : S \rightarrow [0, 1]$

$$f_n(s) = \sum_{x \in S} Q_S^n(s, x).$$

(i) Prove that $f_1 \geq f_2 \geq \dots \geq 0$. Set

$$f = \lim_{n \rightarrow \infty} f_n.$$

(ii) Prove that for any $s \in S$

$$f(s) = \mathbb{P} [X_n \in S, \quad \forall n \in \mathbb{N} \mid X_0 = s].$$

(iii) Prove that for any $s \in S$

$$f(s) = \sum_{x \in S} Q_S(s, x) f(x).$$

(iv) Prove that if $h : S \rightarrow [0, 1]$ satisfies the above equation,

$$\forall s \in S, \quad h(s) = \sum_{x \in S} Q_S(s, x) h(x),$$

then $h(s) \leq f(s), \forall s \in S$.

(v) Prove that the function f satisfies the dichotomy:

$$\text{either } f(s) = 0, \forall s \in S, \text{ or } \sup_{s \in S} f(s) = 1.$$

(vi) Construct an example of HMC $(\mathfrak{X}, Q)$ and subset $S \subset \mathfrak{X}$ such that f_S is nonconstant.

□

Exercise 4.30. Suppose that $\{X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathfrak{X}\}_{n \geq 0}$ is an irreducible, recurrent HMC with state space $\mathfrak{X}$ and transition matrix Q defined on a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Fix $x \in \mathfrak{X}$, assume $\mathbb{P}[X_0 = x] = 1$ and denote by T_k the time of k -th return to x . More precisely

$$T_0 = 0, \quad T_1 := \min \{n > 0; X_n = x\}, \quad T_{k+1} = \min \{n > T_k; X_n = x\}.$$

We set $Y_k := (X_{T_k}, X_{T_k+1}, \dots, X_{T_{k+1}-1})$.

(i) Realize the quantities Y_k as random maps $Y_k : \Omega \rightarrow \mathcal{Y}$ where $\mathcal{Y}$ is a countable set.

(ii) Show that the resulting random maps are i.i.d..

□

Exercise 4.31. Consider a positively recurrent HMC $(X_n)_{n \geq 0}$ with state space $\mathfrak{X}$, transition matrix Q and stationary distribution π . Suppose that T is a stopping time adapted to $(X_n)_{n \geq 0}$ and let $x \in \mathfrak{X}$ be such that $\mathbb{E}_x[T] < \infty$. We denote by $\mathcal{G}_T(x, y)$ the expected number of visits to y before T , when started at x , i.e.,

$$\mathcal{G}_T(x, y) = \mathbb{E}_x[N_{x,y}^T], \quad N_{x,y}^T = \#\{n \geq 0; X_0 = x, X_n = y, n \leq T\}.$$

Prove that $\mathcal{G}_T(x, y) = \pi[y] \mathbb{E}_x[T]$.

□

Exercise 4.32. Consider a positively recurrent HMC $(X_n)_{n \geq 0}$ with state space $\mathfrak{X}$ and transition matrix Q . Denote by π the stationary distribution. For $x \in \mathfrak{X}$ we denote by T_x the first return time to x and for $y \in \mathfrak{X}$ we set

$$N_{x,y} := \{n \in \mathbb{N}; n \leq T_x, X_n = y\}, \quad \mathcal{G}(x, y) := \mathbb{E}_x[N_{x,y}].$$

In other words, $\mathcal{G}(x, y)$ is the expected number of visits to y before returning to x .

(i) Prove that $\mathcal{G}(x, y) = \pi[y] \mathbb{E}_x[T_x]$.

(ii) Prove that

$$\mathbb{P}_x[T_y < T_x] = \frac{1}{\pi[y] (\mathbb{E}_x[T_y] + \mathbb{E}_y[T_x])}.$$

□

Exercise 4.33. Prove the claims in Remark 4.4.11. $\square$

Exercise 4.34 (LeCam). Suppose that $(X_r)_{1 \leq r \leq n}$ is a family of independent Bernoulli random variables with success probabilities p_r and $(Y_r)_{1 \leq r \leq n}$ a family of independent Poisson random variables $Y_r \sim \text{Poi}(p_r)$. Set

$$S_n := \sum_{r=1}^n X_r, \quad \lambda := p_1 + \cdots + p_n.$$

Fix independent optimal couplings $(\hat{X}_r, \hat{Y}_r)$ of (X_r, Y_r) ; see Remark 4.4.11.

(i) Show that the distribution of $(\hat{X}_r, \hat{Y}_r)$ is the measure λ_r on $\mathbb{N}_0 \times \mathbb{N}_0$ given by

$$\lambda_r[(m, n)] = \begin{cases} 1 - p_r, & m = n = 0, \\ e^{-p_r} - 1 + p_r, & m = 1, n = 0, \\ \frac{p_r^n}{n!} e^{-p_r}, & m = 1, n \geq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

(ii) Prove that $d_v(\text{Bin}(p_r), \text{Poi}(p_r)) \leq p_r^2, \forall r = 1, \dots, n$.

(iii) Prove that

$$d_v(S_n, \text{Poi}(\lambda)) \leq \sum_r \mathbb{P}[\hat{X}_r \neq \hat{Y}_r] \leq \sum_{r=1}^n p_r^2.$$

$\square$

Exercise 4.35. Let X, Y be two random variables defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. We say that Y *pointwisely dominates* X if $Y \geq X$ a.s.. We say that Y *stochastically dominates* X and we denote this $Y \stackrel{d}{\geq} X$, if $F_Y(t) \leq F_X(t), \forall t \in \mathbb{R}$, where F_X and F_Y are the cumulative distribution functions of X and respectively Y .

(i) Prove that

$$X \leq Y \Rightarrow X \stackrel{d}{\leq} Y.$$

(ii) Let $(\hat{X}, \hat{Y})$ be any coupling of X and Y . Prove that

$$X \stackrel{d}{\leq} Y \iff \hat{X} \stackrel{d}{\leq} \hat{Y}.$$

(iii) Let $0 < p_0 < p_1 < 1$. Prove that $\text{Bin}(n, p_0) \stackrel{d}{\leq} \text{Bin}(n, p_1), \forall n \in \mathbb{N}$. **Hint.** Let $(U_n)_{n \in \mathbb{N}}$ be an i.i.d. sequence of $\text{Unif}([0, 1])$ random variables. For $p \in (0, 1)$ and $n \in \mathbb{N}$ set $Y_{n,p} := \sum_{j=1}^n \mathbf{I}_{[0,p]}(U_j)$.

(iv) We denote by Q_X and Q_Y the quantiles of X and respectively Y ; see Example 1.2.22. Prove that

$$X \stackrel{d}{\leq} Y \iff Q_X(U) \leq Q_Y(U),$$

where $U \sim \text{Unif}([0, 1])$. The pair $(Q_X(U), Q_Y(U))$ is called the *quantile coupling* of X and Y .

(v) Let $X_0 \sim \text{Ber}(p_0), X_1 \sim \text{Ber}(p_1)$ be two independent Bernoulli random variables with $0 < p_0 < p_1 < 1$. Set $p_2 := \frac{p_1 - p_0}{1 - p_0}$ and let $X_2 \sim \text{Ber}(p_2)$ be independent of X_0, X_1 . Set $Y_1 := \max(X_0, X_2)$. Show that (X_0, Y_1) is a coupling of (X_0, X_1) such that $X_0 \leq Y_1$, a.s.. $\square$

Exercise 4.36 (Card shuffles). Think of a permutation φ of $\mathbb{I}_m = \{1, 2, \dots, m\}$ as an arrangement of a deck of m cards, with $\varphi(1)$ the top-of-the-deck card. A *top-to-random shuffle* consists of moving the top card, with equal probability, below one of the m -cards of the deck. Leaving the top card in place is one of the m equiprobable options. Algebraically, if $k \in \mathbb{I}_m$ and $\sigma_k \in \mathfrak{S}_m$ represents the permutation

$$2, 3, \dots, k, 1, k+1, \dots, m,$$

then we can describe a top-to-random shuffle as a transformation $\varphi \rightarrow \varphi \circ \sigma_K$, where K is a random variable uniformly distributed on $\mathbb{I}_m$. Fix a sequence $(K_n)_{n \in \mathbb{N}}$ of independent random variables uniformly distributed on $\mathbb{I}_m$. Consider the random walk on $\mathfrak{S}_m$ started at $\Phi_0 = \mathbb{1} \in \mathfrak{S}_m$ and transition rule $\Phi_n = \Phi_{n-1} \circ \sigma_{K_n}$. Set

$$T_m := 1 + S_m, \quad S_m = \min \{n \in \mathbb{N}; \Phi_n(1) = \Phi_0(m) = m\}.$$

In other words, T_m is the first moment when the card that initially was at the bottom appears on top.

- (i) Prove that T_m has the same distribution as the coupon collector random variable, i.e., the minimum number of samplings with replacements of m objects until each one of them was sampled.
- (ii) Prove that Φ_{T_m} is uniformly distributed on $\mathfrak{S}_m$ and it is independent of T_m . **Hint.** Prove by induction on k that given that at a moment t there are k cards under the bottom card, then all of the $k!$ possible arrangements of these cards are equally.
- (iii) Prove that

$$\mathbb{P}[T_m \leq n, X_n = \varphi] = \frac{1}{m!} \mathbb{P}[T_m \leq n].$$

- (iv) Prove that for any $A \subset \mathfrak{S}_m$ we have

$$|\mathbb{P}[\Phi_n \in A] - \pi_m[A]| \leq 2\mathbb{P}[T_m > n] \leq \frac{2m \log m}{n}.$$

□

Exercise 4.37. Let $(X_n)_{n \geq 0}$ be an irreducible Markov chain with finite state space $\mathfrak{X}$, transition matrix $\mathcal{Q}$ and invariant probability measure $\mu \in \text{Prob}(\mathfrak{X})$. Assume that the initial distribution is also μ , i.e., $\mathbb{P}_{X_0} = \mu$. For $n \in \mathbb{N}_0$ we set (see Exercise 2.61 for notation)

$$H_n = \frac{1}{n+1} \text{Ent}_2[X_0, X_1, \dots, X_n], \quad L_n = \text{Ent}_2[X_n | X_{n-1}, \dots, X_0].$$

- (i) Prove that the sequence $(L_n)_{n \geq 0}$ is non-increasing and nonnegative. Denote by L its limit.
- (ii) Prove that

$$H_n = \frac{1}{n+1} \sum_{k=0}^n L_k$$

- (iii) Prove that the sequence (H_n) is convergent and its limit is L .
- (iv) Prove that

$$L = - \sum_{x \in \mathfrak{X}} \mu[x] \text{Ent}_2[Q_{x,-}] = - \sum_{x,y \in \mathfrak{X}} \mu[x] Q_{x,y} \log_2 Q_{x,y}.$$

The number L is called *entropy rate* of the irreducible Markov chain. We denote it by $\text{Ent}_2[\mathfrak{X}, \mathcal{Q}]$.

□

Exercise 4.38. Suppose that $(\mathfrak{X}, Q, \mu_0)$ is an irreducible, positively recurrent HMC with state space $\mathfrak{X}$, initial distribution μ_0 and equilibrium (limit) distribution μ_∞ . For $n \in \mathbb{N}$ set

$$\mu_n = \mu \cdot Q^n \in \text{Prob}(\mathfrak{X}).$$

(i) Prove that for any $n \in \mathbb{N}$

$$\mathbb{D}_{KL}(\mu_{n-1} \parallel \pi) \geq \mathbb{D}_{KL}(\mu_n \parallel \pi),$$

where $\mathbb{D}_{KL}$ is the KL-divergence

$$\mathbb{D}_{KL}(\mu \parallel \pi) = \sum_{x \in \mathfrak{X}} \mu[x] \log_2 \frac{\mu[x]}{\pi[x]}.$$

(ii) Prove that for any $n \in \mathbb{N}$

$$\text{Ent}_2[X_n | X_0] \leq \text{Ent}_2[X_{n+1} | X_0],$$

where $\text{Ent}_2[X | Y]$ denotes the relative entropy defined in Exercise 2.61.

□

Exercise 4.39. Let Q denote the $n \times n$ transition matrix describing the random walk on a complete graph with n vertices. Find the spectrum of Q . □

Exercise 4.40 (Doebelin). Suppose that $(X_n)_{\geq 0}$ is an HMC with state space $\mathfrak{X}$, initial distribution μ and transition matrix Q satisfying the *Doebelin condition*

$$\exists \varepsilon > 0, \exists x_0 \in \mathfrak{X} : Q_{x, x_0} > \varepsilon, \forall x \in \mathfrak{X}.$$

Denote $\mathcal{M}$ the space of finite signed measures ρ on $\mathfrak{X}$. For $\rho \in \mathcal{M}$ we set

$$\|\rho\|_1 := \sum_{x \in \mathfrak{X}} |\rho_x| < \infty, \quad \rho_x := \rho[\{x\}].$$

(i) Prove that for any $\rho \in \mathcal{M}$ we have $\rho Q \in \mathcal{M}$ and

$$\sum_{x \in \mathfrak{X}} \rho_x = \sum_{y \in \mathfrak{X}} (\rho Q)_y.$$

(ii) Prove that if $\rho \in \mathcal{M}$ and

$$\sum_{x \in \mathfrak{X}} \rho_x = 0,$$

then

$$\|\rho Q\|_1 \leq (1 - \varepsilon) \|\rho\|_1$$

(iii) Set $\mu_n := \mu \cdot Q^n$. Prove that

$$\|\mu_n - \mu_m\|_1 \leq 2(1 - \varepsilon)^m, \quad \forall n \geq m \geq 1.$$

(iv) Prove that the HMC is irreducible, positively recurrent and the unique invariant probability measure π satisfies

$$\|\mu_n - \pi\|_1 \leq 2(1 - \varepsilon)^n, \quad \forall n \in \mathbb{N}.$$

□

Exercise 4.41. Suppose that $(X_n)_{n \geq 0}$ is an HMC with state space $\mathfrak{X}$, initial distribution μ and transition matrix Q . For each $n \in \mathbb{N}$ we set

$$A_n := \frac{1}{n+1} \sum_{k=0}^n Q^k$$

Suppose that there exist $N \in \mathbb{N}$, $x_0 \in \mathfrak{X}$ and $\varepsilon > 0$ such that

$$(A_N)_{x, x_0} > \varepsilon, \quad \forall x \in \mathfrak{X}.$$

Prove that the HMC is irreducible, positively recurrent and the unique invariant probability measure π satisfies

$$\|\mu A_n - \pi\|_1 \leq \frac{N}{(n+1)\varepsilon}, \quad \forall n \in \mathbb{N}.$$

□

Exercise 4.42. Prove Lemma 4.5.2. □

Exercise 4.43. Suppose that $(\mathfrak{X}, E, c)$ is a finite connected electric network and x_+, x_- are distinct vertices. The *commute time* between x_+, x_- is the quantity

$$K_{x_+, x_-} = \mathbb{E}_{x_+} [T_{x_-}] + \mathbb{E}_{x_-} [T_{x_+}].$$

Set

$$C(\mathfrak{X}) := \sum_{e \in E} c(e).$$

(i) Prove that

$$K_{x_+, x_-} = 2C E_{x_+, x_-}$$

where E_{x_+, x_-} denotes the energy of the Kirchhoff flow with source x_+ and sink x_- ; see (4.5.26). **Hint.** Use Exercise 4.32.

(ii) Consider the Ehrenfest urn with B balls defined in Example 4.1.7. Use (i) to compute $E_0 [T_B]$. In other words, given that initially all the B balls were in the right chamber, find the expected time until all of them move in the left chamber. □

Exercise 4.44. Consider a HMC $(X_n)_{n \geq 0}$, with state space $\mathfrak{X}$, transition function Q . Fix $N \in \mathbb{N}$ and denote by $\mathcal{T}_N$ the set of all stopping times T adapted to the process $(X_n)_{n \geq 0}$ such that $0 \leq T \leq N$ a.s.. Fix a *reward function*

$$R : \{0, 1, \dots, N\} \times \mathfrak{X} \rightarrow [0, \infty).$$

We define inductively $R_N^*(x) = R(N, x)$,

$$R_{m-1}^*(x) = \max \left(R(m-1, x), \sum_{y \in \mathfrak{X}} Q_{x,y} R_m^*(y) \right), \quad 1 \leq m \leq N.$$

$$T_* = \inf \{ k, \quad R_k^*(X_k) = R(k, X_k) \}.$$

- (i) Prove that $U_k := R_k^*(X_k)$ is a super-martingale.
- (ii) Prove that $V_k = U_{k \wedge T_*}$ is a martingale.
- (iii) Show that

$$\mathbb{E}_x [R(T, X_T)] \leq R_0^*(x) = \mathbb{E}_x [R(T^*, X_{T_*})], \quad \forall T \in \mathcal{T}_N.$$

□

Elements of Ergodic Theory

Ergodic theory is a rather eclectic subject with applications in many areas of mathematics, including probability. The ergodicity feature first appeared in the works of L. Boltzmann on statistical mechanics, [30]. The modern formulation of this hypothesis, due to Y. Sinai, came much later, in 1963, and it took a few more decades to be adjudicated mathematically.

Our rather modest goal in this chapter is to describe enough of the fundamentals of this theory so we can shed new light on some of the fundamental limit theorems we have proved in the previous chapters. For more details we refer to [4, 17, 46, 122, 173, 217] that served as our main sources of inspiration.

5.1. The ergodic theorem

5.1.1. Measure preserving maps and invariant sets. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space. A measurable map $T : (\Omega, \mathcal{S}) \rightarrow (\Omega, \mathcal{S})$ is said to be *measure preserving* if $T_{\#}\mathbb{P} = \mathbb{P}$, i.e.,

$$\mathbb{P}[T^{-1}(S)] = \mathbb{P}[S], \quad \forall S \in \mathcal{S}. \quad (5.1.1)$$

The measure preserving map T is called an *automorphism* of the probability space if it is bijective, and its inverse is also measure preserving.

Proposition 1.2.4 shows that (5.1.1) is satisfied if and only if there exists a π -system $\mathcal{C}$ that generates $\mathcal{S}$ such that

$$\mathbb{P}[T^{-1}(C)] = \mathbb{P}[C], \quad \forall C \in \mathcal{C}. \quad (5.1.2)$$

Example 5.1.1. (a) Let $\mathbb{P}_{S^1}$ denote the Euclidean probability measure on S^1 , the unit circle in $\mathbb{R}^2$, i.e. (see Example 1.2.66)

$$\mathbb{P}_{S^1}[d\theta] = \frac{1}{2\pi}d\theta.$$

We denote by R_φ the counterclockwise rotation by an angle φ about the center of this circle. Then R_φ is measure preserving. If we think of S^1 as the set of complex numbers of norm 1,

$$S^1 := \{z \in \mathbb{C}; |z| = 1\},$$

then $R_\varphi(z) = e^{i\varphi}z$.

(b) Consider the n -dimensional torus $\mathbb{T}^n := \mathbb{R}^n/\mathbb{Z}^n$. Set $I = [0, 1]$ and observe that the natural projection $\pi : I^n \rightarrow \mathbb{T}^n$ is Borel measurable. We denote by $\mathbb{P}_{\mathbb{T}^n}$ the push-forward by π of the Lebesgue measure on I^n . The resulting probability space is isomorphic to the product of n copies of $(S^1, \mathcal{B}_{S^1}, \mathbb{P}_{S^1})$.

Suppose that $A \in \text{SL}_n(\mathbb{Z})$, i.e., A is an $n \times n$ matrix with integer coefficients and determinant 1. Then $A(\mathbb{Z}^n) = \mathbb{Z}^n$ and thus we have a well defined induced map

$$T_A : \mathbb{R}^n/\mathbb{Z}^n \rightarrow \mathbb{R}^n/\mathbb{Z}^n.$$

This map is clearly a homeomorphism, hence Borel measurable. It is also measure preserving since $\det A = 1$.




Figure 5.1. Arnold's cat map.

In [4] Arnold and Avez memorably depicted the action of the map T_A for

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$$

as in Figure 5.1. This map is popularly known as *Arnold's cat map*.

(c) In the previous examples, the maps were automorphisms of the corresponding probability spaces. Here is an example of a measure preserving map that is not bijective. More precisely, define

$$Q : S^1 \rightarrow S^1, \quad Q(z) = z^2.$$

Then the Lebesgue measure $\frac{1}{2\pi}d\theta$ is Q -invariant. If we identify S^1 with $\mathbb{R} \bmod \mathbb{Z}$, then we can describe Q as the map $Q : [0, 1) \rightarrow [0, 1)$ given by

$$Q(x) = 2x \bmod 1.$$

If $x \in [0, 1)$ has binary expansion

$$x = \sum_{n=1}^{\infty} \frac{\epsilon_n}{2^n}, \quad \epsilon_n \in \{0, 1\},$$

then

$$Q(x) = \sum_{n=1}^{\infty} \frac{\epsilon_{n+1}}{2^n}.$$

(d) Consider the *tent map* $T : [0, 1] \rightarrow [0, 1]$, $T(x) = \min(2x, 2 - 2x)$. Equivalently, this is the unique continuous map such that $T(0) = T(1) = 0$, $T(1/2) = 1$ and it is linear on each of the intervals $[0, 1/2]$ and $[1/2, 1]$. Its graph looks like a tent with vertices $(0, 0)$, $(1/2, 1)$ and $(1, 0)$.

This map preserves the Lebesgue measure. Indeed, if $I \subset [0, 1]$ is a compact interval then $T^{-1}(I)$ consists of two intervals $I_{\pm}$, symmetrically located with respect to the midpoint $1/2$ of $[0, 1]$, and each having half the size of I .

(e) Suppose that X is a compact metric space and $T : X \rightarrow X$ is a continuous map. Denote by $\text{Prob}(X)$ the set of Borel probability measures on X . Then map T induces a push-forward map $T_{\#} : \text{Prob}(X) \rightarrow \text{Prob}(X)$. The T -invariant measures are precisely the fixed points of $T_{\#}$. One can show (see Exercise 5.2) that the set $\text{Prob}_T(X)$ of T -invariant measures is nonempty, convex and closed with respect to the weak convergence. $\square$

Example 5.1.2 (Stationary sequences). Let $(\mathbb{X}, \mathcal{F})$ be a measurable space and suppose that

$$X_n : \Omega \rightarrow \mathbb{X}, \quad n \in \mathbb{N},$$

is a sequence of random maps defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. The sequence is said to be *stationary* if for any $m, k \in \mathbb{N}$ the random vectors

$$(X_1, \dots, X_m) : \Omega \rightarrow \mathbb{X}^m \quad \text{and} \quad (X_{k+1}, \dots, X_{k+m}) : \Omega \rightarrow \mathbb{X}^m$$

have the same distribution.

- (i) For example, a sequence of i.i.d. random variables is stationary. More generally, an exchangeable sequence of random variables is stationary.
- (ii) Suppose that $(X_n)_{n \geq 0}$ is an HMC with state space $\mathfrak{X}$ and transition matrix Q and initial distribution μ . The sequence $(X_n)_{n \geq 0}$ is stationary if and only if μ is an invariant distribution, i.e., $\mu = \mu \cdot Q$.

To any stationary sequence $X_n : \Omega \rightarrow \mathbb{X}$, we can canonically associate a measure preserving map as follows. Consider the *path space* $\mathbb{U} = \mathbb{X}_{\mathbb{N}} := \mathbb{X}^{\mathbb{N}}$. It consists of sequences of points in $\mathbb{X}$, $\underline{u} = (u_1, u_2, \dots)$. We have natural coordinate maps $U_n : \mathbb{U} \rightarrow \mathbb{X}$, $U_n(\underline{u}) = u_n$.

For $n \in \mathbb{N}$ denote by $\mathcal{U}_n$ the sigma-algebra generated by $U_1, \dots, U_n$ and we set

$$\mathcal{U} = \bigvee_{n=1}^{\infty} \mathcal{U}_n.$$

Note that we have a $\mathcal{U}$ -measurable map *shift map*

$$\Theta : \mathbb{U} \rightarrow \mathbb{U}, \quad \Theta(u_1, u_2, \dots) = (u_2, u_3, \dots).$$

A sequence of random variables $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{F})$, $n \in \mathbb{N}$, defines a measurable map

$$\vec{X} : (\Omega, \mathcal{S}) \rightarrow (\mathbb{U}, \mathcal{U}), \quad \omega \mapsto (X_1(\omega), X_2(\omega), \dots).$$

The distribution of this sequence is the push-forward probability measure $\mathbb{P}_{\vec{X}} := \vec{X}_{\#} \mathbb{P}$. Note that

$$X_n = U_n \circ \vec{X} \quad \text{and} \quad U_{k+1} = U_1 \circ \Theta^k, \quad \forall n, k \in \mathbb{N}.$$

Since the measure $\mathbb{P}_{\bar{X}}$ is uniquely determined by its restrictions to the sigma-subalgebras $\mathcal{U}_n$ we deduce that the sequence $(X_n)_{n \in \mathbb{N}}$ is stationary iff the shift Θ preserves the distribution $\mathbb{P}_{\bar{X}}$ on the path space.

When $\mathbb{X}$ is finite or countable, and the sequence $(X_n)_{n \in \mathbb{N}}$ is i.i.d., the resulting shift is known as *Bernoulli shift*.

Conversely, if T is a measurable map $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\Omega, \mathcal{S}, \mathbb{P})$, then it is measure preserving if and only if, for any measurable function $f : \Omega \rightarrow \mathbb{R}$ the sequence of random variables

$$f, f \circ T, f \circ T^2, \dots$$

is stationary. □

Definition 5.1.3. Suppose that $(\Omega, \mathcal{S})$ is a measurable space and

$$T : (\Omega, \mathcal{S}) \rightarrow (\Omega, \mathcal{S})$$

is a measurable map.

- (i) A measurable function $f : \Omega \rightarrow \mathbb{R}$ is called *T-invariant* if $f \circ T = f$.
- (ii) A measurable set $S \in \mathcal{S}$ is called *T-invariant* if its indicator $\mathbf{I}_S$ is an invariant function.
- (iii) We denote by $\mathcal{I} = \mathcal{I}_T$ the collection of invariant sets. □

Remark 5.1.4. (a) Note the definition of $\mathcal{I}_T$ involves no measure on $\mathcal{S}$.

(b) Note that if $S \in \mathcal{S}$, then $\mathbf{I}_S \circ T = \mathbf{I}_{T^{-1}(S)}$ so the set S is *T-invariant* iff $S = T^{-1}(S)$. Observe that

$$S \subset T^{-1}(S) \iff T(S) \subset S, \quad (5.1.3a)$$

$$T^{-1}(S) \subset S \iff \forall \omega \in \Omega, \quad T(\omega) \in S \Rightarrow \omega \in S. \quad (5.1.3b)$$

We can give a dynamic description of invariance. For $\omega \in \Omega$ we denote by $\mathcal{O}_T(\omega)$ the orbit of ω with respect to the action of T

$$\mathcal{O}_T(\omega) := \{ \omega, T(\omega), T^2(\omega), \dots \}.$$

A set S is invariant if and only if

$$\omega \in S \Rightarrow \mathcal{O}_T(\omega) \subset S \text{ and } \omega \in \Omega \setminus S \Rightarrow \mathcal{O}_T(\omega) \subset \Omega \setminus S.$$

In the universal case $(\mathbb{U}_{\mathbb{X}}, \mathcal{U})$, a subset $S \in \mathcal{U}_{\mathbb{X}}$ is Θ -invariant if

$$\underline{s} = (s_1, s_2, \dots) \in S \Rightarrow (s_2, s_3, \dots) \in S,$$

$$(s_2, s_3, \dots) \in S \Rightarrow \forall s_1 \in \mathbb{X} : (s_1, s_2, s_3, \dots) \in S.$$

Note that if T is an automorphism, then a set S is invariant iff $T(S) = S$. □

Proposition 5.1.5. Suppose that $(\Omega, \mathcal{S})$ is a measurable space and

$$T : (\Omega, \mathcal{S}) \rightarrow (\Omega, \mathcal{S})$$

is a measurable map. Then the following hold.

- (i) The collection $\mathcal{I} = \mathcal{I}_T$ of *T-invariant measurable sets* is a sigma-subalgebra of $\mathcal{S}$.
- (ii) An $\mathcal{S}$ -measurable function $f : \Omega \rightarrow \mathbb{R}$ is *T-invariant* if and only if it is $\mathcal{I}_T$ -measurable.

Proof. (i) Thus follows from the fact that $S \in \mathcal{I}_T$ if and only if $S = T^{-1}(S)$.

(ii) Suppose that f is T -invariant. Then for any $x \in \mathbb{R}$ the set $S = \{f \leq x\}$ is T -invariant since $\mathbf{I}_S \circ T = \mathbf{I}_{\{f \circ T \leq x\}} = \mathbf{I}_S$.

Conversely, if f is $\mathcal{I}_T$ -measurable, then $f^{-1}(\{y\}) \in \mathcal{I}_T, \forall y \in \mathbb{R}$ and

$$(f \circ T)^{-1}(\{y\}) = T^{-1}(f^{-1}(\{y\})) = f^{-1}(\{y\}).$$

If $f(x) = y$, then $x \in f^{-1}(\{y\}) = (f \circ T)^{-1}(\{y\})$ so that $f \circ T(x) = y = f(x)$. $\square$

Remark 5.1.6. Consider the path space $\mathbb{U}_{\mathbb{X}} = \mathbb{X}^{\mathbb{N}}$. We have the *tail* sigma-subalgebra

$$\mathcal{T}_{\infty} = \bigcap_{m \geq 1} \mathcal{T}_m, \quad \mathcal{T}_m = \sigma(U_m, U_{m+1}, \dots).$$

For $\underline{x} = (x_1, x_1, \dots) \in \mathbb{U}_{\mathbb{X}}$ and $x_0 \in \mathbb{X}$ we define the concatenation

$$x_0 * \underline{x} = (x_0, x_1, \dots).$$

Then S is Θ -invariant iff

$$\underline{x} \in S \iff x_0 * \underline{x} \in S, \quad \forall x_0 \in \mathbb{X},$$

i.e.,

$$S = \mathbb{X} * S = \{x_0 * \underline{x}; \quad x_0 \in \mathbb{X}, \quad \underline{x} \in S\} \subset \mathcal{T}_2.$$

Iterating we deduce that

$$S = \mathbb{X}^m * S = \mathbb{X} * (\mathbb{X}^{m-1} * S) \in \mathcal{T}_{m+1}, \quad \forall m \geq 0.$$

Hence $S \in \mathcal{T}_{\infty}$. Hence, in the universal case $\mathcal{I} = \mathcal{I}_{\Theta} \subset \mathcal{T}_{\infty}$.

Fix an element $x_* \in \mathbb{X}$ and consider the set

$$S_{x_*} = \{\underline{x} \in \mathbb{X}^{\mathbb{N}}; \quad \exists N \quad \forall n > N \quad x_n = x_*\}.$$

Thus $\underline{x} \in S_{x_*}$ iff $\underline{x}$ eventually coincides with the constant sequence $(x_*, x_*, \dots)$. Thus S_{x_*} is a tail event.

Observe that the sigma-algebras $\mathcal{I}_{\Theta}$ and $\mathcal{T}_{\infty}$ do not depend on any choice of invariant probability measure on $\mathbb{U}_{\mathbb{X}}$. $\square$

Definition 5.1.7. Suppose that $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\Omega, \mathcal{S}, \mathbb{P})$ is a measure preserving map. A measurable function $f : (\Omega, \mathcal{S}) \rightarrow \mathbb{R}$ is said to be *quasi-invariant* if

$$f = f \circ T \quad \mathbb{P} - \text{a.s.}$$

A subset $S \in \mathcal{S}$ is said to be *quasi-invariant* if $\mathbf{I}_S$ is quasi-invariant, i.e.,

$$\mathbb{P}[S \Delta T^{-1}S] = 0,$$

where $A \Delta B := (A \setminus B) \cup (B \setminus A)$ is the symmetric difference of two sets. We denote by $\mathcal{J} = \mathcal{J}_T$ the collection of T -quasi-invariant sets. $\square$

Proposition 5.1.8. Suppose that $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\Omega, \mathcal{S}, \mathbb{P})$ is a measure preserving map. Then the following hold.

(i) The collection $\mathcal{J}_T$ of quasi-invariant sets is a sigma-algebra.

(ii) The $\mathbb{P}$ -completions of $\mathcal{I}$ and $\mathcal{J}$ coincide, i.e., for any $S \in \mathcal{J}$ there exists $S' \in \mathcal{I}$ such that $\mathbb{P}[S \Delta S'] = 0$.

(iii) A measurable function is T -quasi-invariant if and only if it is $\mathcal{I}_T$ -measurable

Proof. (i) The fact that $\mathcal{I}_T$ is a sigma-algebra follows immediately from the definition of a quasi-invariant.

(ii) Denote by $\bar{\mathcal{I}}$ the $\mathbb{P}$ -completion of $\mathcal{I}$. Let $\bar{S} \in \bar{\mathcal{I}}$. There exists $S \in \mathcal{I}$ such that $\mathbb{P}[\bar{S} \Delta S] = 0$. Since T is measure preserving we deduce

$$0 = \mathbb{P}[T^{-1}(\bar{S} \Delta S)] = \mathbb{P}[T^{-1}(\bar{S}) \Delta S]$$

and thus

$$\begin{aligned} \mathbb{P}[T^{-1}(\bar{S}) \Delta \bar{S}] &= \mathbb{E}[|I_{T^{-1}(\bar{S})} - I_{\bar{S}}|] \\ &\leq \mathbb{E}[|I_{T^{-1}(\bar{S})} - I_S|] + \mathbb{E}[|I_S - I_{\bar{S}}|] = 0. \end{aligned}$$

Conversely, if $S \in \mathcal{I}$ define

$$\bar{S} := \bigcap_{n \in \mathbb{N}} S_n, \quad S_n := \bigcup_{k \geq n} T^{-k}(S).$$

Note that $\bar{S} = T^{-1}(\bar{S})$ so that $\bar{S}$ is invariant. Since $S_1 \supset S_2 \supset \dots$, we have

$$I_{\bar{S}} = \lim_{n \rightarrow \infty} I_{S_n}.$$

On the other hand

$$I_{S_n} = \sup_{k \geq n} I_{T^{-k}(S)} = \sup_{k \geq n} I_S \circ T^k = I_S \text{ a.s.}$$

since S is quasi-invariant and thus $I_S \circ T^k = I_S$ a.s.. Hence $I_{\bar{S}} = I_S$ a.s., so that $S \in \bar{\mathcal{I}}$.

(iii) Clearly, if f is quasi-invariant, then so are the sublevel sets $\{f \leq x\}$, $\forall x \in \mathbb{R}$ and thus f is $\mathcal{I}_T$ -measurable.

Conversely if f is $\mathcal{I}_T$ -measurable, then so are $f^\pm$ and it suffices to show that if $f \geq 0$ is $\mathcal{I}$ -measurable, then f is quasi-invariant. Clearly any $\mathcal{I}$ -measurable elementary function is quasi-invariant. Since f is an increasing limit of $\mathcal{I}$ -measurable elementary functions, it is therefore an increasing limit of quasi-invariant elementary functions and thus it is quasi-invariant. $\square$

Example 5.1.9. Suppose that $\vec{X} = (X_n)_{n \in \mathbb{N}}$ is an exchangeable sequence of random variables. We identify them with the coordinate maps on the universal space $\mathbb{U} := \mathbb{R}^{\mathbb{N}}$. Let $\mathbb{P} \in \text{Prob}(\mathbb{U}, \mathcal{U})$ be their distribution.

The probability measure $\mathbb{P}$ is shift invariant. Moreover, we have three remarkable sigma-subalgebras: the sigma-subalgebra $\mathcal{I}_\Theta$ of shift-invariant events, the tail sigma-subalgebra $\mathcal{I}_\infty$ and the sigma-subalgebra of $\mathcal{S}_\infty$ of exchangeable events. Moreover we have inclusions

$$\mathcal{I}_\Theta \subset \mathcal{I}_\infty \subset \mathcal{S}_\infty.$$

Denote by $\tilde{\mathcal{I}}_\Theta$, $\tilde{\mathcal{I}}_\infty$ and $\tilde{\mathcal{S}}_\infty$ the $\mathbb{P}$ -completions of $\mathcal{I}_\Theta$, $\mathcal{I}_\infty$ and respectively $\mathcal{S}_\infty$. One can show that

$$\tilde{\mathcal{I}}_\Theta = \tilde{\mathcal{I}}_\infty = \tilde{\mathcal{S}}_\infty. \quad (5.1.4)$$

For a proof I refer to [163, Thm. (1)]. In particular the above equality implies Theorem 3.2.62. The proof in [163] of the above equality *does not rely* on the Law of Large numbers for exchangeable sequences as was the case in our proof of Theorem 3.2.62. $\square$

Definition 5.1.10. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be probability space. A measure preserving map $T : \Omega \rightarrow \Omega$ is said to be *ergodic* if any T -invariant set has measure 0 or 1, i.e., the sigma-algebra of invariant sets is a zero-one algebra. $\square$

From Proposition 5.1.8 we deduce the following equivalent characterization of ergodicity.

Proposition 5.1.11. *The map T is ergodic if and only if any $\mathbb{P}$ -quasi-invariant set is a zero-one event, i.e., has measure 0 or 1.* $\square$

Remark 5.1.12. Suppose that T is an ergodic automorphism of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. If $S \in \mathcal{S}$, then the set

$$\widehat{S} = \bigcup_{n \geq 0} T^n(S)$$

is quasi-invariant. Indeed $\widehat{S} \subset T^{-1}(\widehat{S})$ and $\mathbb{P}[T^{-1}(\widehat{S})] = \mathbb{P}[\widehat{S}]$. If $\mathbb{P}[S] > 0$, then $\mathbb{P}[\widehat{S}] > 0$ and the ergodicity of T implies that

$$\mathbb{P}[\Omega \setminus \widehat{S}] = 0.$$

The set $\widehat{S}$ is a union of orbits $\mathcal{O}_T(\omega) = \{T^n(\omega)\}_{n \geq 0}$ of the dynamical system on Ω determined by the iterates of T . The ergodicity shows that the orbits originating in a set S of positive measure reach almost any point in Ω ; the unreachable ones form a negligible set. This shows that the dynamics of an ergodic automorphism is quite chaotic: orbits want to fill the space. $\square$

Definition 5.1.13. Suppose that $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{F})$, $n \in \mathbb{N}$ is a sequence of measurable maps. We say that $(X_n)_{n \in \mathbb{N}}$ is a *Kolmogorov sequence* if its tail algebra

$$\mathcal{T}_\infty := \bigcap_{m \in \mathbb{N}} \mathcal{T}_m, \quad \mathcal{T}_m := \sigma(X_n, n \geq m)$$

is a zero-one algebra. $\square$

If $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. random variables, then Kolmogorov's zero-one theorem shows that it is a Kolmogorov sequence.

Proposition 5.1.14. *If $(X_n)_{n \in \mathbb{N}}$ is a stationary Kolmogorov sequence, then the shift map Θ on the associated path space $\mathbb{X}^{\mathbb{N}}$ is ergodic.*

Proof. As shown in Remark 5.1.6 the sigma-algebra $\mathcal{I}$ of Θ -invariant sets is contained in the tail algebra. For a Kolmogorov sequence the tail algebra is by definition, a zero-one. $\square$

Example 5.1.15. Consider the map $Q : [0, 1) \rightarrow [0, 1)$ discussed in Example 5.1.1(iii). The interval $[0, 1)$ embeds in $\{0, 1\}^{\mathbb{N}}$

$$[0, 1) \ni x = \sum_{n=1}^{\infty} \frac{\epsilon_n}{2^n} \mapsto (\epsilon_1, \epsilon_2, \dots) \in \{0, 1\}^{\mathbb{N}}.$$

The image of the map is a shift-invariant subset of $\{0, 1\}^{\mathbb{N}}$. Its complement is negligible with respect to the product measure on $\{0, 1\}^{\mathbb{N}}$ and the restriction of the product measure on the image of this embedding coincides with the Lebesgue measure; see Exercise 1.6 (vii). The space $\{0, 1\}^{\mathbb{N}}$ equipped with the product measure is the path space corresponding to an i.i.d. sequence of Bernoulli random

variables with success probability $\frac{1}{2}$. Hence the shift map is ergodic, proving that the map Q is also ergodic. $\square$

We have the following characterization of Kolmogorov sequences due to Blackwell and Freedman [20].

Theorem 5.1.16. *Suppose that $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{F})$, $n \in \mathbb{N}$ is a sequence of measurable maps. The following are equivalent.*

- (i) *The sequence is a Kolmogorov sequence.*
- (ii) *For any $A \in \mathcal{S}$*

$$\lim_{m \rightarrow \infty} \sup_{B \in \mathcal{T}_m} |\mathbb{P}[A \cap B] - \mathbb{P}[A]\mathbb{P}[B]| = 0, \quad (5.1.5)$$

where we recall that $\mathcal{T}_m = \sigma(X_n, n \geq m)$.

Proof. (i) $\Rightarrow$ (ii) Then for any $B \in \mathcal{T}_\infty$, $A \in \mathcal{F}$ we have

$$\begin{aligned} |\mathbb{P}[A \cap B] - \mathbb{P}[A]\mathbb{P}[B]| &= |\mathbb{E}[\mathbf{I}_A \mathbf{I}_B] - \mathbb{E}[\mathbf{I}_A]\mathbb{E}[\mathbf{I}_B]| \\ &= \left| \int_B (\mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_m] - \mathbb{P}[A]) \right| \leq \int_\Omega |\mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_m] - \mathbb{P}[A]|. \end{aligned}$$

Hence

$$\sup_{B \in \mathcal{T}_m} |\mathbb{P}[A \cap B] - \mathbb{P}[A]\mathbb{P}[B]| \leq \int_\Omega |\mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_m] - \mathbb{P}[A]|. \quad (5.1.6)$$

The Backwards Martingale Convergence Theorem implies that

$$\mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_m] \rightarrow \mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_\infty] \text{ a.s. and } L^1.$$

Since $\mathcal{T}_\infty$ is a zero-one algebra, we deduce that

$$\lim_{m \rightarrow \infty} \mathbb{E}[\mathbf{I}_A \mid \mathcal{T}_m] = \mathbb{E}[\mathbf{I}_A] = \mathbb{P}[A].$$

Using this in (5.1.6) we obtain (5.1.5).

(ii) $\Rightarrow$ (i) Let $A \in \mathcal{T}_\infty$. Then, $\forall m$, $A \in \mathcal{T}_m$ and thus

$$\begin{aligned} 0 &\leq \mathbb{P}[A] - \mathbb{P}[A]^2 = |\mathbb{P}[A \cap A] - \mathbb{P}[A]\mathbb{P}[A]| \\ &\leq \sup_{B \in \mathcal{T}_m} |\mathbb{P}[A \cap B] - \mathbb{P}[A]\mathbb{P}[B]| \rightarrow 0. \end{aligned}$$

Hence $\mathbb{P}[A] = \mathbb{P}[A]^2$ so that $\mathbb{P}[A] \in \{0, 1\}$ so that $\mathcal{T}_\infty$ is a zero-one algebra. $\square$

Example 5.1.17. Suppose that X is a compact metric space and $T : X \rightarrow X$ is a continuous map. Denote by $\text{Prob}_T(X)$ the set of T -invariant Borel probability measures on X . As mentioned in Example 5.1.1, $\text{Prob}_T(X)$ is a nonempty, convex subset of $\text{Prob}(X)$ that it is closed with respect to the weak topology. A measure $\mu \in \text{Prob}_T(X)$ is ergodic iff it is an extreme point of $\text{Prob}_T(X)$; see e.g. [102, Prop.8.5]. Krein-Milman Theorem [194, Thm.8.14] guarantees that such extreme points exist. $\square$

5.1.2. Ergodic theorems. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be probability space and suppose that $T : \Omega \rightarrow \Omega$ is measure preserving map. For any measurable function $f : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ we denote by $\hat{T}f$ its pullback by T

$$\hat{T}f := f \circ T.$$

This is a measurable function and, since T is measure preserving, we deduce from the change-in-variables formula (Theorem 1.2.54) that

$$\int_{\Omega} \hat{T}f d\mathbb{P} = \int_{\Omega} f dT_{\#}\mathbb{P} = \int_{\Omega} f d\mathbb{P}, \quad \forall f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P}).$$

Note also that

$$(\hat{T}f)^p = \hat{T}f^p \quad \forall f \in \mathcal{L}_+^0(\Omega, \mathcal{S}), \quad p \geq 1.$$

We will denote by $\| \cdot \|_p$ the norm of $L^p(\Omega, \mathcal{S}, \mathbb{P})$. We deduce that $\hat{T}$ defines isometries

$$\hat{T} : L^p(\Omega, \mathcal{S}, \mathbb{P}) \rightarrow L^p(\Omega, \mathcal{S}, \mathbb{P}), \quad \forall p \geq 1.$$

The operator $\hat{T}$ is referred to as the *Koopman operator*.

We denote by $\mathcal{J} = \mathcal{J}_T$ the σ -subalgebra of quasi-invariant measurable subsets. Thus $S \in \mathcal{J}$ if and only if $\hat{T}\mathbf{I}_S = \mathbf{I}_S$, $\mathbb{P}$ -a.s..

More generally, if $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ and $S \in \mathcal{J}$, then

$$\hat{T}\mathbf{I}_S \in L^1(\Omega, \mathcal{S}, \mathbb{P}) \quad \text{and} \quad \hat{T}(f\mathbf{I}_S) = (\hat{T}f) \cdot (\hat{T}\mathbf{I}_S) = (\hat{T}f) \cdot \mathbf{I}_S,$$

so that

$$\int_{\Omega} f\mathbf{I}_S d\mathbb{P} = \int_{\Omega} \hat{T}(f\mathbf{I}_S) d\mathbb{P} = \int_{\Omega} (\hat{T}f) \cdot \mathbf{I}_S d\mathbb{P}, \quad \forall S \in \mathcal{J}, \quad f \in L^1(\Omega, \mathcal{S}, \mathbb{P}). \quad (5.1.7)$$

For each $p \geq 1$ we set

$$\mathcal{Q}_{T,p} := \{ f \in L^p(\Omega, \mathcal{S}, \mathbb{P}); \quad \hat{T}f = f \} \quad (5.1.8)$$

In other words, $\mathcal{Q}_{T,p}$ consists of quasi-invariant L^p -functions, i.e.,

$$\mathcal{Q}_{T,p} = L^p(\Omega, \mathcal{J}, \mathbb{P}) = L^p(\Omega, \mathcal{J}, \mathbb{P}).$$

We set

$$\mathcal{Q}_T := \mathcal{Q}_{T,2} = L^2(\Omega, \mathcal{J}, \mathbb{P}) \quad (5.1.9)$$

and we denote by P_T the orthogonal projection onto $\mathcal{Q}_T$. In the proof of Theorem 1.4.8 we have shown that

$$P_T f = \mathbb{E}[f | \mathcal{J}].$$

The space $\mathcal{Q}_T$ contains the constant functions so $\dim \mathcal{Q}_T \geq 1$.

Proposition 5.1.18. *Suppose that $T : (\Omega, \mathcal{J}, \mathbb{P}) \rightarrow (\Omega, \mathcal{J}, \mathbb{P})$ is a measure preserving map. Then the following statements are equivalent.*

- (i) *The map T is ergodic.*
- (ii) *For any $p \geq 1$ $\dim \mathcal{Q}_{T,p} = 1$*
- (iii) *There exists $p \geq 1$ such that $\dim \mathcal{Q}_T^p = 1$.*

Proof. (i) $\Rightarrow$ (ii) Assume that T is ergodic so $\mathcal{J}$ is a zero-one sigma-subalgebra. Hence, any $\mathcal{J}$ -measurable elementary function is constant. Hence any L^p function must be a.s.-constant as a limit of elementary functions.

Clearly (ii) $\Rightarrow$ (iii). To prove the implication (iii) $\Rightarrow$ (i) note any $\mathcal{J}$ -measurable function belongs to any L^p and thus must be a.s. constant. $\square$

To summarize

$$\boxed{T \text{ is ergodic} \iff \dim \mathcal{Q}_T = 1.} \quad (5.1.10)$$

For each n we denote by A_n the n -th temporal average/mean operator

$$f \mapsto A_n f = \frac{1}{n} (1 + \hat{T} + \hat{T}^2 + \dots + \hat{T}^{n-1}) f.$$

Note that A_n defines linear operators

$$A_n : L^p(\Omega, \mathcal{S}, \mathbb{P}) \rightarrow L^p(\Omega, \mathcal{S}, \mathbb{P}), \quad p \geq 1$$

satisfying

$$\|A_n f\|_p \leq \|f\|_p, \quad \forall f \in L^p. \quad (5.1.11)$$

Remark 5.1.19. Let me briefly explain the intuition of the temporal averages $A_n(f)$. Think of Ω as the space of states of a physical system that evolves in discrete time. Thus, if the system was initially in the state ω , it will be in the state $T^n(\omega)$ after n units of time.

A function $f : \Omega \rightarrow \mathbb{R}$ can be viewed as a macroscopic quantity that associates to each state ω a measurable numerical quantity $f(\omega)$. Note that for each $n \in \mathbb{N}$ and each $\omega \in \Omega$ we have

$$(A_{n+1}f)(\omega) = \frac{f(\omega) + f(T\omega) + \dots + f(T^n\omega)}{n+1}$$

is the average value of the macroscopic quantity f as the system evolves for n units of time. $\square$

We have the following *mean ergodic theorem* due to John von Neumann,

Theorem 5.1.20 (L^2 -Mean ergodic theorem). *Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $T : \Omega \rightarrow \Omega$ is a measure preserving map. Then, $\forall f \in L^2(\Omega, \mathcal{S}, \mathbb{P})$, the temporal averages $A_n f$ converge in L^2 to the orthogonal projection of f onto the space $\mathcal{Q}_T$ of quasi-invariant functions, i.e.,*

$$\frac{1}{n} (1 + \hat{T} + \hat{T}^2 + \dots + \hat{T}^{n-1}) f \rightarrow P_T f = \mathbb{E}[f | \mathcal{J}].$$

In particular, if T is ergodic we have

$$\frac{1}{n} (1 + \hat{T} + \hat{T}^2 + \dots + \hat{T}^{n-1}) f \rightarrow \mathbb{E}[f] \mathbf{I}_\Omega \text{ in } L^2.$$

Proof. Denote by $\mathfrak{X}_2$ the collection of functions $f \in L^2(\Omega, \mathcal{S}, \mathbb{P})$ such that $A_n f$ converges in L^2 to some function $A_\infty f$. Clearly $\mathfrak{X}_2$ is a vector space. We will gradually show that $\mathfrak{X}_2 = L^2(\Omega, \mathcal{S}, \mathbb{P})$ and $A_\infty = P_T$.

1. $\mathcal{Q}_T \subset \mathfrak{X}_2$ and $A_\infty f = f, \forall f \in \mathcal{Q}_T$.

Indeed $A_n f = f, \forall f \in \mathcal{Q}_T$.

2. $\forall f \in \mathfrak{X}_2$, we have $\hat{T}f \in \mathfrak{X}_2$ and $A_\infty f \in \mathcal{Q}_T$.

Let $f \in \mathfrak{X}_2$. Note first that $\widehat{T}$ commutes with A_n . Since $\widehat{T}$ is continuous we deduce

$$\lim_{n \rightarrow \infty} A_n \widehat{T} f = \lim_{n \rightarrow \infty} \widehat{T} A_n f = \widehat{T} A_\infty f,$$

i.e., $\widehat{T} f \in \mathfrak{X}_2$ and $A_\infty \widehat{T} f = \widehat{T} A_\infty f$.

On the other hand,

$$n A_n \widehat{T} f = (n+1) A_{n+1} f - f \implies A_n \widehat{T} f = \frac{n+1}{n} A_{n+1} f - \frac{1}{n} f,$$

so that

$$\widehat{T} A_\infty f = A_\infty \widehat{T} f = \lim_{n \rightarrow \infty} \frac{n+1}{n} A_{n+1} f = A_\infty f.$$

Hence $A_\infty f \in \mathcal{Q}_T$.

3. $\widehat{T} f - f \in \mathfrak{X}_2, \forall f \in L^2(\Omega, \mathcal{S}, \mathbb{P})$.

Indeed

$$A_n(\widehat{T} f - f) = \frac{1}{n}(\widehat{T}^n f - f)$$

and, since $\widehat{T}$ is unitary, we deduce that

$$\|A_n(\widehat{T} f - f)\|_2 \leq \frac{1}{n}(\|\widehat{T} f\|_2 + \|f\|_2) = \frac{2}{n}\|f\|_2 \rightarrow 0.$$

4. $\forall f \in L^2(\Omega, \mathcal{S}, \mathbb{P}), \forall k \in \mathbb{N}$ we have $\widehat{T}^k f - f \in \mathcal{Q}_T^\perp$.

We first prove the claim for $k = 1$. Indeed, for any $g \in \mathcal{Q}_T$ we have $\widehat{T} g = g$ and

$$(\widehat{T} f - f, g) = (\widehat{T} f, g) - (f, g) = (\widehat{T} f, \widehat{T} g) - (f, g) = 0,$$

where at the last step we used the fact that $\widehat{T}$ is unitary. In general

$$\widehat{T}^k f - f = \sum_{j=1}^k (\widehat{T}^j f - \widehat{T}^{j-1} f) = \sum_{j=1}^k (\widehat{T} f_j - f_j), \quad f_j := (\widehat{T}^{j-1} f),$$

and $\widehat{T} f_j - f_j \in \mathcal{Q}_T^\perp, \forall j$.

5. $A_\infty f = P_T f, \forall f \in \mathfrak{X}_2$.

We have

$$f - A_n f = \frac{1}{n} \sum_{k=1}^{n-1} (f - \widehat{T}^k f) \in \mathcal{Q}_T^\perp.$$

Letting $n \rightarrow \infty$ and using the fact that $\mathcal{Q}_T^\perp$ is a closed subspace of L^2 we deduce from **4** $f - A_\infty f \in \mathcal{Q}_T^\perp$ so that $A_\infty f = P_T f$.

6. $\mathfrak{X}_2$ is closed.

Let $(f_k)_{k \in \mathbb{N}}$ be a sequence in $\mathfrak{X}_2$ that converges in L^2 to f . To show that $f \in \mathfrak{X}_2$ we will show that the sequence $A_n f$ is Cauchy. Fix $\varepsilon > 0$. We have

$$\|A_n f - A_m f\|_2 \leq \|A_n f - A_n f_k\|_2 + \|A_n f_k - A_m f_k\|_2 + \|A_m f_k - A_m f\|_2$$

(use (5.1.11), i.e., $\|A_n\| \leq 1$ as operator $L^2 \rightarrow L^2$)

$$\leq \|f - f_k\|_2 + \|A_n f_k - A_m f_k\|_2 + \|f - f_k\|_2.$$

Hence

$$\|A_n f - A_m f\|_2 \leq 2\|f - f_k\|_2 + \|A_n f_k - A_m f_k\|_2, \quad \forall k, m, n.$$

Fix k such that

$$\|f - f_k\|_2 < \frac{\varepsilon}{3}.$$

The sequence $(A_n f_k)_{n \in \mathbb{N}}$ is convergent since $f_k \in \mathfrak{X}_2$. It is thus Cauchy, so there exists $N = N(\varepsilon, k)$ such that $\forall m, n > N$

$$\|A_n f_k - A_m f_k\|_2 < \frac{\varepsilon}{3}.$$

Hence

$$\|A_n f - A_m f\|_2 \leq \varepsilon, \quad \forall m, n > N(\varepsilon, k).$$

7. $\mathfrak{X}_2 = L^2(\Omega, \mathcal{S}, \mathbb{P})$.

We know that $\mathcal{Q}_T \subset \mathfrak{X}_2$ and

$$\text{Range}(\hat{T} - 1) \subset \mathcal{Q}_T^\perp \cap \mathfrak{X}_2.$$

At this point we invoke a classical result of functional analysis: if $S : H \rightarrow H$ is a bounded linear operator on a Hilbert space, then the closure of the range of S is $(\ker S^*)^\perp$; see e.g. [29, Cor. 2.18].

The operator $\hat{T}$ is unitary, so that $\hat{T}^* = \hat{T}^{-1}$. Hence if we let $S = \hat{T} - 1$, then

$$S^* = \hat{T}^* - 1 = \hat{T}^{-1} - 1,$$

since $\hat{T}$ is unitary. We deduce

$$\text{closure}(\text{Range}(\hat{T} - 1)) = (\ker(\hat{T}^{-1} - 1))^\perp = \mathcal{Q}_T^\perp.$$

Since $\mathfrak{X}_2$ is closed we deduce $\mathcal{Q}_T^\perp \subset \mathfrak{X}_2$. Hence $\mathcal{Q}_T \oplus \mathcal{Q}_T^\perp \subset \mathfrak{X}_2$. This completes the proof of Theorem 5.1.20. $\square$

Corollary 5.1.21. *Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $T : \Omega \rightarrow \Omega$ is a measure preserving map. Then, $\forall f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ the temporal averages $A_n f$ converge in L^1 to $\mathbb{E}[f \mid \mathcal{J}]$, i.e.,*

$$\frac{1}{n}(1 + \hat{T} + \hat{T}^2 + \dots + \hat{T}^{n-1})f \rightarrow \mathbb{E}[f \mid \mathcal{J}] \text{ in } L^1 \text{ as } n \rightarrow \infty.$$

Proof. Denote by $\mathfrak{X}_1$ the collection of functions $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that $A_n f$ converges in L^1 to some function $A_\infty f$. Since $\|\cdot\|_1 \leq \|\cdot\|_2$ we deduce that $\mathfrak{X}_2 \subset \mathfrak{X}_1$. The argument in Step 6. in the proof of Theorem 5.1.20 extends without change to the L^1 since, according to (5.1.11), the operators A_n are contractions $A_n : L^1 \rightarrow L^1$. This proves that $\mathfrak{X}_1$ is a closed subspace of L^1 that contains L^2 and thus $\mathfrak{X}_1 = L^1$.

From (5.1.7) we deduce that for any $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$, and any $S \in \mathcal{J}$ we have

$$\int_\Omega f \mathbf{I}_S d\mathbb{P} = \int_\Omega (A_n f) \mathbf{I}_S d\mathbb{P}.$$

Letting $n \rightarrow \infty$ we deduce

$$\int_\Omega f \mathbf{I}_S d\mathbb{P} = \int_\Omega (A_\infty f) \mathbf{I}_S d\mathbb{P}, \quad \forall S \in \mathcal{J}.$$

Hence (see Definition 1.4.3) $A_\infty f = \mathbb{E}[f \mid \mathcal{J}]$, $\square$

We can now formulate and prove *Birkhoff's ergodic theorem*

Theorem 5.1.22 (Birkhoff's ergodic theorem). *Let $(\Omega, \mathcal{S}, \mathbb{P})$ be probability space. Suppose that $T : \Omega \rightarrow \Omega$ a measure preserving map. If $f \in \mathcal{L}^1(\Omega, \mathcal{S}, \mathbb{P})$, then the temporal averages*

$$A_n(f) = \frac{1}{n+1} \left(f + f \circ T + \cdots + f \circ T^n \right) = \frac{1}{n+1} \left(f + \widehat{T}f + \cdots + \widehat{T}^n f \right)$$

converge a.s. to $\mathbb{E}[f \mid \mathcal{I}]$.

Proof. Denote by $\mathfrak{X}_0$ the set of functions $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that $A_n f$ converges a.s. to a function $A_\infty f$. Corollary 5.1.21 shows that in this case $A_\infty f = \mathbb{E}[f \mid \mathcal{I}] \in L^1$. Clearly $\mathfrak{X}_0$ is a vector subspace of $L^1(\Omega, \mathcal{S}, \mathbb{P})$.

We will show that $\mathfrak{X}_0 = L^1(\Omega, \mathcal{S}, \mathbb{P})$ in two steps.

- (i) The set $\mathfrak{X}_0$ is a closed subspace of $L^1(\Omega, \mathcal{S}, \mathbb{P})$.
- (ii) $\widehat{T}f - f \in \mathfrak{X}_0, \forall f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$.

The claim (i) is the difficult one. Assuming temporarily its validity, I will show how it implies (ii) and the conclusion of the theorem.

Proof of (ii) assuming (i). Observe that for any $f \in L^1$ we have

$$A_n(\widehat{T}f - f) = \frac{1}{n} (\widehat{T}^n f - f)$$

In particular, if $f \in L^\infty$ we deduce

$$\|A_n(\widehat{T}f - f)\|_\infty \leq \frac{2}{n} \|f\|_\infty$$

so $A_n(\widehat{T}f - f) \rightarrow 0$ a.s. so $(\widehat{T}f - f) \in \mathfrak{X}_0$ if $f \in L^\infty$.

Suppose now that $f \in L^1$ then $f = f_+ - f_-$ and $(\widehat{T}f)_\pm = \widehat{T}f_\pm$. Thus it suffices to show that $\widehat{T}f - f \in \mathfrak{X}_0$ if $f \in L^1$ and $f \geq 0$ a.s..

In this case we can find a sequence of elementary functions f_n such that $f_n \nearrow f$. Hence

$$\widehat{T}f_n - f_n \rightarrow \widehat{T}f - f \text{ in } L^1.$$

Since the functions f_n are bounded, so are the functions $\widehat{T}f_n - f_n$ we deduce that $\widehat{T}f_n - f_n \in \mathfrak{X}_0$. We know from (i) that $\mathfrak{X}_0$ is L^1 -closed. This proves (ii).

From (ii) we deduce that $\widehat{T}f - f \in \mathfrak{X}_0, \forall f \in L^2 \subset L^1$. Since $\mathfrak{X}_0$ is closed in L^1 we deduce from the proof of Theorem 5.1.20 that

$$\text{closure}_{L^2}(\text{range}(\widehat{T} - 1)) \subset \text{closure}_{L^1}(\text{range}(\widehat{T} - 1)) \subset \mathfrak{X}_0.$$

On the other hand, $Q_{T,2} := Q_T \cap L^2 \subset \mathfrak{X}_0$, so that

$$L^2 = \mathfrak{X}_2 = Q_{T,2} + Q_{T,2}^\perp = Q_{T,2} + \text{closure}_{L^2}(\text{range}(\widehat{T} - 1)) \in \mathfrak{X}_0.$$

Since L^2 is dense in L^1 , and $\mathfrak{X}_0$ is closed in L^1 we conclude that $\mathfrak{X}_0 = L^1$.

Proof of (i) The proof of this result is based on a technical inequality similar in spirit to Doob's maximal inequality (3.2.36). For $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ we define $\mathbb{M}[f] \in \mathcal{L}^0(\Omega, \mathcal{S})$,

$$\mathbb{M}[f](\omega) := \sup_{n \geq 1} A_n f(\omega) = \sup_{n \geq 1} \frac{1}{n} \left(f(\omega) + f(T\omega) + \cdots + f(T^{n-1}\omega) \right).$$

Lemma 5.1.23 (Maximal Ergodic Lemma). $\forall \lambda > 0, \forall f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$

$$\forall \lambda > 0, f \in L^1(\Omega, \mathcal{S}, \mathbb{P}) : \lambda \mathbb{P}[\{\mathbb{M}[|f|] > \lambda\}] \leq \|f\|_1. \quad (5.1.12)$$

□

Lemma 5.1.24. *The Maximal Ergodic Lemma implies the claim (i), i.e., the set $\mathfrak{X}_0$ of functions $f \in L^1(\Omega, \mathcal{S}, \mathbb{P})$ such that $A_n f$ converges a.s. is closed in L^1 .*

Proof. Suppose that the sequence (f_k) in $\mathfrak{X}_0$ converges in L^1 to a function f . We want to show that the sequence $A_n(f)$ is a.s. Cauchy, i.e.,

$$\mathcal{C}[f] := \limsup_{m, n \rightarrow \infty} |A_n(f) - A_m(f)| = \lim_{N \rightarrow \infty} \sup_{m, n \geq N} |A_n(f) - A_m(f)| = 0 \text{ a.s..}$$

We have $\mathcal{C}[f_k] = 0$ a.s., for all k . Hence

$$\mathcal{C}[f] = \mathcal{C}[f] - \mathcal{C}[f_k] \leq \mathcal{C}[f - f_k] \leq 2\mathbb{M}[|f - f_k|].$$

Hence, for any $\varepsilon > 0$ we have

$$\mathbb{P}[\mathcal{C}[f] \geq \varepsilon] \leq \mathbb{P}[2\mathbb{M}[|f - f_k|] > \varepsilon] \leq \frac{2\|f - f_k\|_1}{\varepsilon}$$

Letting $k \rightarrow \infty$ we deduce

$$\mathbb{P}[\mathcal{C}[f] \geq \varepsilon] = 0, \quad \forall \varepsilon > 0$$

so $\mathcal{C}[f] = 0$, a.s. □

Proof of the Maximal Ergodic Lemma. Let us observe that the inequality (5.1.12) follows from

$$\int_{\{\mathbb{M}[g] > 0\}} g d\mathbb{P} \geq 0 \quad \forall g \in L^1(\Omega, \mathcal{S}, \mathbb{P}). \quad (5.1.13)$$

Indeed, if in (5.1.13) we let $g = f - \lambda$, $\lambda > 0$, then

$$\|f\|_1 \geq \int_{\{\mathbb{M}[f] > \lambda\}} f d\mathbb{P} \stackrel{(5.1.13)}{\geq} \lambda \mathbb{P}[\{\mathbb{M}[f] > \lambda\}].$$

We will present two proofs of (5.1.13).

The first proof, due to F. Riesz, is a bit longer but a more intuitive. The second proof, due to A. Garsia [84] is a lot shorter, but less intuitive.

Set

$$X := \{\mathbb{M}[g] > 0\} \subset \Omega.$$

Define

$$S_n(g) := \sum_{j=0}^{n-1} g \circ T^j, \quad \mathbb{M}_n[g] := \max_{1 \leq k \leq n} S_k(g), \quad (5.1.14)$$

$$X_k := \{\mathbb{M}_k[g] > 0\}.$$

First proof of (5.1.13). Note that $X_k \subset X_{k+1}$ and

$$\int_X g d\mathbb{P} = \lim_{n \rightarrow \infty} \int_{X_n} g d\mathbb{P} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \int_{X_k} g d\mathbb{P}.$$

At the last step we used the fact that the Cèsaro means of a convergent sequence have the same limit as the sequence; see Exercise 2.6 with $p_{n,k} = \frac{1}{n}$. Thus, it suffices to show that

$$\sum_{k=1}^n \int_{X_k} g d\mathbb{P} \geq 0, \quad \forall n \geq 0. \quad (5.1.15)$$

Fix n . We have

$$\sum_{k=1}^n \int_{X_k} g d\mathbb{P} = \sum_{j=0}^{n-1} \int_{X_{n-j}} g d\mathbb{P} = \sum_{j=0}^{n-1} \int_{T^{-j}(X_{n-j})} g \circ T^j d\mathbb{P},$$

where at the last step we used the change-in-variables formula (1.2.25) and the fact that T is measure preserving. We set $Y_j := T^{-j}(X_{n-j})$. Hence

$$\sum_{k=1}^n \int_{X_k} g d\mathbb{P} = \int_{\Omega} \left(\sum_{j=0}^{n-1} g(T^j \omega) \mathbf{I}_{Y_j}(\omega) \right) \mathbb{P}[d\omega].$$

We will prove the stronger fact

$$h(\omega) := \sum_{j=0}^{n-1} g(T^j \omega) \mathbf{I}_{Y_j}(\omega) \geq 0, \quad \forall \omega \in \Omega. \quad (5.1.16)$$

Let $\omega \in \Omega$. Set $x_j = x_j(\omega) := g(T^j \omega)$. Note that $\omega \in Y_j$ if and only if $T^j(\omega) \in X_{n-j}$, i.e., at least one of the numbers

$$x_j, x_j + x_{j+1}, \dots, x_j + x_{j+1} + \dots + x_{n-1}$$

is positive. The inequality (5.1.16) is a special case of the following cute combinatorial lemma of F. Riesz [178].

Lemma 5.1.25. *Suppose are given a finite sequence of real numbers*

$$\underline{x} := x_0, \dots, x_{n-1}$$

We say that x_j is a leading term of $\underline{x}$ if there exists $\ell \geq j$ such that $x_j + \dots + x_\ell > 0$. Then the sum of the leading terms is ≥ 0 .

Proof. The lemma is easily proved by induction on n . For $n = 1$ this is obviously true. Assume that it is true for any $m < n$ and any sequence of m real numbers. Denote by L the set of indices $j = 0, 1, \dots, n-1$ such that x_j is a leading terms. If $L = \emptyset$ the conclusion is trivially true.

Suppose $L \neq \emptyset$, set $j_0 := \min L$ and denote by ℓ_0 the smallest $\ell \geq j_0$ such that

$$x_{j_0} + \dots + x_{\ell_0} > 0.$$

If $\ell_0 = j_0$, then $x_{j_0} > 0$. Suppose that $\ell_0 > j_0$. The minimality of ℓ_0 implies that for any j , such that $j_0 \leq j < \ell_0$ we have $x_{j_0} + \dots + x_j < 0$ so that, for $j_0 < k \leq \ell_0$ we have

$$x_k + \dots + x_{\ell_0} > 0.$$

This proves that each of the terms $x_{j_0}, x_{j_0+1}, \dots, x_{\ell_0}$ is a leading term. Their sum is obviously nonnegative.

Consider now the (shorter) sequence

$$\underline{y} : y_0 = x_{\ell_0+1}, \dots, y_{m-1} := x_n, \quad m := n - 1 - \ell_0 < n - 1.$$

The induction assumption implies that the sum of the leading terms of $\underline{y}$ is ≥ 0 . The minimality of j_0 implies that the leading terms of $\underline{x}$ are $x_{j_0}, \dots, x_{\ell_0}$ together with the leading terms of $\underline{y}$. This proves Lemma 5.1.25 and completes the proof of Theorem 5.1.22. $\square$

Second proof of (5.1.13). We continue using the notations (5.1.14). Set $G_n := \mathbb{M}_n[g]$.

Since $X_n \nearrow X$, it suffices to show that

$$\int_{X_n} g d\mathbb{P} \geq 0, \quad \forall n.$$

The operator $f \mapsto \hat{T}f$ is monotone, i.e., $f_0 \leq f_1 \Rightarrow \hat{T}f_0 \leq \hat{T}f_1$, and we deduce that for $1 \leq k \leq m$ we have

$$S_{k-1}(g) \leq \max_{1 \leq j \leq m-1} S_j(g) = G_{m-1} \leq G_{m-1}^+$$

and

$$S_k(g) = g + \hat{T}S_{k-1}(g) \leq g + \hat{T}G_{m-1} \leq g + \hat{T}G_{m-1}^+$$

so that

$$G_{m-1} \leq G_m \leq g + \hat{T}G_{m-1}^+, \quad \forall m \in \mathbb{N},$$

or equivalently

$$g \geq G_n - \hat{T}G_n^+, \quad \forall n.$$

We deduce

$$\begin{aligned} \int_{X_n} g &\geq \int_{X_n} G_n - \int_{X_n} \hat{T}G_n^+ \\ (\hat{T}G_n^+ &\geq 0 \text{ on } \Omega, G_n = G_n^+ \text{ on } X_n, G_n^+ = 0 \text{ on } \Omega \setminus X_n) \\ &\geq \int_{X_n} G_n^+ - \boxed{\int_{\Omega} \hat{T}G_n^+} = \int_{\Omega} G_n^+ - \boxed{\int_{\Omega} G_n^+} = 0 \end{aligned}$$

where, at the last step, the equality of the boxed terms is due to the fact that T is measure preserving. $\square$

Remark 5.1.26. (a) For a shorter proof of the pointwise ergodic theorem I refer to [112].

(b) Step (i) is equivalent to some form of maximal inequality similar to (5.1.12). This is often referred to as the Banach principle. For details I refer to [71, Sec. 11.2] or [85, Sec.1.1]. $\square$

Remark 5.1.27. In Remark 5.1.12 we suggested that the ergodicity condition points to a chaotic behavior of the dynamics of the iterates of T . The ergodic theorem makes this much more precise.

Suppose that T is a measure preserving self-map of the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. If T is ergodic, then for any subset $S \in \mathcal{S}$ there exists a negligible subset $\mathcal{N} \in \mathcal{S}$ such that

$$\forall \omega \in \Omega \setminus \mathcal{N}, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \# \{ k; T^k \omega \in S, \quad 0 \leq k < n \} = \mathbb{P}[S]. \quad (5.1.17)$$

Indeed, the left-hand-side of (5.1.17) is the temporal average $A_n[\mathbf{I}_S](\omega)$ so (5.1.17) follows from the Ergodic Theorem 5.1.22. Observe that (5.1.17) states that for most ω , the orbit $\mathcal{O}_T(\omega)$ spends equal amounts of time in sets of equal measures. In other words, most orbits are equidistributed. The equidistribution phenomenon characterizes ergodicity.

Let us observe that, conversely, if a measure preserving map T satisfies the above equidistribution property, then it has to be ergodic. Indeed, suppose that S is a T -invariant set. Let $\mathcal{N}$ be a negligible set as in (5.1.17). Then for any $\omega \in (\Omega \setminus S) \setminus \mathcal{N}$ the orbit $\mathcal{O}_T(\omega)$ does not intersect S since S is

invariant. In this case the left-hand-side of (5.1.17) is equal to zero so $\mathbb{P}[S] = 0$. Thus if S is invariant and its complement $\Omega \setminus S$ is not negligible, then S must be so. Hence T is ergodic.

If we partition Ω into a finite number of measurable sets $S_1, \dots, S_N$, $p_k = \mathbb{P}[S_k] > 0$, $\forall k$, then there exists a negligible set $N \subset \Omega$ so that for any $\omega \in \Omega \setminus N$ the orbit $\mathcal{O}_T(\omega)$ will be located at each moment of time in one of the chambers S_k of this partition. Moreover, it spends a fraction p_k of the time in the chamber S_k . From this point of view, we can regard the dynamics as hopscotching randomly from one chamber to another, and each chamber is frequented as often as its size. We want to warn that this hopscotching need not have a Markovian nature. $\square$

5.2. Applications

Ergodicity is the unifying principle behind some of the limit theorems we have discussed in the previous chapters and it is the source of many interesting non-probabilistic results.

5.2.1. Limit theorems. The Strong Law of Large Numbers is a consequence of the Ergodic Theorem.

Example 5.2.1 (I.i.d. random variables). Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of i.i.d. integrable random variables defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Kolmogorov's zero-one theorem shows that this is a Kolmogorov family, thus ergodic. Consider the coordinate maps on the path space

$$U_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}, \quad U_m(u_1, u_2, \dots) = u_n$$

The Ergodic Theorem implies

$$\frac{1}{n}(U_1 + \dots + U_n) \rightarrow \mathbb{E}[U_1] \quad \mu - \text{a.s.}$$

Observing that $X_n = U_n \circ \vec{X}$ we deduce the Strong Law of Large Numbers. $\square$

Example 5.2.2 (Exchangeable sequences variables). Suppose that $(X_n)_{n \geq 1}$ is an exchangeable sequence of integrable real valued random variables defined on the same probability space $(\Omega, \mathcal{S}, \mathbb{P})$. $\square$

Consider the coordinate maps on the path space

$$U_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}, \quad U_m(u_1, u_2, \dots) = u_n$$

Denote by $\mathcal{I}_{\Theta}$ the sigma-subalgebra of $\mathbb{R}^{\mathbb{N}}$ consisting of shift-invariant events and by $\mathcal{S}_{\infty}$ the sigma-subalgebra of exchangeable events. The Ergodic Theorem implies

$$\frac{1}{n}(U_1 + \dots + U_n) \rightarrow \mathbb{E}[U_1 \mid \mathcal{I}_{\Theta}]$$

On the other hand the equality (5.1.4) implies

$$\mathbb{E}[U_1 \mid \mathcal{I}_{\Theta}] = \mathbb{E}[U_1 \mid \mathcal{S}_{\infty}] = \mathbb{E}[U_1 \mid \mathcal{T}_{\infty}].$$

Thus the Law of Large Numbers for exchangeable sequences is also a special case of the Ergodic Theorem. $\square$

Example 5.2.3 (Markov chains). Consider an irreducible, recurrent HMC $(X_n)_{n \geq 0}$ with state space $\mathfrak{X}$, transition matrix Q and initial distribution μ . The path space of this Markov chain (see Theorem 4.1.3) is the probability space

$$\mathbb{U} = \mathbb{U}_{\mu} = (\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_{\mu}),$$

where

$$U_n(u_0, u_1, u_2, \dots) = u_n.$$

Recall that for any $x \in \mathfrak{X}$ we set $\mathbb{P}_x = \mathbb{P}_{\delta_x}$ where δ_x is the Dirac measure on $\mathfrak{X}$ concentrated at x then

$$\mathbb{P}_\mu = \sum_{x \in \mathfrak{X}} \mu_x \mathbb{P}_x, \quad \mu_x = \mu[\{x\}]. \quad (5.2.1)$$

Denote by $\mathcal{J} \subset \mathcal{E}$ the sigma-algebra of sets that are Θ -invariant sets, where Θ denotes the shift on $\mathbb{U}$. Fix $x \in \mathfrak{X}$ and let $A \in \mathcal{J}$.

The Markov property (4.1.16) implies that

$$\mathbb{E}_x[\mathbf{I}_A \circ \Theta^n \mid \mathcal{E}_n] = \mathbb{E}_{X_n}[\mathbf{I}_A] = \mathbb{P}_{X_n}[A], \quad \mathcal{E}_n = \sigma(X_0, \dots, X_n).$$

Since A is invariant we deduce $\mathbf{I}_A = \mathbf{I}_A \circ \Theta^n$ a.s. so

$$\mathbb{E}_x[\mathbf{I}_A \mid \mathcal{E}_n] = \mathbb{E}_{X_n}[\mathbf{I}_A].$$

We deduce from Lévy's 0-1 theorem that

$$\mathbb{E}_x[\mathbf{I}_A \mid \mathcal{E}_n] \rightarrow \mathbf{I}_A \text{ a.s..}$$

Hence

$$\mathbb{P}_{X_n}[A] = \mathbb{E}_{X_n}[\mathbf{I}_A] \rightarrow \mathbf{I}_A \text{ a.s..}$$

Note that

$$\mathbb{P}_{X_n}[A] = \mathbb{P}_x[A \mid X_n = x] \mathbb{P}_x[X_n = x] + \sum_{x' \in \mathfrak{X} \setminus \{x\}} \mathbb{P}_x[A \mid X_n = x'] \mathbb{P}_x[X_n = x'].$$

On the other hand, $\mathbb{P}_x[X_n = x \text{ i.o.}] = 1$, since the chain is recurrent. Thus

$$\mathbb{P}\left[\left\{\mathbb{P}_{X_n}[A] = \mathbb{P}_x[A] \text{ i.o.}\right\}\right] = 1.$$

Hence $\mathbf{I}_A = \mathbb{P}_x[A]$ a.s. so $\mathbb{P}_x[A] \in \{0, 1\}$. Using (5.2.1) we deduce that $\mathbb{P}_\mu[A] \in \{0, 1\}$ for any initial distribution μ .

If the chain is positively recurrent and π_∞ is the invariant distribution, then Θ is measure preserving and we deduce that $\mathcal{J}$ is a zero-one algebra so Θ is ergodic. For a description of the sigma-subalgebra of shift-invariant subsets of a positively recurrent HMC we refer to [20].

We see that the Ergodic Theorem for Markov chains (Corollary 4.4.3) is a special case of Birkhoff's Ergodic Theorem because any $f \in L^1(\mathfrak{X}, \pi)$ induces a function $\bar{f} = f \circ U_0 \in L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathbb{P}_{\pi_\infty})$.

The fact that the shift map is ergodic allows us to state results stronger than Corollary 4.4.3. For any finite set $B \subset \mathfrak{X} \times \mathfrak{X}$ we obtain a function $F_B \in L^1(\mathfrak{X}^{\mathbb{N}_0}, \mathbb{P}_{\pi_\infty})$

$$F_B(u_0, u_1, u_2, \dots) = \mathbf{I}_B(u_0, u_1)$$

and a corresponding Law of Large Numbers

$$\frac{1}{n} \sum_{k=0}^{n-1} \mathbf{I}_B(X_k, X_{k+1}) = \mathbb{E}_{\pi_\infty}[F_B] \rightarrow \sum_{(x_0, x_1) \in B} \pi_\infty[x_0] Q_{x_0, x_1}, \quad (5.2.2)$$

One should think of B as a collection of directed edges, a "bridge". In the left-hand-side we have the fraction of time a path of the Markov chain "crosses the bridge B ".

Here is an amusing simple illustration of this result. Consider the graph G obtained from two disjoint connected graphs G_0, G_1 with a single edge from a vertex u_0 in G_0 to a vertex u_1 in G_1 . For a vertex v_i of G_i we denote by $\deg_i(v_i)$ its degree in G_i . We denote by E_i the number of edges of G_i .

Let B be the set consisting of the single oriented edge (u_0, u_1) . In this case

$$Q_{u_0, u_1} = \frac{1}{\deg_0(u_0) + 1}, \quad \pi_\infty[u_0] = \frac{\deg_0(u_0) + 1}{2E_0 + 2E_1 + 2}.$$

Formula (5.2.2) shows that the standard random walk on G crosses the bridge from u_0 to u_1 roughly a fraction $\frac{1}{2E_0 + 2E_1 + 2}$ of the time. $\square$

Example 5.2.4 (Weyl's equidistribution theorem). Fix $\varphi \in (0, 2\pi)$ and denote by R_φ the planar counterclockwise of angle φ about the origine. This induces a transformation of the unit circle

$$S^1 := \{z \in \mathbb{C}; |z| = 1\}.$$

This preserves the canonical probability measure μ on S^1

$$\mu[d\theta] = \frac{1}{2\pi}d\theta.$$

As in the previous section this induces a unitary operator

$$\widehat{R}_\varphi : L^2(S^1, \mu) \rightarrow L^2(S^1, \mu), \quad \widehat{R}_\varphi f(\theta) = f(\theta + \varphi).$$

Above the functions in $L^2(S^1, \mu)$ are *complex valued*. For $n \in \mathbb{Z}$ we set

$$e_n(\theta) = e^{in\theta} \in L^2(S^1, \mu).$$

Note that $\widehat{R}_\varphi e_n = e^{in\varphi} e_n$. Since the collection $(e_n)_{n \in \mathbb{Z}}$ is a complete orthonormal sistem we deduce that the eigenspace corresponding to the eigenvalue 1 of $\widehat{R}_\varphi$

$$\ker(1 - \widehat{R}_\varphi) = \text{span} \left\{ e_n; \frac{n\varphi}{2\pi} \in \mathbb{Z} \right\}.$$

We deduce that $\ker(1 - \widehat{R}_\varphi)$ is 1-dimensional iff $\frac{\varphi}{2\pi}$ is irrational. In this case $\widehat{R}_\varphi$ is ergodic and we deduce from (5.1.17) that if $A \subset S^1$ then for almost any $\theta \in S^1$ we have the asymptotic equidistribution equality

$$\frac{1}{n} \sum_{k=0}^{n-1} \mathbf{I}_k(\theta + k\varphi) = \frac{\theta_1 - \theta_0}{2\pi}, \quad \text{a.s.} \quad (5.2.3)$$

With a little bit more work one can show that (5.2.3) holds for any θ . This is *Weyl's equidistribution theorem*, [221]. The reader interested in more details on the equidistribution problem can consult [124]. $\square$

5.2.2. Mixing. Suppose that T is a measure preserving transformation of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Note that if T is ergodic, then the L^2 ergodic theorem implies that

$$\frac{1}{n} \sum_{k=0}^{n-1} f \circ T^k \xrightarrow{L^2} \mathbb{E}[f] \mathbf{I}_\Omega, \quad \forall f \in L^2(\Omega, \mathcal{S}, \mathbb{P}).$$

If we take the inner product with $g \in L^2$ of both sides in the above equality we deduce

$$\frac{1}{n} \sum_{k=0}^{n-1} \int_{\Omega} (f \circ T^k) g \, d\mathbb{P} \rightarrow \mathbb{E}[f] \mathbb{E}[g], \quad \forall f, g \in L^2(\Omega, \mathcal{S}, \mathbb{P}) \quad (5.2.4)$$

In particular, if we let $f = \mathbf{I}_A$, $g = \mathbf{I}_B$, $A, B \in \mathcal{S}$, we deduce

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{P}[T^{-k}(A) \cap B] = \mathbb{P}[A] \mathbb{P}[B]. \quad (5.2.5)$$

Let us observe that the above condition is equivalent with ergodicity. Indeed if we let A quasi-invariant and $B = X \setminus A$, then $\mathbb{P}[T^{-k}(A) \cap B] = 0$, $\forall k$ and we deduce

$$\mathbb{P}[A](1 - \mathbb{P}[A]) = 0$$

so any quasi-invariant set has measure 0 or 1.

Since convergent sequences are also Cèsaro convergent we deduce that condition (5.2.5) follows from the stronger requirement

$$\lim_{n \rightarrow \infty} \mathbb{P}[T^{-n}(A) \cap B] = \mathbb{P}[A] \mathbb{P}[B], \quad \forall A, B \in \mathcal{S}. \quad (5.2.6)$$

A measure preserving map T satisfying this condition is said to be *mixing*.

When T is an automorphism one can give a more visual interpretation of the mixing condition. In this case mixing is also equivalent to the condition

$$\lim_{n \rightarrow \infty} \mathbb{P}[A \cap T^n(B)] = \mathbb{P}[A] \mathbb{P}[B], \quad \forall A, B \in \mathcal{S}. \quad (5.2.7)$$

Assume that the region B is occupied molecules of black ink in a glass of crystalline water. These molecules occupy a fraction $\mathbb{P}[B]$ of the entire space. Flow the black region B using T . Thus, $T^n(B)$ represents the location of the black region after n units of time.

The mixing condition shows that after a while, the fraction $\mathbb{P}[A \cap T^n(B)] / \mathbb{P}[A]$ of a region A occupied by these moving molecules of black ink is equal to $\mathbb{P}[B]$. Thus in the long run, all the regions will have the same fraction of black ink. To use a very apt analogy in Arnold and Avez [4], this is what happens when we mix well a cocktail.

The mixing condition (5.2.6) can be rewritten as

$$\lim_{n \rightarrow \infty} (\widehat{T^n} \mathbf{I}_A, \mathbf{I}_B) = \lim_{n \rightarrow \infty} \mathbb{E}[(\mathbf{I}_A \circ T^n) \cdot \mathbf{I}_B] = \mathbb{E}[\mathbf{I}_A] \cdot \mathbb{E}[\mathbf{I}_B], \quad \forall A, B \in \mathcal{S}. \quad (5.2.8)$$

This implies that for any elementary functions $f, g \in \text{Elem}(\Omega, \mathcal{S})$ we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} (f \circ T^n) g \, d\mathbb{P} = \left(\int_{\Omega} f \, d\mathbb{P} \right) \left(\int_{\Omega} g \, d\mathbb{P} \right).$$

Since $\text{Elem}(\Omega, \mathcal{S})$ is dense in $L^2(\Omega, \mathcal{S}, \mathbb{P})$ we deduce that if T is mixing, then

$$\forall f, g \in L^2(\Omega, \mathcal{S}, \mathbb{P}) : \lim_{n \rightarrow \infty} \int_{\Omega} (f \circ T^n) g \, d\mathbb{P} = \left(\int_{\Omega} f \, d\mathbb{P} \right) \left(\int_{\Omega} g \, d\mathbb{P} \right). \quad (5.2.9)$$

Clearly, if a measure preserving map satisfies (5.2.9), then it is mixing. The above argument has the following immediate generalization.

Proposition 5.2.5. *Suppose that $T : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\Omega, \mathcal{S}, \mathbb{P})$ is a measure preserving map and*

$$\mathcal{C} \subset L^2(\Omega, \mathcal{S}, \mathbb{P})$$

is a collection of functions such that $\text{span}(\mathcal{C})$ is dense in $L^2(\Omega, \mathcal{S}, \mathbb{P})$. Then the following are equivalent.

- (i) *The map T is mixing.*

(ii) For every $f, g \in \mathcal{C}$,

$$\lim_{n \rightarrow \infty} (\hat{T}^n f, g)_{L^2(\Omega)} = \left(\int_{\Omega} f d\mathbb{P} \right) \left(\int_{\Omega} g d\mathbb{P} \right). \quad (5.2.10)$$

□

Remark 5.2.6. Note that condition (5.2.10) can be rephrased as

$$\lim_{n \rightarrow \infty} \mathbb{E}[\hat{T}^n f \cdot g] = \mathbb{E}[f] \mathbb{E}[g].$$

Observing that

$$\mathbb{E}[f] = \mathbb{E}[\hat{T}^n f], \quad \forall n$$

we can rewrite (5.2.10) as

$$\lim_{n \rightarrow \infty} \left(\mathbb{E}[\hat{T}^n f \cdot g] - \mathbb{E}[\hat{T}^n f] \mathbb{E}[g] \right),$$

or equivalently

$$\lim_{n \rightarrow \infty} \text{Cov}[\hat{T}^n f, g] = 0, \quad \forall f, g \in \mathcal{C}. \quad (5.2.11)$$

Thus, the mixing condition is a weak form of asymptotic independence. □

Let us give a few examples of mixing maps.

Proposition 5.2.7. Suppose that $X_n : (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow (\mathbb{X}, \mathcal{F})$, $n \in \mathbb{N}$, is a Kolmogorov stationary sequence of measurable maps. Then the shift map on the path space is mixing.

Proof. We will show that the shift map Θ satisfies (5.2.6). Denote by U_n the coordinate maps on the path space $U_n : \mathbb{X}^{\mathbb{N}} \rightarrow \mathbb{X}$, and set

$$\mathcal{T}_n := \sigma(U_n, U_{n+1}, \dots).$$

For $B \in \mathcal{F}$ and $m \in \mathbb{N}$ we set

$$\varepsilon_m(B) := \sup_{S \in \mathcal{T}_m} |\mathbb{P}[S \cap B] - \mathbb{P}[S] \mathbb{P}[B]|.$$

Since $(X_n)_{n \in \mathbb{N}}$ is a Kolmogorov sequence we deduce from Theorem 5.1.16 $\varepsilon_m(B) \rightarrow 0$ as $m \rightarrow \infty$.

Observe that if $A \in \mathcal{F}$, then $T^{-m}(A) \in \mathcal{T}_n$ so that

$$|\mathbb{P}[T^{-m}(S) \cap B] - \mathbb{P}[T^{-m}(A)] \mathbb{P}[B]| \leq \varepsilon_m(B) \rightarrow 0.$$

This implies (5.2.6) since T is measure preserving so $\mathbb{P}[T^{-m}(A)] = \mathbb{P}[A]$. □

Proposition 5.2.8. Suppose that $(X_n)_{n \geq 0}$ is an irreducible, positively recurrent HMC with state space $\mathfrak{X}$, transition matrix Q and stationary distribution π . Then the following are equivalent.

- (i) The HMC is aperiodic.
- (ii) The shift map on the path space $(\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_{\pi})$ is mixing.

Proof. I follow the approach in [28, Sec. 16.1.2].

(i) $\Rightarrow$ (ii) Suppose that our HMC is aperiodic. Consider the path space $\mathbb{U} = \mathfrak{X}^{\mathbb{N}_0}$. Denote by $\mathcal{C}$ the collection of cylindrical subsets of $\mathbb{U}$ of the form

$$C_{x_{i_1}, \dots, x_{i_k}} := \{\underline{u} \in \mathbb{U}; \quad u_{i_j} = x_{i_j} \in \mathfrak{X}, \quad \forall 1 \leq j \leq k\}, \quad 0 \leq i_1 < \dots < i_k, \quad k \in \mathbb{N}.$$

In view of Proposition 5.2.5 it suffices to show that (5.2.6) is satisfied for any $A, B \in \mathcal{C}$. Suppose that

$$A = C_{x_{i_1}, \dots, x_{i_k}}, \quad B = C_{x_{j_1}, \dots, x_{j_m}}.$$

For $n > j_m$ we have

$$\Theta^{-n}(A) \cap B = C_{x_{j_1}, \dots, x_{j_m}, x_{i_1+n}, \dots, x_{i_k+n}}, \quad x_{i_j+n} := x_{i_j},$$

and

$$\begin{aligned} \mathbb{P}_\pi[\Theta^{-n}(A) \cap B] &= \pi[x_{j_1}] Q_{x_{j_1}, x_{j_2}}^{j_2-j_1} \cdots Q_{x_{j_{m-1}}, x_{j_m}}^{j_m-j_{m-1}} \boxed{Q_{x_{j_m}, x_{i_1}}^{n+i_1-j_m}} \times \\ &\quad \times Q_{x_{i_1}, x_{i_2}}^{i_2-i_1} \cdots Q_{x_{i_{k-1}}, x_{i_k}}^{i_k-i_{k-1}}. \end{aligned} \quad (5.2.12)$$

Since the HMC is aperiodic we deduce from (4.4.9) we deduce that

$$\lim_{n \rightarrow \infty} Q_{x_{j_m}, x_{i_1}}^{n+i_1-j_m} = \pi[x_{i_1}].$$

Using this in (5.2.12) we deduce that

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}_\pi[\Theta^{-n}(A) \cap B] &= \underbrace{\pi[x_{j_1}] Q_{x_{j_1}, x_{j_2}}^{j_2-j_1} \cdots Q_{x_{j_{m-1}}, x_{j_m}}^{j_m-j_{m-1}}}_{\mathbb{P}_\pi[B]} \times \\ &\quad \times \underbrace{\pi[x_{i_1}] Q_{x_{i_1}, x_{i_2}}^{i_2-i_1} \cdots Q_{x_{i_{k-1}}, x_{i_k}}^{i_k-i_{k-1}}}_{\mathbb{P}_\pi[A]}. \end{aligned}$$

(ii) $\Rightarrow$ (i) Suppose that the shift map is mixing. To prove that it is aperiodic we argue by contradiction and assume the period d is bigger than 1. As in Proposition 4.2.14 consider the communication classes of Q^d ,

$$C_1, C_2, \dots, C_d \subset \mathfrak{X}.$$

Hence

$$\mathbb{P}[X_{n+1} \in C_{i+1 \bmod d} \mid X_n \in C_i \bmod d] = 1, \quad \forall n \geq 0, \quad i = 1, \dots, d.$$

Consider the sets

$$A_i = \{ \underline{u} \in \mathfrak{X}^{\mathbb{N}_0}; \quad u_0 \in C_i \}, \quad i = 1, 2, \dots.$$

Then $\Theta^{-n}(A_i) = A_{i+n \bmod d}$, $A_i \cap A_j = \emptyset$ if $i \not\equiv j \bmod d$. We deduce that for any $n \in \mathbb{N}$ we have

$$\mathbb{P}_\pi[\Theta^{-nd}(A_0) \cap A_1] = 0, \quad \mathbb{P}_\pi[\Theta^{-nd-1}(A_0) \cap A_1] = \mathbb{P}_\pi[A_1] = \pi[A_1] \neq 0.$$

This contradicts the fact that Θ is mixing. $\square$

Remark 5.2.9. Suppose that $(X_n)_{n \geq 0}$ is an HMC as in the above proposition. We know that if the sequence $(X_n)_{n \geq 0}$ is Kolmogorov, then it is mixing. A theorem of Blackwell and Freedman [20] shows that the converse is also true so $(X_n)_{n \geq 0}$ is mixing if and only if it is Kolmogorov. In fact, the following properties are equivalent.

- (i) The HMC $(X_n)_{n \geq 0}$ is aperiodic
- (ii) For any probability measures $\mu, \nu \in \text{Prob}(\mathfrak{X})$

$$\lim_{n \rightarrow \infty} d_v(\mu Q^n - \nu Q^n) = 0, .$$

- (iii) The HMC $(X_n)_{n \geq 0}$ is mixing.
- (iv) The HMC $(X_n)_{n \geq 0}$ is Kolmogorov.

For a proof I refer to [109, Thm. 26.10]. $\square$

Example 5.2.10 (The tent map). Consider the tent map $T : [0, 1] \rightarrow [0, 1]$ introduced in Example 5.1.1(d). Recall that T is the continuous map $[0, 1] \rightarrow [0, 1]$ such that $T(0) = 0 = T(1)$, $T(1/2) = 1$ and T is linear on each of the intervals $[0, 1/2]$ and $[1/2, 1]$. We want to show that T is mixing.

Consider the Haar orthonormal basis of $L^2([0, 1])$. Recall its definition. It consists of the *Haar functions*

$$\begin{aligned} H_{-1} &= 1, \quad H_1 = H_{0,0} = \mathbf{I}_{[0,1/2]} - \mathbf{I}_{[1/2,1]} \\ H_{n,k}(x) &= 2^{n/2} H_{0,0}(2^n x - k) \\ &= 2^{n/2} \mathbf{I}_{\left[\frac{k}{2^n}, \frac{k}{2^n} + \frac{1}{2^{n+1}}\right)} - 2^{n/2} \mathbf{I}_{\left[\frac{k}{2^n} + \frac{1}{2^{n+1}}, \frac{k+1}{2^n}\right)}, \quad 0 \leq k < 2^n. \end{aligned}$$

Define

$$\begin{aligned} \mathbf{H}_{-1} &= \text{span } \mathbf{I}_{[0,1]}, \quad \mathbf{H}_n := \text{span } \{ H_{n,k}; \quad 0 \leq k < 2^n \}, \quad n \geq 0, \\ \mathcal{H} &= \{ H_0 \} \cup \{ H_{n,k}; \quad n \geq 0, \quad 0 \leq k < 2^n \}. \end{aligned}$$

The subspaces $\mathbf{H}_n$ are mutually orthogonal and the collection $\mathcal{H}$ spans a dense subspace of $L^2([0, 1])$; see [31, Sec. 9.2]. Moreover

$$\widehat{T} \mathbf{H}_n \subset \mathbf{H}_{n+1}, \quad \forall n \geq 0.$$

Thus if $m, n \geq 0$, $0 \leq j \leq 2^m$, $0 \leq k \leq 2^n$ we have

$$(\widehat{T}^\ell H_{m,j}, H_{n,k})_{L^2} = 0 = \left(\int_0^1 H_{m,j}(x) dx \right) \left(\int_0^1 H_{n,k}(x) dx \right), \quad \forall \ell > n - m.$$

Clearly $\widehat{T} H_0 = H_0$. Proposition 5.2.5 applied to the collection $\mathcal{H}$ implies that T is mixing, hence ergodic. $\square$

Example 5.2.11 (Arnold's cat). We consider a slightly more general situation. Let $d > 1$ and denote by $\mathbb{T}^d$ the d -dimensional torus

$$\mathbb{T}^m = \underbrace{S^1 \times \cdots \times S^1}_m$$

equipped with the invariant probability measure

$$\mu[d\theta] = \frac{1}{(2\pi)^d} d\theta^1 \cdots d\theta_d,$$

where θ_i are the standard angular coordinates on the torus. Suppose that $A \in \text{SL}_d(\mathbb{Z})$, i.e., A is a $d \times d$ matrix with integral entries and determinant 1. Since $A\mathbb{Z}^d = \mathbb{Z}^d$ we deduce that A defines a measure preserving map of $\mathbb{T}^d$

$$\theta = \begin{bmatrix} \theta_1 \\ \vdots \\ \theta_m \end{bmatrix} \mapsto T_A \theta := A \cdot \theta \bmod (2\pi\mathbb{Z})^d.$$

Denote by $\langle -, - \rangle$ the canonical inner product in $\mathbb{R}^d$ and by A^* the transpose of A . Clearly

$$A^* \in \text{SL}_d(\mathbb{Z}) \quad \text{and} \quad A^*(\mathbb{Z}^d) = \mathbb{Z}^d.$$

For each $\vec{m} \in \mathbb{Z}^d$ we denote by $\mathcal{O}_{\vec{m}}$ the orbit of the action of A^* on $\mathbb{Z}^d$, i.e., the set

$$\mathcal{O}_{\vec{m}} = \{ (A^*)^n \vec{m}; \quad n \geq 0 \}.$$

To any $\vec{m} \in \mathbb{Z}^d$ we associate the *character*¹ $\chi_{\vec{m}} \in L^2(\mathbb{T}^d, \mu)$

$$\chi_{\vec{m}}(\theta) = e^{i\langle \vec{m}, \theta \rangle} = e^{i(m_1\theta_1 + \dots + m_d\theta_d)}, \quad i := \sqrt{-1}.$$

The set of characters

$$\mathcal{C}_d := \{\chi_{\vec{m}}; \vec{m} \in \mathbb{Z}^d\} \subset L^2(\mathbb{T}^d, \mu) \quad (5.2.13)$$

is an orthonormal family that spans a vector subspace dense in $L^2(\mathbb{T}^d, \mu)$.

The unitary operator $\hat{T}_A : L^2(\mathbb{T}^d, \mu) \rightarrow L^2(\mathbb{T}^d, \mu)$ has the explicit description

$$\hat{T}_A f(\theta) = f(A\theta).$$

In particular,

$$\hat{T}_A \chi_{\vec{m}}(\theta) = e^{i\langle \vec{m}, A\theta \rangle} = e^{i\langle A^* \vec{m}, \theta \rangle} = \chi_{A^* \vec{m}}(\theta).$$

We have the following result.

Theorem 5.2.12. *Let $A \in \text{SL}_d(\mathbb{Z})$, $d > 1$. The following are equivalent.*

- (i) *The map $A : \mathbb{T}^d \rightarrow \mathbb{T}^d$ is ergodic.*
- (ii) *For any $\vec{m} \in \mathbb{Z}^d \setminus \{0\}$ the orbit $\mathcal{O}_{\vec{m}}$ is infinite.*
- (iii) *The map $A : \mathbb{T}^d \rightarrow \mathbb{T}^d$ is mixing.*

Proof. I follow the approach in [46, Sec. 4.3]. We only need to prove (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii).

(i) $\Rightarrow$ (ii) We argue by contradiction. Suppose there exists $\vec{m} \in \mathbb{Z}^d \setminus \{0\}$ such that $\mathcal{O}_{\vec{m}}$ is finite. Denote by n the smallest $n \in \mathbb{N}$ such that $(A^*)^n \vec{m} = \vec{m}$. Then the function

$$f = \chi_{\vec{m}} + \dots + \chi_{(A^*)^{n-1} \vec{m}}$$

is $\hat{T}_A$ -invariant and nonconstant since the functions $1, \chi_{\vec{m}} \dots \chi_{(A^*)^{n-1} \vec{m}}$ are linearly independent. Hence A is not ergodic.

(ii) $\Rightarrow$ (iii) We apply Proposition 5.2.5 to the set of characters $\mathcal{C}_d$ in (5.2.13). Note that if

$$\int_{\mathbb{T}^d} \chi_{\vec{m}} d\mu = \begin{cases} 1, & \vec{m} = 0, \\ 0, & \vec{m} \neq 0. \end{cases}$$

Clearly if $f = g = 1$, then (5.2.10) holds trivially. Suppose $f \neq 1$. Then

$$\left(\int_{\Omega} f d\mathbb{P} \right) \left(\int_{\Omega} g d\mathbb{P} \right) = 0.$$

Assumption (ii) implies that $\hat{T}_A^n f$ is a character different from g for all n sufficiently large and thus

$$(\hat{T}_A^n f, g)_{L^1} = 0, \quad \forall n \gg 0.$$

We deduce that A is mixing. □

The condition (ii) above holds if and only if none of the eigenvalues of A are roots of 1. Observe that if one eigenvalue of $A \in \text{SL}_2(\mathbb{Z})$ is a root of 1 then all eigenvalues are roots of 1. We deduce that the only matrices in $\text{SL}_2(\mathbb{Z})$ are

$$\pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \pm \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

¹Any continuous group morphism $\chi : \mathbb{T}^d \rightarrow S^1$ has the form $\chi_{\vec{m}}$ for some $\vec{m} \in \mathbb{Z}^d$.

In particular, this shows that Arnold's cat map is mixing. □

Remark 5.2.13. There is another condition that intermediates between mixing and ergodicity. More precisely, a measure preserving self-map of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ is called *weakly mixing* if,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} | \mathbb{P}[T^{-k}(A) \cap B] - \mathbb{P}[A] \mathbb{P}[B] | = 0 \quad (5.2.14)$$

for any $A, B \in \mathcal{S}$. Clearly (5.2.14) implies (5.2.5) so weakly mixing are ergodic.

Since convergent sequences are Cèsaro convergent we deduce that (5.2.6) implies (5.2.14) so mixing maps are weakly mixing.

It turn out that most weakly mixing automorphisms of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$ are not mixing. More precisely the mixing operators form a meagre (first Baire category) subset in the set of weakly mixing automorphisms, [99, p.77].

5.3. Exercises

Exercise 5.1. Suppose that $(\Omega, \mathcal{S})$ is a measurable space and $T : (\Omega, \mathcal{S}) \rightarrow (\Omega, \mathcal{S})$ a measurable map. Denote by $\text{Prob}_T(\Omega, \mathcal{S})$ the set of T -invariant probability measures $\mathbb{P} : \mathcal{S} \rightarrow [0, 1]$.

- (i) Prove that $\text{Prob}_T(\Omega, \mathcal{S})$ is a convex subset of the space of finite measures on $\mathcal{S}$.
- (ii) Prove that T is ergodic with respect to a probability measure $\mathbb{P}$ if and only if $\mathbb{P}$ is an extremal point of $\text{Prob}_T(\Omega, \mathcal{S})$ i.e., $\mathbb{P}$ cannot be written as a convex combination

$$\mathbb{P} = (1 - t)\mathbb{P}_0 + t\mathbb{P}_1, \quad t \in (0, 1), \quad \mathbb{P} \neq \mathbb{P}_0, \mathbb{P}_1.$$

□

Exercise 5.2. Suppose that (X, d) is a compact metric spaces and $T : X \rightarrow X$ is a continuous map. Let $\mathbb{P}$ be a Borel probability measure on X . For $n \in \mathbb{N}$ we set

$$\mathbb{P}_n := \frac{1}{n} \sum_{k=0}^{n-1} T^k_{\#} \mathbb{P}.$$

- (i) Prove that the sequence $(\mathbb{P}_n)_{n \in \mathbb{N}}$ contains a subsequence $(\mathbb{P}_{n_k})$ that converges weakly to a Borel probability measure $\mathbb{P}_*$ on X , i.e.,

$$\lim_{k \rightarrow \infty} \int_X f(x) \mathbb{P}_{n_k} [dx] = \int_{\mathbb{P}_*} f(x) \mathbb{P}_* [dx], \quad \forall f \in C(X).$$

Hint. Use Banach-Alaoglu compactness theorem.

- (ii) Prove that $\mathbb{P}_*$ is T -invariant.
- (iii) Prove that the set $\text{Prob}_T(X)$ of T -invariant Borel probability measures on X is convex and closed with respect to the weak convergence.

□

Exercise 5.3. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and $T : \Omega \rightarrow \Omega$ a measure preserving map. We say that T is *quasi-mixing* if there exist $c_1, c_2 > 0$ such that $\forall A, B \in \mathcal{S}$

$$c_1 \mathbb{P}[A] \mathbb{P}[B] \leq \mathbb{P}[T^{-1}(A) \cap B] \leq c_2 \mathbb{P}[A] \mathbb{P}[B]. \quad (5.3.1)$$

- (i) Suppose that $\mathcal{A} \subset \mathcal{S}$ is a collection of measurable subsets that generates $\mathcal{S}$, $\sigma(\mathcal{A}) = \mathcal{S}$. Show that T is quasi-mixing if (5.3.1) holds for all $A, B \in \mathcal{A}$.
- (ii) Prove that if T is quasi-mixing, then it is ergodic.

□

Exercise 5.4. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space and $T : \Omega \rightarrow \Omega$ a measure preserving map. Suppose that $(\mathcal{F}_n)_{n \geq 1}$ is a filtration of sigma -subalgebras with the following properties.

- (i)

$$\bigvee_{n \geq 1} \mathcal{F}_n = \mathcal{S}.$$

- (ii) $T^{-1}(\mathcal{F}_n) \subset \mathcal{F}_n, \forall n \in \mathbb{N}$.

- (iii) For any $k \in \mathbb{N}$ the intersection

$$\bigcap_{n \geq 1} (T^k)^{-1}(\mathcal{F}_n)$$

is a 0-1-sigma subalgebra.

Prove that T is mixing. □

Exercise 5.5. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a probability space, $T : \Omega \rightarrow \Omega$ a measure preserving map and $g \in L^1(\Omega, \mathcal{S}, \mathbb{P})$. Prove that the following are equivalent.

- (i) The function g is T -invariant, i.e., $g \circ T = g$ a.s..
- (ii) For any $f \in L^\infty(\Omega, \mathcal{F}, \mathbb{P})$, $\mathbb{E}[gf] = \mathbb{E}[g(f \circ T)]$. □

Exercise 5.6 (Poincaré). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $T : \Omega \rightarrow \Omega$ is a measure preserving measurable map. Prove that for any $S \in \mathcal{S}$ such that $\mathbb{P}[S] > 0$ we have

$$\mathbb{P}[\{\omega \in \Omega; T^n \omega \in S \text{ i.o.}\}] = 1.$$

□

Exercise 5.7 (Kac). Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a probability space and $T : \Omega \rightarrow \Omega$ is a measure preserving measurable map. For $S \in \mathcal{S}$ such that $\mathbb{P}[S] > 0$ we define the first return map

$$T_S : \Omega \rightarrow \mathbb{N} \cup \{\infty\}, \quad T_S(\omega) = \min \{n \in \mathbb{N} : T^n \omega \in S\}$$

Set

$$\Omega_S := \{\omega \in \Omega \setminus S; T^n \omega \notin S, \forall n \geq 1\}.$$

- (i) Prove that

$$\int_S T_S(\omega) \mathbb{P}[d\omega] = 1 - \mathbb{P}[\Omega_S].$$

- (ii) Prove that if T is ergodic then $\mathbb{P}[\Omega_S] = 0$. □

Exercise 5.8. Consider an irreducible HMC $(X_n)_{n \geq 0}$ with finite state space $\mathfrak{X}$, transition matrix Q and whose initial distribution is the stationary distribution μ . The path space of this Markov chain is (see Theorem 4.1.3)

$$\mathbb{U}_\mu = (\mathfrak{X}^{\mathbb{N}_0}, \mathcal{E}, \mathbb{P}_\mu).$$

For $n \in \mathbb{N}_0$ we denote by U_n the n -th coordinate map $U_n(u_0, u_1, \dots) = u_n$. Let

$$f : \mathbb{U}_\mu \rightarrow \mathbb{R}, \quad f(u_0, u_1, \dots) = -\log_2 Q_{u_0, u_1}$$

- (i) Prove that

$$\int_{\mathbb{U}_\mu} f(\underline{u}) \mathbb{P}_\mu[d\underline{u}] = \text{Ent}_2[\mathfrak{X}, Q],$$

where $\text{Ent}_2[\mathfrak{X}, Q]$ denotes the entropy rate of the Markov chain described in Exercise 4.37.

- (ii) Prove that

$$-\lim_{n \rightarrow \infty} \frac{1}{n} \log_2 (Q_{U_0, U_1} \cdots Q_{U_{n-1}, U_n}) = \text{Ent}_2[\mathfrak{X}, Q] \text{ a.s..}$$

□

Exercise 5.9 (Kac). Consider the map $Q : [0, 1) \rightarrow [0, 1)$, $x \mapsto Qx := 2x \bmod 1$. Show that Q is mixing with respect to the Lebesgue measure. **Hint.** See Example 5.1.15. □

Exercise 5.10. Consider the tent map $T : [0, 1] \rightarrow [0, 1]$, $T(x) = \min(2x, 2 - 2x)$ and the logistic map $L : [0, 1] \rightarrow [0, 1]$, $L(x) = 4x(1 - x)$.

- (i) Prove that the map $\Phi : [0, 1] \rightarrow [0, 1]$, $\Phi(x) = \frac{1}{2}(1 - \cos(\pi x))$ is a homeomorphism and $L \circ \Phi = \Phi \circ T$.
- (ii) Describe the measure $\mu := \Phi_{\#}\lambda$, where λ is the Lebesgue measure on $[0, 1]$.
- (iii) Prove that the logistic map preserves μ and it is mixing with respect to this measure. $\square$

Exercise 5.11. Fix $m \in \mathbb{N}$, $m \geq 2$. For any $\vec{\epsilon} \in \{0, 1\}^m$ define

$$F_{\vec{\epsilon}} : [0, 1] \rightarrow [0, 1], \quad F_{\vec{\epsilon}}(x) = \begin{cases} (-1)^{\epsilon_k} m \left(x - \frac{k-1}{m} \right) + \epsilon_k, & \frac{k-1}{m} \leq x < \frac{k}{m}, \\ 0, & x = 1. \end{cases}$$

Prove that $F_{\vec{\epsilon}}$ is mixing for any $\vec{\epsilon} \in \{0, 1\}^m$. $\square$

Exercise 5.12. Consider the Haar functions $H_{n,k}$ used in Example 5.2.10. We define the *Rademacher functions*,

$$R_n : [0, 1] \rightarrow \mathbb{R}, \quad R_n = 2^{-n/2} \sum_{0 \leq k < 2^n} H_{n,k}, \quad n \geq 0$$

- (i) Prove that

$$\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} R_n(x) = 1 - 2x, \quad \forall x \in [0, 1].$$

- (ii) Prove that the functions $(R_n)_{n \geq 0}$, viewed as random variables defined on the probability space $([0, 1], \lambda)$, are i.i.d.. $\square$

Exercise 5.13. Suppose that X is a finite set and π is a probability measure on X given by the weights $\pi_x := \pi[\{x\}] > 0$, $\forall x \in X$. Consider the Cartesian product $\mathbb{U}_{\pi} := X^{\mathbb{Z}}$ equipped with the product sigma-algebra and product measure $\pi_{\infty} := \pi^{\otimes \mathbb{Z}}$. The elements of $\mathbb{X}$ are functions $\mathbf{u} : \mathbb{Z} \rightarrow X$. Consider the shift $\Theta : \mathbb{U}_{\pi} \rightarrow \mathbb{U}_{\pi}$, $\Theta \mathbf{u}(n) = \mathbf{u}(n+1)$.

- (i) Prove that Θ is mixing with respect to the measure π_{∞} .
- (ii) Denote for $S \subset X$ by $\mathcal{N}_{\pi}^S$ the subset of $\mathbb{U}_{\pi}$ consisting of functions $\mathbf{u} : \mathbb{Z} \rightarrow X$ such that $\lim_{n \rightarrow \infty} \mathbf{u}(n)$ exists. Prove that $\pi_{\infty}[\mathcal{N}_{\pi}^S] = 0$ and that the complement $\mathbb{U}_{\pi}^S$ is Θ -invariant. $\square$

Exercise 5.14. Consider the *baker's transform* $B : [0, 1]^2 \rightarrow [0, 1]^2$,

$$B(x, y) = \begin{cases} (q(2x), q(y/2)), & x \leq 1/2, \\ (q(2x), q((y+1)/2)), & x > 1/2, \end{cases}$$

where $q(t)$ denotes the fractional part of the real number t , $q(t) = t - [t]$. Prove that B is mixing with respect to the Lebesgue measure. Consider the map $\Phi : \{0, 1\}^{\mathbb{Z}} \rightarrow [0, 1]^2$ given by $\Phi(\mathbf{u}) = (x(\mathbf{u}), y(\mathbf{u}))$,

$$x(\mathbf{u}) = \sum_{n=0}^{\infty} \frac{\mathbf{u}(-n)}{2^{n+1}}, \quad y(\mathbf{u}) = \sum_{n=1}^{\infty} \frac{\mathbf{u}(n)}{2^n}.$$

Denote by π the uniform measure on $\{0, 1\}$ and by π_{∞} the induced product measure on $\{0, 1\}^{\mathbb{Z}}$.

- (i) Prove that $\Phi_{\#}\pi_{\infty} = \lambda$, where λ is the Lebesgue measure on the square $[0, 1]^2$.

(ii) Show that $B \circ \Phi = \Phi \circ \Theta$, where Θ is the shift defined in Exercise 5.13.

(iii) Prove that the baker's transform is mixing with respect to the Lebesgue measure. $\square$

Exercise 5.15 (Gauss). Consider the map $G : [0, 1] \rightarrow [0, 1]$ given by

$$G(x) = \begin{cases} 0, & x = 0, \\ \frac{1}{x} - \left\lfloor \frac{1}{x} \right\rfloor, & x \in (0, 1]. \end{cases}$$

For $k \in \mathbb{N}$ we set $I_k := (1/(k+1), 1/k)$. Any $x \in (0, 1]$ has a continuous fraction decomposition

$$x = [0 : a_1 : a_2 : \cdots] := 0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}, \quad a_n = a_n(x) \in \mathbb{N}_0, \quad \forall n \in \mathbb{N}.$$

(The number x is rational if and only if $a_n = 0$ for all n sufficiently large.) Set

$$[0, 1]_* := [0, 1] \setminus \mathbb{Q}.$$

(i) Let $x = [0 : a_1 : a_2 : \cdots] \in [0, 1]_*$. Prove that $G(x) = [0 : a_2 : a_3 : \cdots]$ and, for any $n \in \mathbb{N}$, we have

$$x = [0 : a_1 : \cdots : a_{n-1} : a_n + G^n(x)] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots + a_{n-1} + \frac{1}{a_n + G^n(x)}}}}.$$

(ii) Let $x \in [0, 1]_*$. Prove that $a_n(x) = k$ iff $G^n(x) \in I_k, \forall k, n \in \mathbb{N}$.

(iii) For each $a \in \mathbb{R}$ we set

$$T_a = \begin{bmatrix} a & 1 \\ 1 & 0 \end{bmatrix}.$$

Prove that

$$[0 : a_1 : \cdots : a_n] = \frac{p_n}{q_n}, \quad \begin{bmatrix} p_n \\ q_n \end{bmatrix} = T_0 \cdot T_{a_1} \cdots T_{a_n} \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (5.3.2)$$

(iv) Let $x := [0 : a_1 : a_2 : \cdots] \in [0, 1]_*$. Prove that for any $n \in \mathbb{N}$ we have

$$x = \frac{p_n(x) + p_{n-1}(x)G^n(x)}{q_n(x) + q_{n-1}(x)G^n(x)}$$

where $p_n(x), q_n(x)$ are defined in terms of the $a_n(x)$'s by (5.3.2).

(v) Prove that $q_n(x) \geq 2^{\frac{n-2}{2}}, \forall x \in [0, 1]_*, n \in \mathbb{N}$.

(vi) Prove that the restriction of G to I_k is a diffeomorphism onto $(0, 1)$.

(vii) Fix $c > 0$ and set $\rho : [0, 1] \rightarrow [0, \infty)$

$$\rho(x) = \frac{c}{x+1}.$$

Prove that for any $x \in [0, 1]_*$ we have

$$\rho(x) = \sum_{G(y)=x} \frac{\rho(y)}{|G'(y)|}$$

(viii) Prove that the probability measure μ on defined by

$$\mu[dx] = \frac{1}{(x+1)\log 2} \lambda[dy]$$

is G -invariant.

(ix) Prove that for any $n \in \mathbb{N}$ the map $A_n : [0, 1]_* \rightarrow \mathbb{N}$, $x \mapsto a_n(x)$ is measurable and the sigma-algebra generated by these random variables coincides with the Borel sigma algebra.

Hint. Show that the set $I_{a_1, \dots, a_m} := \{A_k = a_k, 1 \leq k \leq m\}$ is an interval with endpoints expressible in terms of the fractions $\frac{p_k}{q_k}$ defined as in (5.3.2).

(x) Show that G is quasi-mixing (see Exercise 5.3) hence ergodic. □

Exercise 5.16. Suppose that T is an automorphism of a probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Define

$$T^{\times 2} : \Omega \rightarrow \Omega \rightarrow \Omega \times \Omega, \quad T^{\times 2}(\omega_1, \omega_2) = (T\omega_1, T\omega_2).$$

Prove that $T^{\times 2}$ is ergodic (with respect to $\mathbb{P}^{\otimes 2}$) if and only if T is weakly mixing, i.e., satisfies (5.2.14). □

An introduction to stochastic calculus

In Example 2.5.9 we associated to a Brownian motion

$$B : [0, \infty) \times (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}, \quad (t, \omega) \mapsto B(t, \omega)$$

a Gaussian white noise, $\mathcal{W} = \mathcal{W}_B$. This is an isometry

$$\mathcal{W} : L^2([0, \infty)) \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P}), \quad f \mapsto \mathcal{W}[f]$$

such that

$$\mathcal{W}[\mathbf{I}_{[0,t]}] = B(t).$$

The Gaussian random variable $\mathcal{W}[f]$ is traditionally denoted as an integral

$$\mathcal{W}[f] = \int_0^\infty f(s)dB(s).$$

For this reason $\mathcal{W}[f]$ is called the *Wiener integral* with respect the Brownian motion B . The details behind construction of $\mathcal{W}$ are outlined in Exercise 2.78. We recall the most salient points.

Choose a sequence of step functions,

$$f_n = \sum_{k \geq 1} c_{k,n} \mathbf{I}_{(s_{kn}, t_{kn}]},$$

where the intervals $(s_{kn}, t_{kn}]$ are disjoint and $c_{kn} = 0$ for all but finitely many k 's, and such that

$$\|f - f_n\|_{L^2} \rightarrow 0.$$

The equality

$$\mathcal{W}[\mathbf{I}_{(s,t]}] = \mathcal{W}[\mathbf{I}_{[0,t]}] - \mathcal{W}[\mathbf{I}_{[0,s]}] = B(t) - B(s)$$

implies that

$$\mathcal{W}[f_n] := \sum_{k \geq 1} c_{kn} (B(t_{kn}) - B(s_{kn})).$$

The map $\mathcal{W}$ is an isometry so $\mathcal{W}[f]$ is the L^2 -limit of $\mathcal{W}[f_n]$. For $0 \leq t_0 < t_1$ we set

$$\int_{t_0}^{t_1} f(s)dB(s) := \mathcal{W}[\mathbf{I}_{[t_0, t_1]}(s)f(s)].$$

The resulting process

$$X_f(t) = \int_0^t f(s)dB(s)$$

is a Gaussian process with independent increments and covariance form

$$K_f(s, t) = \int_0^\infty \mathbf{I}_{[0, t]}(\tau)\mathbf{I}_{[0, t]}(\tau)f(\tau)^2 d\tau = \int_0^{t \wedge s} f(\tau)^2 d\tau.$$

It is thus a martingale with respect to the filtration

$$\mathcal{F}_t^X = \sigma(X(s), 0 \leq s \leq t).$$

One can prove that the process X_f is also a martingale with respect to the Brownian filtration. In particular, it is adapted to the Brownian filtration.

The Gaussian random variable $X(t) - X(s)$, $s < t$, has mean zero and variance

$$V(s, t) = \int_s^t f(\tau)^2 d\tau,$$

so

$$\mathbb{E}[|X(t) - X(s)|^q] = C_q V(s, t)^{q/2}, \quad C_q = (2\pi)^{-1/2} \int_{\mathbb{R}} |x|^q e^{-x^2/2} dx.$$

If $f \in L_{\text{loc}}^{2p}([0, \infty))$, $p > 1$, then $X(t)$ satisfies the assumptions of Kolmogorov's continuity theorem so $X(t)$ admits a continuous modification. The process X_f has a continuous modification even when f is only L^2 , but it takes a bit more work to show this.

The Itô integral extends Wiener's integrals

$$L^2([0, \infty)) \ni f \mapsto \int_0^t f(s)dB(s) = \mathcal{W}[f\mathbf{I}_{[0, t]}]$$

to integrals

$$F \mapsto \mathcal{J}_t[F] = \int_0^t F(s)dB(s),$$

where $F : [0, \infty) \times (\Omega, \mathcal{S}, \mathbb{P}) \rightarrow \mathbb{R}$ is a *stochastic process*. For suitable processes F , the resulting process $t \mapsto \mathcal{J}_t[F]$ is a continuous L^2 -martingale.

You might ask whether this is a mathematician's penchant for generalizations, or there is a deeper motivation. Like all fundamental objects in mathematics, the stochastic integral too is rooted in fundamental problems. To put it briefly, Itô's original motivation was to get a better understanding of diffusion processes. At that time these were being investigated by Feller and Kolmogorov using analytic methods and Itô was seeking an approach closer to the physical intuition. I refer the reader to the very informative historical survey [106] for more details. Future developments cemented the importance, usefulness and versatility of this concept.

This chapter is devoted to explaining the basics of this process and describing a few of its numerous applications.

My presentation is inspired from many sources, but the ones I have consulted frequently are [67, 132, 174, 177, 181, 199, 202, 211].

6.1. Itô's integral with respect to the Brownian motion

To better appreciate some of the sophistication of the modern theory of stochastic integration I think that it helps to first see how this looked in its original form, i.e., Itô's integral of (certain) random processes with respect to the Gaussian white noise on $[0, \infty)$. This approach will be superseded by the more structured approach in the next section due to Kunita and Watanabe. In my experience, the novel ideas in the Kunita-Watanabe approach are best grasped by comparing it with Itô's original approach that we are about to describe. That is why I chose to be a bit repetitive in this chapter.

6.1.1. Itô integrals of progressive processes. Fix a Brownian motion $(B(t))_{t \geq 0}$ defined on a complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Denote by $\mathcal{F}_\bullet^B = (\mathcal{F}_t)_{t \geq 0}$ the completed Brownian filtration

$$\mathcal{F}_t^B = \sigma(B(s), 0 \leq s \leq t) \vee \sigma(\mathcal{N}),$$

where $\mathcal{N} \subset \mathcal{S}$ denotes the collection of $\mathbb{P}$ -negligible events. As shown in Theorem 3.4.2, this filtration is also right-continuous.

Fix a B -admissible (or nonanticipating) filtration $\mathcal{F}_\bullet$; see Definition 3.4.6. Denote by $\mathbb{L}^p = \mathbb{L}^p(\mathcal{F}_\bullet)$ the space of processes

$$X \in L^p([0, \infty) \times \Omega, \mathcal{B}_{[0, \infty)} \otimes \mathcal{S}, \lambda \otimes \mathbb{P}), \quad X = X(t, \omega)$$

that are *progressively measurable* with respect to the filtration $\mathcal{F}_\bullet$. Recall that this means that for any $t \geq 0$, the restriction of X to $[0, t] \times \Omega$ is measurable with respect to the sigma-algebra $\mathcal{B}_{[0, t]} \otimes \mathcal{F}_t$. In particular, the progressive processes are adapted to the filtration $\mathcal{F}_\bullet$. One advantage of the progressive over the simply adapted ones is that integral quantities such that

$$\omega \mapsto \int_0^t f(X(t, \omega)) dt$$

are $\mathcal{F}_t$ -measurable. This is a simple application of Fubini-Tonnelli Theorem 1.3.41.

We denote by $c\mathbb{L}^p = c\mathbb{L}^p(\mathcal{F}_\bullet)$ the subspace of $\mathbb{L}^p$ consisting of processes with *continuous* paths. We set

$$\mathbb{L}^p(0, T) := \{X \in \mathbb{L}^p; \forall t > T \quad X(t) = 0 \text{ a.s.}\}.$$

Note that $\mathbb{L}^p(0, T)$ is a closed subspace of $\mathbb{L}^p$. When $p = 2$, the orthogonal projection onto $\mathbb{L}^2(0, T)$ is given by $X \mapsto \mathbf{I}_{[0, T]} X$. We will frequently identify processes in $\mathbb{L}^p(0, T)$ with processes parametrized by $[0, T]$. Such a process admits an extension by 0 to a process over $[0, \infty)$.

Define

$$c\mathbb{L}^p(0, T) := \mathbf{I}_{[0, T]} c\mathbb{L}^p \subset \mathbb{L}^p(0, T).$$

Thus $X \in \mathbb{L}^p(0, T)$ if $X \in c\mathbb{L}^p$, $X(t) = 0, \forall t > T$ and the paths of X are continuous on $[0, T]$.

Observe that for any $0 \leq s_0 < s_1$, and any $\mathcal{F}_{s_0}$ -measurable *bounded* random variable X , the process

$$(t, \omega) \mapsto X(\omega) \mathbf{I}_{(s_0, s_1]}(t)$$

belongs to $\mathbb{L}^p$. We denote by $\mathcal{E}_{\text{pred}}$ the vector subspace of $\mathbb{L}^p$ spanned by such processes. Note that

$$\mathcal{E}_{\text{pred}} \subset \bigcap_{p \geq 1} \mathbb{L}^p.$$

We will refer to the processes in $\mathcal{E}_{\text{pred}}$ as *elementary predictable processes*. We set

$$\mathcal{E}_{\text{pred}}(0, T) := \mathbf{I}_{[0, T]} \mathcal{E}_{\text{pred}}.$$

Any $X \in \mathcal{E}_{\text{pred}}$ can be described *non-uniquely* as a sum

$$X = \sum_{k=1}^n X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}, \quad (6.1.1)$$

where $0 \leq t_0 < t_1 < \dots < t_n < \infty$, $X_{k-1} \in L^\infty(\Omega, \mathcal{F}_{t_{k-1}}, \mathbb{P})$. We will refer to such a description as *convenient*, and to the points t_k as the *nodes* of the convenient description.

Any convenient description as above can be refined by replacing a term $X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}$ with a sum

$$X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]} \rightarrow X_{k-1} \mathbf{I}_{(t_{k-1}, \tau]} + X_{k-1} \mathbf{I}_{(\tau, t_k]}, \quad \tau \in (t_{k-1}, t_k).$$

Note that since $t_{k-1} < \tau$ we have $X_{k-1} \in \mathcal{F}_\tau$, so both terms in the right-hand side above are progressive. If $X \in \mathcal{E}_{\text{pred}}(0, T)$, then it admits a convenient description (6.1.1) with $t_n = T$.

Using refinements, we can find convenient descriptions of two processes $X, Y \in \mathcal{E}_{\text{pred}}$ with identical nodes

$$0 \leq t_0 < t_1 < \dots < t_n < \infty \quad \text{and} \quad X_{k-1}, Y_{k-1} \in L^2(\Omega, \mathcal{F}_{t_{k-1}}, \mathbb{P})$$

such that

$$X = \sum_{k=1}^n X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}, \quad Y = \sum_{k=1}^n Y_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}. \quad (6.1.2)$$

When this happens we say that the convenient descriptions of X and Y are *compatible*.

If

$$X = \sum_{k=1}^n X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]} \in \mathcal{E}_{\text{pred}}$$

we set

$$\mathcal{J}[X] := \int_0^\infty X(t) dB(t) := \sum_{k=1}^n X_{k-1} (B(t_k) - B(t_{k-1})).$$

I leave to the reader the simple task of verifying that $\mathcal{J}[X]$ is independent of the choice of convenient description of X .

Let us observe that the resulting map

$$\mathcal{J} : \mathcal{E}_{\text{pred}} \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$$

is linear. Indeed, if $X, Y \in \mathcal{E}_{\text{pred}}$, choose compatible convenient descriptions of X and Y to verify that

$$\mathcal{J}[X + Y] = \mathcal{J}[X] + \mathcal{J}[Y].$$

Moreover

$$\forall X \in \mathcal{E}_{\text{pred}}, \quad \mathbb{E}[\mathcal{J}[X]] = 0. \quad (6.1.3)$$

Indeed if

$$X = C \mathbf{I}_{(s, t]}, \quad s < t, \quad C \in L^2(\Omega, \mathcal{F}_s, \mathbb{P}),$$

then $\mathcal{J}[X] = C(B(t) - B(s))$ and

$$\mathbb{E}[\mathcal{J}[X]] = \mathbb{E}\left[\mathbb{E}[C(B(t) - B(s)) \mid \mathcal{F}_s]\right] = \mathbb{E}[B(t) - B(s)] \mathbb{E}[C] = 0.$$

The general equality (6.1.3) now follows from the linearity of $\mathcal{J}$.

Lemma 6.1.1 (Itô isometry). *For any $X \in \mathcal{E}_{\text{pred}}$*

$$\| \mathcal{I}[X] \|_{L^2(\Omega)}^2 = \| X \|_{L^2(\mathbb{R}_{\geq 0} \times \Omega)}^2 = \mathbb{E} \left[\int_0^\infty X(\tau)^2 d\tau \right] = \int_0^\infty \mathbb{E} [X(\tau)^2] d\tau.$$

Proof. Let

$$X = \sum_{k=1}^n X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]} \in \mathcal{E}_{\text{pred}}.$$

Since the mean-zero random variable $B(t_k) - B(t_{k-1})$ is independent of $\mathcal{F}_{t_{k-1}}$ we deduce that for any $j < k$

$$\mathbb{E} \left[(B(t_k) - B(t_{k-1})) \underbrace{X_{k-1} X_{j-1} (B(t_j) - B(t_{j-1}))}_{\mathcal{F}_{t_{k-1}}} \right] = 0.$$

Hence

$$\| \mathcal{I}[X] \|_{L^2}^2 = \sum_{k=1}^n \mathbb{E} [X_{k-1}^2 (B(t_k) - B(t_{k-1}))^2]$$

$$(B(t_k) - B(t_{k-1})) \perp X_{k-1})$$

$$= \sum_{k=1}^n \| X_{k-1} \|_{L^2}^2 \cdot \| B(t_k) - B(t_{k-1}) \|_{L^2}^2 = \sum_{k=1}^n (t_k - t_{k-1}) \| X_{k-1} \|_{L^2}^2.$$

On the other hand, $\forall j < k$,

$$X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]} \cdot X_{j-1} \mathbf{I}_{(t_{j-1}, t_j]} = 0,$$

so

$$\begin{aligned} \| X \|_{L^2(\mathbb{R}_{\geq 0} \times \Omega)}^2 &= \sum_{k=1}^n \int_{t_{k-1}}^{t_k} \left(\int_{\Omega} X_{k-1}(\omega)^2 \mathbb{P}[d\omega] \right) dt \\ &= \sum_{k=1}^n (t_k - t_{k-1}) \| X_{k-1} \|_{L^2}^2. \end{aligned}$$

□

Let $\bar{\mathcal{E}}_{\text{pred}}$ denote the closure of $\mathcal{E}_{\text{pred}}$ in $\mathbb{L}^2$. Since

$$\mathcal{I} : \mathcal{E}_{\text{pred}} \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$$

is an isometry, it extends in a unique fashion to an isometry

$$\mathcal{I} : \bar{\mathcal{E}}_{\text{pred}} \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P}).$$

Theorem 6.1.2 (Approximation theorem).

$$\bar{\mathcal{E}}_{\text{pred}} = \mathbb{L}^2.$$

Proof. The Dominated Convergence Theorem shows that for any $X \in \mathbb{L}^2$

$$\lim_{n \rightarrow \infty} \| \mathbf{I}_{[0, n]} X - X \|_{L^2} = 0$$

so it suffices to show that

$$\forall T > 0, \quad \mathbb{L}^2(0, T) \subset \bar{\mathcal{E}}_{\text{pred}}. \quad (6.1.4)$$

Denote by $b\mathbb{L}^2(0, T)$ the subspace of $\mathbb{L}^2(0, T)$ consisting of *bounded* processes, i.e., processes X such that

$$\exists M > 0, \quad \forall (t, \omega) \in [0, T] \times \Omega, \quad |X_\omega(t)| \leq M$$

Let us observe that $b\mathbb{L}^2(0, T)$ is dense in $\mathbb{L}^2(0, T)$ since $\mathbf{I}_{\{|X| \leq n\}} X \rightarrow X$ in L^2 and $\mathbf{I}_{\{|X| \leq n\}}$ is a progressive process. Hence it suffices to show that

$$\forall T > 0, \quad b\mathbb{L}^2(0, T) \subset \bar{\mathcal{E}}_{\text{prog}}. \quad (6.1.5)$$

Set

$$bc\mathbb{L}^2(0, T) := b\mathbb{L}^2(0, T) \cap c\mathbb{L}^2(0, T).$$

We first show that

$$bc\mathbb{L}^2(0, T) \in \bar{\mathcal{E}}_{\text{prog}}. \quad (6.1.6)$$

For any continuous function $f : [0, T] \rightarrow \mathbb{R}$ we define $S_n[f] : [0, T] \rightarrow \mathbb{R}$ by

$$S_n[f](t) = \sum_{k=1}^n f((k-1)T/n) \mathbf{I}_{((k-1)T/n, kT/n]}.$$

Note that $S_n[f](0) = 0$ and $S_n[f](t) \rightarrow f(t)$, $\forall t \in (0, T]$.

Let $X \in bc\mathbb{L}^2(0, T)$. Set $M = \|X\|_{L^\infty}$. Note that

$$\forall n, \quad S_n[X] \in \mathcal{E}_{\text{pred}}(0, T), \quad \|S_n[X]\|_{L^\infty} \leq M,$$

and

$$\lim_{n \rightarrow \infty} S_n[X_\omega](t) = X_\omega(t), \quad \forall (t, \omega) \in (0, T] \times \Omega.$$

The Dominated Convergence Theorem implies that $S_n[X] \rightarrow X$ in L^2 . This proves (6.1.6). Therefore it suffices to show that $bc\mathbb{L}^2(0, T)$ is dense in $b\mathbb{L}^2(0, T)$.

For any $f \in L^\infty(0, T)$ we define $R_n[f] \in L^\infty(0, T)$,

$$R_n[f](t) = n \int_{(t-1/n)_+}^t f(s) ds.$$

Note that $R_n[f]$ is continuous and $\|R_n[f]\|_{L^\infty} \leq \|f\|_{L^\infty}$. Moreover, the Lebesgue differentiation theorem [207, Prop. 11.11] implies that $R_n[f] \rightarrow f$ a.e. so

$$\|R_n[f] - f\|_{L^2} \rightarrow 0.$$

Let $X \in b\mathbb{L}^2(0, T)$. Since X is progressive, the process $R_n[X]$ is adapted to the filtration $\mathcal{F}_t$. On the other hand, $R_n[X]$ is continuous, and thus it is progressive. Moreover, $\forall \omega \in \Omega$,

$$\int_0^T |R_n[X_\omega](t) - X_\omega(t)|^2 dt \rightarrow 0.$$

Since $\|R_n[X]\|_{L^\infty} \leq \|X\|_{L^\infty}$, the Dominated Convergence Theorem implies that

$$\int_\Omega \left(\int_0^T |R_n[X_\omega](t) - X_\omega(t)|^2 dt \right) \mathbb{P}[d\omega] \rightarrow 0.$$

This proves that $bc\mathbb{L}^2(0, T)$ is dense in $b\mathbb{L}^2(0, T)$ and concludes the proof of Theorem 6.1.2. $\square$

Remark 6.1.3. The same proof shows that $\mathcal{E}_{\text{pred}}$ is dense in $\mathbb{L}^p$, $\forall p \in [1, \infty)$. $\square$

Hence, the map $\mathcal{J} : \mathcal{E}_{\text{pred}} \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$ extends to a linear isometry

$$\mathcal{J} : \mathbb{L}^2 \rightarrow L^2(\Omega, \mathcal{S}, \mathbb{P})$$

called the *Itô integral*. Moreover

$$\forall X \in \mathbb{L}^2, \quad \mathbb{E}[\mathcal{J}[X]] = 0. \quad (6.1.7)$$

We will use the notation

$$\int_0^\infty X(t)dB(t) := \mathcal{I}[X], \quad \forall X \in \mathbb{L}^2.$$

For $t > 0$ and $X \in \mathbb{L}^2$ we set

$$\mathcal{I}_t[X] = \int_0^t X dB := \mathcal{I}[X \mathbf{I}_{[0,t]}] \in L^2(\Omega, \mathcal{F}_t, \mathbb{P}).$$

The above equality describes $\mathcal{I}_t[X]$ *not as a function* on Ω , but as an *equivalence class of functions* on Ω modulo a.s. equality.

Remark 6.1.4. Was all the contortionism in the proof of the Approximation Theorem really necessary? Could have we defined $\int_0^t X dB$ as a limit of Riemann sums over shrinking partitions of $[0, t]$

$$\int_0^t X dB = \lim_{n \rightarrow \infty} \sum_{k=1}^n X(t_{k-1})(B(t_k) - B(t_{k-1}))?$$

As explained in [174, Sec. I.8], the answer is no.

The culprit is the fact that, according to Proposition 2.5.17, the Brownian does not have finite variation. This fact coupled with the Banach-Steinhaus theorem implies that this Riemann sum type approach will not work for *all* X . As we will see a bit later in Corollary 6.1.20, the Riemann sum approximation method does work for a large class of processes X .

This situation is similar with the one encountered the theory of Fourier series. In that case, the Fourier series of a continuous function f need not converge pointwisely to the function. It does so if, e.g., the function f is Lipschitz. $\square$

For any measurable subset $A \subset \mathcal{S}$ we set

$$\mathbb{L}_A^2 := \{ X \in \mathbb{L}^2 : X|_{[0, \infty) \times A} = 0 \}.$$

In other words, $X \in \mathbb{L}_A^2$ iff $X \in \mathbb{L}^2$ and its A -paths are trivial, i.e., for any $\omega \in A$, the path $t \mapsto X_\omega(t)$ is identically zero. We denote by $\mathcal{E}_{\text{pred}, A}$ the vector subspace of $\mathcal{E}_{\text{pred}}$ consisting of processes with vanishing A -paths. Note that if $X \in \mathbb{L}_A^2$, then $\mathbf{I}_{[0, T]}X \in \mathbb{L}_A^2, \forall T > 0$.

Proposition 6.1.5. *If $X \in \mathbb{L}_A^2$, then $\mathbf{I}_A \mathcal{I}[X] = 0$ a.s. on A .*

Proof. The statement is clearly true if $X \in \mathcal{E}_{\text{pred}}$ and $X|_{[0, \infty) \times A} = 0$, then $\mathcal{I}[X] = 0$ a.s. on A .

The linear operators S_n and T_n used in the proof of the Approximation Theorem are path-wise defined so they map processes with vanishing A -paths to processes with vanishing A -paths. This shows that $\mathcal{E}_{\text{pred}, A}$ is dense in $\mathbb{L}_A^2$. The conclusion follows by observing that the linear map

$$\mathbb{L}^2 \ni X \mapsto \mathbf{I}_A X \in L^2(\Omega, \mathcal{S}, \mathbb{P})$$

is continuous and vanishes on of a subspace dense in $\mathbb{L}_A^2$. $\square$

Corollary 6.1.6. *Let $X, Y \in \mathbb{L}^2$ and $A \subset \mathcal{S}$. If*

$$X(t, \omega) = Y(t, \omega), \quad \forall (t, \omega) \in [0, \infty) \times A$$

the $\mathcal{I}[X] = \mathcal{I}[Y]$ a.s. on A . $\square$

6.1.2. Itô's integral as a stochastic process. For $p \in [1, \infty]$ we denote by $\mathbb{L}_{\text{loc}}^p$ the space of measurable processes

$$X : ([0, \infty) \times \Omega, \mathcal{B}_{[0, \infty)} \otimes \mathcal{S}) \rightarrow \mathbb{R}$$

such that $\mathbf{I}_{[0, t]} X \in \mathbb{L}^p, \forall t \geq 0$. In particular, each $X \in \mathbb{L}_{\text{loc}}^p$ is progressively measurable. Note that $\mathcal{E}_{\text{pred}} \subset \mathbb{L}^2 \subset \mathbb{L}_{\text{loc}}^2$.

To each stochastic process $X \in \mathbb{L}_{\text{loc}}^2$ we associate the $\mathcal{F}_\bullet$ -adapted stochastic process

$$\mathcal{J}_t[X] = \int_0^t X dB := \mathcal{J}[\mathbf{I}_{[0, t]} X], \quad \forall t \geq 0.$$

We set

$$\int_s^t X dB := \int_0^\infty \mathbf{I}_{(s, t]}(\tau) X(\tau) dB(\tau) = \mathcal{J}[\mathbf{I}_{(s, t]} X].$$

Since $\mathbf{I}_{[0, t]} = \mathbf{I}_{[0, s]} + \mathbf{I}_{(s, t]}$ we deduce

$$\int_0^t X(\tau) dB(\tau) = \int_0^s X(\tau) dB(\tau) + \int_s^t X(\tau) dB(\tau).$$

Lemma 6.1.7. *Let $Z \in \mathcal{E}_{\text{pred}}$. Then the random process*

$$t \mapsto \mathcal{J}_t[Z] = \int_0^t Z(\tau) dB(\tau)$$

admits a modification as a continuous martingale adapted to the filtration $\mathcal{F}_\bullet$. Moreover, for any $s < t$

$$\mathbb{E}[\mathcal{J}[\mathbf{I}_{(s, t]} Z]^2 \mid \mathcal{F}_s] = \mathbb{E}\left[\int_s^t Z(\tau)^2 d\tau \mid \mathcal{F}_s\right]. \quad (6.1.8)$$

Proof. We can write Z in the form

$$Z = \sum_{k=1}^n Z_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}$$

where $0 = t_0 < t_1, \dots, t_n$, $Z_{k-1} \in L^2(\Omega, \mathcal{F}_{t_{k-1}}, \mathbb{P})$. Then

$$\mathcal{J}_t[Z] = \sum_{k=1}^n Z_{k-1} (B(t_k \wedge t) - B(t_{k-1} \wedge t)) \quad (6.1.9)$$

showing that for any $\omega \in \Omega$

$$t \mapsto \mathcal{J}_t[Z](\omega)$$

is continuous. We will next show that it is also a martingale adapted to the filtration $\mathcal{F}_\bullet$.

Let us observe that the terms in the sum (6.1.9) such that $t_{k-1} \wedge t = t$ are identically zero since $t_k \wedge t = t$ in this case. Thus in the sum (6.1.9) there exists at most one nonzero term such that $t_k \wedge t = t$. This proves that the process is adapted to $\mathcal{F}_t$ since $\mathcal{F}_{t_{k-1}} \subset \mathcal{F}_t$ whenever the interval $(t_{k-1} \wedge t, t_k \wedge t]$ is nonempty.

Let $0 \leq s < t$. Choose $t_{n+1} > t$. Then

$$Z = \sum_{k=1}^n Z_{k-1} \mathbf{I}_{(t_{k-1}, t_k]} + 0 \cdot \mathbf{I}_{(t_n, t_{n+1}]}$$

Using refinements we can produce a convenient description of Z

$$Z = \sum_{k=1}^N Z_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}$$

that has s and t among its nodes $s = t_\ell, t = t_m, \ell < m$. Then

$$\begin{aligned} I_{[0,t]} Z &= \underbrace{\sum_{1 \leq k \leq \ell} Z_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}}_{=: I_{[0,s]} Z} + \underbrace{\sum_{\ell < k \leq m} Z_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}}_{=: I_{(s,t]} Z}, \\ \mathcal{I}_t[Z] &= \mathbb{E}[I_{[0,s]} Z] + \mathcal{I}[I_{(s,t]} Z] = \mathcal{I}_s[Z] \\ &\quad + \sum_{\ell < k \leq m} Z_{k-1} (B(t_k) - B(t_{k-1})). \end{aligned}$$

Hence

$$\mathbb{E}[\mathcal{I}_t[Z] \parallel \mathcal{F}_s] = \mathcal{I}_s[Z] + \mathbb{E}\left[\sum_{\ell < k \leq m} Z_{k-1} (B(t_k) - B(t_{k-1})) \parallel \mathcal{F}_s\right].$$

For $k > \ell$ we have

$$\mathbb{E}[Z_{k-1} (B(t_k) - B(t_{k-1})) \parallel \mathcal{F}_{t_{k-1}}] = Z_{k-1} \mathbb{E}[(B(t_k) - B(t_{k-1})) \parallel \mathcal{F}_{t_{k-1}}] = 0.$$

Since $\mathcal{F}_s \subset \mathcal{F}_{t_{k-1}}$ we deduce

$$\mathbb{E}[Z_{k-1} (B(t_k) - B(t_{k-1})) \parallel \mathcal{F}_s] = 0, \quad \forall k > \ell.$$

The proof of (6.1.8) is similar to the proof of Itô's isometry Lemma 6.1.1. We have

$$\begin{aligned} \int_s^t Z(\tau)^2 d\tau &= \sum_{\ell < k \leq m} Z_{k-1}^2 (t_k - t_{k-1}), \\ \mathbb{E}\left[\int_s^t Z(\tau)^2 d\tau \parallel \mathcal{F}_s\right] &= \sum_{\ell < k \leq m} \mathbb{E}[Z_{k-1}^2 \parallel \mathcal{F}_s] (t_k - t_{k-1}), \\ \mathcal{I}[I_{(s,t]} Z]^2 &= \underbrace{\sum_{k > \ell} Z_{k-1}^2 (B(t_k) - B(t_{k-1}))^2}_{=: T_1} \\ &\quad + 2 \underbrace{\sum_{k > j > \ell} Z_{j-1} Z_{k-1} (B(t_j) - B(t_{j-1})) (B(t_k) - B(t_{k-1}))}_{=: T_2}. \end{aligned}$$

Arguing as in the proof of Lemma 6.1.1 we deduce that $\mathbb{E}[T_2 \parallel \mathcal{F}_2] = 0$ and

$$\mathbb{E}[T_1 \parallel \mathcal{F}_2] = \sum_{\ell < k \leq m} \mathbb{E}[Z_{k-1}^2 \parallel \mathcal{F}_s] (t_k - t_{k-1}).$$

□

Proposition 6.1.8. For any $X \in \mathbb{L}_{\text{loc}}^2$ the stochastic process

$$t \mapsto \mathcal{J}_t[X], \quad t \geq 0$$

is an L^2 -martingale that admits a modification with continuous paths. Moreover, for any $t > 0$

$$\|\mathcal{J}_t[X]^2\| = \int_0^t \mathbb{E}[X_s^2] ds. \quad (6.1.10)$$

Proof. Fix $T > 0$ and sequence of elementary processes $Z_n \in \mathcal{E}_{\text{pred}}$ such that

$$\mathcal{J}[\mathbf{I}_{[0,T]}Z_n] \rightarrow \mathcal{J}[\mathbf{I}_{[0,T]}X] \quad \text{in } L^2.$$

We deduce that

$$\forall t \in [0, T], \quad \mathcal{J}_t[Z_n] = \mathcal{J}[\mathbf{I}_{[0,t]}Z_n] \rightarrow \mathcal{J}[\mathbf{I}_{[0,t]}X] = \mathcal{J}_t[X] \quad \text{in } L^2.$$

In particular, for any $0 \leq s < t$,

$$\mathbb{E}[\mathcal{J}_t[Z_n] \mid \mathcal{F}_s] \rightarrow \mathbb{E}[\mathcal{J}_t[X] \mid \mathcal{F}_s].$$

This proves that $\mathcal{J}_t[X]$ is an L^2 -martingale.

By extracting a subsequence we can assume that

$$\int_0^T \mathbb{E}[(Z_{n+1}(s) - Z_n(s))^2] dt < \frac{1}{2^{n+1}}, \quad \forall n \in \mathbb{N}$$

so

$$\int_0^T \mathbb{E}[(X(s) - Z_n(s))^2] dt < \frac{1}{2^n}, \quad \forall n \in \mathbb{N}.$$

The process $(\mathcal{J}_t[Z_{n+1}] - \mathcal{J}_t[Z_n])_{t \in [0, T]}$ is a continuous L^2 martingale and Doob's maximal inequality (3.5.12) implies

$$\begin{aligned} \mathbb{P}\left[\|\mathcal{J}_\bullet[Z_{n+1}] - \mathcal{J}_\bullet[Z_n]\|_{C([0, T])} > \frac{1}{n^2}\right] &= \mathbb{P}\left[\sup_{0 \leq t \leq T} |\mathcal{J}_t[Z_{n+1}] - \mathcal{J}_t[Z_n]| > \frac{1}{n^2}\right] \\ &= \mathbb{P}\left[\sup_{0 \leq t \leq T} |\mathcal{J}_t[Z_{n+1}] - \mathcal{J}_t[Z_n]|^2 > \frac{1}{n^4}\right] \leq n^4 \mathbb{E}[(\mathcal{J}_T[Z_{n+1}] - \mathcal{J}_T[Z_n])^2] \\ &= n^4 \int_0^T \mathbb{E}[(Z_{n+1}(t) - Z_n(t))^2] dt < \frac{n^4}{2^{n+1}}. \end{aligned}$$

The Borel-Cantelli Lemma implies that

$$\mathbb{P}\left[\|Z_{n+1} - Z_n\|_{C([0, T])} > \frac{1}{n^2} \text{ i.o.}\right] = 0.$$

This shows that the sequence $(\mathcal{J}_t[Z_n])_{t \in [0, T]}$ is a.s. *uniformly* convergent on $[0, T]$.

Thus, the process

$$t \mapsto \mathcal{J}_t[X], \quad t \in [0, T],$$

admits a continuous modification. By choosing a sequence $T_n \nearrow \infty$ we deduce that the process

$$t \mapsto \mathcal{J}_t[X], \quad t \geq 0,$$

admits a continuous modification. □

☞ In the sequel, for any $X \in \mathbb{L}_{\text{loc}}^2$, $\mathcal{I}_t[X]$ will denote a continuous version of the process $\mathcal{I}_t[X]$. Any two such versions are modifications of each other.

Arguing exactly as in the above proposition we obtain the following result.

Corollary 6.1.9. Let $(X_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{L}_{\text{loc}}^2$. Suppose that there exists $T > 0$ and $X \in \mathbb{L}_{\text{loc}}^2$ such that

$$\lim_{n \rightarrow \infty} \int_0^T \mathbb{E}[(X_n(t) - X(t))^2] dt = 0.$$

Then

$$\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T} |\mathcal{I}_t[X_n] - \mathcal{I}_t[X]| = 0 \quad \text{a.s.}$$

□

Example 6.1.10. Observe that $B \in \mathbb{L}_{\text{loc}}^2$ since

$$\int_0^t \mathbb{E}[B(s)^2] ds = \int_0^t \text{Var}[B(s)] ds = \int_0^t s ds = \frac{t^2}{2}.$$

Let us compute

$$\int_0^t B(s) dB(s) := \int_0^\infty \mathbf{I}_{[0,t]} B(s) dB(s).$$

Set

$$X_n := R_n[\mathbf{I}_{[0,t]} B](t) = \sum_{k=1}^n B\left(\frac{(k-1)t}{n}\right) \mathbf{I}_{((k-1)t/n, kt/n]} \in \mathcal{E}_{\text{pred}}.$$

Note that for $\tau \in ((k-1)T/n, kT/n]$ we have $0 \leq \tau - \frac{(k-1)t}{n} \leq \frac{t}{n}$ and

$$\|B(\tau) - X_n(\tau)\|_{L^2}^2 \leq \frac{\tau}{n}.$$

We deduce

$$\int_0^t \|B(s) - X_n(s)\|_{L^2}^2 ds \leq \frac{s}{n} \sum_{k=1}^n \frac{1}{n} = \frac{s}{n}.$$

Hence $X_n \rightarrow \mathbf{I}_{[0,t]} B$ in $L^2(\mathbb{R}_{\geq 0} \times \Omega)$ so that

$$\mathcal{I}[X_n] \xrightarrow{L^2} \int_0^t B(s) dB(s).$$

For simplicity we set $B_k := B(kt/n)$, $k = 0, 1, \dots, n$. We have

$$\mathcal{I}[X_n] = \sum_{k=1}^n B_{k-1} (B_k - B_{k-1}).$$

Using Abel's trick¹ we deduce that

$$\sum_{k=1}^n B_k (B_k - B_{k-1}) = B_n^2 - \sum_{k=1}^n B_{k-1} (B_k - B_{k-1}) = B_n^2 - \mathcal{I}[X_n].$$

Hence

$$B_n^2 - 2\mathcal{I}[X_n] = \sum_{k=1}^n B_k (B_k - B_{k-1}) - \mathcal{I}[X_n] = \sum_{k=1}^n (B_k - B_{k-1})^2.$$

¹I.e., discrete integration by parts.

Note that $B_n^2 = B(t)^2$. Proposition 2.5.17 on the quadratic variation of the Brownian motion shows that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n (B_k - B_{k-1})^2 = t \text{ in } L^2.$$

We deduce that

$$\int_0^t B(s) dB(s) = \frac{1}{2} (B(t)^2 - t). \quad (6.1.11)$$

The process $B(t)^2 - t$ is one of the martingale presented in Example 3.5.3. $\square$

For $p \in [1, \infty)$ we denote by $\mathcal{M}_c^p = \mathcal{M}_c^p(\mathcal{F}_\bullet)$ the space of continuous L^p -martingales $(Y_t)_{t \geq 0}$ adapted to $\mathcal{F}_\bullet$. In other words (Y_t) is a martingale with continuous paths and $Y_t \in L^p(\Omega, \mathcal{F}_t, \mathbb{P})$ for any $t \geq 0$. We denote by $b\mathcal{M}_c^p$ the subspace of $\mathcal{M}_c^p$ consisting of continuous L^p -bounded martingales (Y_t) , i.e.,

$$\sup_{t \geq 0} \|Y_t\|_{L^p} < \infty.$$

The Itô integral induces a map

$$\mathcal{I}_\bullet : \mathbb{L}_{\text{loc}}^2 \rightarrow \mathcal{M}_c^2, \quad X_\bullet \mapsto \mathcal{I}_\bullet[X_\bullet],$$

such that

$$\|\mathcal{I}_t[X]\|_{L^2}^2 = \mathbb{E} \left[\int_0^t |X(s)|^2 ds \right] = \int_0^t \mathbb{E}[X(s)^2] ds, \quad \forall t \geq 0.$$

Moreover, $\mathcal{I}_\bullet[\mathbb{L}^2] \subset b\mathcal{M}_c^2$.

6.1.3. Localization. The process of stochastic integration interacts nicely with the process of localization with respect to a stopping time, leading to rather strong and useful conclusions.

Proposition 6.1.11. *Let $X \in \mathbb{L}_{\text{loc}}^2$ and $T : \Omega \rightarrow [0, \infty]$ be a stopping time adapted to the filtration $\mathcal{F}_\bullet$. Then, the stopped martingale $(\mathcal{I}[X]^T)_t = \mathcal{I}_{t \wedge T}[X]$ is also described by an Itô integral. More precisely, for any $\tau \geq 0$ we have*

$$\int_0^{\tau \wedge T} X dB = \int_0^\tau X \mathbf{I}_{[[0, T]]} dB. \quad (6.1.12)$$

Above, $\mathbf{I}_{[[0, T]]}$ is the predictable stochastic process introduced in Proposition 3.1.23, namely

$$\mathbf{I}_{[[0, T]]}(t, \omega) = \begin{cases} 1, & t \leq T(\omega), \\ 0, & t > T(\omega). \end{cases}$$

Proof. The following proof is inspired from [9, Thm.7.5]. Fix $\tau > 0$. Replacing the stopping time T with $\tau \wedge T$ we can assume $T \leq \tau$.

Step 1. We prove the result for $X \in \mathcal{E}_{\text{pred}}(0, \tau)$ and the stopping time T is has a finite range.

After refining a convenient description of X we can assume that the range of T is among the nodes of this decomposition

$$X = \sum_{k=1}^n X_{k-1} \mathbf{I}_{(t_{k-1}, t_k]}, \quad 0 = t_0 < t_1 < \dots < t_n = \tau.$$

Note that for any $j = 1, \dots, n$ $\{T \geq t_j\} = \Omega \setminus \{T \leq t_{j-1}\} \in \mathcal{F}_{t_{j-1}}$.

$$\begin{aligned} XI_{[[0,T]]}(t, \omega) &= \sum_{j=1}^n I_{[[0,T]]}(t, \omega) X_{j-1}(\omega) I_{(t_{j-1}, t_j]}(t) \\ &= \sum_{j=1}^n \underbrace{I_{\{t_j \leq T\}}(\omega) X_{j-1}(\omega)}_{=: Z_{j-1}(\omega)} I_{(t_{j-1}, t_j]}(t). \end{aligned}$$

This proves that $XI_{[[0,T]]} \in \mathcal{E}_{\text{pred}}(0, \tau)$ and the last term in the above equality is a convenient description.

We have

$$\begin{aligned} \mathcal{J}_T[X](\omega) &= \sum_k I_{\{T=t_k\}}(\omega) \mathcal{J}_{t_k}[X](\omega) \\ &= \sum_k I_{\{T=t_k\}}(\omega) \mathcal{J}_{t_k}[X](\omega) \\ &= \sum_{k=1}^n I_{\{T=t_k\}} \sum_{j=1}^k X_{j-1}(\omega) (B(t_j, \omega) - B(t_{j-1}, \omega)) \\ &= \sum_{j=1}^n \left(\sum_{k \geq j} I_{\{T=t_k\}} \right) X_{j-1}(\omega) (B(t_j, \omega) - B(t_{j-1}, \omega)) \\ &= \sum_{j=1}^n I_{\{t_j \leq T\}} X_{j-1}(\omega) (B(t_j, \omega) - B(t_{j-1}, \omega)) \\ &= \sum_{j=1}^n Z_{j-1}(\omega) (B(t_j, \omega) - B(t_{j-1}, \omega)) = \mathcal{J}_\tau[XI_{[[0,T]]}]. \end{aligned}$$

Step 2. We prove (6.1.12) when $X \in \mathcal{E}_{\text{pred}}(0, t)$, but T is an arbitrary stopping time such that $T \leq \tau$. For $n \in \mathbb{N}$ we set

$$T_n(\omega) := 2^{-n} \lceil 2^n T(\omega) \rceil$$

where $\lceil x \rceil$ is the smallest integer $\geq x$. Thus

$$T_n(\omega) = \frac{k}{2^n} \iff \frac{k-1}{2^n} < T(\omega) \leq \frac{k}{2^n}.$$

We deduce $T_{n+1} \leq T_n$. Since

$$T_n - \frac{1}{2^n} < T \leq T_n,$$

we deduce $T_n \searrow T$ as $n \rightarrow \infty$. Note that T_n is also a stopping time adapted to $\mathcal{F}_\bullet$. Indeed,

$$\{T_n \leq t\} = \{T \leq 2^{-n} \lceil 2^n t \rceil\} \in \mathcal{F}_{2^{-n} \lceil 2^n t \rceil} \subset \mathcal{F}_t.$$

Then

$$\mathcal{J}_{T_n \wedge \tau}[X] = \mathcal{J}_\tau[XI_{[0, T_n]}], \quad \forall n.$$

Since the process $[0, \tau] \ni t \rightarrow \mathcal{J}_t[X]$ is a.s. continuous we deduce

$$\lim_{n \rightarrow \infty} \mathcal{J}_{T_n \wedge \tau}[X] = \mathcal{J}_{T \wedge \tau}[X] \quad \text{a.s.}$$

On the other hand, the Dominated Convergence Theorem implies that

$$XI_{[0, T_n]} I_{[0, \tau]} \rightarrow XI_{[[0, T]]} I_{[0, \tau]} \quad \text{in } \mathbb{L}^2(0, \tau)$$

and Corollary 6.1.9 implies that

$$\mathcal{I}_s[X \mathbf{I}_{[[0, T_n]]}] \rightarrow \mathcal{I}_s[X \mathbf{I}_{[[0, T]]}] \text{ uniformly in } s \in [0, \tau] \text{ a.s..}$$

Hence $\mathcal{I}_{T_n \wedge \tau}[X] = \mathcal{I}_\tau[X \mathbf{I}_{[[0, T_n]]}] \rightarrow \mathcal{I}_\tau[X \mathbf{I}_{[[0, T]]}]$, a.s..

Step 3. The general case. Let $X \in \mathbb{L}_{\text{loc}}^2$ and T a stopping time such that $T \leq \tau$. Choose a sequence $X_n \in \mathcal{E}_{\text{pred}}(0, \tau)$ such that

$$X_n \rightarrow X \mathbf{I}_{[0, \tau]} \text{ in } \mathbb{L}^2(0, \tau).$$

Then

$$X_n \mathbf{I}_{[[0, T]]} \rightarrow X \mathbf{I}_{[[0, T]]} \mathbf{I}_{[0, \tau]} \text{ in } \mathbb{L}^2(0, \tau).$$

We know from Step 2 that $\mathcal{I}_{\tau \wedge T}[X_n] = \mathcal{I}_\tau[X_n \mathbf{I}_{[[0, T]]}]$ and we deduce

$$\mathcal{I}_{\tau \wedge T}[X_n] = \mathcal{I}_\tau[X_n \mathbf{I}_{[[0, T]]}] \rightarrow \mathcal{I}_\tau[X \mathbf{I}_{[[0, T]]}] \text{ in } L^2(\Omega).$$

On the other hand, according to Corollary 6.1.9, $\mathcal{I}_t[X_n] \rightarrow \mathcal{I}_t[X]$ uniformly in $t \in [0, \tau]$ a.s.. We deduce that

$$\mathcal{I}_{t \wedge T}[X_n] \rightarrow \mathcal{I}_{t \wedge T}[X] \text{ a.s..}$$

Thus $\mathcal{I}_{\tau \wedge T}[X] = \mathcal{I}_\tau[X \mathbf{I}_{[0, T]}]$ a.s.. □

One consequence of the above result is the Persistence of Identity Principle, [199, Thm. 6,4].

Corollary 6.1.12 (Persistence of identity principle). *If $X, Y \in \mathbb{L}^2$ and T is a stopping time such that*

$$X(t, \omega) = Y(t, \omega), \text{ if } 0 \leq t \leq T(\omega),$$

then

$$\mathcal{I}_t[X](\omega) = \mathcal{I}_t[Y](\omega),$$

for almost all ω such that $t \leq T(\omega)$.

Proof. For any t we have the a.s. equalities in ω

$$\begin{aligned} \mathcal{I}_{t \wedge T}[X](\omega) &= \int_0^t X(s) \mathbf{I}_{[[0, T]]}(s, \omega) ds \\ &= \int_0^t Y(s) \mathbf{I}_{[[0, T]]}(s, \omega) ds = \mathcal{I}_{t \wedge T}[Y](\omega). \end{aligned}$$

□

Using the above localization result, we will extend the Itô integral to integrands that are not necessarily in $\mathbb{L}_{\text{loc}}^2$.

Definition 6.1.13. For $p \in [1, \infty)$ we denote by $\mathcal{L}^p = \mathcal{L}^p(\mathcal{F}_\bullet)$ the space of $\mathcal{F}_\bullet$ -progressive processes

$$X : [0, \infty) \times \Omega \ni (t, \omega) \mapsto X(t, \omega) \in \mathbb{R}$$

such that

$$\mathbb{P}\left[\left\{\omega; \int_0^\infty |X(t, \omega)|^p dt < \infty\right\}\right] = 1.$$

We denote by $\mathcal{L}_{\text{loc}}^p$ the space of $\mathcal{F}_\bullet$ -progressive processes X such that

$$\forall n \in \mathbb{N}, \quad \mathbb{P}\left[\left\{\omega; \int_0^n |X(t, \omega)|^p dt < \infty\right\}\right] = 1.$$

For $X \in \mathcal{L}_{\text{loc}}^2 = \mathcal{L}_{\text{loc}}^2(\mathcal{F}_\bullet)$ and $t > 0$ we denote by $A_t(X)$ the random variable

$$A_t(X)_\omega := \int_0^t X(t, \omega)^2 dt.$$

□

Observe that

$$\mathbb{L}^2 \subset \mathcal{L}^2, \quad \mathbb{L}_{\text{loc}}^2 \subset \mathcal{L}_{\text{loc}}^2.$$

Remark 6.1.14. The spaces $\mathbb{L}^p$ and $\mathbb{L}_{\text{loc}}^p$ depend on the choice of probability measure $\mathbb{P}$. On the other hand, the space $\mathcal{L}_{\text{loc}}^p$ does not change if we replace $\mathbb{P}$ with an equivalent probability measure $\mathbb{P}'$, i.e., $\mathbb{P}[S] = 0$ iff $\mathbb{P}'[S] = 0$. □

Lemma 6.1.15. Let $X \in \mathcal{L}_{\text{loc}}^p$, $1 \leq p < \infty$. Then there exists a sequence of stopping times (T_n) with the following properties

- (i) $T_n \leq T_{n+1}$, $\forall n$.
- (ii) $T_n \rightarrow \infty$ as $n \rightarrow \infty$,
- (iii) For any $n \in \mathbb{N}$, $\mathbf{I}_{[0, T_n]} X \in \mathbb{L}^2$.

$$\int_{\Omega} \left(\int_0^{T_n(\omega)} |X(t, \omega)|^p dt \right) \mathbb{P}[d\omega] < \infty.$$

A sequence (T_n) with above properties is called a localizing sequence for X .

Proof. Define

$$T_n(\omega) = n \wedge \inf \left\{ t \geq 0; \int_0^t |X(s)|^p ds \geq n \right\},$$

where we use the convention $\inf \emptyset = \infty$. Clearly $T_n \leq T_{n+1}$ and

$$\int_0^{T_n(\omega)} |X(t, \omega)|^p dt \leq n \quad \text{a.s..}$$

Since $X \in \mathcal{L}^{p, \text{loc}}$ we deduce $T_n \nearrow \infty$ a.s.. □

Proposition 6.1.16. Let $X \in \mathcal{L}_{\text{loc}}^2$ and suppose that (T_n^1) and (T_n^2) are localizing sequence. For $n \in \mathbb{N}$ define the continuous L^2 -martingales

$$Z_n^i(t) := \int_0^t X \mathbf{I}_{[0, T_n^i]}(s) ds, \quad i = 1, 2.$$

Then there exists a continuous process $Z_\infty(t) \in \mathcal{L}_{\text{loc}}^2$, adapted to the filtration $\mathcal{F}_\bullet$, and a negligible set $\mathcal{N} \subset \mathcal{S}$ such that, $\forall \omega \in \Omega \setminus \mathcal{N}$, $\forall i \in \{1, 2\}$, and any $t \geq 0$.

$$\lim_{n \rightarrow \infty} Z_n^i(t, \omega) = Z_\infty(t, \omega).$$

Proof. Assume first that $T_n^1 \leq T_n^2$, $\forall n$. Fix a negligible set $\mathcal{N} \subset \mathcal{S}$ such that $Z_n^i(-, \omega)$ are continuous for any $\omega \in \Omega \setminus \mathcal{N}$ and any $i \in \{1, 2\}$. For $\omega \in \mathcal{N}$ define $Z_\infty^i(t, \omega) = 0$. For $\omega \in \Omega \setminus \mathcal{N}$ and $t \geq 0$ define

$$Z_\infty^i(t, \omega) := Z_n^i(t, \omega),$$

for some n such that $T_n^i(\omega) > t$. Note that if $n > m$ and $T_m^i(\omega) > t$, then

$$Z_m^i(t) = Z_n^i(t \wedge T_m^i),$$

and the Persistence of Identity Principle (Corollary 6.1.12) implies that

$$Z_m^i(t, \omega) = Z_n^i(t, \omega), \quad T_n^i(\omega) \geq T_m^i(\omega) > t.$$

Thus $Z_\infty^i(t)$ is well defined. Clearly $Z_\infty^i(t)$ has continuous paths and

$$\lim_{n \rightarrow \infty} Z_n^i(t, \omega) = Z_\infty^i(t, \omega), \quad \forall t \in [0, \infty) \times \mathcal{N}^c.$$

Since $T_n^1 \leq T_n^2$,

$$Z_n^1(t) = Z_n^2(t) \quad \text{if } t < T_n^1(\omega) \leq T_n^2(\omega).$$

Hence in this case $Z_\infty^1 = Z_\infty^2$.

To prove the equality in general consider the localizing sequence $T_n^0 = T_n^1 \wedge T_n^2$, define Z_∞^0 as above and conclude as before that $Z_\infty^0 = Z_\infty^1 = Z_\infty^2$. $\square$

Definition 6.1.17. Let $X \in \mathcal{L}_{\text{loc}}^2$. The continuous process Z_∞ constructed in Proposition 6.1.16 is called the *Itô integral* of X and it is denoted by

$$\int_0^t X(s) dB(s).$$

$\square$

For any $X \in \mathcal{L}_{\text{loc}}^2$ and any $t > 0$ we set

$$\mathcal{A}_t(X) := \int_0^t X(s)^2 ds.$$

Theorem 6.1.18 (Approximation of Itô integrals). *Let*

$$(X_n), \quad X_\infty \in \mathcal{L}_{\text{loc}}^2$$

such that, for any $t > 0$

$$\mathcal{A}_t(X_n - X_\infty) := \int_0^t (X_n(s) - X_\infty(s))^2 ds \rightarrow 0, \quad \text{in probability.}$$

Then

$$\int_0^t X_n(s) dB(s) \rightarrow \int_0^t X_\infty(s) dB(s), \quad \text{in probability.} \quad (6.1.13)$$

Proof. The proof is inspired from [9, Prop.7.3]. Replacing X_n by $X_n - X_\infty$ we can assume that $X_\infty = 0$. The desired conclusion is a consequence of the following technical result.

Lemma 6.1.19. *Let $X \in \mathcal{L}_{\text{loc}}^2$ and set*

$$Y(t) := \int_0^t X(s) dB(s).$$

Then for any $t, \varepsilon, \delta > 0$, we have

$$\mathbb{P}[|Y(t)| > \varepsilon] \leq \mathbb{P}[\mathcal{A}_t(X) > \delta] + \frac{\delta}{\varepsilon^2}. \quad (6.1.14)$$

Proof. Let

$$T_\delta := \inf \{ t \geq 0; \mathcal{A}_t(X) > \delta \}.$$

We have

$$\begin{aligned} \mathbb{P}[|Y(t)| > \varepsilon] &= \mathbb{P}[|Y(t)| > \varepsilon, T_\delta < t] + \mathbb{P}[|Y(t)| > \varepsilon, T_\delta \geq t] \\ &\leq \mathbb{P}[T_\delta < t] + \mathbb{P}[|Y(t)| > \varepsilon, T_\delta \geq t] \\ &\leq \mathbb{P}[\mathcal{A}_t(X) > \delta] + \mathbb{P}[|Y(t)| > \varepsilon, T_\delta \geq t]. \end{aligned}$$

Observe that

$$\begin{aligned} \mathbb{P}[|Y(t)| > \varepsilon, T_\delta \geq t] &= \mathbb{P}[Y(t)^2 > \varepsilon^2, T_\delta \geq t] \\ &= \mathbb{P}\left[\left|\int_0^t X(s) \mathbf{I}_{[0, T_\delta]} ds\right|^2 > \varepsilon^2\right] \end{aligned}$$

(use Markov's inequality)

$$\leq \frac{1}{\varepsilon^2} \left\| \int_0^t X(s) \mathbf{I}_{[0, T_\delta]} ds \right\|_{L^2(\Omega)}^2.$$

The punchline is the fact that $X(s) \mathbf{I}_{[0, T_\delta]} \in \mathbb{L}^2$ and we can use Itô's isometry to conclude that

$$\left\| \int_0^t X(s) \mathbf{I}_{[0, T_\delta]} ds \right\|_{L^2(\Omega)}^2 = \mathbb{E} \left[\underbrace{\int_0^t (X(s) \mathbf{I}_{[0, T_\delta]}(s))^2 ds}_{=A_{t \wedge T_\delta}(X)} \right] \leq \delta.$$

□

We can now complete the proof of (6.1.13). Set

$$Y_n(t) = \int_0^t X_n(s) dB(s).$$

We have to show that $Y_n(t) \rightarrow 0$ in probability. Fix $\varepsilon > 0$, $k \in \mathbb{N}$, and set $\delta = \delta_k(\varepsilon) = 2^{-k-1}\varepsilon^2$.

Since $A_t(X_n) \rightarrow 0$ in probability there exists $N = N_k(\varepsilon) > 0$ such that,

$$\mathbb{P}[A_t(X_n) > \delta_k(\varepsilon)] < 2^{-k-1}, \quad \forall n \geq N_k(\varepsilon).$$

Using (6.1.14) we deduce that

$$\forall n \geq N_k(\varepsilon), \quad \mathbb{P}[|Y_n(t)| > \varepsilon] \leq 2^{-k-1} + \frac{\delta_k(\varepsilon)}{\varepsilon^2} = 2^{-k}.$$

Hence, for any $k \in \mathbb{N}$

$$0 \leq \liminf_{n \rightarrow \infty} \mathbb{P}[|Y_n(t)| > \varepsilon] \leq \limsup_{n \rightarrow \infty} \mathbb{P}[|Y_n(t)| > \varepsilon] \leq 2^{-k}.$$

□

Corollary 6.1.20. Suppose that $X : [0, \infty) \times \Omega \rightarrow \mathbb{R}$ is a process with continuous paths adapted to the filtration $\mathcal{F}_\bullet$. For any partition $\mathbf{P} = (t_0, \dots, t_N)$ of $[0, T]$ we set

$$X(\mathbf{P}) := \sum_{k=1}^N X(t_{k-1}) \mathbf{I}_{(t_{k-1}, t_k]},$$

$$\mathcal{R}_X(\mathbf{P}) = \sum_{k=1}^N X(t_{k-1}) (B(t_k) - B(t_{k-1})).$$

If $(\mathbf{P}_n)$ is a sequence of partitions of $[0, T]$ such that $\|\mathbf{P}_n\| \rightarrow 0$, then

$$\mathcal{R}_X(\mathbf{P}_n) \rightarrow \int_0^T X(s)dB(s) \text{ in probability.}$$

Proof. Since X has continuous paths we deduce that

$$\mathcal{A}_T(X(\mathbf{P}_n) - X) \rightarrow 0 \text{ a.s.}$$

and thus in probability. The claim now follows from Theorem 6.1.18. $\square$

6.1.4. Local martingales. Processes produced by Itô integrals of processes in $\mathcal{L}_{\text{loc}}^2$ have special features. This subsection describes a few of them

Definition 6.1.21. A *continuous local martingale* adapted to $\mathcal{F}_\bullet$ is a continuous process $X(t)$, $t \geq 0$, adapted to $\mathcal{F}_\bullet$, such that there exists a sequence of stopping times $(T_n)_{n \in \mathbb{N}}$ with the following properties

- (i) $T_N \leq T_{n+1}$, $\forall n$,
- (ii) as such that $T_n \nearrow \infty$ as $n \rightarrow \infty$ and,
- (iii) the stopped processes $X^{T_n}(t) - X_0 = X(t \wedge T_n) - X_0$ are UI martingales.

Such a sequence of stopping times is called a *localizing sequence* for the local continuous martingale and its said to *reduce* the local martingale. We denote by $\mathcal{M}^{c, \text{loc}}$ the space of continuous local martingales. $\square$

Remark 6.1.22. The UI requirement in (iii) of the above definition can be dropped. If T_n is an increasing sequence of stopping times such that $T_n \nearrow \infty$ and X^{T_n} is a continuous martingale, then the sequence $T'_n = n \wedge T_n$ is localizing in the sense of Definition 6.1.21. Indeed, each T'_n is a bounded stopping time and thus the stopped processes $X^{T'_n}$ are UI. $\square$

Example 6.1.23. (a) The continuous martingales are also continuous local martingales. To see this note that the stopping times $T_n = n$ form a localizing sequence.

(b) The Itô integral of a process $X \in \mathcal{L}_{\text{loc}}^2$ is a continuous local martingale.

(c) If X is a continuous local martingale, then for any stopping time T , the stopped process X^T is also a continuous local martingale. $\square$

Proposition 6.1.24. Let $X = (X(t))_{t \geq 0}$ be continuous local martingale such that $X(0) \in L^1$.

- (i) If $X(t) \geq 0$, $\forall t \geq 0$, then X is a supermartingale.
- (ii) If there exists $Z \in L^1$ such that $|X(t)| \leq Z$, $\forall t$, then X is an UI-martingale.
- (iii) The sequence of stopping times

$$T_n = \inf \{ t \geq 0; |X(t)| \geq n \}$$

is a localizing sequence. The stopped martingales X^{T_n} are bounded i.e.,

$$\sup_{\substack{t \geq 0, n \in \mathbb{N}, \\ \omega \in \Omega}} |X_{T_n(\omega) \wedge t}(\omega)| < \infty.$$

In particular, they are continuous L^2 -bounded martingales.

Proof. I follow the approach in [132, Prop. 4.7]

(i) Set $Y(t) := X(t) - X(0)$. Choose a localizing sequence T_n . Then, $\forall s \leq t$ we have

$$Y[(s \wedge T_n)] = \mathbb{E}[Y[(t \wedge T_n) \mid \mathcal{F}_s]].$$

Since $X(0) \in L^1(\Omega, \mathcal{F}_s, \mathbb{P})$, $\forall s \geq 0$, we deduce

$$X[(s \wedge T_n)] = \mathbb{E}[X(t \wedge T_n) \mid \mathcal{F}_s]$$

Letting $n \rightarrow \infty$ and invoking the continuity of X and the conditional Fatou lemma Theorem 1.4.12(vii) we deduce

$$X(s) \geq \mathbb{E}[X(t) \mid \mathcal{F}_s]$$

(ii) Arguing as in (i) we deduce

$$X(s \wedge T_n) = \mathbb{E}[X(t \wedge T_n) \mid \mathcal{F}_s]$$

Using the conditional Dominated Convergence Theorem Theorem 1.4.12(viii) we let $n \rightarrow \infty$ and we deduce

$$X(s) = \mathbb{E}[X(t) \mid \mathcal{F}_s]$$

Thus X is a martingale. It is UI since it is dominated by Z .

(iii) In this case we have $|Y^{T_n}(t)| \leq n$ so Y^{T_n} is an UI martingale. The stopped process X^{T_n} is dominated by $n + |X(0)|$ so it is a UI martingale. $\square$

6.1.5. Finite variation processes. We will soon be forced to deal with finite variation processes and I need to mention a few facts about them and their connections to local martingales.

We begin by recalling a few facts of classical real analysis.

Definition 6.1.25. Let $T \in (0, \infty)$.

(i) Given $a : [0, T] \rightarrow \mathbb{R}$ and $t \in (0, T]$ we define the *total variation* of a on $[0, t]$ to be

$$V(a, t) := \sup_{\underline{t}} \sum_{i=1}^m |a(t_i) - a(t_{i-1})|,$$

where the supremum is over the partitions

$$P = \{0 = t_0 < t_1 < \cdots < t_{k-1} < t_k = t\}$$

of $[0, t]$.

(ii) A function $a : [0, T] \rightarrow \mathbb{R}$ is said to have *bounded variation* if $V(a, T) < \infty$. We denote by $BV([0, T])$ the space of R -functions of bounded variation on $[0, T]$.

(iii) A function $a : [0, \infty) \rightarrow \mathbb{R}$ is said to have *finite* or *locally bounded* variation if $V(a, T) < \infty$ for any $T > 0$.

$\square$

To $a \in BV([0, T])$ we associate a signed Borel measure μ_a defined

$$a(t) = \mu_a([0, t]).$$

The measure μ_a admits a canonical decomposition

$$\mu_a = \mu_a^+ - \mu_a^-,$$

where $\mu_a^\pm$ are positive measures supported on disjoint Borel subsets of $[0, T]$. If we set

$$|\mu_a| := \mu_a^+ + \mu_a^-,$$

then

$$|\mu_a|([0, t]) = \mathbf{V}(a, t).$$

The traditional notation is

$$da := \mu_a, \quad |da| := |\mu_a|.$$

The following result is classical. For a proof we refer to [13, §14.3].

Proposition 6.1.26. *If $a \in BV([0, T])$, and $f : [0, T] \rightarrow \mathbb{R}$ is measurable and*

$$\int_0^T |f(s)| |da(s)| < \infty,$$

then the function

$$t \mapsto \int_0^t f(s) da(s)$$

is continuous and has bounded variation. $\square$

Proposition 6.1.27. *Suppose that $f : [0, T] \rightarrow \mathbb{R}$ is continuous and $a \in BV([0, T])$. Fix $t \in (0, T]$. If*

$$P^\nu = \{0 = t_0 < t_1 < \cdots < t_{k_\nu} = t\}$$

is a sequence of partitions of $[0, t]$ with mesh converging to 0 as $\nu \rightarrow \infty$, then

$$\int_0^t f(s) da(s) = \lim_{\nu \rightarrow \infty} \sum_{j=1}^{k_\nu} f(t_{j-1}) (a(t_j) - a(t_{j-1})).$$

$\square$

Definition 6.1.28. Fix a filtered probability space $(\Omega, (\mathcal{F}_t)_{t \geq 0}, \mathcal{F}, \mathbb{P})$. A *finite variation process* is a process $(A_t)_{t \geq 0}$ adapted to the filtration $\mathcal{F}_t$ such that

- $A_0 \equiv 0$ and
- all its sample paths are continuous functions with finite variation on $[0, \infty)$.

We will denote by $\mathcal{V}$ or $\mathcal{V}(\mathcal{F}_\bullet)$ the space of finite variation processes adapted to $\mathcal{F}_\bullet$. If additionally the sample paths of A_t are nondecreasing, then we say that the process is nondecreasing. $\square$

Note that if $(A_t)_{t \geq 0}$ is a finite variation process, then its variation

$$V_t = \int_0^t |dA_s|$$

is a nondecreasing process. Writing

$$A_t := \frac{1}{2}(A_t + V_t) - \frac{1}{2}(V_t - A_t)$$

we see that any finite variation process is the difference of two nondecreasing processes.

Proposition 6.1.29. *Let $A = (A_t)_{t \geq 0}$ be a finite variation process and $H = (H_t)_{t \geq 0}$ be a progressive process such that*

$$\forall t \geq 0, \forall \omega \in \Omega : \int_0^t |H_s(\omega)| \cdot |dA_s(\omega)| < \infty. \quad (6.1.15)$$

Then the process $H \cdot A$ defined by

$$(H \cdot A)_t(\omega) := \int_0^t H_s(\omega) dA_s(\omega) \quad (6.1.16)$$

is also a finite variation process.

Proof. Proposition 6.1.26 implies that $H \cdot A$ is a finite variation process. We only have to show that it is adapted to the filtration. For this it suffices to show that if $h : \Omega \times [0, 1]$ is measurable with respect to the product σ -algebra $\mathcal{F}_t \otimes \mathcal{B}_{[0,t]}$ and if

$$\int_0^t |h(\omega, s)| |dA_s(\omega)| < \infty, \quad \forall \omega \in \Omega, /$$

then the random variable

$$\omega \mapsto \int_0^t h(\omega, s) dA_s(\omega)$$

is $\mathcal{F}_t$ -measurable. Indeed, iff

$$h(\omega, s) = \mathbf{I}_{(u,v]}(s) \mathbf{I}_\Gamma(\omega), \quad \Gamma \in \mathcal{F}_t$$

then the claim is obvious: the integral $\int_0^t h(\omega, s) dA_s(\omega)$ is a limit of $\mathcal{F}_t$ -measurable functions. A monotone class argument then allows us to extend the claim to functions of the form $f(\omega, s) = \mathbf{I}_G(\omega, s)$, $G \in \mathcal{F}_t \otimes \mathcal{B}_{[0,t]}$. This implies the general case since any $\mathcal{F}_t \otimes \mathcal{B}_{[0,t]}$ -measurable function is a pointwise limit of step functions h_n such that $|h_n| \leq |h|$. Then

$$\lim_{n \rightarrow \infty} \int_0^t h_n(\omega, s) dA_s(\omega) = \int_0^t h(\omega, s) dA_s(\omega),$$

by the dominated convergence theorem. □

Remark 6.1.30. If the filtration $\mathcal{F}_t$ is complete we can replace the assumption (6.1.15) by the weaker one: for any

$$\forall t > 0, \int_0^t |h(\omega, s)| |dA_s(\omega)| < \infty \quad \text{a.s.}$$

□

Proposition 6.1.31 (Associativity). *Suppose that A is a finite variation process and H^1, H^2 are two progressive processes such that*

$$\forall t \geq 0, \forall \omega \in \Omega : \int_0^t |H_s^i(\omega)|^2 \cdot |dA_s(\omega)| < \infty, \quad i = 1, 2. \quad (6.1.17)$$

Then

$$H^1 \cdot (H^2 \cdot A) = (H^1 H^2) \cdot A$$

Proof. Clearly it suffices to consider only the deterministic case (when Ω consists of a single point) and A is non-increasing so that $|dA| = dA$. Denote by μ the (random) signed Borel measure dA so that

$$A(t) = \mu[[0, t]].$$

The signed measure $d(H^2 \cdot A)_2$ is $H^2(s)\mu[ds]$ so

$$\begin{aligned} H^1 \cdot (H^2 \cdot A) &= \int_{[0, t]} H^1(s) d(H^2 \cdot A)(s) \\ &= \int_{[0, 1]} H^1(s) H^2(s) \mu[ds] = (H^1 H^2) \cdot A_t. \end{aligned}$$

□

Proposition 6.1.32. *Let Y be a continuous local martingale adapted to the filtration $\mathcal{F}_\bullet$ that satisfies the usual conditions. If $Y(t)$ is also bounded variation process such that $Y(0) = 0$, then $\mathbb{P}[\forall t, Y(t) = 0] = 1$.*

Proof. Set

$$T_n := \inf \left\{ t \geq 0; \int_0^t |dY_s| \geq n \right\}.$$

We deduce from Example 3.3.11 applied to the process $V(Y, t)$ that T_n is a stopping time. Fix $n \in \mathbb{N}$ and set

$$N := M^{T_n}.$$

Then N is a local martingale starting at 0 satisfying

$$\int_0^\infty |dN_s| \leq n.$$

In particular,

$$|N_t| \leq \int_0^t |dN_s| \leq n, \quad \forall t \geq 0.$$

We deduce from Proposition 6.1.24 that N is a UI martingale. Fix a partition

$$P = \{0 = t_0 < t_1 < \dots < t_p = t\}$$

of $[0, t]$. Let us observe that for any $0 < i < j$ we have

$$\begin{aligned} \mathbb{E}[(N_{t_j} - N_{t_{j-1}})(N_{t_i} - N_{t_{i-1}})] &= \mathbb{E}[\mathbb{E}[(N_{t_j} - N_{t_{j-1}})(N_{t_i} - N_{t_{i-1}}) \mid \mathcal{F}_{t_i}]] \\ &= \mathbb{E}[(N_{t_i} - N_{t_{i-1}}) \underbrace{\mathbb{E}[(N_{t_j} - N_{t_{j-1}}) \mid \mathcal{F}_{t_i}]}_{=0}] = 0. \end{aligned}$$

We deduce

$$\begin{aligned} \mathbb{E}[N_t^2] &= \sum_{i=1}^p \mathbb{E}[(N_{t_i} - N_{t_{i-1}})^2] \leq \sum_{i=1}^p \mathbb{E}[(\sup_{1 \leq i \leq p} |N_{t_i} - N_{t_{i-1}}|) |N_{t_i} - N_{t_{i-1}}|] \\ &\leq n \mathbb{E} \underbrace{\sup_{1 \leq i \leq p} [|N_{t_i} - N_{t_{i-1}}|]}_{=: \Delta(N, P)} \end{aligned}$$

Now choose a sequence of partitions P^ν with mesh going to zero. Since N is continuous, it is uniformly continuous on $[0, t]$ and thus

$$\lim_{\nu \rightarrow \infty} \Delta(N, P^\nu) = 0.$$

Since N is bounded we deduce from the dominated convergence theorem that

$$\lim_{\nu \rightarrow \infty} \mathbb{E}[\Delta(N, P)] = 0$$

so that $N_t = 0$, i.e., $Y_{t \wedge T_n} = 0$. Letting $n \rightarrow \infty$ we deduce $Y_t = 0$. $\square$

6.1.6. Itô calculus. Suppose that $B^1, \dots, B^{N_0}$ are independent Brownian motions defined on the probability space $(\Omega, \mathcal{S}, \mathbb{P})$. Set

$$\mathbf{B} : [0, \infty) \times \Omega \rightarrow \mathbb{R}^{N_0}, \quad \mathbf{B}(t, \omega) = (B^1(t, \omega), \dots, B^{N_0}(t, \omega)),$$

$$\mathcal{F}_t^{\mathbf{B}} := \bigvee_{k=1}^{N_0} \mathcal{F}_t^{B^k}, \quad t \geq 0.$$

Fix a filtration $\mathcal{F}_\bullet$ that is non-anticipating (Definition 3.4.6) with respect to each of the Brownian motions B^k , $1 \leq k \leq N_0$. Thus

$$\mathcal{F}_t \supset \mathcal{F}_t^{\mathbf{B}}, \quad \forall t \geq 0,$$

and, $\forall t > s$, $(\mathbf{B}(t) - \mathbf{B}(s)) \perp \mathcal{F}_s$.

Definition 6.1.33. A $\mathcal{F}_\bullet$ -adapted vector-valued process

$$\mathbf{X} = (X^\alpha(t))_{1 \leq \alpha \leq N_1}$$

is said to be an *Itô process* with respect to $\mathbf{B}$ if there exist

$$F^\alpha \in \mathcal{L}_{\text{loc}}^1(\mathcal{F}_\bullet), \quad G_j^\alpha \in \mathcal{L}_{\text{loc}}^2(\mathcal{F}_\bullet), \quad 1 \leq \alpha \leq N_1, \quad 1 \leq j \leq N_0$$

such that

$$\begin{aligned} X^\alpha(t) &= X^\alpha(0) + \int_0^t F^\alpha(s) ds \\ &+ \sum_{j=1}^{N_0} \int_0^t G_j^\alpha(s) dB^j(s), \quad \forall 1 \leq \alpha \leq N_1. \end{aligned} \tag{6.1.18}$$

We write this in compact form

$$\mathbf{X}(t) = \mathbf{X}(0) + \mathcal{J}_t[\mathbf{F}, \mathbf{G}].$$

Let us observe that an Itô process is the sum of a continuous finite variation process and a continuous local martingale. We formally define the *stochastic differential* of the Itô process X^α to be

$$dX^\alpha = F^\alpha(s)ds + \sum_{j=1}^{N_0} G_j^\alpha(s)dB^j(s), \quad \forall 1 \leq \alpha \leq N_1.$$

$\square$

Remark 6.1.34. The stochastic differential is expressed in terms of multidimensional Brownian motion $\mathbf{B} = (B^1, \dots, B^{N_0})$. The Itô isometry and Proposition 6.1.32 show that the stochastic differential is well defined, i.e., independent of the description of $\mathbf{X}$ as a sum between a finite variation process and an Itô integral.

More precisely, if $\mathbf{X}$, $\hat{\mathbf{X}}$ are Itô processes defined by two equalities of the type (6.1.18),

$$\mathbf{X} = \mathbf{X}(0) + \mathcal{I}_t[\mathbf{F}, \mathbf{G}] \quad \text{and} \quad \hat{\mathbf{X}} = \hat{\mathbf{X}}(0) + \mathcal{I}_t[\hat{\mathbf{F}}, \hat{\mathbf{G}}],$$

then

$$\hat{\mathbf{X}}(t) - \mathbf{X}(t) = \hat{\mathbf{X}}(0) - \mathbf{X}(0), \quad \forall t \geq 0 \iff \mathbf{F} = \hat{\mathbf{F}}, \quad \mathbf{G} = \hat{\mathbf{G}},$$

and $d\mathbf{X} = \mathbf{F}dt + \mathbf{G}dB$. I will refer to the contribution $\mathbf{G}dB$ as the *noisy part of the stochastic differential*. $\square$

Theorem 6.1.35 (Itô's formula). *Suppose that $\mathbf{X} = (X^\alpha(t))_{1 \leq \alpha \leq N_1}$ is a $\mathcal{F}_\bullet$ -adapted Itô process with respect to $\mathbf{B} = (B^1, \dots, B^{N_0})$,*

$$\mathbf{X}(t) = \mathbf{X}(0) + \mathcal{I}_t[\mathbf{F}, \mathbf{G}].$$

Then, for any $u : [0, \infty) \times \mathbb{R}^{N_1} \rightarrow \mathbb{R}$ such that $u(t, \mathbf{x})$ is C^1 in the t -variable and C^2 in the $\mathbf{x}$ -variable, the process $u(t, \mathbf{X}(t))$ is also an Itô process with respect to $(B^1, \dots, B^{N_0})$. More precisely, we have Itô's formula

$$\begin{aligned} u(t, \mathbf{X}(t)) &= u(0, \mathbf{X}(0)) \\ &+ \sum_{j=1}^{N_0} \int_0^t \underbrace{\left(\sum_{\alpha=1}^{N_1} u'_{x_\alpha}(s, \mathbf{X}) G_j^\alpha \right)}_{=: \hat{G}_j} dB^j(s) \\ &+ \underbrace{\int_0^t \left(u'_s(s, \mathbf{X}) + \sum_{\alpha=1}^{N_1} u'_{x_\alpha}(s, \mathbf{X}) F^\alpha + \frac{1}{2} \sum_{j=1}^{N_0} \sum_{\alpha, \beta=1}^{N_1} u''_{x_\alpha x_\beta}(s, \mathbf{X}) G_j^\alpha G_j^\beta \right)}_{=: \hat{F}} ds. \end{aligned} \tag{6.1.19}$$

Hence

$$du(t, \mathbf{X}(t)) = \hat{F}ds + \sum_j \hat{G}_j dB_j.$$

Proof. I will prove (6.1.19) gradually, in increasing degree of generality. The approach is a blend of the approaches in [202, Sec. 3.3] and [211, Sec. 4.4]. Fix $t > 0$. We set

$$\mathbf{F}(t) := (F^\alpha(t))_{1 \leq \alpha \leq N_1}, \quad \mathbf{G}(t) := (G_j^\alpha(t))_{1 \leq j \leq N_0, 1 \leq \alpha \leq N_1},$$

and we will denote by $\mathcal{R}_t(u, \mathbf{F}, \mathbf{G})$ the right-hand side of (6.1.19),

Step 1. We prove (6.1.19) in the special case when $F^\alpha, G_j^\alpha \in \mathcal{E}_{\text{pred}}$ and $u \in C_b^3([0, \infty) \times \mathbb{R}^{N_1})$. Analytically, this is the most demanding step.

We can assume that the elementary processes $\mathbf{I}_{[0, t]} F^\alpha, \mathbf{I}_{[0, t]} G_j^\alpha$ admit convenient descriptions with the same sets of nodes

$$0 = t_0 < t_1 < \dots < t_N = t.$$

For every $\nu \in \mathbb{N}$, $k = 0, \dots, N-1$, $n = 0, 1, \dots, 2^\nu$ we set

$$t_{k,n}^\nu := t_k + n \frac{t_{k+1} - t_k}{2^\nu}.$$

Note that $t_k = t_{k,0}^\nu = t_{k-1,2^\nu}^\nu$. Geometrically, the points $t_{k,n}^\nu$ are obtained by dividing the interval $[t_k, t_{k+1}]$ into 2^ν equal parts. We have

$$\begin{aligned} u(t, \mathbf{X}(t)) - u(0, \mathbf{X}(0)) &= \sum_{k=1}^N \left(u(t_k, \mathbf{X}(t_k)) - u(t_{k-1}, \mathbf{X}(t_{k-1})) \right) \\ &= \sum_{k=1}^N \left(\underbrace{\sum_{n=1}^{2^\nu} \left(u(t_{k-1,n}^\nu, \mathbf{X}(t_{k-1,n}^\nu)) - u(t_{k-1,n-1}^\nu, \mathbf{X}(t_{k-1,n-1}^\nu)) \right)}_{=: S_{k,\nu}} \right). \end{aligned}$$

We investigate the behavior of $S_{k,\nu}$ for k fixed and $\nu \rightarrow \infty$. To simplify the notation we set

$$\begin{aligned} \Delta_k t &= t_k - t_{k-1}, \quad t_n^\nu := t_{k-1,n}^\nu, \\ \mathbf{X}_{\nu,n} &:= \mathbf{X}(t_n^\nu), \quad \Delta_{\nu,n} \mathbf{X} := \mathbf{X}_{\nu,n} - \mathbf{X}_{\nu,n-1}. \end{aligned}$$

We have

$$S_{k,\nu} := \sum_{n=1}^{2^\nu} \left(u(t_n^\nu, \mathbf{X}_{\nu,n}) - u(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) \right).$$

Set $\Delta_k t := t_k - t_{k-1}$. Using Taylor's formula with integral remainder we deduce that

$$\begin{aligned} u(t_n^\nu, \mathbf{X}_{\nu,n}) - u(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) &= \underbrace{u'_s(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) 2^{-\nu} \Delta_k t}_{a_{\nu,n}} \\ &+ \frac{1}{2} \underbrace{u''_s(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) 2^{-2\nu} (\Delta_k t)^2}_{b_{\nu,n}} + \sum_{\alpha} \underbrace{u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) \Delta_{\nu,n}^\alpha}_{c_{\alpha|\nu,n}} \\ &+ \frac{1}{2} \sum_{\alpha,\beta} \underbrace{u''_{x_\alpha x_\beta}(t_{n-1}^\nu, \mathbf{X}_{\nu,n-1}) \Delta_{\nu,n}^\alpha \Delta_{\nu,n}^\beta}_{d_{\alpha\beta|\nu,n}} + R_{\nu,n}. \end{aligned} \tag{6.1.20}$$

The remainder $R_{\nu,n}$ is a *random variable*² that satisfies an upper bound

$$|R_{\nu,n}| \leq C(2^{-3\nu} (\Delta_k t)^3 + \|\Delta_{\nu,n-1}\|^3),$$

where C is a deterministic positive constant depending only on $\|u\|_{C_b^3}$. We have

$$\begin{aligned} S_{k,\nu} &= \sum_{n=1}^{2^\nu} a_{\nu,n} + \frac{1}{2} \sum_{n=1}^{2^\nu} b_{\nu,n} + \sum_{\alpha} \sum_{n=1}^{2^\nu} c_{\alpha|\nu,n} + \frac{1}{2} \sum_{\alpha,\beta} \sum_{n=1}^{2^\nu} d_{\alpha\beta|\nu,n} \\ &\quad + \sum_{n=1}^{2^\nu} R_{\nu,n}. \end{aligned}$$

Now observe that

$$\Delta_{\nu,n-1}^\alpha = \int_{t_{n-1}^\nu}^{t_n^\nu} F^\alpha(s) ds + \sum_{j=1}^{N_0} \int_{t_{n-1}^\nu}^{t_n^\nu} G_j^\alpha(s) dB^j(s)$$

²The measurability of $R_{\nu,n}$ follows from the representation of $R_{\nu,n}$ as an integral of a progressive process.

At this point we invoke the fact that the processes F^α and G_j^α are elementary and independent of t on each of the intervals $(t_{k-1}, t_k]$. We deduce that

$$\Delta_{\nu, n-1}^\alpha = 2^{-\nu} \underbrace{F^\alpha(t_{k-1}) \Delta_k t}_{\phi^\alpha} + \underbrace{\sum_j G_j^\alpha(t_{k-1}) \left(B^j(t_n^\nu) - B^j(t_{n-1}^\nu) \right)}_{v_{\nu n}^\alpha}.$$

Set

$$\Delta_{\nu, n} B^j := B^j(t_n^\nu) - B^j(t_{n-1}^\nu).$$

We deduce from (6.1.20) that the sums

$$\mathcal{A}_\nu := \sum_{n=1}^{2^\nu} a_{\nu, n}, \quad \mathcal{B}_\nu := \left[\frac{2^\nu}{\Delta_k t} \right] \cdot \sum_{n=1}^{2^\nu} b_{\nu, n}$$

are Riemann sums. As $\nu \rightarrow \infty$ they converge a.s. to

$$\int_{t_{k-1}}^{t_k} u'_s(s, \mathbf{X}(s)) ds \quad \text{and respectively} \quad \frac{1}{2} \int_{t_{k-1}}^{t_k} u''_s(s, \mathbf{X}(s)) ds.$$

Hence

$$\lim_{\nu \rightarrow \infty} \sum_{n=1}^{2^\nu} b_{\nu, n} = 0 \quad \text{a.s..}$$

Note that

$$\begin{aligned} c_{\alpha|\nu, n} &= u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu, n-1}) F^\alpha(t_{k-1}) 2^{-\nu} \\ &+ \sum_j G_j^\alpha(t_{k-1}) u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu, n-1}) \Delta_{\nu, n} B^j. \end{aligned}$$

As $\nu \rightarrow \infty$ we have

$$\sum_{n=1}^{2^\nu} u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu, n-1}) F^\alpha(t_{k-1}) 2^{-\nu} \rightarrow \int_{t_{k-1}}^{t_k} u'_{x_\alpha}(s, \mathbf{X}) F^\alpha(s) ds \quad \text{a.s.}$$

since the term on the left-hand side is a Riemann sum approximation for the integral on the right-hand side. Here we have tacitly used the fact that $F^\alpha(s) = F^\alpha(t_{k-1})$ for $s \in (t_{k-1}, t_k]$.

For fixed j , the sum

$$\sum_{n=1}^{2^\nu} G_j^\alpha(t_{k-1}) u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu, n-1}) \Delta_{\nu, n} B^j$$

converges in L^2 to

$$\int_{t_{k-1}}^{t_k} u'_{x_\alpha}(s, \mathbf{X}(s)) G_j^\alpha(s) dB^j(s)$$

The reason is that the process

$$Y^\nu = \sum_{n=1}^{2^\nu} u'_{x_\alpha}(t_{n-1}^\nu, \mathbf{X}_{\nu, n-1}) G_j^\alpha(t_{k-1}^\nu) \mathbf{I}_{(t_{n-1}^\nu, t_n^\nu]}$$

converges in $\mathbb{L}^2$ to the process $\mathbf{I}_{(t_{k-1}, t_k]} u'_{x_\alpha}(s, \mathbf{X}(s)) G_j^\alpha(s)$. More precisely

$$\|Y^\nu - \mathbf{I}_{(t_{k-1}, t_k]} u'_{x_\alpha}(s, \mathbf{X}(s)) G_j^\alpha(s)\|_{\mathbb{L}^2} = O(2^{-\nu})$$

since $u \in C_b^3$. The Borel-Cantelli lemma implies that the convergence is also a.s.. We deduce that

$$\begin{aligned} \lim_{\nu \rightarrow \infty} \sum_{n=1}^{2^\nu} c_{\alpha|\nu,n} &= \int_{t_{k-1}}^{t_k} u'_{x_\alpha}(s, \mathbf{X}) F^\alpha(s) ds \\ &+ \int_{t_{k-1}}^{t_k} u'_{x_\alpha}(s, \mathbf{X}(s)) G_j^\alpha(s) dB^j(s) \quad \text{a.s..} \end{aligned}$$

Note that

$$\mathbb{E}[|R_{\nu,n}|] \leq C \left(2^{-3\nu} (\Delta_k t)^3 + \mathbb{E}[|B^j(2^{-\nu} \Delta_k t)|^3] \right) = O(2^{-3\nu/2}).$$

We deduce that

$$\mathbb{E}\left[\left|\sum_n R_{\nu,n}\right|\right] = O(2^{-\nu/2}) \quad \text{as } \nu \rightarrow \infty$$

so

$$\lim_{\nu \rightarrow \infty} \sum_n R_{\nu,n} = 0 \quad \text{in } L^1 \text{ and a.s..}$$

We need to elucidate the behavior as $\nu \rightarrow \infty$ of

$$\mathcal{D}_{\alpha\beta|\nu} := \sum_{n=1}^{2^\nu} d_{\alpha\beta|\nu,n},$$

Set

$$Z(t) := u''_{x_\alpha x_\beta}(t, \mathbf{X}(t))$$

Then

$$\begin{aligned} d_{\alpha\beta|\nu,n} &= Z(t_{n-1}^\nu) \Delta_{\nu,n}^\alpha \Delta_{\nu,n}^\beta = Z(t_{n-1}^\nu) (2^{-\nu} \phi^\alpha + v_{\nu n}^\alpha) (2^{-\nu} \phi^\beta + v_{\nu n}^\beta) \\ &= Z(t_{n-1}^\nu) \left(2^{-2\nu} \phi^\alpha \phi^\beta + 2^{-\nu} (\phi^\alpha v_{\nu n}^\beta + \phi^\beta v_{\nu n}^\alpha) + v_{\nu n}^\alpha v_{\nu n}^\beta \right). \end{aligned}$$

Note that

$$\begin{aligned} \|Z(t_{n-1}^\nu)\|_{L^\infty} &\leq \|u\|_{C_b^3}, \quad \|\phi^\alpha\|_{L^\infty} \leq \|F^\alpha\|_\infty \Delta_k t \\ \|v_{\nu n}^\alpha\|_{L^2} &\leq \sum_j \|G_j^\alpha\|_{L^\infty} \underbrace{\|B^j(t_n^\nu) - B^j(t_{n-1}^\nu)\|_{L^2}}_{=(2^{-\nu} \Delta_k t)^{1/2}} = O(2^{-\nu/2}). \end{aligned}$$

We deduce that

$$\begin{aligned} \sum_{n=1}^{2^\nu} d_{\alpha\beta|\nu,n} &= 2^{-2\nu} \underbrace{\sum_{n=1}^{2^\nu} Z(t_{n-1}^\nu) \phi^\alpha \phi^\beta}_{=T_\nu^1} + 2^{-\nu} \underbrace{\sum_{n=1}^{2^\nu} (\phi^\alpha v_{\nu n}^\beta + \phi^\beta v_{\nu n}^\alpha)}_{=T_\nu^2} \\ &\quad + \underbrace{\sum_{n=1}^{2^\nu} Z(t_{n-1}^\nu) v_{\nu n}^\alpha v_{\nu n}^\beta}_{=T_\nu^3}. \end{aligned}$$

Note that

$$\|T_\nu^1\|_{L^\infty} = O(2^{-\nu}), \quad \|T_\nu^2\|_{L^2} = O(2^{-\nu/2}) \quad \text{as } \nu \rightarrow \infty.$$

On the other hand

$$T_\nu^3 = \sum_{i,j} G_i^\alpha(t_{k-1}) G_j^\beta(t_{k-1}) Q_\nu(B^i, B^j)$$

$$Q_\nu(B^i, B^j) = \sum_{n=1}^{2^\nu} Z(t_{n-1}^\nu) \Delta_{\nu,n} B^i \Delta_{\nu,n} B^j.$$

The term $Q_\nu(B^i, B^j)$ is symmetric and bilinear in B^i, B^j and we deduce from the polarization formula that

$$Q_\nu(B^i, B^j) = \frac{1}{2} (Q_\nu(B^i + B^j, B^i + B^j) - Q_\nu(B^i, B^i) - Q_\nu(B^j, B^j)).$$

Note that $2^{-1/2}(B^i + B^j)$ is a Brownian motion if $i \neq j$ so it suffices to understand the behavior of $Q_\nu(B, B)$ where B is a Brownian motion and $\mathcal{F}_\bullet$ is non-anticipating with respect to B . We have

$$Q_\nu(B, B) = \sum_{n=1}^{2^\nu} Z(t_{n-1}^\nu) (\Delta_{\nu,n} B)^2,$$

Set

$$I_n^\nu := Z(t_n^\nu) (t_n^\nu - t_{n-1}^\nu) = Z(t_n^\nu) 2^{-\nu} \Delta_k t.$$

Note that

$$I_\nu := \sum_{n=1}^{2^\nu} I_n^\nu$$

is a Riemann sum that converges a.s. to

$$\int_{t_{k-1}}^{t_k} Z(s) ds$$

as $\nu \rightarrow \infty$. Since

$$\sup_{0 \leq s \leq t} \|Z(s)\|_{L^\infty(\Omega)} < \infty.$$

the convergence holds in any $L^p(\Omega)$, $1 \leq p < \infty$. We have

$$Q_\nu(B) - I_\nu = \sum_{n=1}^{2^\nu} \Xi_n^\nu$$

where

$$\Xi_n^\nu := Z(t_{n-1}^\nu) (\Delta_{\nu,n} B)^2 - I_n^\nu = Z(t_{n-1}^\nu) (\Delta_{\nu,n} B)^2 - 2^{-\nu} \Delta_k t.$$

Note that

$$\mathbb{E}[(\Delta_{\nu,n} B)^2] = 2^{-\nu} \Delta_k t,$$

since $\Delta_{\nu,n} B$ is a normal random variable with mean 0 and variance $2^{-\nu} \Delta_k t$. For $m < n$ we have

$$\begin{aligned} \Xi_m^\nu \Xi_n^\nu &= \underbrace{Z(t_{m-1}^\nu) Z(t_{n-1}^\nu) ((\Delta_{\nu,m} B)^2 - 2^{-\nu} \Delta_k t)}_I \\ &\quad \times \underbrace{((\Delta_{\nu,n} B)^2 - 2^{-\nu} \Delta_k t)}_{II}. \end{aligned}$$

Observing that I and II are independent and $\mathbb{E}[II] = 0$ we deduce

$$\mathbb{E}[\Xi_m^\nu \Xi_n^\nu] = 0, \quad \forall 1 \leq m < n \leq 2^\nu.$$

Hence

$$\mathbb{E}[(Q_\nu(B, B) - I_\nu)^2] = \mathbb{E}\left[\left(\sum_n \Xi_n^\nu\right)^2\right] = \sum_n \mathbb{E}[(\Xi_n^\nu)^2]$$

$$\begin{aligned}
&= \sum_n \mathbb{E} [Z(t_{n-1}^\nu)^2 (\Delta_{\nu,n} B)^2 - 2^{-\nu} \Delta_k t)^2] \\
&= \sum_n \mathbb{E} [Z(t_{n-1}^\nu)^2] \mathbb{E} [(\Delta_{\nu,n} B)^2 - 2^{-\nu} \Delta_k t)^2] = 2(2^{-\nu} \Delta_k)^2 \sum_n \mathbb{E} [Z(t_{n-1}^\nu)^2],
\end{aligned}$$

where at the last step we use the fact that if N is normal random variable with mean zero and variance v , then

$$\mathbb{E} [(N^2 - v)^2] = 2v^2.$$

Since Z is bounded we deduce that

$$\mathbb{E} \left[\left(\sum_n \Xi_n^\nu \right)^2 \right] = \sum_n \mathbb{E} [(\Xi_n^\nu)^2] = O(2^{-\nu}) \text{ as } \nu \rightarrow \infty.$$

We deduce that

$$\|Q_\nu - I_\nu\|_{L^2}^2 = O(2^{-\nu}) \text{ as } \nu \rightarrow \infty.$$

Hence

$$\lim_{\nu \rightarrow \infty} Q_\nu(B, B) = \int_{t_{k-1}}^{t_k} Z(t) dt \text{ a.s. and } L^2. \quad (6.1.21)$$

From the polarization formula we deduce that

$$\lim_{\nu \rightarrow \infty} Q_\nu(B^i, B^j) = \delta_{ij} \int_{t_{k-1}}^{t_k} Z(t) dt \text{ a.s. and } L^2,$$

where δ_{ij} is Kronecker's symbol. Since $G_i^\alpha(t), G_j^\beta(t) \in L^\infty$ are independent of t on $(t_{k-1}, t_k]$ we deduce

$$\begin{aligned}
\lim_{\nu \rightarrow \infty} T_\nu^3 &= \lim_{\nu \rightarrow \infty} G_i^\alpha G_j^\beta Q_\nu(B^i, B^j) \\
&= \delta_{ij} \int_{t_{k-1}}^{t_k} G_i^\alpha(t) G_j^\beta(t) u''_{x_\alpha x_\beta}(t, \mathbf{X}(t)) dt,
\end{aligned}$$

a.s. and in L^2 . This concludes **Step 1**.

Step 2. We prove (6.1.19) when $u \in C_b^3([0, \infty \times \mathbb{R}^{N_1}])$, $F^\alpha \in \mathbb{L}_{\text{loc}}^1$, $G_j^\alpha \in \mathbb{L}_{\text{loc}}^2$. In this case we can find $F_n^\alpha, G_{j,n}^\alpha \in \mathcal{E}_{\text{pred}}$ such that

$$F_n^\alpha \rightarrow F^\alpha \text{ in } \mathbb{L}^1(0, t) \text{ and a.s., } G_{j,n}^\alpha \rightarrow G_j^\alpha \text{ in } \mathbb{L}^2(0, t) \text{ and a.s.}$$

We let $n \rightarrow \infty$ in (6.1.19) for u, F_n^α and $G_{j,n}^\alpha$ and observe that $\mathcal{R}_t(u, \mathbf{F}_n, \mathbf{G}_n) \rightarrow \mathcal{R}_t(u, \mathbf{F}, \mathbf{G})$ a.s..

Step 3. We prove (6.1.19), when $F^\alpha \in \mathcal{L}_{\text{loc}}^1$, $G_j^\alpha \in \mathcal{L}_{\text{loc}}^2$ and $u \in C_b^3$. Using a localizing sequence T_n that works for both $\mathbf{F}$ and $\mathbf{G}$ (see Lemma 6.1.15) we obtain sequences

$$\mathbf{F}_n = \mathbf{F} \mathbf{I}_{[[0, T_n]]} \in \mathbb{L}_{\text{loc}}^1, \quad \mathbf{G}_n = \mathbf{G} \mathbf{I}_{[[0, T_n]]} \in \mathbb{L}_{\text{loc}}^2$$

such that

$$\begin{aligned}
\mathbf{F}_n(t, \omega) &\rightarrow \mathbf{F}(t, \omega) \quad \mathbf{G}_n(t, \omega) \rightarrow \mathbf{G}(t, \omega), \quad \forall t, \omega, \\
|\mathbf{F}_n(t, \omega)| &\leq |\mathbf{F}(t, \omega)|, \quad |\mathbf{G}_n(t, \omega)| \leq |\mathbf{G}(t, \omega)|, \quad \forall n, t, \omega.
\end{aligned}$$

We let $n \rightarrow \infty$ and we observe that the Dominated Convergence Theorem implies $\mathcal{R}_t(u, \mathbf{F}_n, \mathbf{G}_n)(\omega)$ converges to $\mathcal{R}_t(u, \mathbf{F}, \mathbf{G})(\omega)$, $\forall \omega$.

Step 4. We prove (6.1.19), when $F^\alpha \in \mathcal{L}_{\text{loc}}^1$, $G_j^\alpha \in \mathcal{L}_{\text{loc}}^2$ and

$$u \in C_b^{1,2}([0, \infty) \times \mathbb{R}^{N_1}), \quad \text{i.e. } u, u'_t, u'_{x_\alpha}, u''_{x_\alpha x_\beta} \in C_b^0([0, \infty) \times \mathbb{R}^{N_1}).$$

This is achieved by selecting a sequences $u_n \in C_b^3([0, \infty) \times \mathbb{R}^{N_1})$ such that

$$u_n, (u_n)'_t, (u_n)'_{x_\alpha}, (u_n)''_{x_\alpha x_\beta}$$

converge sufficiently fast uniformly on compacts to $u, u'_t, u'_{x_\alpha}, u''_{x_\alpha x_\beta}$. Since $\mathbf{X}(t)$ has continuous paths we deduce that

$$\mathcal{R}_t(u_n, \mathbf{F}, \mathbf{G}) \rightarrow \mathcal{R}_t(u, \mathbf{F}, \mathbf{G}) \text{ a.s..}$$

Step 5. We prove (6.1.19) in general, when $F \in \mathcal{L}^{1, \text{loc}}, G \in \mathcal{L}_{\text{loc}}^2, u \in C^{1,2}([0, \infty) \times \mathbb{R}^{N_1})$.

Fix a smooth cutoff function $\rho : \mathbb{R} \rightarrow [0, 1]$ such that

$$\rho(s) = \begin{cases} 1, & s \leq 0, \\ 0, & s \geq 1 \end{cases}.$$

Set $\rho_n(s) = \rho(s - n)$, and define

$$\psi_n : [0, \infty) \times \mathbb{R}^{N_1} \rightarrow [0, 1], \quad \psi_n(t, \mathbf{x}) = \rho_n(t) \rho_n(|\mathbf{x}|),$$

where $|\cdot|$ denotes the Euclidean norm on $\mathbb{R}^{N_1}$. Set $u_n = \psi_n u$. Then $u_n \in C_{\text{cpt}}^{1,2} \subset C_b^{1,2}$ and $u_n \rightarrow u, C^{1,2}$ on compacts. Since $\mathbf{X}$ has continuous paths we deduce that

$$\mathcal{R}_t(u_n, \mathbf{F}, \mathbf{G}) \rightarrow \mathcal{R}_t(u, \mathbf{F}, \mathbf{G}) \text{ on } \Omega.$$

□

Remark 6.1.36 (Itô calculus rules). Suppose $u = u(t, x^1, \dots, x^n)$ is a $C^{1,2}$ function and

$$\mathbf{X} = (X^1, \dots, X^n) = (X^\alpha)_{1 \leq \alpha \leq n}$$

is an n -dimensional Itô process. Then $u(t, X^1, \dots, X^n)$ is an Itô process as well and

$$\begin{aligned} du(t, X^1, \dots, X^n) &= u'_t dt + \sum_{\alpha} u'_{x^\alpha}(\mathbf{X}) dX^\alpha \\ &\quad + \frac{1}{2} \sum_{\alpha, \beta} u''_{x^\alpha x^\beta}(\mathbf{X}) dX^\alpha dX^\beta \\ &= u'_t dt + \sum_{\alpha} u'_{x^\alpha}(\mathbf{X}) dX^\alpha \\ &\quad + \frac{1}{2} \sum_{\alpha} u''_{x^\alpha x^\alpha}(\mathbf{X}) dX^\alpha dX^\alpha + \sum_{\alpha < \beta} u''_{x^\alpha x^\beta}(\mathbf{X}) dX^\alpha dX^\beta. \end{aligned} \tag{6.1.22}$$

Above, the differentials dX^α are linear combinations of the elementary differentials dt and dB^j and the multiplication of these elementary differentials is given by the multiplication table 6.1

Table 6.1. Itô's multiplication rules

$\times$	dt	dB^i
dt	0	0
dB^j	0	$\delta_{ij} dt$

Formally, the equality (6.1.22) is obtained as follows. First, consider the 2nd (reduced) jet of u

$$\begin{aligned} & u'_t(t, x^\alpha)dt + \sum_{\alpha} u'_{x^\alpha}(t, x^\alpha)dx^\alpha + \frac{1}{2}u''_{tt}(t, x^\alpha)dt^2 \\ & + \frac{1}{2} \sum_{\alpha} u''_{x^\alpha x^\alpha}(t, x^\alpha)(dx^\alpha)^2 + \sum_{\alpha < \beta} u''_{x^\alpha x^\beta}(t, x^\alpha)dx^\alpha dx^\beta. \end{aligned}$$

Next, in the above expression make the replacements $x^\alpha \rightarrow X^\alpha$ and then apply the multiplication rules in Table 6.1. If

$$dX^\alpha = F^\alpha dt + \sum_j G_j^\alpha dB^j$$

Then $du(t, \mathbf{X})$ will have the form

$$du(t, \mathbf{X}) = Lu(t, \mathbf{X})dt + \sum_j \left(\sum_{\alpha} u'_{x^\alpha}(t, \mathbf{X})G_j^\alpha \right) dB^j, \quad (6.1.23)$$

where L is the second order partial differential operator

$$L = \partial_t + \frac{1}{2} \sum_{\alpha} \left(\sum_j G_j^\alpha G_j^\alpha \right) \partial_{x^\alpha}^2 + \sum_{\alpha} F^\alpha \partial_{x^\alpha}.$$

□

6.1.7. Some computational examples. The best way to get a feel of Itô's formula is to see it at work in a few concrete situations. This subsection collects a few and I believe revealing examples.

Example 6.1.37. Let $B(t)$ be a $\mathcal{F}_\bullet$ -Brownian motion. Fix

$$f = f(t, x) \in C_b^{1,2}([0, \infty) \times \mathbb{R}).$$

Set $X(t) = f(t, B(t))$. Itô's formula implies

$$dX(t) = f'_x(B(t))dB + (Lf)(B(t))dt, \quad Lf = \partial_t f + \frac{1}{2}\partial_{xx}^2 f.$$

Hence

$$f(t, B(t)) - f(0) - \frac{1}{2} \int_0^t f''_{xx}(s, B(s))ds = \int_0^t f'_x(s, B(s))dB(s)$$

Since f'_x is bounded, we deduce that the term in the right hand side is an L^2 -bounded martingale. This shows that Proposition 3.5.5 is a special case of Itô's formula.

More generally, suppose that $\mathbf{B} = (B^1(t), \dots, B^n(t))$ is a Brownian motion on $\mathbb{R}^n$ started at the origin and $f \in C^2(\mathbb{R}^n)$ such that

$$e^{-|\mathbf{x}|} f(\mathbf{x}) \in C_b^2(|\mathbf{x}| \geq 1).$$

Then

$$f(\mathbf{B}(t)) = f(0) + \frac{1}{2} \int_0^t \Delta f(\mathbf{B}(s))ds + \sum_k \int_0^t \partial_{x_k} f(\mathbf{B}(s))dB^k(s),$$

where Δ denotes the *physicists'* Laplacian,³ $\Delta = \sum_k \partial_{x_k}^2$. Note that if f is subharmonic, i.e., $\Delta f \geq 0$, then $f(\mathbf{B}(t))$ is a submartingale. □

³The geometers' Laplacian differs by a sign.

Example 6.1.38. Fix a Brownian motion B and a B -admissible filtration $\mathcal{F}_\bullet$. Let $G \in \mathcal{L}_{\text{loc}}^2(\mathcal{F}_\bullet)$ and set

$$X(t) := \int_0^t G(s)dB(s).$$

In this case $L = \frac{1}{2}G^2\partial_{xx}^2$. Itô's formula shows that for any $f \in C^2(\mathbb{R})$ we have

$$df(X) = \frac{1}{2}G^2f''(X)dt + f'(X)dB.$$

When $f(x) = x^2$ we get

$$dX^2 = G^2dt + 2XGdB,$$

i. e.,

$$X^2(t) - \int_0^t G(s)^2ds = 2 \int_0^t X(s)G(s)dB(s). \quad (6.1.24)$$

Consider now the Itô process

$$Y(t) = \int_0^t G(s)^2ds.$$

For $\lambda \in \mathbb{R}$ set $f_\lambda(x, y) = e^{\lambda x - \frac{\lambda^2}{2}y}$. We want to compute $df_\lambda(X, Y)$. In this case

$$L = \frac{1}{2}G^2\partial_{xx}^2 + G^2\partial_y,$$

$$de^{\lambda X - \frac{\lambda^2}{2}Y} = Lf_\lambda(X, Y)dt + G\partial_x f_\lambda(X, Y)dB$$

($Lf_\lambda = 0$)

$$= \frac{G\lambda^2}{2}e^{\lambda X - \frac{\lambda^2}{2}Y}dB.$$

Hence $f_\lambda(X, Y) = e^{\lambda X - \frac{\lambda^2}{2}Y}$ is a local martingale. $\square$

Example 6.1.39 (The stochastic exponential and Hermite polynomials). Recall the n -th Hermite polynomial H_n

$$H_n(y) = (-1)^n e^{y^2/2} \frac{d^n}{dy^n} e^{-y^2/2} = n! \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^r}{2^r r! (n-2r)!} y^{n-2r}. \quad (6.1.25)$$

The generating function of the Hermite polynomials is given by (1.6.6),

$$\sum_{n \geq 0} H_n(y) \frac{\lambda^n}{n!} = e^{\lambda y - \lambda^2/2}. \quad (6.1.26)$$

Set

$$H_n(t, x) := t^{n/2} H_n(xt^{-1/2}) = (-t)^n e^{\frac{x^2}{2t}} \frac{d^n}{dx^n} e^{-\frac{x^2}{2t}}.$$

I will show the following identity of K. Itô, [104, Thm 3.1].

$$H_{n+1}(t, B(t)) = \int_0^t H_n(s, B(s))dB(s), \quad \forall n \geq 0. \quad (6.1.27)$$

I will follow the strategy in [150, Sec. 2.7]. Consider the exponential martingale (3.5.2)

$$X(t) = f_\theta(t, B(t)), \quad f_\theta(t, x) = \exp(\theta x - \theta^2 t/2)$$

Note that

$$\partial_t f_\theta + \partial_x^2 f_\theta = 0.$$

Itô's formula implies

$$dX(t) = \partial_x f_\theta(t, B(t))dB = \theta f_\theta(t, B(t))dB = \theta X(t)dB(t).$$

Hence

$$X(t) = 1 + \theta \int_0^t X(s)dB(s) \quad (6.1.28)$$

If we set $y = xt^{-1/2}$, $\lambda = \theta t^{1/2}$ in (6.1.25) we deduce

$$f_\theta(t, x) = \sum_{n \geq 0} H_n(t, x) \frac{\theta^n}{n!}. \quad (6.1.29)$$

From the equality (6.1.25) we deduce that

$$|H_n(y)| \leq \sum_{m=0}^n \binom{n}{m} |y|^{n-m} \left(\frac{1}{2}\right)^{m/2} = \left(\frac{1}{\sqrt{2}} + |y|\right)^n,$$

so

$$|H_n(t, x)| \leq \left(\sqrt{\frac{t}{2}} + |x|\right)^n. \quad (6.1.30)$$

Set

$$f_{\theta,n}(t, x) = \sum_{k=0}^n H_k(t, x) \frac{\theta^k}{k!}.$$

The estimate (6.1.30) implies that for any continuous function $u : [0, \infty) \rightarrow \mathbb{R}$ we have

$$\lim_{n \rightarrow \infty} \int_0^t \left(f_{\theta,n}(s, u(s)) - f_\theta(s, u(s)) \right)^2 ds = 0, \quad \forall t \geq 0.$$

Using the Approximation Theorem 6.1.18 we deduce that

$$\int_0^t X(s)dB(s) = \int_0^t f_\theta(s, B(s))dB(s) = \sum_{n \geq 0} \frac{\theta^n}{n!} \int_0^t H_n(s, B(s))dB(s),$$

where the last series converges in probability. Using this in the integral equation (6.1.28) we deduce

$$\begin{aligned} \sum_{n \geq 0} H_n(t, B(t)) \frac{\theta^n}{n!} &= X(t) = 1 + \theta \int_0^t X(s)dB(s) \\ &= 1 + \sum_{n \geq 1} \left(\int_0^t H_{n-1}(s, B(s))dB(s) \right) \frac{\theta^n}{n!}. \end{aligned}$$

The equalities (6.1.27) are obtained by identifying the coefficients of the above power series. $\square$

Example 6.1.40 (Stochastic Integration by parts). Suppose that X, Y are one-dimensional Itô processes

$$dX = udt + FdB, \quad dY = vdt + GdB.$$

Then XY is also an Itô process and

$$d(XY) = XdY + YdX + dXdY = XdY + YdX + FGdt$$

The integral form of this identity is an integration-by parts formula

$$X(t)Y(t) - X(0)Y(0) = \int_0^t X_s dY_s + \int_0^t Y_s dX_s + \int_0^t F(s)G(s)ds \quad (6.1.31)$$

Note that the noisy parts of the stochastic differential introduce a correction to the usual integration by parts formula for deterministic functions. For example, if $X = Y = B$ then we get

$$2 \int_0^t B(s) dB(s) = B(t)^2 - t.$$

Observe that $B^2 \in \mathbb{L}_{\text{loc}}^2$ so its Itô integral is a continuous L^2 -martingale, something we already knew. $\square$

Example 6.1.41. Fix $n \in \mathbb{N}$ and a point $\mathbf{x}_0 = (x_0^1, \dots, x_0^n) \in \mathbb{R}^n$. Denote by $|\cdot|$ the canonical Euclidean norm on $\mathbb{R}^n$. Consider a Brownian motion $\mathbf{B}$ in $\mathbb{R}^n$ starting at 0 and set

$$\mathbf{B}_{\mathbf{x}_0}(t) := \mathbf{x}_0 + \mathbf{B} = (x_0^1 + B^1(t), \dots, x_0^n + B^n(t)).$$

For any open set $D \subset \mathbb{R}^n$ and any $\mathbf{x} \in D$ we set

$$T_{D,\mathbf{x}} = \inf \{ t \geq 0; \mathbf{x} + \mathbf{B}(t) \in D^c := \mathbb{R}^n \setminus D \}.$$

When D is the open ball $\mathbb{B}_r = \{|\mathbf{x}| < r\}$ we use the simpler notation $T_{r,\mathbf{x}} := T_{\mathbb{B}_r,\mathbf{x}}$. Note that if $\sup_{\mathbf{x} \in D} |\mathbf{x}| \leq r$, then $T_{D,\mathbf{x}} \leq T_{r,\mathbf{x}}$.

For any $r > 0$ we define $f_r \in C_b^2(\mathbb{R}^n)$ by $f_r(\mathbf{x}) = -e^{-\frac{|\mathbf{x}|^2}{4r^2}}$. Note that

$$\begin{aligned} \partial_{x^j} f_r &= -\frac{x^j}{2r^2} f_r, \quad \partial_{x^j}^2 f_r = -\frac{1}{2r^2} f_r + \frac{(x^j)^2}{4r^4} f_r \\ \Delta f_r &= -\left(\frac{|\mathbf{x}|^2}{4r^4} - \frac{n}{2r^2}\right) f_r = \left(\frac{n}{2r^2} - \frac{|\mathbf{x}|^2}{4r^4}\right) e^{-\frac{|\mathbf{x}|^2}{4r^2}}. \end{aligned}$$

Note that

$$\inf_{|\mathbf{x}| < r} \Delta f_r(|\mathbf{x}|) = \underbrace{\frac{2n-1}{4r^2} e^{-1/4}}_{=: \lambda}.$$

Writing $X^\alpha = x_0^\alpha + \delta_j^\alpha B_j$, $1 \leq \alpha, j \leq n$ we deduce from (6.1.23)

$$df_r(\mathbf{B}_{\mathbf{x}_0}(t)) = \frac{1}{2} \Delta f_r(\mathbf{B}_{\mathbf{x}_0}(t)) dt + \sum_j \partial_{x^j} f_r(\mathbf{B}_{\mathbf{x}_0}(t)) dB^j$$

Hence

$$\begin{aligned} f_r(\mathbf{B}_{\mathbf{x}_0}(t)) - f_t(\mathbf{x}_0) &= \frac{1}{2} \int_0^t \Delta f_r(\mathbf{B}_{\mathbf{x}_0}(s)) ds \\ &\quad + \underbrace{\sum_j \int_0^t \partial_{x^j} f_r(\mathbf{B}_{\mathbf{x}_0}(s)) dB^j(s)}_{=: Y(t)}. \end{aligned}$$

The process $Y(t)$ is an L^2 -martingale with $Y(0) = 0$. For any $N \in \mathbb{N}$ we have

$$\begin{aligned} |\mathbf{B}_{\mathbf{x}_0}(T_{r,\mathbf{x}_0} \wedge n)| &\leq r, \quad \Delta f_r(\mathbf{B}_{\mathbf{x}_0}(T_{r,\mathbf{x}_0} \wedge n)) \geq \lambda \\ 1 &\geq f_r(\mathbf{B}_{\mathbf{x}_0}(T_{r,\mathbf{x}_0} \wedge N)) - f_t(\mathbf{x}_0) \\ &= \frac{1}{2} \int_0^{T_{r,\mathbf{x}_0} \wedge N} \Delta f_r(\mathbf{B}_{\mathbf{x}_0}(s)) ds + Y(T_{r,\mathbf{x}_0} \wedge N) \\ &\geq \frac{\lambda}{2} (T_{r,\mathbf{x}_0} \wedge N) + Y(T_{r,\mathbf{x}_0} \wedge N). \end{aligned}$$

By taking expectations and using the fact that $N \rightarrow Y(T_{r,x_0} \wedge N)$ is a martingale we deduce

$$1 \leq \frac{\lambda}{2} \mathbb{E}[T_{r,x_0} \wedge N].$$

Letting $N \rightarrow \infty$ we deduce

$$\mathbb{E}[T_{r,x_0}] \leq \frac{2}{\lambda} < \infty.$$

We deduce that $\mathbb{E}[T_{D,x_0}] < \infty$.

Suppose now that D is bounded with smooth boundary. It is known (see [87, Chap. 4]) that for any function $g \in C^{0,\alpha}(\partial D)$, there exists a unique function $u \in C^{0,\alpha}(\bar{D}) \cap C^{2,\alpha}(D)$ satisfying the boundary value problem,

$$\Delta u = 0 \text{ in } D, \quad u|_{\partial D} = g.$$

For $\varepsilon \geq 0$ we set

$$D_\varepsilon := \{x \in D; \text{dist}(x, \partial D) > \varepsilon\}.$$

This is a bounded domain such that $\bar{D}_\varepsilon \subset D$ for $\varepsilon > 0$. Moreover, for $\varepsilon > 0$ sufficiently small, the boundary ∂D_ε is smooth. Note that $D = D_0$. For simplicity we set

$$T_{\varepsilon,x} = T_{D_\varepsilon,x} = \inf \{t > 0; B_x(t) \in \mathbb{R}^n \setminus D_\varepsilon\}.$$

Since ∂D_ε is smooth for $\varepsilon > 0$ small, we deduce that, for such ε , there exists a function $u_\varepsilon \in C_{\text{cpt}}^{2,\alpha}(\mathbb{R}^n)$ such that $u_\varepsilon(x) = u(x)$, $\forall x \in \bar{D}_\varepsilon$. Itô's formula implies that for any $x \in D_\varepsilon$ we have

$$\begin{aligned} u_\varepsilon(B_x(t)) - u_\varepsilon(x) - \frac{1}{2} \int_0^t \Delta u_\varepsilon(B_x(s)) ds \\ = \underbrace{\sum_{j=1}^n \int_0^t \partial_{x^j} u_\varepsilon(B_x(s)) dB^j(s)}_{=: X(t)}. \end{aligned}$$

The process $X(t)$ is an L^2 -martingale and thus $u_\varepsilon(B_x(t)) - u_\varepsilon(x)$ is an L^2 martingale. Observing that $\Delta u_\varepsilon(B_x(s)) = 0$ for $s < T_{\varepsilon,x}$ we deduce that

$$Y(t) = u_\varepsilon(B_x(t \wedge T_{\varepsilon,x})) - u(x)$$

is an L^2 -martingale. Since $\mathbb{E}[T_\varepsilon] < \infty$ we conclude that

$$\mathbb{E}[Y(T_{\varepsilon,x})] = \mathbb{E}[Y(0)] = 0,$$

$$\mathbb{E}[u_\varepsilon(B_x(T_{\varepsilon,x}))] = \mathbb{E}[u_\varepsilon(x + B_x(T_{\varepsilon,x}))] = u(x).$$

Letting $\varepsilon \searrow 0$ we deduce from the Dominated Convergence Theorem that

$$u(x) = \mathbb{E}[u(B_x(T_{D,x}))] = \mathbb{E}[g(B_x(T_{D,x}))]$$

We have obtained a probabilistic description of the harmonic function u in terms of its boundary values g , assuming a classical existence result concerning existence for the Dirichlet problem.

It turns out that if the boundary of D is “not too wild”, then we can prove entirely by probabilistic methods that the function

$$\bar{D} \ni x \mapsto u(x) := \mathbb{E}[g(B_x(T_{D,x}))]$$

is the unique solution of the boundary value problem

$$u \in C^2(D) \cap C^0(\bar{D}), \quad \Delta u = 0 \text{ in } D, \quad u|_{\partial D} = g.$$

In particular, we obtain a purely probabilistic proof for the existence and uniqueness of solutions of this boundary value problem. For details I refer to [185, Sec.8.4]. $\square$

Example 6.1.42 (Stochastic differential equations). A *stochastic differential equation* (SDE) is an equation of the form

$$dX_t = \mu(t, X_t)dt + \sigma(t, X_t)dB_t, \quad X_0 = Z, \quad t \in [0, T]. \quad (6.1.32)$$

Above, Z is a random variable and μ, σ are functions of appropriate regularity.

A *weak* solution of this equation consists of

- (i) a Brownian motion $(B_t)_{t \geq 0}$,
- (ii) a complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$ together with a B -admissible filtration

$$\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0},$$

and

- (iii) an $\mathcal{F}_\bullet$ -adapted continuous process (X_t) satisfying the stochastic integral equation

$$X_t - Z = \int_0^t \mu(s, X_s)ds + \int_0^t \sigma(s, X_s)dB_s, \quad t \in [0, 1].$$

The solution is called *strong* if the filtration $\mathcal{F}_\bullet$ is the (completed) filtration determined by the Brownian motion B . Note that since the initial condition Z is $\mathcal{F}_0$ -measurable it is independent of the Brownian motion since the filtration $\mathcal{F}_\bullet$ is nonanticipating (or B -admissible).

There is a rich theory of SDEs and I refer to curious reader to [161] or [181, Chap.V]. I have only one modest goal, a uniqueness result. I will describe a simple condition on μ and σ that guarantees that (6.1.32) has at most one solution.

More precisely, I will assume that μ, σ are continuous and that there exist $C, L > 0$ such that, $\forall t \in [0, T], \forall x, y \in \mathbb{R}$,

$$|\mu(t, x)| + |\sigma(t, x)| \leq C(1 + |x|), \quad (6.1.33a)$$

$$|\mu(t, x) - \mu(t, y)|, |\sigma(t, x) - \sigma(t, y)| \leq L|x - y|. \quad (6.1.33b)$$

Suppose that B is a Brownian motion, $\mathcal{F}_\bullet$ is a B -admissible filtration, and $(X_t)_{t \in [0, T]}, (Y_t)_{t \in [0, T]}$ are $\mathcal{F}_\bullet$ -adapted continuous processes in satisfying (6.1.32), (6.1.33a), (6.1.33b) and $X_0 = Y_0$, a.s..

I will show that if μ and σ satisfy (6.1.33b), then the processes X and Y are indistinguishable.

For each $n \in \mathbb{N}$ consider the stopping time

$$T_n := \inf \{ t > 0; \max(|X_t|, |Y_t|) \geq n \}$$

and define

$$U(t) := (\mu(t, X_t) - \mu(t, Y_t)), \quad V(t) = (\sigma(t, X_t) - \sigma(t, Y_t)).$$

Since X and Y are solutions of (6.1.32), we deduce

$$X_t - Y_t = \int_0^t U_s ds + \int_0^t V_s dB_s, \quad \forall t \in [0, T].$$

Localizing with respect to T_n in (6.1.32) we deduce

$$X_t^{T_n} - Y_t^{T_n} = \int_0^{t \wedge T_n} U_s ds + \int_0^{t \wedge T_n} V_s dB_s.$$

Hence

$$\mathbb{E}[(X_t^{T_n} - Y_t^{T_n})^2] = \mathbb{E}\left[\left(\int_0^{t \wedge T_n} U(s)ds + \int_0^{t \wedge T_n} V(s)dB_s\right)^2\right]$$

(use the elementary inequality $(a+b)^2 \leq 2(a^2 + b^2)$)

$$\leq 2\mathbb{E}\left[\left(\int_0^{t \wedge T_n} U(s)ds\right)^2\right] + 2\mathbb{E}\left[\left(\int_0^{t \wedge T_n} V(s)dB_s\right)^2\right].$$

Observe that

$$|\mu(s, X_s) - \mu(s, Y_s)| \leq L|X_s - Y_s|, \quad \forall s \in [0, T],$$

so

$$\begin{aligned} \mathbb{E}\left[\left(\int_0^{t \wedge T_n} U(s)ds\right)^2\right] &\leq L^2\mathbb{E}\left[\left(\int_0^T ds\right) \int_0^{t \wedge T_n} (X_s - Y_s)^2 ds\right] \\ &= TL^2\mathbb{E}\left[\int_0^{t \wedge T_n} (X_s - Y_s)^2 ds\right] \leq TL^2\mathbb{E}\left[\int_0^t (X_{s \wedge T_n} - Y_{s \wedge T_n})^2 ds\right] \\ &= TL^2 \int_0^t \mathbb{E}[(X_s^{T_n} - Y_s^{T_n})^2] ds. \end{aligned}$$

Set $V_n := \mathbf{I}_{[0, T_n]} V \in \mathbb{L}^2(0, T)$. We have

$$\mathbb{E}\left[\left(\int_0^{t \wedge T_n} V(s)dB_s\right)^2\right] = \mathbb{E}\left[\left(\int_0^t V_n(s)dB_s\right)^2\right]$$

(use Itô's isometry (6.1.10))

$$= \mathbb{E}\left[\int_0^t V_n(s)^2 ds\right] \leq \mathbb{E}\left[\int_0^{t \wedge T_n} (X_s - Y_s)^2 ds\right] \leq L^2 \int_0^t \mathbb{E}[(X_s^{T_n} - Y_s^{T_n})^2] ds.$$

Hence

$$\mathbb{E}[(X_t^{T_n} - Y_t^{T_n})^2] \leq 2L^2(1+T) \int_0^t \mathbb{E}[(X_s^{T_n} - Y_s^{T_n})^2] ds.$$

Thus, if we set

$$f_n(s) = \mathbb{E}[(X_s^{T_n} - Y_s^{T_n})^2] < \infty,$$

then f_n satisfies the integral inequality

$$f_n(0) = 0, \quad f(t) \leq 2L^2(1+T) \int_0^t f_n(s) ds.$$

Gronwall's Lemma [12, Cor. 1.1] implies $f_n(s) = 0, \forall n$. Thus

$$\forall t \in [0, T], \quad \forall n, \quad X_t^{T_n} = Y_t^{T_n} \text{ a.s..}$$

Letting $n \rightarrow \infty$ we deduce that

$$\forall t \in [0, T], \quad X_t = Y_t \text{ a.s..}$$

Since X and Y are continuous processes we deduce that X and Y are indistinguishable. $\square$

Example 6.1.43 (The Ornstein-Uhlenbeck process). We have seen that the stochastic differential equation

$$dX(t) = \theta X(t)dB(t)$$

has one solution of the form $X(t) = f(t, B(t))$ namely

$$X(t) = \exp(\theta B(t) - \theta t^2/2).$$

I would like to investigate a very simple linear stochastic ODE

$$dX = -\alpha X dt + \sigma dB(t) \quad (6.1.34)$$

where $\alpha, \sigma \in (0, \infty)$. This is an example of a *Langevin equation*.

One should regard the above equality as a “noisy” perturbation of the simplest linear o.d.e., $x' = \alpha x$. The solutions of the *deterministic* perturbation of this equation,

$$x' = \alpha x + f(t),$$

are given by Duhamel formula

$$x(t) = x(0)e^{-\alpha t} + \int_0^t t e^{-\alpha(t-s)} f(s) ds.$$

Using this as inspiration we try the following Ansatz for a solution of (6.1.34)

$$\begin{aligned} X(t) &= X(0)e^{-\alpha(t-t_0)} + \int_{t_0}^t \sigma e^{-\alpha(t-s)} dB(s) \\ &= X(0)e^{-\alpha t} + e^{-\alpha t} \int_{t_0}^t \sigma e^{\alpha s} dB(s). \end{aligned}$$

where $X(t)$ is adapted to the completed filtration $\mathcal{F}_t^B$ determined by B . In particular this implies that $X(0)$ is a.s. constant since $\mathcal{F}_0^B$ is a 0 – 1 sigma subalgebra. Set $x_0 = X(0)$.

Itô’s formula shows that this Ansatz is correct and describes a solution of (6.1.34). The discussion in the previous example shows that it is the unique solution satisfying the initial condition $X(0) = x_0 \in \mathbb{R}$. The stochastic process $X(t)$ is known as the *Ornstein-Uhlenbeck process*.

The Wiener integral above describes a Gaussian process so $X(t)$ is a Gaussian process with mean

$$\mathbb{E}[X(t)] = x_0 e^{-\alpha t}$$

and variance

$$\text{Var}[X(t)] = \sigma^2 e^{-2\alpha t} \int_0^t e^{2\alpha s} ds = \frac{\sigma^2}{2\alpha} (1 - e^{-2\alpha t}).$$

It can be characterized as the unique Gaussian process that is simultaneously time-homogeneous and Markov, [60].

Just as the Brownian motion can be represented as a scaling limit of the standard random walk on $\mathbb{Z}$, so does the Ornstein-Uhlenbeck process admits a description as a scaling limit of Bernoulli-Laplace urn; see [105] or [162, Sec.3]. For the physical interpretation of this process we refer to [22, Sec. IV.16].

Using (6.1.26) we deduce

$$e^{\alpha s} = e^{\alpha^2/2} \sum_{n \geq 0} \frac{\alpha^n}{n!} H_n(s).$$

We deduce

$$\begin{aligned} \int_0^t e^{-\alpha s} dB(s) &= e^{\alpha^2/2} \sum_{n \geq 0} \frac{\alpha^n}{n!} \int_0^t H_n(s) dB(s) \\ &\stackrel{(6.1.27)}{=} e^{\alpha^2/2} \sum_{n \geq 0} \frac{\alpha^n}{n!} H_{n+1}(t, B(t)). \end{aligned}$$

Note that for any $f \in C^1(\mathbb{R})$ the stochastic process $f(X(t))$ satisfies the stochastic differential equation

$$\begin{aligned} df(X) &= f'(X)dX + \frac{1}{2}f''(X)dt = f'(X)(-\alpha Xdt + \sigma dB) \\ &= \left(\frac{1}{2}f''(X) - \alpha Xf'(X)\right)dt + \sigma f'(X)dB. \end{aligned}$$

□

6.2. A more general stochastic integral

The above construction of the Itô integral can be generalized and streamlined revealing deeper structures. This extension is similar to the generalization of the integral with respect to Lebesgue measure to an integral with respect to an abstract measure.

In the stochastic case, the role of abstract measures is played by the so called *integrators* and the functions we integrate are called *integrands*. In the case of Itô integral discussed in the previous section the integrator is the Brownian motion and the integrands are the predictable processes in $\mathcal{L}_{\text{loc}}^2$.

The integrators are the so called *continuous semimartingales*. The continuous local martingales form an important subclass of continuous semimartingales. The Itô processes are, essentially, all the semimartingales. The integrands are progressive processes with certain integrability properties.

Given the stochastic integral of a progressive process H with respect to a semimartingale M is a process

$$H \cdot M(t) = \int_0^t H(s)dM(s).$$

One important feature of the stochastic integral is that its output $H \cdot M$ has the same nature as M . For example, if M is a continuous L^2 -martingale, then so is $H \cdot M$. I refer to [181, Sec.IV.1] for a very illuminating bird's-eye-view of this stochastic integral.

There are different ways of constructing the stochastic integral, each requiring to bypass certain technical obstacles.

One can define $H \cdot M(t)$ as an integral of $H(s, \omega)$ over $[0, t] \times \Omega$ with respect a certain measure λ_M on $[0, \infty) \times \Omega$. This is the point of view adopted in [40] and the major obstacle is defining the measure λ_M .

We adopt an approach pioneered by Kunita and Watanabe [125] that mimics the construction of the original Itô integral in the previous section. Recall that it consisted of three main steps.

- (i) Construct the integrals of sufficiently many elementary *predictable* processes.
- (ii) Extend by continuity the integral to many more processes.
- (iii) Localization with respect to stopping times.

We performed Step (ii) by relying on Itô's isometry. There exists an Itô isometry in the more general case as well, defined on an appropriate Hilbert space. The major new difficulty is describing explicitly this "appropriate Hilbert space". This description is based of the key concept *quadratic variation* of a continuous local martingale. It generalizes the equality (6.1.21) that we needed in the proof of Itô's formula.

Conceptually, the quadratic variation is something intimately related to the Doob decomposition of submartingales.

Recall that, in the discrete case, for any submartingale $(S_n)_{n \geq 0}$ such that $S_0 = 0$, there exists a unique *predictable* increasing process $(C_n)_{n \geq 0}$ such that $C_0 = 0$ and $S_n - C_n$ is a martingale. In particular, if $(M_n)_{n \geq 0}$ is an L^2 -martingale such that $M_0 = 0$, then there exists a unique *predictable* increasing process $(Q_n)_{n \geq 0}$ such that $Q_0 = 0$ and $M_n^2 - Q_n$ is a martingale. The process Q is called the quadratic variation of the discrete L^2 -martingale M ; see Example 3.1.16.

In the continuous time case, the existence of a Doob decomposition is a rather delicate matter. That includes the special case of quadratic variation. There are different ways of constructing the quadratic variation, but each involves considerable effort. We chose an approach that avoids certain foundational issues, but requires a bit more analytical effort.

In the remainder of this section we will work with a complete filtered probability space

$$(\Omega, \mathcal{S}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$$

and we will assume that the filtration satisfies the usual conditions.

6.2.1. Quadratic variation of a local martingale. The next result plays a fundamental role in the theory of stochastic integral.

Theorem 6.2.1. *Let $M = (M_t)_{t \geq 0}$ be a continuous local martingale such that $M_0 = 0$. There exists a, unique up to indistinguishability, increasing continuous process $[M] = ([M]_t)_{t \geq 0}$, such that $M^2 - [M]$ is a local martingale. Moreover, for any $T > 0$ we have*

$$[M]_T = \lim_{\|P\| \rightarrow 0} Q_T(M, P) \text{ in probability,} \quad (6.2.1)$$

where for any subdivision $P = \{0 = t_0 < t_1 < \dots < t_p = T\}$ of $[0, T]$ we denote by $\|P\|$ its mesh and we denoted by $Q_T(M, P)$ the quadratic variation of M along P , i.e.,

$$Q_T(M, P) := \sum_{i=1}^p (M_{t_i} - M_{t_{i-1}})^2.$$

The process $[M]$ is called the quadratic variation process associated to the continuous local martingale M .

Proof. This approach is inspired by the presentations in [67, §2.3] and [132, §4.3]. We carry the proof in several steps.

Step 0. The uniqueness of $[M]_t$ follows from Proposition 6.1.32.

Step 1.⁴ We prove the existence of $[M]$ in the special case when $M_0 = 0$ and M is a bounded continuous martingale. In this case M is a UI continuous martingale.

Fix $T > 0$. Given a subdivision

$$P = \{0 = t_0 < t_1 < \dots < t_p = T\}$$

of $[0, T]$, we set

$$\|P\| := \max_{1 \leq k \leq p} (t_k - t_{k-1}),$$

and

$$X_t^P = X_t^P(M) := \sum_{i=1}^p X_t^i, \quad X_t^i = M_{t_{i-1} \wedge t} (M_{t \wedge t_i} - M_{t \wedge t_{i-1}}), \quad t \geq 0.$$

⁴This is the meatiest part of the proof.

More precisely,

$$X_t^i = \begin{cases} M_{t_{i-1}}(M_t - M_{t_{i-1}}), & t \geq t_i, \\ M_{t_{i-1}}(M_t - M_{t_{i-1}}), & t \in [t_{i-1}, t_i), \\ 0, & t < t_{i-1}. \end{cases}$$

The next elementary result describe the key property of X_t^i .

Lemma 6.2.2. *The process X_t^i is a bounded martingale for any $i = 1, \dots, p$.*

Proof. Let $0 \leq s < t$. The following hold.

- If $s \geq t_i$, then X_t^i is $\mathcal{F}_s$ measurable and thus

$$\mathbb{E}[X_t^i \mid \mathcal{F}_s] = M_{t_{i-1}}(M_t - M_{t_{i-1}}) = X_s^i.$$

- If $s \in [t_{i-1}, t_i)$, then $M_{t_{i-1} \wedge t} = M_{t_{i-1}}$ is $\mathcal{F}_s$ -measurable and thus

$$\mathbb{E}[X_t^i \mid \mathcal{F}_s] = M_{t_{i-1}} \mathbb{E}[(M_t - M_{t_{i-1}}) \mid \mathcal{F}_s] = M_{t_{i-1}}(M_s - M_{t_{i-1}}) = X_s^i.$$

- If $s < t < t_{i-1}$, then $X_s^i = X_t^i = 0$.
- If $s < t_{i-1} \leq t$, then

$$\begin{aligned} \mathbb{E}[X_t^i \mid \mathcal{F}_s] &= \mathbb{E}[\mathbb{E}[X_t^i \mid \mathcal{F}_{t_{i-1}}] \mid \mathcal{F}_s] \\ &= \mathbb{E}\left[M_{t_{i-1}} \underbrace{\mathbb{E}[(M_{t_i \wedge t} - M_{t_{i-1}}) \mid \mathcal{F}_{t_{i-1}}]}_{=0} \mid \mathcal{F}_s\right] = 0 = X_s^i. \end{aligned}$$

□

This shows that X_t^P is a *martingale*. Observe that if $t \in (t_{k-1}, t_k]$, then

$$X_t^P(M) = \sum_{i=1}^{k-1} M_{t_{i-1}}(M_{t_i} - M_{t_{i-1}}) + M_{t_{k-1}}(M_t - M_{t_{k-1}}).$$

For any adapted process A_t and any partition

$$P = \{t_0 < t_1 < \dots < t_p = T\}$$

of $[0, T]$ we define the process

$$Q_t^P(A) := \sum_{i=1}^p Q_t^i(A), \quad Q_t^i(A) := (A_{t \wedge t_i} - A_{t \wedge t_{i-1}})^2.$$

Note that for any $t \in (0, T]$, the partition P of $[0, T]$ induces a partition P_t of $[0, t]$,

$$P_t = t_0 < t_1 < \dots < t_{k(t)} < t \leq T, \quad k(t) := \max \{k \leq p : t_k < t\}.$$

Then

$$Q_t^P(A) = Q_t(A, P_t).$$

A simple computation shows that, $\forall t \in [0, T]$ we have

$$M_t^2 - Q_t^P(M) = 2X_t^P(M), \tag{6.2.2a}$$

$$\mathbb{E}[Q_t^P(M)] = \mathbb{E}[M_t^2]. \tag{6.2.2b}$$

In particular, for $T \geq t > s$ we have

$$\begin{aligned}\mathbb{E}[Q_t^P(M) - Q_s^P(M) \mid \mathcal{F}_s] &= \mathbb{E}[M_t^2 - M_s^2 \mid \mathcal{F}_s] \\ &= \mathbb{E}[(M_t - M_s)^2 \mid \mathcal{F}_s].\end{aligned}\tag{6.2.3}$$

Lemma 6.2.3. *If $|M_t| \leq K$ for any t , then for any $T > 0$ and any partition P of $[0, T]$ we have*

$$\mathbb{E}[Q_T^P(M)^2] \leq 12K^4.\tag{6.2.4}$$

Proof of Lemma 6.2.3. Suppose

$$P = \{0 = t_0 < t_1 < \cdots < t_p = T\}.$$

Then

$$\begin{aligned}Q_T^P(M)^2 &= \left(\sum_{k=1}^p (M_{t_k} - M_{t_{k-1}})^2 \right)^2 \\ &= \sum_{k=1}^p (M_{t_k} - M_{t_{k-1}})^4 \\ &\quad + 2 \sum_{k=1}^{p-1} (M_{t_k} - M_{t_{k-1}})^2 (Q_T^P(M) - Q_{t_k}^P(M)).\end{aligned}\tag{6.2.5}$$

We have

$$\begin{aligned}\mathbb{E}\left[\sum_{k=1}^p (M_{t_k} - M_{t_{k-1}})^4 \right] &\leq 4K^2 \mathbb{E}\left[\sum_{k=1}^p (M_{t_k} - M_{t_{k-1}})^2 \right] \\ &= 4K^2 \sum_{k=1}^p \mathbb{E}[M_{t_k}^2 - M_{t_{k-1}}^2] = 4K^2 \mathbb{E}[M_T^2] \leq 4K^4.\end{aligned}$$

Next,

$$\begin{aligned}&\mathbb{E}\left[(M_{t_k} - M_{t_{k-1}})^2 (Q_T^P(M) - Q_{t_k}^P(M)) \right] \\ &= \mathbb{E}\left[\mathbb{E}\left[(M_{t_k} - M_{t_{k-1}})^2 (Q_T^P(M) - Q_{t_k}^P(M)) \mid \mathcal{F}_{t_k} \right] \right] \\ &= \mathbb{E}\left[(M_{t_k} - M_{t_{k-1}})^2 \mathbb{E}\left[(Q_T^P(M) - Q_{t_k}^P(M)) \mid \mathcal{F}_{t_k} \right] \right] \\ &\stackrel{(6.2.3)}{=} \mathbb{E}\left[(M_{t_k} - M_{t_{k-1}})^2 \mathbb{E}\left[(M_T^2 - M_{t_k}^2) \mid \mathcal{F}_{t_k} \right] \right] \\ &\leq 4K^2 \mathbb{E}\left[(M_{t_k} - M_{t_{k-1}})^2 \right] = \mathbb{E}[M_{t_k}^2 - M_{t_{k-1}}^2].\end{aligned}$$

Hence

$$\begin{aligned}&\mathbb{E}\left[\sum_{k=1}^{p-1} (M_{t_k} - M_{t_{k-1}})^2 (Q_T^P(M) - Q_{t_k}^P(M)) \right] \\ &\leq 4K^2 \sum_{k=1}^{p-1} \mathbb{E}[M_{t_k}^2 - M_{t_{k-1}}^2] = 4K^2 \mathbb{E}[M_{t_{p-1}}^2] \leq 4K^4.\end{aligned}$$

This proves the lemma. $\square$

Lemma 6.2.4. *Let $T > 0$. If $(P^\nu)_{\nu \in \mathbb{N}}$ is a sequence of partitions of $[0, T]$ such that $\|P^\nu\| \rightarrow 0$, then the sequence $Q_T^{P^\nu}$ converges in L^2 .*

Proof of Lemma 6.2.4. For simplicity, in the sequel I will write Q^P instead of Q_T^P . If P, P' are two partitions of $[0, T]$ we denote by $P \vee P'$ the partition whose set of nodes is the union of the sets of nodes of P and P' .

From (6.2.2a) we deduce that for any two partitions P, P' of $[0, T]$ the difference

$$Y_t := Q_t^P(M) - Q_t^{P'}(M)$$

is a bounded martingale. By construction, $Y_t^2 - Q_t^{P \vee P'}(Y)$ is also a bounded martingale. Moreover

$$\mathbb{E} \left[(Q_t^P(M) - Q_t^{P'}(M))^2 \right] = \mathbb{E} [Y_t^2] \stackrel{(6.2.2b)}{=} \mathbb{E} [Q_t^{P \vee P'}(Y)]. \quad (6.2.6)$$

Note that for any two adapted processes A, B and any partition

$$P = \{0 = t_0 < t_1 < \dots < t_p = T\}$$

we have

$$\begin{aligned} Q_t^P(A+B) &= \sum_{i=1}^p Q_t^i(A+B) \\ Q_t^i(A+B) &= ((A+B)_{t \wedge t_i} - (A+B)_{t \wedge t_{i-1}})^2 \\ &= \left((A_{t \wedge t_i} - A_{t \wedge t_{i-1}}) + (B_{t \wedge t_i} - B_{t \wedge t_{i-1}}) \right)^2 \\ &\leq 2(A_{t \wedge t_i} - A_{t \wedge t_{i-1}})^2 + 2(B_{t \wedge t_i} - B_{t \wedge t_{i-1}})^2 = 2Q_t^i(A) + 2Q_t^i(B). \end{aligned}$$

Hence

$$Q_t^P(A+B) \leq 2Q_t^P(A) + 2Q_t^P(B). \quad (6.2.7)$$

If we write Q^P for the process $t \mapsto Q_t^P(M)$ we deduce

$$\begin{aligned} Q_t^{P \vee P'}(Y) &= Q_t^{P \vee P'} \left(\underbrace{Q^P(M) - Q^{P'}(M)}_{=Y} \right) \\ &\stackrel{(6.2.7)}{\leq} 2Q_t^{P \vee P'}(Q^P) + 2Q_t^{P \vee P'}(Q^{P'}). \end{aligned} \quad (6.2.8)$$

Using (6.2.6) we deduce

$$\mathbb{E} \left[(Q_t^P(M) - Q_t^{P'}(M))^2 \right] \leq 2\mathbb{E} [Q_t^{P \vee P'}(Q^P)] + 2\mathbb{E} [Q_t^{P \vee P'}(Q^{P'})]. \quad (6.2.9)$$

Thus, to prove the lemma, it suffices to show that

$$\lim_{\|P\| + \|P'\| \rightarrow 0} \mathbb{E} [Q_T^{P \vee P'}(Q^P)] = 0 = \lim_{\|P\| + \|P'\| \rightarrow 0} \mathbb{E} [Q_T^{P \vee P'}(Q^{P'})]. \quad (6.2.10)$$

Let s_k be a node of $P \vee P'$ and t_j a node of P such that

$$t_j \leq s_k < s_{k+1} \leq t_{j+1}.$$

Observe that

$$\begin{aligned} Q_{s_{k+1}}^P(M) - Q_{s_k}^P(M) &= (M_{s_{k+1}} - M_{t_j})^2 - (M_{s_k} - M_{t_j})^2 \\ &= (M_{s_{k+1}} - M_{s_k})(M_{s_{k+1}} + M_{s_k} - 2M_{t_j}). \end{aligned}$$

Summing the squares we obtain

$$\begin{aligned} Q_T^{P \vee P'}(Q^P) &= \sum_{s_k \in P \vee P'} (Q_{s_{k+1}}^P(M) - Q_{s_k}^P(M))^2 \\ &\leq Q_T^{P \vee P'}(M) \sup_k (M_{s_{k+1}} + M_{s_k} - 2M_{t_{j(k)}})^2, \end{aligned}$$

where

$$j(k) := \max \{ j; t_j \leq s_k \}.$$

Using the Cauchy-Schwartz inequality we deduce

$$\begin{aligned} \mathbb{E} \left[Q_T^{P \vee P'} (Q^P) \right] &\leq \mathbb{E} \left[Q_T^{P \vee P'} (M)^2 \right]^{1/2} \cdot \mathbb{E} \left[\sup_k (M_{s_{k+1}} + M_{s_k} - 2M_{t_{j(k)}})^4 \right]^{1/2} \\ &\stackrel{(6.2.4)}{\leq} \sqrt{12K^4} \cdot \mathbb{E} \left[\sup_k (M_{s_{k+1}} + M_{s_k} - 2M_{t_{j(k)}})^4 \right]^{1/2}. \end{aligned}$$

The second factor goes to zero as $\|P\| + \|P'\| \rightarrow 0$ since M_t is bounded and uniformly continuous on $[0, T]$. $\square$

Consider now the sequence P^ν of partitions of $[0, T]$. For any $\nu > \mu$ the difference $Q_t^{P^\nu}(M) - Q_t^{P^\mu}(M)$ is a martingale and Doob's L^2 -maximal inequality (3.5.13) implies

$$\mathbb{E} \left[\sup_{t \leq T} |Q_t^{P^\nu}(M) - Q_t^{P^\mu}(M)|^2 \right] \leq 4\mathbb{E} \left[|Q_T^{P^\nu}(M) - Q_T^{P^\mu}(M)|^2 \right]. \quad (6.2.11)$$

Lemma 6.2.5. *There exists a subsequence $\nu(k)$ such that $Q_t^{P^{\nu(k)}}$ converges in L^2 uniformly on $[0, T]$.*

Proof. Using Lemma 6.2.4 we can find an increasing sequence $\nu(k)$ such that

$$4\mathbb{E} \left[|Q_T^{P^\nu}(M) - Q_T^{P^{\nu(k)}}(M)|^2 \right] \leq \frac{1}{2^k}, \quad \forall \nu \geq \nu(k).$$

Using Chebyshev's inequality we deduce from (6.2.11) that

$$\mathbb{P} \left[\sup_{t \leq T} |Q_t^{P^{\nu(k+1)}}(M) - Q_t^{P^{\nu(k)}}(M)| > \frac{1}{k^2} \right] \leq \frac{k^4}{2^k}.$$

Using the Borel-Cantelli lemma we deduce that the sequence $Q_t^{P^{\nu(k)}}$ converges in L^2 uniformly on $[0, T]$ to a process A_t . $\square$

Thus any subsequence of $Q_t^{P^\nu}$ has a sub-subsequence that converges in L^2 uniformly on $[0, T]$ proving that the entire sequence $Q_t^{P^\nu}$ converges in L^2 uniformly on $[0, T]$.

By letting $T = N \in \mathbb{N}$ and using a diagonal procedure we can find a subsequence of partitions P^ν of $[0, \infty)$ of the form

$$\frac{k}{2^\nu}, \quad k = 0, 1, \dots,$$

$Q_t^{P^\nu}(M)$ converges in L^2 to a process A_t , uniformly on $[0, N]$, $\forall N \in \mathbb{N}$. In particular, the process A_t is continuous.

The processes $Q_t^{P^\nu}$ are not increasing but become so when restricted to the set of nodes of P^ν . Since the union of the nodes of P^ν is dense in $[0, \infty)$ we deduce that A_t is also increasing. The sequence of martingales

$$Z_t^\nu := M_t^2 - Q_t^{P^\nu}$$

converges in L^2 to $Z_t = M_t^2 - A_t$. Since the map $X \mapsto \mathbb{E}[X | \mathcal{F}_s]$ is $L^p \rightarrow L^p$ continuous we deduce that Z_t is a martingale. Thus when M_t is a bounded martingale we have $[M]_t = A_t$.

The equality (6.2.1) is now a consequence of the above convergences. In this case we have

$$[M]_T = \lim_{\|P\| \rightarrow 0} Q_T(M, P) \text{ in } L^2.$$

If M does not vanish at 0, then we set

$$[M] = [M'], \quad M' = M - M_0.$$

Step 2. Suppose that M is a local martingale. The existence of the quadratic variation process $[M]_t$ is a consequence of the following localization result.

Lemma 6.2.6. *Suppose that M is a bounded martingale and T is a stopping time. Then*

$$[M^T]_t = [M]_t^T = [M]_{T \wedge t}.$$

Proof. From the optional stopping time theorem we deduce that

$$(M^2 - [M])^T = (M^T)^2 - [M]^T$$

is a martingale. The result is now a consequence of the uniqueness of the quadratic variation process. $\square$

Consider now a sequence of stopping times (T_n) increasing to ∞ such that

$$Y^n = M^{T_n} \mathbf{I}_{\{T_n > 0\}}$$

is a bounded martingale. According to **Step 1** there exists a unique increasing continuous process A_t^n such that $(Y_t^n)^2 - A_t^n$ is a martingale. From the uniqueness result in Proposition 6.1.32 we deduce that

$$A_t^n = A_t^{n+1} \quad \forall t \leq T_n.$$

We can define unambiguously

$$[M]_t = A_t^n, \quad t \leq T_n.$$

Using Lemma 6.2.6 we deduce that

$$M_{T_n \wedge t}^2 \mathbf{I}_{\{T_n > 0\}} - [M]_{T_n \wedge t} = M_{T_n \wedge t}^2 \mathbf{I}_{\{T_n > 0\}} - [M]_{T_n \wedge t}$$

is a bounded martingale proving that $M^2 - [M]$ is a local martingale.

Finally, let us prove (6.2.1) when M is a local martingale. More precisely, we will prove that if M_t is a continuous local martingale, $t > 0$, and P^ν is a sequence of partitions of $[0, t]$ such that $\|P^\nu\| \rightarrow 0$, then

$$\lim_{\nu \rightarrow \infty} \sup_{s \leq t} |Q_t^{P^\nu}(M) - [M]_s| = 0 \quad \text{in probability.} \quad (6.2.12)$$

To prove this we fix $\varepsilon, \delta > 0$ arbitrarily. We can then find a stopping time S such that X^S is a bounded martingale and $\mathbb{P}[S \leq t] < \delta$. Indeed, define

$$S_n := \inf \{ s > 0; |M_s| > n \}.$$

Then $S_n \rightarrow \infty$ a.s.. In particular, $\mathbb{P}[S_n \leq t] \rightarrow 0$. Thus S_n with n large enough has the desired property.

Lemma 6.2.6 shows that $[M^S] = [M]$ on $[0, S]$.

$$\mathbb{P} \left[\sup_{s \leq t} |Q_t^{P^\nu}(M) - [M]_s| > \varepsilon \right] \leq \delta + \mathbb{P} \left[\sup_{s \leq t} |Q_t^{P^\nu}(M^S) - [M^S]_s| > \varepsilon \right].$$

The second term on the right goes to zero as ν goes to ∞ since $Q_t^{P^\nu}(M^S)$ converges in L^2 uniformly on $[0, T]$ to $[M^S]_t$. This completes the proof of Theorem 6.2.1. $\square$

Example 6.2.7 (The quadratic variation of the Brownian motion). Let $(B_t)_{t \geq 0}$ be a standard Brownian motion. From Proposition 2.5.17 we deduce

$$[B]_t = t, \quad \forall t \geq 0.$$

□

Example 6.2.8 (The quadratic variation of an Itô process). Fix a Brownian motion B and a B -admissible filtration and consider the Itô process $X = GdB$. It is a continuous local martingale. The equality (6.1.24) shows that

$$X^2(t) - \int_0^t G(s)^2 ds = 2 \int_0^t X(s)G(s)dB(s).$$

Since the right-hand-side is a continuous local martingale we deduce that the quadratic variation of X is the increasing and continuous process

$$t \mapsto \int_0^t G(s)^2 ds.$$

□

Proposition 6.2.9. Suppose that $(M_t)_{t \geq 0}$ is a continuous local martingale such that $M_0 = 0$. Then $[M] = 0$ if and only if $M = 0$.

Proof. Suppose that $[M] = 0$. Note that $M^2 - [M]$ is a nonnegative local martingale and in particular it is a supermartingale. Hence

$$\mathbb{E}[M_t^2] \leq \mathbb{E}[M_0^2] = 0, \quad \forall t \geq 0.$$

□

Theorem 6.2.10. Suppose that M is a continuous local martingale and $M_0 \in L^2$.

(i) The following are equivalent.

(a) M_t is a L^2 -bounded martingale, i.e., $\sup_{t \geq 0} \mathbb{E}[M_t^2] < \infty$.

(b) $\mathbb{E}[M]_\infty] < \infty$.

Moreover, if the above properties hold, then the process $M_t^2 - [M]_t$ is a UI martingale and

$$\mathbb{E}[M_\infty^2] - \mathbb{E}[M_0^2] = \mathbb{E}[M]_\infty].$$

(ii) The following are equivalent.

(a) M is an L^2 -martingale.

(b) $\mathbb{E}[M]_t] < \infty, \forall t > 0$.

Moreover, if the above properties hold, then the process $M_t^2 - [M]_t$ is a true martingale.

Proof. (i) (a) $\Rightarrow$ (b) Assume that M is a martingale bounded in L^2 . By replacing M with $M - M_0$ we can assume $M_0 = 0$.

Doob's L^2 -inequality implies

$$\mathbb{E}\left[\sup_{t \leq T} M_t^2\right] \leq 4\mathbb{E}[M_T^2].$$

Letting $T \rightarrow \infty$ we deduce

$$\mathbb{E} \left[\sup_{0 \leq t} M_t^2 \right] \leq 4\mathbb{E} [M_\infty^2] =: C < \infty.$$

Set

$$S_n := \inf \{ t \geq 0 : [M]_t \geq n \}.$$

The local martingale $M_{s \wedge S_n}^2 - [M]_{s \wedge S_n}$ is dominated by the integrable random variable

$$n + \sup_{s \geq 0} M_s^2$$

and thus it is a UI martingale. Hence

$$\mathbb{E} [[M]_{s \wedge S_n}] = \mathbb{E} [M_{s \wedge S_n}^2] \leq \mathbb{E} \left[\sup_{s \geq 0} M_s^2 \right] \leq C.$$

By letting $n \rightarrow \infty$ and then $t \rightarrow \infty$ we deduce from the monotone convergence theorem that $\mathbb{E} [[M]_\infty] < \infty$.

(b) $\Rightarrow$ (a) Assume that $\mathbb{E} [[M]_\infty] < \infty$. Define

$$T_n := \inf \{ t \geq 0 : |M_t| \geq n \}.$$

The local martingale

$$M_{t \wedge T_n}^2 - [M]_{t \wedge T_n}$$

is dominated by the integrable random variable $n^2 + [M]_\infty$ and thus it is a UI martingale. Hence

$$\mathbb{E} [M_{t \wedge T_n}^2] = \mathbb{E} [[M]_{t \wedge T_n}] \leq \mathbb{E} [[M]_\infty] < \infty. \quad (6.2.13)$$

Letting $n \rightarrow \infty$ we deduce from Fatou's lemma that

$$\mathbb{E} [M_t^2] < \infty, \quad \forall t \geq 0.$$

It remains to prove that M_t is a genuine martingale. From (6.2.13) we deduce that the family $(M_{t \wedge T_n})_{n \in \mathbb{N}}$ is uniformly integrable and thus $M_{t \wedge T_n}$ converge a.s. and L^1 to M_t as $n \rightarrow \infty$. The local martingale M^{T_n} is bounded and thus it is a martingale. We deduce that

$$M = \lim_{n \rightarrow \infty} M^{T_n}$$

is also a martingale.

Finally, if (a) and (b) hold, then the local martingale

$$M^2 - [M]$$

is dominated by the integrable random variable

$$\sup_{t \geq 0} M_t^2 + [M]_\infty$$

and thus it is a UI martingale.

Part (ii) of the theorem follows by applying part (i) to the local martingales

$$M^n := (M_{t \wedge n})_{t \geq 0}, \quad n \in \mathbb{N}.$$

□

6.2.2. The bracket of two local martingales. Let $(\Omega, \mathcal{S}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a filtered complete probability space.

Definition 6.2.11. If M, N are two continuous local martingales, then we define their *bracket* or *covariance* to be the *finite variation* continuous process $\langle M, N \rangle$ defined by

$$\langle M, N \rangle_t := \frac{1}{2} \left([M + N]_t - [M]_t - [N]_t \right)$$

□

Note that

$$[M]_t = [M]_t.$$

The following result is an immediate consequence of the definition of the bracket and the properties of the quadratic variation of a local martingale.

Proposition 6.2.12. *Let M, N be two continuous local martingales. Then the following hold.*

- (i) *The bracket $\langle M, N \rangle$ is, up to indistinguishability, the unique continuous finite variation process such that $M_t N_t - \langle M, N \rangle_t$ is a local martingale.*
- (ii) *The application $(M, N) \mapsto \langle M, N \rangle$ is symmetric and bilinear.*
- (iii) *If $P^\nu = \{0 = t_0^\nu < t_1^\nu < \dots < t_{p(\nu)}^\nu = T\}$ is a sequence of partitions of $[0, T]$ such that $\|P^\nu\| \rightarrow 0$, then*

$$\lim_{\nu \rightarrow \infty} \sum_{j=1}^{p(\nu)} (M_{t_j} - M_{t_{j-1}})(N_{t_j} - N_{t_{j-1}}) = \langle M, N \rangle_T \text{ in probability.} \quad (6.2.14)$$

- (iv) *For any stopping time T we have*

$$\langle M^T, N^T \rangle = \langle M^T, N \rangle = \langle M, N \rangle^T. \quad (6.2.15)$$

- (v) *If M, N are two martingales bounded in L^2 then $MN - \langle M, N \rangle$ is a UI martingale. In particular $\langle M, N \rangle_\infty$ is well defined as a.s limit of $\langle M, N \rangle_t$ as $t \rightarrow \infty$ and*

$$\mathbb{E}[M_\infty N_\infty] - \mathbb{E}[M_0 N_0] = \mathbb{E}[\langle M, N \rangle_\infty].$$

□

Example 6.2.13. Let

$$X_i(t) = \int_0^t G_i(s) dB(s), \quad i = 1, 2,$$

be two Itô processes. The computation in Example 6.2.8 shows that

$$\langle X_1, X_2 \rangle_t = \int_0^t G_1(s) G_2(s) ds.$$

□

Theorem 6.2.14 (Kunita-Watanabe Inequality). *Let M, N be two continuous local martingales and H, K two measurable processes. Then, a.s. we have*

$$\int_0^\infty |H_s| |K_s| |d\langle M, N \rangle_s| \leq \left(\int_0^\infty H_s^2 d[M]_s \right)^{\frac{1}{2}} \left(\int_0^\infty K_s^2 d[N]_s \right)^{\frac{1}{2}}. \quad (6.2.16)$$

Proof. We set

$$\langle M, N \rangle_s^t := \langle M, N \rangle_t - \langle M, N \rangle_s, \quad s \leq t.$$

From the approximation result (6.2.14) and the Cauchy-Schwartz inequality we deduce that a.s., for all rational $s \leq t$ we have

$$|\langle M, N \rangle_s^t| \leq \sqrt{[M]_s^t} \cdot \sqrt{\langle N, N \rangle_s^t}.$$

From the continuity of M and N we deduce that this holds a.s. for all $s \leq t$. Using the above inequality for s, t consecutive nodes of a subdivision of $[u, v]$ we deduce

$$\int_u^v |d\langle M, N \rangle_\tau| \leq \sqrt{[M]_u^v} \cdot \sqrt{\langle N, N \rangle_u^v}.$$

We deduce that for any Borel set $B \subset [0, \infty)$ we have

$$\int_B |d\langle M, N \rangle_\tau| \leq \sqrt{\int_B d[M]_u} \cdot \sqrt{\int_B d\langle N, N \rangle_u}$$

If

$$h = \sum_i \lambda_i \mathbf{I}_{A_i}, \quad k = \sum_i \mu_i \mathbf{I}_{A_i}$$

are two step functions, then

$$\begin{aligned} \int h(s)k(s) |d\langle M, N \rangle_s| &= \sum_i \lambda_i \mu_i \int_{A_i} |d\langle M, N \rangle_s| \\ &\leq \left(\sum_i \lambda_i^2 \int_{A_i} d[M]_s \right)^{\frac{1}{2}} \left(\sum_i \mu_i^2 \int_{A_i} d\langle N, N \rangle_s \right)^{\frac{1}{2}} \\ &= \left(\int h(s) d[M]_s \right)^{\frac{1}{2}} \left(\int k(s) d\langle N, N \rangle_s \right)^{\frac{1}{2}}. \end{aligned}$$

This proves (6.2.16) for step functions. The general case follows from this via a limiting procedure. $\square$

☞ If M is a continuous local martingale, non necessarily vanishing at 0, then we define its quadratic variation to be

$$[M]_t = [M']_t, \quad M'_t = M_t - M_0.$$

6.2.3. Continuous semimartingales. A continuous process $(X_t)_{t \geq 0}$ is called a *semimartingale* if it has the form

$$X_t = X_0 + M_t + A_t,$$

where M is a continuous local martingale and A is finite variation continuous process, both vanishing at 0.

From Proposition 6.1.32 we deduce that the above decomposition is unique. The finite variation part is sometimes called the *compensator* of X . We will denote by $\mathcal{SM}$ the space of semimartingales.

If $Y_t = M'_t + A'_t$ is another continuous semimartingale, then we define the covariance bracket

$$\langle X, Y \rangle_t := \langle M, M' \rangle_t.$$

Proposition 6.2.15. Suppose that $P^\nu = \{0 = t_0 < t_1^\nu < \dots < t_{p_\nu}^\nu = t\}$ is a sequence of subdivisions of $[0, t]$ such that $\|P^\nu\| \rightarrow 0$. Then

$$\langle X, Y \rangle_t = \lim_{\nu \rightarrow \infty} \sum_{i=1}^{p_\nu} (X_{t_i^\nu} - X_{t_{i-1}^\nu})(Y_{t_i^\nu} - Y_{t_{i-1}^\nu})$$

in probability.

Proof. It suffices to consider only the case $X = Y$. We have

$$\begin{aligned} \sum_{i=1}^{p_\nu} (X_{t_i^\nu} - X_{t_{i-1}^\nu})^2 &= \sum_{i=1}^{p_\nu} (M_{t_i^\nu} - M_{t_{i-1}^\nu})^2 \\ &+ 2 \sum_{i=1}^{p_\nu} (M_{t_i^\nu} - M_{t_{i-1}^\nu})(A_{t_i^\nu} - A_{t_{i-1}^\nu}) + \sum_{i=1}^{p_\nu} (A_{t_i^\nu} - A_{t_{i-1}^\nu})^2. \end{aligned}$$

The first sum converges to $\langle M, M \rangle_t$ by (6.2.14).

$$\begin{aligned} \sum_{i=1}^{p_\nu} (A_{t_i^\nu} - A_{t_{i-1}^\nu})^2 &\leq \left(\max_i |A_{t_i^\nu} - A_{t_{i-1}^\nu}| \right) \sum_{i=1}^{p_\nu} |A_{t_i^\nu} - A_{t_{i-1}^\nu}| \\ &\leq \left(\max_i |A_{t_i^\nu} - A_{t_{i-1}^\nu}| \right) \int_0^t |dA_s|. \end{aligned}$$

The last term converges to zero due to the continuity of A_s . Finally

$$\begin{aligned} &\left| \sum_{i=1}^{p_\nu} (M_{t_i^\nu} - M_{t_{i-1}^\nu})(A_{t_i^\nu} - A_{t_{i-1}^\nu}) \right| \\ &\leq \left(\max_i |M_{t_i^\nu} - M_{t_{i-1}^\nu}| \right) \cdot \sum_{i=1}^{p_\nu} |A_{t_i^\nu} - A_{t_{i-1}^\nu}| \\ &\leq \left(\max_i |M_{t_i^\nu} - M_{t_{i-1}^\nu}| \right) \int_0^t |dA_s|. \end{aligned}$$

The last term converges to 0 due to the continuity of M . □

We will now construct the stochastic integral in several stages. As in the case of Itô integral discussed in the previous section the construction is based on two fundamental techniques: approximation and localization.

We fix a complete probability space $(\Omega, \mathcal{S}, \mathbb{P})$ and a filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. In the sequel all semimartingales will be adapted to $\mathcal{F}_\bullet$.

6.2.4. The stochastic integral with respect to an L^2 -bounded martingale. We denote by

$$b\mathcal{M}^2 = b\mathcal{M}^2(\mathcal{F}_\bullet),$$

the spaces of L^2 -bounded continuous martingales adapted to $\mathcal{F}_\bullet$. Thus $M \in b\mathcal{M}^2$ iff

$$\sup_{t \geq 0} \mathbb{E}[M_t^2] < \infty.$$

We denote $b\mathcal{M}_0^2$ the subspace of $b\mathcal{M}^2$ consisting of martingales vanishing at 0

We deduce from Proposition 6.2.12(v) that if $M, N \in b\mathcal{M}^2$, then $\langle M, N \rangle_\infty$ exists and it is integrable. Moreover

$$\mathbb{E}[\langle M, N \rangle_\infty] = \mathbb{E}[M_\infty N_\infty].$$

We have a bilinear, symmetric, nonnegative definite form on $b\mathcal{M}^2$

$$(M, N)_{b\mathcal{M}^2} := \mathbb{E}[\langle M, N \rangle_\infty] = \mathbb{E}[M_\infty N_\infty].$$

Since $[M]_t$ is a nondecreasing, nonnegative process we deduce that

$$[M]_\infty = 0 \Rightarrow [M] = 0 \Rightarrow M = 0.$$

Thus the above bilinear form defines an inner product on $b\mathcal{M}^2$ with norm

$$\|M\|_{b\mathcal{M}^2} = \sqrt{\mathbb{E}[[M]_\infty]} = \sqrt{\mathbb{E}[M_\infty^2]}.$$

Proposition 6.2.16. *The space $b\mathcal{M}_0^2$ with the above inner product is complete, and thus a Hilbert space.*

Proof. Suppose $(M^n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $b\mathcal{M}^2$,

$$\lim_{m, n \rightarrow \infty} \mathbb{E}[(M_\infty^m - N_\infty^n)] = 0.$$

From Doob's L^2 -inequality we deduce

$$\lim_{m, n \rightarrow \infty} \mathbb{E}\left[\sup_{t \geq 0} (M_t^m - M_t^n)^2\right] = 0.$$

Hence, for any $t \geq 0$ the sequence M_t^n converges in L^2 to a random variable M_t^∞ . We need to show that the process M^∞ is continuous.

We can find a subsequence (n_k) such that

$$\mathbb{E}\left[\sup_{t \geq 0} |M_t^{n_{k+1}} - M_t^{n_k}| \right] \sqrt{\mathbb{E}\left[\sup_{t \geq 0} (M_t^{n_{k+1}} - M_t^{n_k})^2\right]} < \frac{1}{2^k}.$$

Hence

$$\mathbb{E}\left[\sum_k \sup_{t \geq 0} |M_t^{n_{k+1}} - M_t^{n_k}| \right] < \infty$$

so that

$$\sum_k \sup_{t \geq 0} |M_t^{n_{k+1}} - M_t^{n_k}|$$

is a.s. finite proving the a.s. uniform convergence on $[0, \infty)$. $\square$

Given $M \in b\mathcal{M}^2$ we denote by $\mathbb{L}^2(M) = \mathbb{L}^2(M, \mathcal{F}^\bullet)$ the collection of progressive random processes H such that

$$\mathbb{E}\left[\int_0^\infty H_s^2 d[M]_s\right] < \infty, \quad [M]_t = [M]_t.$$

Alternatively

$$\mathbb{L}^2(M) = L^2(\Omega \times \mathbb{R}_+, \mathcal{F}_{\text{prog}} \otimes \mathcal{B}_{\mathbb{R}_+}, \mathbb{P} \otimes d[M]).$$

This is clearly a Hilbert space with inner product

$$(H, K)_{\mathbb{L}^2(M)} := \mathbb{E}\left[\int_0^\infty H_s K_s d[M]_s\right].$$

Let $\mathcal{E} = \mathcal{E}_{\text{pred}}$ denote the space of elementary predictable processes.

Proposition 6.2.17. *If $M \in b\mathcal{M}_0^2$, then the space $\mathcal{E}$ is dense in $\mathbb{L}^2(M)$.*

Proof. It suffices to show that if $K \in \mathbb{L}^2(M)$ is orthogonal to $\mathcal{E}$, then $K = 0$. Let $K \in \mathbb{L}^2(M)$, $K \perp \mathcal{E}$. Set

$$X_t := \int_0^t K_s d[M]_s, \quad t \geq 0.$$

From the Cauchy-Schwartz inequality we deduce

$$\mathbb{E} \left[\int_0^t |K_s| d[M]_s \right] \leq \mathbb{E} \left[\int_0^t |K_s|^2 d[M]_s \right]^{1/2} \mathbb{E} [M_\infty]^{1/2}.$$

We deduce that $X_t \in L^1$ and

$$\forall t \geq 0 \quad \int_0^t |K_s| d[M]_s < \infty, \quad \text{a.s.}$$

We deduce from Proposition 6.1.29 that X_t is a continuous, bounded variation process and $X_0 = 0$.

Let $0 \leq s < t$ and $F \in L^\infty(\Omega, \mathcal{F}_s)$. Set

$$H := F \mathbf{I}_{(s,t]} \in \mathcal{E}.$$

We have

$$0 = (H, K)_{\mathbb{L}^2(M)} = \mathbb{E} \left[F \int_s^t K_s d[M]_s \right] = \mathbb{E} [F(X_t - X_s)].$$

Since F is an arbitrary bounded $\mathcal{F}_s$ -measurable function we deduce

$$\mathbb{E} [X_t | \mathcal{F}_s] = X_s, \quad \forall t > s \geq 0.$$

Thus X_t is a bounded variation continuous martingale and Proposition 6.1.32 implies $X_t = 0, \forall t > 0$. Hence $K_s = 0$ a.s. with respect to $d[M]_s$, i.e., $K = 0$ in $\mathbb{L}^2(M)$. $\square$

Remark 6.2.18. We should contrast the above proof with the proof of the analogous result, the Approximation Theorem 6.1.2.

In that case, the proof was constructive in the sense that given $T > 0$ (deterministic) and $X \in \mathbb{L}^2(B^T)$ we have constructed an explicit sequence of elementary predictable processes converging to X . The above nonconstructive proof uses in a crucial manner Proposition 6.1.32.

There exist however, more laborious, constructive proofs of Proposition 6.2.17, e.g., the proof of Lemma [110, Lemma 3.2.4]. $\square$

Theorem 6.2.19. *Let $M \in b\mathcal{M}_0^2$. For each $H \in \mathcal{E}$ of the form*

$$H(\omega, t) = \sum_{i=1}^p H_i(\omega) \mathbf{I}_{(t_{i-1}, t_i]}(t), \quad H_i \in L^\infty(\Omega, \mathcal{F}_{t_{i-1}}),$$

we define $H \cdot M \in b\mathcal{M}_0^2$ by setting

$$(H \cdot M)_t := \sum_{i=1}^p H_i (M_{t \wedge t_i} - M_{t \wedge t_{i-1}}). \quad (6.2.17)$$

The map $H \mapsto H \cdot M$ extends to an isometry $\mathbb{L}^2(M) \rightarrow b\mathcal{M}_0^2$. Moreover, the process $H \cdot M$ is uniquely determined by the equality

$$\langle H \cdot M, N \rangle = H \cdot \langle M, N \rangle, \quad \forall N \in b\mathcal{M}_0^2, \quad (6.2.18)$$

where the term in the right-hand side is a Stieltjes integral with respect to the bounded variation process $\langle M, N \rangle$. If T is a stopping time, then

$$(\mathbf{I}_{[[0, T]]} H) \cdot M = (H \cdot M)^T = H \cdot M^T.$$

We often write

$$\int_0^t H_s dM_s := (H \cdot M)_t,$$

and we will refer to $H \cdot M$ the stochastic integral of H with respect to M . Using this notation the characteristic equality (6.2.18) reads

$$\left\langle \int_0^\bullet H_s dM_s, N_\bullet \right\rangle_t = \int_0^t H_s d\langle M, N \rangle_s.$$

Proof. We first prove that the linear map

$$\mathcal{E} \ni H \mapsto H \cdot M \in b\mathcal{M}_0^2$$

is an isometry. We have

$$H(s) = \sum_{i=1}^p H_i \mathbf{I}_{(t_{i-1}, t_i]}(s), \quad H_i \in L^\infty(\Omega, \mathcal{F}_{t_i})$$

so $X = H \cdot M$ is a sum of L^2 -bounded continuous martingales of the form

$$X_t^i = H_i (M_{t \wedge t_i} - M_{t \wedge t_{i-1}}), \quad 0 = t_0 < t_1 < \dots < t_p.$$

Moreover

$$X_\infty^i = H_i (M_{t_i} - M_{t_{i-1}}).$$

For $i < j$ we have

$$\begin{aligned} & \mathbb{E} [H_i H_j (M_{t_i} - M_{t_{i-1}}) (M_{t_j} - M_{t_{j-1}})] \\ &= \mathbb{E} [H_i (M_{t_i} - M_{t_{i-1}}) \mathbb{E} [H_j (M_{t_j} - M_{t_{j-1}}) \mid \mathcal{F}_i]] \\ &= \mathbb{E} [X_{t_i}^i \mathbb{E} [X_{t_j}^j \mid \mathcal{F}_i]] = \mathbb{E} [X_{t_i}^i X_{t_i}^j] = 0, \end{aligned}$$

since $X_{t_i}^j = 0$. Using (6.2.1) we deduce

$$\langle X^i, X^i \rangle_t = H_i^2 [M]_{t_i \wedge t} - [M]_{t_{i-1} \wedge t}.$$

Thus

$$\begin{aligned} \langle H \cdot M, H \cdot M \rangle_t &= \sum_{i=1}^p H_i^2 ([M]_{t_i \wedge t} - [M]_{t_{i-1} \wedge t}) \\ &= \int_0^t H(s)^2 d[M]_s. \end{aligned}$$

Hence

$$\begin{aligned} \|H \cdot M\|_{b\mathcal{M}^2}^2 &= \mathbb{E} [\langle H \cdot M, H \cdot M \rangle_\infty] \\ &= \mathbb{E} \left[\int_0^\infty H(s)^2 d[M]_s \right] = \|H\|_{\mathbb{L}^2(M)}^2. \end{aligned}$$

Since $\mathcal{E}$ is dense in $\mathbb{L}^2(M)$ and $b\mathcal{M}_0^2$ is a Hilbert space, the map

$$\mathcal{E} \ni H \mapsto H \cdot M \in b\mathcal{M}_0^2$$

extends uniquely to an isometry of $\mathbb{L}^2(M)$ into $b\mathcal{M}^2$.

To prove the equality (6.2.18) we observe that the Kunita-Watanabe inequality (6.2.16) implies that

$$\mathbb{E} \left[\int_0^\infty |H_s| d|\langle M, N \rangle_s| \right] \leq \|H_s\|_{\mathbb{L}^2(M)} \|N\|_{b\mathcal{M}^2}$$

and thus the random variable

$$(H \cdot \langle M, N \rangle)_\infty = \int_0^\infty H_s d\langle M, N \rangle_s$$

is well defined and integrable.

Let us first prove the equality (6.2.18) in the special case when $H \in \mathcal{E}$. We have

$$H_s = \sum_{i=1}^p H_i \mathbf{I}_{(t_{i-1}, t_i]}(s), \quad H_i \in L^\infty(\Omega, \mathcal{F}_{t_i})$$

so

$$\langle H \cdot M, N \rangle_t = \sum_{i=1}^p H_i \langle M_t^{t_i} - M_t^{t_{i-1}}, N \rangle_t$$

(use Proposition 6.2.12(iv))

$$\begin{aligned} & \sum_{i=1}^p H_i (\langle M^{t_i}, N^{t_i} \rangle_t - \langle M^{t_{i-1}}, N^{t_{i-1}} \rangle_t) \\ &= \sum_{i=1}^p H_i (\langle M, N \rangle_{t_i \wedge t} - \langle M, N \rangle_{t_{i-1} \wedge t}) \\ &= \int_0^t H_s d\langle M, N \rangle_s = (H \cdot \langle M, N \rangle)_t. \end{aligned}$$

This proves (6.2.18) when $H \in \mathcal{E}$.

The map

$$b\mathcal{M}^2 \ni X \mapsto \langle X, N \rangle_\infty \in L^1$$

is also continuous. Indeed, the Kunita-Watanabe inequality (6.2.16) shows that

$$\mathbb{E} [|\langle X, N \rangle_\infty|] \leq \mathbb{E} [[X]_\infty]^{1/2} \mathbb{E} [[N]_\infty]^{1/2} = \|N\|_{b\mathcal{M}^2} \|X\|_{b\mathcal{M}^2}.$$

Next, observe that the map

$$L^2(M) \ni H \mapsto (H \cdot \langle M, N \rangle)_\infty \in L^1,$$

is also continuous. Indeed, the Kunita-Watanabe inequality shows that

$$\begin{aligned} \mathbb{E} \left[\left| \int_0^\infty H d\langle M, N \rangle \right| \right] &\leq \mathbb{E} \left[\int_0^\infty H_s^2 d[M]_s \right]^{1/2} \mathbb{E} [[N]_\infty]^{1/2} \\ &= \|N\|_{b\mathcal{M}^2} \|H\|_{\mathbb{L}^2(M)}. \end{aligned}$$

If now H^n is a sequence in $\mathcal{E}$ that converges to $H \in \mathbb{L}^2(M)$, then

$$\begin{aligned} \langle H \cdot M, N \rangle_\infty &= \lim_{n \rightarrow \infty} \langle H^n \cdot M, N \rangle_\infty \\ &= \lim_{n \rightarrow \infty} (H^n \cdot \langle M, N \rangle)_\infty = (H \cdot \langle M, N \rangle)_\infty. \end{aligned}$$

If we replace N with the stopped process N^t in the above equality we obtain (6.2.18).

It is not hard to see that the equality (6.2.18) determines $H \cdot M$ uniquely. Indeed, if X is another martingale satisfying the same property we deduce

$$\langle H \cdot M - X, N \rangle = 0.$$

Letting $N = H \cdot M - X$ we deduce $\langle N, N \rangle = 0$. Proposition 6.2.9 implies that $N = 0$, so $X = H \cdot M$.

We only need to verify the stopping property. We have

$$\begin{aligned} \langle (H \cdot M)^T, N \rangle_t &= (\langle H \cdot M, N \rangle)_{T \wedge t} \\ &= (H \cdot \langle M, N \rangle)_{T \wedge t} = (\mathbf{I}_{[0, T]} H \cdot \langle M, N \rangle)_t, \end{aligned}$$

showing that $\mathbf{I}_{[0, T]} H \cdot M$ satisfies the characterizing equality (6.2.18). The equality

$$H \cdot M^T = \mathbf{I}_{[0, T]} H \cdot M$$

is proved in a similar fashion. $\square$

Corollary 6.2.20. *If $M \in b\mathcal{M}^2$ and $H \in \mathbb{L}^2(M)$, then the quadratic variation of $H \cdot M \in b\mathcal{M}^2$ is*

$$[H \cdot M]_t = \langle H \cdot M, H \cdot M \rangle_t = \int_0^t H_s^2 d[M]_s$$

Proof. Using (6.2.18) with $N = H \cdot M$ we deduce

$$\langle H \cdot M, H \cdot M \rangle = H \cdot \langle M, H \cdot M \rangle.$$

Using (6.2.18) with $N = M$ we deduce

$$\langle H \cdot M, M \rangle = H \cdot \langle M, M \rangle.$$

Proposition 6.1.31 implies that

$$\langle H \cdot M, H \cdot M \rangle = H \cdot (H \cdot [M]) = H^2 \cdot [M].$$

$\square$

Proposition 6.2.21. *If $K \in \mathbb{L}^2(M)$ and $H \in \mathbb{L}^2(K \cdot M)$, then $HK \in \mathbb{L}^2(M)$ and*

$$H \cdot (K \cdot M) = (HK) \cdot M, \tag{6.2.19}$$

i.e.

$$\int_0^t H_s d(K \cdot M)_s = \int_0^t H_s K_s dM_s.$$

Proof. Using Corollary 6.2.20 we deduce that

$$[K \cdot M]_t = \int_0^t K_s^2 d[M]_s.$$

Hence

$$\int_0^r H_s^2 d[K \cdot M]_s = \int_0^r H_s^2 K_s^2 d[M]_s$$

so $HK \in L^2(M)$. To prove (6.2.19) observe that for $N \in b\mathcal{M}_0^2$ we have

$$\langle (KH) \cdot M, N \rangle \stackrel{(6.2.18)}{=} (KH) \cdot \langle M, N \rangle.$$

Using Proposition 6.1.31 we deduce that

$$(KH) \cdot \langle M, N \rangle = K \cdot (H \cdot \langle M, N \rangle)$$

$$\stackrel{(6.2.18)}{=} K \cdot \langle H \cdot M, N \rangle \stackrel{(6.2.18)}{=} \langle K \cdot (H \cdot M), N \rangle.$$

Hence, $\forall N \in b\mathcal{M}^2$

$$\langle (KH) \cdot M, N \rangle = \langle K \cdot (H \cdot M), N \rangle.$$

The equality (6.2.19) now follows from the uniqueness part of Theorem 6.2.19. $\square$

6.2.5. Stochastic integration with respect to local martingales. If M is a continuous local martingale we denote by $\mathbb{L}^2(M)$ the space of progressive processes H such that

$$\mathbb{E} \left[\int_0^\infty H_s^2 d[M]_s \right],$$

and by $\mathcal{L}_{\text{loc}}^2(M)$ the space of progressive processes H such that

$$\mathbb{P} \left[\int_0^t H_s^2 d[M]_s < \infty, \forall t \geq 0 \right] = 1.$$

We denote by $c\mathcal{M}_{\text{loc}}$ the space of continuous local martingales and $c\mathcal{M}_{\text{loc}}^0$ the space of continuous local martingales vanishing at 0.

Theorem 6.2.22. *Let $M \in c\mathcal{M}_{\text{loc}}^0$. For any $H \in \mathcal{L}_{\text{loc}}^2(M)$ there exist a local martingale $H \cdot M \in c\mathcal{M}_{\text{loc}}^0$ uniquely determined by the requirement*

$$\langle H \cdot M, N \rangle = H \cdot \langle M, N \rangle, \quad \forall N \in c\mathcal{M}_{\text{loc}}^0, \quad (6.2.20)$$

where the stochastic integral on the right-hand-side is a pathwise Stieltjes integral, i.e.,

$$\langle H \cdot M, N \rangle_t(\omega) = \int_0^t H_s(\omega) d\langle M, N \rangle_s(\omega). \quad (6.2.21)$$

Moreover, the following hold.

- (i) If $M \in b\mathcal{M}_0^2$, and $H \in \mathbb{L}^2(M)$, then $H \cdot M$ coincides with the stochastic integral defined in the previous subsection.
- (ii) For any stopping time T we have

$$(\mathbf{I}_{[0,T]} H) \cdot M = (H \cdot M)^T = H \cdot M^T. \quad (6.2.22)$$

- (iii) If $H \in \mathcal{L}_{\text{loc}}^2(M)$ and K is a progressive process such that $K \in \mathcal{L}_{\text{loc}}^2(H \cdot M)$, then $KH \in \mathcal{L}_{\text{loc}}^2(M)$ and

$$(KH) \cdot M = K \cdot (H \cdot M). \quad (6.2.23)$$

Proof. We can assume that for any $\omega \in \Omega$

$$\forall t > 0, \quad \int_0^t H_s(\omega)^2 d[M]_s(\omega) < \infty.$$

For $n \in \mathbb{N}$ we define the stopping time

$$T_n = \inf \left\{ t > 0; \int_0^t (1 + H_s^2) d[M]_s \right\}.$$

The sequence (T_n) is increasing and

$$\lim_{n \rightarrow \infty} T_n = \infty.$$

Since

$$[M^{T_n}]_t = [M]_{t \wedge T_n} \leq n$$

we deduce $M^{T_n} \in b\mathcal{M}^2$. Moreover

$$\int_0^\infty H_t^2 d[M^{T_n}]_t = \int_0^{T_n} H_s^2 d[M]_s \leq n$$

so $H \in \mathbb{L}^2(M^{T_n})$. Hence we can define $H \cdot M^{T_n}$ in the sense of Theorem 6.2.19. The localization property (6.2.22) shows that if $N > n$ we have

$$H \cdot M^{T_n} = (H \cdot M^{T_N})^{T_n}.$$

We deduce that there exists a unique continuous process X such that

$$\forall n \in \mathbb{N}, \quad X^{T_n} = H \cdot M^{T_n}.$$

Set $H \cdot M := X$. Since the processes $H \cdot M^{T_n}$ are martingales in $b\mathcal{M}^2$, we deduce that $H \cdot M$ is a local martingale.

To verify the characterization (6.2.20), let $N \in c\mathcal{M}_{\text{loc}}^0$. Set

$$T'_n := \inf \{ t > 0; |N_t| \geq n \}, \quad S_n := T_n \wedge T'_n.$$

We have $N^{T'_n} \in \mathcal{M}^{2,b}$ and

$$\begin{aligned} \langle H \cdot M, N \rangle^{S_n} &\stackrel{(6.2.15)}{=} \langle (H \cdot M)^{T_n}, N^{T'_n} \rangle = \langle H \cdot M^{T_n}, N^{T'_n} \rangle \\ &= H \cdot \langle M^{T_n}, N^{T'_n} \rangle = H \cdot \langle M, N \rangle^{S_n} = (H \cdot \langle M, N \rangle)^{S_n}. \end{aligned}$$

Letting $n \rightarrow \infty$ we deduce

$$\langle H \cdot M, N \rangle = H \cdot \langle M, N \rangle.$$

The same argument as in the proof of Theorem 6.2.19 shows Proposition 6.2.9 implies that (6.2.20) determines $H \cdot M$.

The properties (ii) f and (iii) follow from (6.2.20) exactly as in the proof of Theorem 6.2.19.

Finally, let us prove (i). Let $M \in b\mathcal{M}_0^2$ and $H \in \mathbb{L}^2(M)$. The equality (6.2.20) shows that

$$\langle H \cdot M, H \cdot M \rangle = H^2[M]$$

showing that $H \cdot M \in b\mathcal{M}^2$.

The characterizing equalities (6.2.18) and (6.2.20) now imply that the definition of stochastic integrals in Theorem 6.2.19 and Theorem 6.2.22 agree. $\square$

Let $M \in c\mathcal{M}_{\text{loc}}^0$ and $H \in \mathcal{L}_{\text{loc}}^2(M)$. Then $H \cdot M \in c\mathcal{M}_{\text{loc}}^0$ and, using (6.2.21) we deduce

$$\begin{aligned} \langle H \cdot M, H \cdot M \rangle_t(\omega) &= \int_0^t H_s(\omega) d\langle M, H \cdot M \rangle_s(\omega) \\ &= \int_0^t H_s(\omega)^2 d\langle M, M \rangle_s(\omega). \end{aligned}$$

In other words, the quadratic variation of $H \cdot M$ is

$$\int_0^t H_s(\omega)^2 d[M]_s(\omega).$$

This generalizes the result in Example 6.2.8 that we obtained as a consequence of Itô's formula for the stochastic integral with respect to the Brownian motion.

By polarization we deduce that $\forall M, N \in c\mathcal{M}_{\text{loc}}^0$ and $H, K \in \mathcal{L}_{\text{loc}}^\infty$ we have

$$\langle H \cdot M, K \cdot N \rangle = (HK) \cdot \langle M, N \rangle, \quad (6.2.24)$$

i.e.,

$$\langle H \cdot M, K \cdot N \rangle_t = \int_0^t H_s K_s d\langle M, N \rangle_s.$$

☞ If M is a continuous local martingale, non necessarily vanishing at 0, and $H \in \mathcal{L}_{\text{loc}}^2$, then we define the stochastic integral $H \cdot M$ to be

$$H \cdot M_t := H \cdot M'_t, \quad M'_t = M_t - M_0 \in c\mathcal{M}_{\text{loc}}^0. \quad (6.2.25)$$

6.2.6. Stochastic integration with respect to semimartingales. Denote by $\mathcal{L}_{\text{loc}}^\infty$ the spaces of locally bounded progressive processes. Thus $H \in \mathcal{L}_{\text{loc}}^\infty$ if

$$\forall \omega, \quad \forall t > 0 \quad \sup_{s \in [0, t]} |H_s(\omega)| < \infty.$$

If X is a continuous semimartingale (see Subsection 6.2.3), then X decomposes uniquely as a sum

$$X = X_0 + M + A,$$

where M is a local martingale vanishing at 0 and A is a (continuous) finite variation process. In other words, the space $\mathcal{SM}^0$ of semimartingales vanishing at 0 admits a direct sum decomposition

$$\mathcal{SM}^0 = c\mathcal{M}_{\text{loc}}^0 \oplus \mathcal{V}^0,$$

where $\mathcal{V}^0$ denotes the space of finite variation processes vanishing at 0.

For $H \in \mathcal{L}_{\text{loc}}^\infty$ and $X = X_0 + M + A \in \mathcal{SM}^0$ we set

$$H \cdot X := H \cdot M + H \cdot A,$$

where $H \cdot A$ is defined *pathwise* by (6.1.16). Note that $H \cdot M \in c\mathcal{M}_{\text{loc}}^0$ and, since $H \in \mathcal{L}_{\text{loc}}^\infty$, $H \cdot A \in \mathcal{V}^0$. This can be viewed as a bilinear map

$$\mathcal{L}_{\text{loc}}^\infty \times \mathcal{SM}^0 \rightarrow \mathcal{SM}^0$$

compatible with the decomposition $\mathcal{SM} = c\mathcal{M}_{\text{loc}}^0 \oplus \mathcal{V}^0$. Additionally, it satisfies the following properties.

(i) For any $H, K \in \mathcal{L}_{\text{loc}}^\infty$ and any $X \in \mathcal{SM}^0$,

$$H \cdot (K \cdot X) = (HK) \cdot X.$$

(ii) For any stopping time T

$$(\mathbf{I}_{[0, T]} H) \cdot X = (H \cdot X)^T = H \cdot X^T. \quad (6.2.26)$$

Proposition 6.2.23. Suppose that $X \in \mathcal{SM}^0$ and H has the form

$$H_s(\omega) = \sum_{i=1}^p H_i(\omega) \mathbf{I}_{(t_{i-1}, t_i]}(s),$$

where $0 = t_0 < t_1 < \dots < t_p$ and H_i is $\mathcal{F}_{t_{i-1}}$ -measurable. Then $H \in \mathcal{L}_{\text{loc}}^\infty$ and

$$(H \cdot X)_t = \sum_{i=1}^p H_i(X_{t \wedge t_i} - X_{t \wedge t_{i-1}}).$$

Proof. By linearity, we can assume that $p = 1$ and

$$H = H_0 \mathbf{I}_{(t_0, t_1]},$$

where H_0 is $\mathcal{F}_{t_0}$ -measurable. The result is clearly true if X is a finite variation process so it suffices to prove it when X is a local martingale. Localizing as in the proof of Theorem 6.2.22 we can reduce this to the case when X is a bounded L^2 -integral, i.e., $X \in b\mathcal{M}^2$.

We still have one issue to deal with namely, the random variable H_0 may not be bounded so we cannot use the definition (6.2.17).

For $n \in \mathbb{N}$ define the stopping time T_n by

$$T_n(\omega) := \inf \{ t > 0; \mid H_t(\omega) \mid > n \} = \begin{cases} \infty, & |H_0(\omega)| \leq n, \\ 0, & |H_0(\omega)| > n. \end{cases}$$

We deduce that

$$\mathbf{I}_{[0, T_n]} \cdot H = H_0^n \cdot \mathbf{I}_{(t_0, t_n]}, \quad H_0^n := \mathbf{I}_{\{|H_0| \leq n\}} H_0.$$

Hence

$$H_0^n (X_{t \wedge t_1} - X_{t \wedge t_0}) = (\mathbf{I}_{[0, T_n]} \cdot H) \cdot X = H_0 \cdot X^{T_n}.$$

We reach the desired conclusion by letting $n \rightarrow \infty$. □

The stochastic integral $H \cdot X$ depends continuously on H in appropriate topologies. The next result, similar in spirit to Theorem 6.1.18, states this in more precise terms.

Theorem 6.2.24 (Approximation theorem). *Consider a continuous semimartingale*

$$X = M + A \in c\mathcal{M}_{\text{loc}}^0 \oplus \mathcal{V}^0$$

and $H \in \mathcal{L}_{\text{loc}}^\infty$. Fix $t > 0$ and suppose that $(H^n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{L}_{\text{loc}}^\infty$ such that,

$$\int_0^t (H_s^n - H_s)^2 d[X]_s \rightarrow 0 \text{ in probability,} \quad (6.2.27a)$$

$$\int_0^t |H_s^n - H_s| |dA_s| \rightarrow 0 \text{ a.s..} \quad (6.2.27b)$$

Then

$$\int_0^t H_s^n dX_s \rightarrow \int_0^t H_s dX_s \text{ in probability.}$$

Proof. We only have to verify that

$$\int_0^t H_s^n dM_s \rightarrow \int_0^t H_s dM_s \text{ in probability.}$$

We have the following technical result generalizing Lemma 6.1.19.

Lemma 6.2.25. *Let $M \in c\mathcal{M}_{\text{loc}}^0$ and $H \in \mathcal{L}_{\text{loc}}^2(M)$. Set*

$$X := H \cdot M.$$

Then, for any $t, \varepsilon, \delta > 0$ we have

$$\mathbb{P}[\mid X_t \mid > \varepsilon] \leq \mathbb{P}[[X]_t > \delta] + \frac{\delta}{\varepsilon^2}$$

Proof. I am following the strategy in the proof of [167, Lemma 10.2.25]. Set

$$T_\delta := \inf \{ t > 0; [X]_t \geq \delta \}.$$

Then

$$[X^{T_\delta}]_t \leq \delta, \quad \forall t$$

so $X^{T_\delta} \in b\mathcal{M}^2$. We have

$$\begin{aligned} \mathbb{P}[|X_t| > \varepsilon] &= \mathbb{P}[|X_t| > \varepsilon, T_\delta \leq t] + \mathbb{P}[|X_t| > \varepsilon, T_\delta > t]. \\ &\leq \mathbb{P}[\langle X, X \rangle_t \geq \delta] + \mathbb{P}[|X_t| > \varepsilon, T_\delta > t]. \end{aligned}$$

On the other hand

$$\begin{aligned} \mathbb{P}[|X_t| > \varepsilon, T_\delta > t] &= \mathbb{P}[|X_t| > \varepsilon, T_\delta > t] \\ &= \mathbb{P}[|X_t^{T_n}| > \varepsilon, T_\delta > t] \leq \mathbb{P}[|X_t^{T_n}| > \varepsilon] \end{aligned}$$

(use Chebysev's inequality)

$$\leq \frac{1}{\varepsilon^2} \mathbb{E}[(X_t^{T_n})^2] = \frac{1}{\varepsilon^2} \mathbb{E}[X_t^{T_n}] \leq \frac{\delta}{\varepsilon^2}.$$

□

Set

$$X^n := (H^n - H) \cdot M.$$

We deduce from Lemma 6.2.25 that

$$\mathbb{P}[|X_t^n| > \varepsilon] \leq \mathbb{P}[X^n]_t > \delta] + \frac{\delta}{\varepsilon^2}$$

The assumption (6.2.27a) shows that

$$\lim_{n \rightarrow \infty} \mathbb{P}[X^n]_t > \delta] = 0.$$

Hence

$$\limsup_{n \rightarrow \infty} \mathbb{P}[|X_t^n| > \varepsilon] \leq \frac{\delta}{\varepsilon^2}, \quad \forall \delta > 0.$$

□

Corollary 6.2.26. Let $M \in c\mathcal{M}_{\text{loc}}^0$ be a continuous local martingale and H a continuous adapted process. Fix $t > 0$. For any partition

$$P: 0 = t_0 < t_1 < \dots < t_n = t$$

of $[0, t]$ consider the Riemann sum

$$S_t(H, M, P) := \sum_{i=1}^n H_{t_{i-1}} (M_{t_i} - M_{t_{i-1}}).$$

If $(P^\nu)_{\nu \geq 1}$ is a sequence of partitions of $[0, t]$ such that

$$\lim_{\nu \rightarrow \infty} \|P^\nu\| = 0,$$

then

$$\lim_{\nu \rightarrow \infty} S_t(H, M, P^\nu) = H \cdot M_t \text{ in probability.} \quad (6.2.28)$$

Proof. Suppose that P^ν is given by

$$0 = t_0^\nu < t_1^\nu < \cdots < t_{N(\nu)}^\nu = t.$$

Set

$$H^\nu = \sum_{i=1}^{N(\nu)} H_{t_i^\nu}^\nu \mathbf{I}_{(t_{i-1}^\nu, t_i^\nu]}.$$

Proposition 6.2.23 shows that

$$H^\nu \cdot M_t = S_t(H, M, P^\nu).$$

Since H is continuous we deduce that

$$\lim_{\nu \rightarrow \infty} \sup_{s \in [0, t]} |H_s^\nu - H_s| = 0 \text{ a.s.}$$

Hence

$$\int_0^t (H_s^\nu - H_s)^2 d[M]_s \leq \left(\sup_{s \in [0, t]} |H_s^\nu - H_s|^2 \right) [M]_t \rightarrow 0, \text{ a.s.}$$

The equality (6.2.28) now follows from Theorem 6.2.24. $\square$

6.2.7. Itô's formula revisited. Itô's formula (6.1.19) generalizes to stochastic integrals with respect to continuous semimartingales. The effort spent constructing these more general stochastic integrals will have an immediate payoff in the form of a much simpler proof.

Theorem 6.2.27 (Itô's formula). *Let $X^1, \dots, X^N$ be continuous semimartingales and $F \in C^2(\mathbb{R}^N)$. Then $F(X^1, \dots, X^N)$ is a continuous semimartingale and*

$$\begin{aligned} F(X_t^1, \dots, X_t^N) - F(X_0^1, \dots, X_0^N) &= \sum_{j=1}^N \int_0^t F'_{x^j}(X_s^1, \dots, X_s^N) dX_s^j \\ &\quad + \frac{1}{2} \sum_{j,k=1}^N \int_0^t F''_{x^j x^k}(X_s^1, \dots, X_s^N) d\langle X^j, X^k \rangle_s. \end{aligned} \quad (6.2.29)$$

Proof. I follow the elegant approach in [67, Sec.2.10] or [177, Sec. IV.3]. The proof is carried out in several steps.

Step 1. We will prove that for any continuous semimartingales X, Y we have an integration by parts formula generalizing (6.1.31)

$$\begin{aligned} X_t Y_t - X_0 Y_0 &= \int_0^t X_s dY_s + \int_0^t Y_s dX_s + \int_0^t d\langle X, Y \rangle_s \\ &= (X \cdot Y)_t + (Y \cdot X)_t + \langle X, Y \rangle_t. \end{aligned} \quad (6.2.30)$$

In differential form this reads

$$d(XY) = X dY + Y dX + d\langle X, Y \rangle. \quad (6.2.31)$$

By polarization it suffices to prove (6.2.30) in the special case $X = Y$, i.e.,

$$X_t^2 - X_0^2 = 2 \int_0^t X_s dX_s + [X]_t. \quad (6.2.32)$$

Fix $t > 0$. Consider a sequence of partitions (P_ν) of $[0, t]$

$$P_\nu : 0 = t_0^\nu < \cdots < t_{n(\nu)}^\nu = t$$

such that $\|P_\nu\| \rightarrow 0$. Then

$$X_t^2 - X_0^2 = 2 \underbrace{\sum_{i=1}^{n(\nu)} X_{t_{i-1}^\nu} (X_{t_i^\nu} - X_{t_{i-1}^\nu})}_{=: I_\nu} + \underbrace{\sum_{i=1}^{n(\nu)} (X_{t_i^\nu} - X_{t_{i-1}^\nu})^2}_{=: Q_\nu}.$$

Corollary 6.2.26 shows that

$$\lim_{\nu \rightarrow \infty} I_\nu = 2 \int_0^t X_s dX_s \text{ in probability}$$

Proposition 6.2.15 shows that

$$\lim_{\nu \rightarrow \infty} Q_\nu = [X]_t \text{ in probability}$$

Step 2. We will show that if $F(X^1, \dots, X^n)$ satisfies (6.2.29), then,

$$G(X^1, \dots, X^N) := X^i F(X^1, \dots, X^n)$$

also satisfies (6.2.29).

Set $\vec{X}_t = (X_t^1, \dots, X_t^N)$. Using the integration-by-parts formula (6.2.30) we deduce

$$G(\vec{X}_t) - G(\vec{X}_0) = X^i \cdot F(\vec{X})_t + F(\vec{X}) \cdot X_t^i + \langle X^i, F(\vec{X}) \rangle_t.$$

Since F satisfies (6.2.29) we deduce

$$\begin{aligned} F(\vec{X})_t &= F(\vec{X}_0) + \sum_j F'_{x^j}(\vec{X}) \cdot X_t^j \\ &\quad + \frac{1}{2} \sum_{j,k} F''_{x^j x^k}(\vec{X}) \cdot \langle X^j, X^k \rangle_t. \end{aligned}$$

Now observe that

$$\left\langle X^i, \boxed{F''_{x^j x^k}(\vec{X}) \cdot \langle X^j, X^k \rangle} \right\rangle_t = 0$$

since the quadratic variation of the boxed term is 0.

On the other hand, using (6.2.24) we deduce that

$$\langle X^i, F'_{x^i}(\vec{X}) \cdot X^j \rangle = F'_{x^i}(\vec{X}) \cdot \langle X^i, X^j \rangle.$$

Moreover

$$X^i \cdot (F'_{x^j}(\vec{X}) \cdot X^j) \stackrel{(6.2.23)}{=} (X^i \cdot F'_{x^j}) \cdot X^j.$$

To conclude note that

$$\begin{aligned} G'_{x^j} &= X^i F'_{x^j} + \delta_{ji} F, \\ G''_{x^j x^k} &= X^i F''_{x^j x^k} + \delta_{ki} F'_{x^j} + \delta_{ji} F'_{x^k}. \end{aligned}$$

Step 3. To deal with the general case note that, localizing we can reduce to the case

$$\sup_{i,t} |X_t^i| < M.$$

Denote by C_M the cube $[-M, M]^N$. There exists a sequence of polynomials $Q_n : \mathbb{R}^N \rightarrow \mathbb{R}$ such that

$$\|Q_\nu - F\|_{C^2(C_N)} \rightarrow 0.$$

Invoking the Approximation Theorem we see that it suffices to prove (6.2.29) in the special case when F is a polynomial. By linearity, it suffices to prove (6.2.29) when F is a monomial. This follows immediately from **Step 2**. $\square$

Remark 6.2.28. For a proof of (6.2.29) similar in spirit to the proof of Theorem 6.1.35 I refer to [125, 132]. $\square$

Corollary 6.2.29. Let $F \in C^{1,2}(\mathbb{R} \times \mathbb{R}^N)$. Suppose that X^0 is a finite variation process and $X^1, \dots, X^N$ are continuous semimartingales. Then

$$\begin{aligned} & F(X_t^0, X_t^1, \dots, X_t^N) - F(X_0^0, X_0^1, \dots, X_0^N) \\ &= \sum_{j=0}^N \int_0^t F'_{x^j}(X_s^0, X_s^1, \dots, X_s^N) dX_s^j \\ &+ \frac{1}{2} \sum_{j,k=1}^N \int_0^t F''_{x^j x^k}(X_s^0, X_s^1, \dots, X_s^N) d\langle X^j, X^k \rangle_s. \end{aligned} \quad (6.2.33)$$

Proof. Since $\langle X^0, X^j \rangle = 0$, $\forall j = 1, \dots, N$ we deduce that if $F \in C^2$, then (6.2.33) is a special case of (6.2.29). The general case follows from this case by localizing and approximating F with C^2 -functions. $\square$

Corollary 6.2.30. Let $F \in C^{1,2}(\mathbb{R} \times \mathbb{R}^N)$. Suppose that $X^1, \dots, X^N$ are continuous semimartingales. Then

$$\begin{aligned} & F(t, X_t^1, \dots, X_t^N) - F(0, X_0^1, \dots, X_0^N) \\ &= \int_0^t F'_t(s, X_s^1, \dots, X_s^N) ds + \sum_{j=1}^N \int_0^t F'_{x^j}(s, X_s^1, \dots, X_s^N) dX_s^j \\ &+ \frac{1}{2} \sum_{j,k=1}^N \int_0^t F''_{x^j x^k}(X_s^0, X_s^1, \dots, X_s^N) d\langle X^j, X^k \rangle_s. \end{aligned} \quad (6.2.34)$$

$\square$

6.3. Some applications

We worked very hard to construct the stochastic integral with respect to a continuous semimartingale and hopefully all this effort has some payoff. That is indeed the case. Given the limited goal of this chapter, I will present only a few of the numerous applications of this theory. I refer to [132, 177, 181] for richer illustrations of the power and reach of this technique.

6.3.1. The Doléans exponential. Suppose that M is a continuous local martingale. For every complex number λ we obtain a continuous semimartingale

$$\text{EXP}(\lambda M)_t := e^{\lambda M_t - \frac{\lambda^2}{2} [M]_t}$$

Proposition 6.3.1. For any $\lambda \in \mathbb{C}$, the semimartingale $\text{EXP}(\lambda M)_t$ is a local martingale. For $\lambda \in \mathbb{R}$ it is also a supermartingale. Moreover, $\text{EXP}(\lambda M)$ satisfies the SDE

$$d \text{EXP}(\lambda M) = \lambda \text{EXP}(\lambda M) dM \quad (6.3.1)$$

i.e.,

$$\text{EXP}(\lambda M)_t = 1 + \lambda \int_0^t \text{EXP}(M)_s dM_s.$$

Proof. Consider function $F \in C^2(\mathbb{R}^2)$ given by $F(x, y) = e^{\lambda y - \frac{\lambda^2}{2}x}$. Note that

$$\text{EXP}(M)_t = F([M]_t, M_t)$$

and Itô's formula implies that

$$dF([M]_t, M_t) = \left(F'_x + \frac{1}{2} F''_{yy} \right) d[M] + F'_y dM_t.$$

Observing that

$$F'_x + \frac{1}{2} F''_{yy} = 0$$

we deduce that $\text{EXP}(M)$ satisfies the stochastic differential equations (6.3.1) and thus it is a continuous local martingale. When $\lambda \in \mathbb{R}$ is a nonnegative and thus also a supermartingale according to Proposition 6.1.24(i). $\square$

The local martingale $\text{EXP}(\lambda M)$ is called the *Doléans exponential* of M . We refer to it as *exponential* since the SDE (6.3.1) resembles the classical equation

$$d \exp(\lambda x) = \lambda \exp(\lambda x) dx.$$

Observe that when M is the Brownian motion, then $\text{EXP}(\lambda M)$ is the L^2 -martingale (3.5.2).

Proposition 6.1.24(ii) shows that $\text{EXP}(\lambda M)$ is an L^2 martingale when λ is purely imaginary. I want to analyze when $\text{EXP}(\lambda M)$ is an L^2 -martingale if λ is real. We already know that $\text{EXP}(\lambda M)$ is a supermartingale since it is a nonnegative continuous local martingale.

Proposition 6.3.2. *Let $(X_t)_{t \geq 0}$ be a continuous supermartingale adapted to a filtration $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions $\mathbb{E}[X_t] = \mathbb{E}[X_0]$, $\forall t \geq 0$, then (X_t) is a martingale. If moreover X_t is nonnegative and*

$$\mathbb{E}[X_\infty] = \mathbb{E}[X_0],$$

then (X_t) is a UI martingale

Proof. Let $s \leq t$. Since (X_t) is a supermartingale we deduce that for any $F \in \mathcal{F}_s$ we have $\mathbb{E}[X_s \mathbf{I}_F] \geq \mathbb{E}[X_t \mathbf{I}_F]$. Since $\mathbb{E}[X_s] = \mathbb{E}[X_t]$ we deduce

$$\mathbb{E}[(X_s \mathbf{I}_F - X_s)] \geq \mathbb{E}[(X_t \mathbf{I}_F - X_t)].$$

Hence, for any $F \in \mathcal{F}_s$

$$\mathbb{E}[X_s \mathbf{I}_{F^c}] \leq \mathbb{E}[X_t \mathbf{I}_{F^c}],$$

i.e., $\mathbb{E}[X_t | \mathcal{F}_s] \geq X_s$, proving that (X_t) is a martingale. The last conclusion follows from the positivity of X and the equivalence (i) $\iff$ (iii) in Theorem 3.2.21. $\square$

Theorem 6.3.3 (Novikov). *Suppose that (M_t) is a continuous local martingale such that*

$$\mathbb{E}[e^{\frac{1}{2}[M]_\infty}] < \infty. \quad (6.3.2)$$

Then $\text{Exp}(M)_t$ is a UI martingale.

Proof. I follow the ingenious approach in [123]. Assume temporarily the validity of the following result which is a weaker version of Theorem 6.3.3.

Lemma 6.3.4. *Suppose that N is a continuous local martingale such that,*

$$\exists \varepsilon > 0 : \mathbb{E} \left[e^{\frac{1+\varepsilon}{2} [N]_\infty} \right] < \infty. \quad (6.3.3)$$

Then $\text{EXP}(N)$ is a UI martingale and thus $\mathbb{E}[\text{EXP}(N)_\infty] = 1$. $\square$

Note that

$$\mathbb{E}[\text{EXP}(M)_\infty] \leq \mathbb{E}[\text{EXP}(M)_t] = \mathbb{E}[\text{EXP}(M)_0] = 1, \quad \forall t > 0$$

so it suffices to show that $\mathbb{E}[\text{EXP}(M)_\infty] \geq 1$.

Condition (6.3.2) shows that for any $\varepsilon > 0$ sufficiently small we have

$$\mathbb{E} \left[e^{\frac{(1+\varepsilon)^2}{2} [(1-\varepsilon)M]_\infty} \right] = \mathbb{E} \left[e^{\frac{(1-\varepsilon)^2}{2} [M]_\infty} \right] \leq \mathbb{E} \left[e^{\frac{1}{2} [M]_\infty} \right] < \infty.$$

Lemma 6.3.4 applied to $N = (1 - \varepsilon)M$ shows that for any sufficiently small $\varepsilon > 0$ we have

$$\mathbb{E}[\text{EXP}((1 - \varepsilon)M)_\infty] = 1.$$

In the sequel, to ease the notation, I will drop the subscript ∞ . Thus $M = M_\infty$, $[M] := [M]_\infty$ etc.

For any $\varepsilon > 0$ and any positive real number $T > 0$ we have

$$\begin{aligned} 1 &= \mathbb{E}[\text{EXP}((1 - \varepsilon)M)] = \mathbb{E}[e^{(1-\varepsilon)(M-[M]/2)} e^{\varepsilon(1-\varepsilon)[M]/2}] \\ &= \mathbb{E}[e^{(1-\varepsilon)(M-[M]/2)} e^{\varepsilon(1-\varepsilon)[M]/2} \mathbf{I}_{[M] < T}] \\ &\quad + \mathbb{E}[e^{(1-\varepsilon)(M-[M]/2)} e^{\varepsilon(1-\varepsilon)[M]/2} \mathbf{I}_{[M] > T}] \end{aligned}$$

(use Hölder's inequality)

$$\begin{aligned} &\leq \mathbb{E}[e^{(M-[M]/2)}]^{1-\varepsilon} \mathbb{E}[e^{(1-\varepsilon)[M]/2} \mathbf{I}_{[M] < T}]^\varepsilon \\ &\quad + \mathbb{E}[e^{(M-[M]/2)} \mathbf{I}_{[M] > T}]^{1-\varepsilon} \mathbb{E}[e^{(1-\varepsilon)[M]/2}]^\varepsilon. \end{aligned}$$

Letting $\varepsilon \searrow 0$ and taking into account (6.3.2) we deduce

$$1 \leq \mathbb{E}[e^{(M-[M]/2)}] + \mathbb{E}[e^{(M-[M]/2)} \mathbf{I}_{[M] > T}].$$

Finally, letting $T \rightarrow \infty$ we deduce $1 \leq \mathbb{E}[\text{EXP}(M)]$. This completes the proof of Theorem 6.3.3. $\square$

Proof of Lemma 6.3.4. For any $r, p > 1$ we have

$$p(N - [N]/2)_t = \underbrace{\left(pN - \frac{rp^2}{2} [N]/2 \right)_t}_{A_t} + \underbrace{\frac{p(rp-1)}{2} [M]_t}_{B_t}.$$

Let T be a stopping time so that N^T is a bounded martingale. We have

$$\begin{aligned} \mathbb{E}[\text{EXP}(N)_{t \wedge T}^p] &= \mathbb{E}[e^{A_{t \wedge T}} \cdot e^{B_{t \wedge T}}] \leq \mathbb{E}[e^{rA_{t \wedge T}}]^\frac{1}{r} \cdot \mathbb{E}[e^{\frac{r}{r-1} B_{t \wedge T}}]^\frac{r-1}{r} \\ &= \mathbb{E}[\text{EXP}(prN)_{t \wedge T}]^\frac{1}{r} \mathbb{E}\left[\exp\left(\frac{pr(pr-1)}{2(r-1)} [N]_{t \wedge T}\right)\right]^\frac{r-1}{r}. \end{aligned}$$

The coefficient

$$c(r, p) := \frac{pr(pr-1)}{2(r-1)}$$

can be made $< \frac{1+\varepsilon}{2}$ if we choose $p = 1 + \delta^2$, $r = 1 + \delta$, with $\delta > 0$ sufficiently small. Fix such p and r .

Since $t \mapsto [M]_t$ is nondecreasing, the condition (6.3.3) shows that

$$\sup_{t>0} \mathbb{E} \left[e^{c(p,r)} [N]_{t \wedge T} \right]$$

is bounded by a universal constant $\mathfrak{Z}$ that depends only p, r and $\mathbb{E} \left[e^{\frac{1+\varepsilon}{2} [N]_\infty} \right]$. The integrand $\text{EXP}(prN)_t$ is a positive supermartingale so that

$$\mathbb{E} \left[\text{EXP}(prN)_{t \wedge T} \right] \leq \mathbb{E} \left[\text{EXP}(prN)_t \right] = 1.$$

Hence

$$\mathbb{E} \left[\text{EXP}(N)_{t \wedge T}^p \right] \leq \mathfrak{Z},$$

for any t and any stopping time T such that N^T is bounded. We can find a localizing sequence (T_n) of such stopping times and we deduce that the submartingale is L^p -bounded. Doob's maximal inequalities (3.5.13) show that

$$\mathbb{E} \left[\sup_{t \geq 0} \text{EXP}(N)_t^p \right] \leq \text{const}(p) \cdot \mathfrak{Z}$$

Thus the submartingale $(\text{EXP}(N))_{t \geq 0}$ is dominated by an integrable random variable. Proposition 6.1.24(ii) shows that it must be a UI martingale. $\square$

Corollary 6.3.5. *Suppose that (M_t) is continuous local martingale such that*

$$\forall t > 0, \quad \mathbb{E} \left[e^{\frac{1}{2} [M]_t} \right] < \infty. \quad (6.3.4)$$

Then $\text{Exp}(M)_t$ is a martingale.

Proof. For every positive real number $t_0 > 0$ the stopped martingale M^{t_0} satisfies (6.3.2) so it is a UI martingale. Hence $(\text{EXP}(M)_t)_{t \in [0, t_0]}$ is a martingale, $\forall t_0 > 0$. $\square$

6.3.2. Lévy's characterization of Brownian motion. We know at least two characterizations of the Brownian motion. The original one is as a continuous process $B_t, t \geq 0$ such that $B_0 = 0$ and for any $t > s \geq 0$ the increment $B_t - B_s$ is a normal random variable with mean zero, variance $t - s$ and independent of $\mathcal{F}_s^B$.

The second characterization is as a continuous Gaussian process $B_t, t \geq 0$, such that

$$\mathbb{E}[B_t] = 0, \quad \mathbb{E}[B_s B_t] = \min(s, t), \quad \forall s, t \geq 0.$$

We have seen that the process $B_t^2 - t$ is a $\mathcal{F}_\bullet^B$ martingale. It turns out that the last condition alone characterizes the Brownian motion. This is essentially the content of the next theorem usually referred to as *Lévy's characterization theorem*.

Theorem 6.3.6. *Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space and $(M_t)_{t \geq 0}$ is a continuous local martingale with respect to a filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. If $M_0 = 0$ and $[M]_t = t$, $\forall t \geq 0$, then M_t is a $\mathcal{F}_\bullet$ -Brownian motion; see Definition 3.4.6.*

Proof. For any real number θ consider the Doléans exponential

$$\text{EXP} \left(i\theta M_t \right) = e^{i\theta M_t + \frac{\theta^2 t}{2}}.$$

It is bounded on every compact interval $[0, a]$ and Proposition 6.1.24(ii) shows that it is a martingale, i.e., both its real and imaginary parts are martingales. Hence, for any $t > s \geq 0$ we have

$$\mathbb{E} \left[e^{i\theta M_t + \frac{\theta^2 t}{2}} \middle| \mathcal{F}_s \right] = e^{i\theta M_s + \frac{\theta^2 s}{2}}.$$

Hence

$$\mathbb{E}[e^{i\theta(M_t - M_s)} \mid \mathcal{F}_s] = e^{-\frac{\theta^2}{2}(t-s)}.$$

In particular,

$$\mathbb{E}[e^{i\theta(M_t - M_s)}] = e^{-\frac{\theta^2}{2}(t-s)},$$

so $M_t - M_s$ has the same characteristic function as normal random variable with mean 0 and variance $(t - s)$, i.e., $M_t - M_s \sim N(0, t - s)$. Hence

$$\mathbb{E}[e^{i\theta(M_t - M_s)} \mid \mathcal{F}_s] = \mathbb{E}[e^{i\theta(M_t - M_s)}].$$

Corollary 2.2.30 implies that $M_t - M_s$ is independent of $\mathcal{F}_s$. This proves that M_t is a $\mathcal{F}_\bullet$ -Brownian motion. $\square$

6.3.3. Martingale representation theorem. Suppose that $(\Omega, \mathcal{S}, \mathbb{P})$ is a complete probability space and $(B_t)_{t \geq 0}$ is a Brownian motion. Denote by $\mathcal{F}_\bullet^B = (\mathcal{F}_t^B)_{t \geq 0}$ the completed Brownian filtration. Recall that $\mathbb{L}^2(B)$ denotes the space of $\mathcal{F}_\bullet^B$ progressive processes $(H_t)_{t \geq 0}$ such that

$$\mathbb{E}\left[\int_0^\infty H_t^2 dt\right] < \infty.$$

We denote by $\mathbb{L}_{\text{loc}}^2(B)$ the space of progressive processes $(H_t)_{t \geq 0}$ such that

$$\mathbb{E}\left[\int_0^t H_s^2 ds\right] < \infty, \quad \forall t > 0.$$

For $T > 0$ we denote by $\mathbb{L}^2([0, T], B)$ the subspace of $\mathbb{L}_{\text{loc}}^2(B)$ consisting of processes that vanish for $t > T$. Equivalently

$$\mathbb{L}^2([0, T], B) = \mathbf{I}_{[0, T]} \mathbb{L}_{\text{loc}}^2(B).$$

If $H \in \mathbb{L}_{\text{loc}}^2(B)$, then the Itô process

$$X_t = \int_0^t H_s dB_s, \quad t > 0,$$

is a L^2 -martingale. In particular, for any $T > 0$,

$$X_T \in L^2(\Omega, \mathcal{F}_T^B, \mathbb{P}), \quad \mathbb{E}[X_T] = 0.$$

Theorem 6.3.7. Let $T > 0$. Suppose that $Y \in L^2(\Omega, \mathcal{F}_T^B, \mathbb{P})$. Then there exists a unique process $H = H^Y \in \mathbb{L}^2([0, T], B)$ such that

$$Y = \mathbb{E}[Y] + \int_0^T H_s dB_s. \quad (6.3.5)$$

Proof. The uniqueness follows immediately from Itô's isometry. Denote by $\mathcal{Y}$ the subspace of $L^2(\Omega, \mathcal{F}_T^B, \mathbb{P})$ consisting of random variables that admit a representation (6.3.5). The map

$$\mathcal{Y} \ni Y \rightarrow H^Y \in \mathbb{L}^2([0, T], B)$$

satisfies

$$\| (Y_1 - \mathbb{E}[Y_1]) - (Y_2 - \mathbb{E}[Y_2]) \|_{L^2} = \| H^{Y_1} - H^{Y_2} \|_{\mathbb{L}^2}.$$

This shows that if (Y_n) is a Cauchy sequence in $\mathcal{Y}$, then (H^{Y_n}) is Cauchy in $\mathbb{L}^2(B)$. This implies that $\mathcal{Y}$ is a closed subspace of $L^2(\Omega, \mathcal{F}_T^B, \mathbb{P})$.

I will prove that $\mathcal{Y}$ contains a dense subset.

Lemma 6.3.8. For any $n \in \mathbb{N}$, any

$$0 = t_0 < t_1 < t_2 < \cdots < t_n \leq T,$$

and any $\theta_1, \dots, \theta_n \in \mathbb{R}$ we have

$$\prod_{j=1}^n e^{i\theta_j(B_{t_j} - B_{t_{j-1}})} \in \mathcal{Y}.$$

Proof. Set

$$X_t = \int_0^t \left(\sum_{j=1}^n \theta_j \mathbf{I}_{(t_{j-1}, t_j]} \right) dB_s.$$

Note that

$$X_T = \sum_{j=1}^n \theta_j (B_{t_j} - B_{t_{j-1}}), \quad [X]_T = \sum_{j=1}^n \theta_j^2 (t_j - t_{j-1})$$

and

$$e^{iX_T + \frac{1}{2}[X]_T} = \text{EXP}(iX)_T = 1 + i \int_0^T \text{EXP}(iX)_t dt.$$

Hence

$$e^{iX_T} = e^{-\frac{1}{2}[X]_T} + i \int_0^T e^{-\frac{1}{2}[X]_t} \text{EXP}(iX)_t dt \in \mathcal{Y}.$$

□

Denote by $\mathcal{Y}_0$ the subspace of $\mathcal{Y}$ spanned by the random variables of the type described in the above lemma.

I will show that $\mathcal{Y}_0$ is dense in $L^2(\Omega, \mathcal{F}_T^B, \mathbb{P})$. More precisely, I will show that if $Z \in L^2(\Omega, \mathcal{F}_T^B, \mathbb{P})$ and

$$\mathbb{E} \left[Z \prod_{j=1}^n e^{i\theta_j(B_{t_j} - B_{t_{j-1}})} \right] = 0, \quad (6.3.6)$$

for any n , any

$$0 = t_0 < t_1 < t_2 < \cdots < t_n \leq T,$$

and any $\theta_1, \dots, \theta_n \in \mathbb{R}$, then $Z = 0$. I will use the very elegant argument in [132, Lemma 5.18].

Fix $0 = t_0 < t_1 < t_2 < \cdots < t_n \leq T$ and consider the finite complex measure $\mu = \mu_{t_1, \dots, t_n}$ on $\mathbb{R}^n$ given by

$$\mu[F] = \mathbb{E} \left[Z \mathbf{I}_F(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}}) \right], \quad F \in \mathcal{B}_{\mathbb{R}^n}.$$

For any bounded Borel measurable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ we have

$$\int_{\mathbb{R}^n} f(x) \mu[dx] = \mathbb{E} \left[Z f(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}}) \right].$$

Condition (6.3.6) reads

$$\int_{\mathbb{R}^n} e^{i\theta_1 x_1 + \cdots + i\theta_n x_n} \mu[dx_1, \dots, dx_n] = 0, \quad \forall \theta_1, \dots, \theta_n \in \mathbb{R}.$$

In other words, the Fourier transform μ is zero so $\mu_{t_1, \dots, t_n}$ is trivial.

Hence, for any for any n , any $0 < t_1 < \cdots < t_n \leq T$ and any $S \in \sigma(B_{t_1}, \dots, B_{t_n})$ we have

$$\mathbb{E}[Z \mathbf{I}_S] = \mu[S] = 0.$$

This implies via the Monotone Class Theorem that

$$\mathbb{E}[Z\mathbf{I}_S] = \mu[S] = 0, \quad \forall S \in \sigma(B_t, 0 \leq t \leq T).$$

Hence $Z = 0 \in L^2\Omega, \mathcal{F}_T^B, \mathbb{P}$.

□

Theorem 6.3.9 (Martingale Representation Theorem). *Suppose that $(M_t)_{t \geq 0}$ is an L^2 R -martingale adapted to the completed Brownian filtration $\mathcal{F}_\bullet^B$. Then there exists a unique $H = H^M \in \mathbb{L}_{\text{loc}}^2(B)$ such that*

$$M_t = M_0 + \int_0^t H_s dB_s, \quad \forall t \geq 0.$$

In particular, M_t is continuous!

Proof. Observe first that since $\mathcal{F}_0^B$ is a 0-1 sigma-subalgebra, the initial value M_0 is a.s. constant. Hence

$$M_0 = \mathbb{E}[M_t], \quad \forall t.$$

Theorem 6.3.7 shows that for any $n \in \mathbb{N}$ there exists a unique $H^n \in \mathbb{L}^2([0, n], B)$ such that

$$M_n = M_0 + \int_0^n H_s^n dB_s.$$

Since

$$(M_t)_{t \geq 0} \quad \text{and} \quad \left(X_t^n = M_0 + \int_0^t H_s^n dB_s \right)_{t \geq 0}$$

are martingales we deduce that for any $t \leq n$ we have

$$M_t = \mathbb{E}[M_n | \mathcal{F}_t] = \mathbb{E}[X_t^n | \mathcal{F}_t] = M_0 + \int_0^t H_s^n dB_s.$$

The uniqueness part of Theorem 6.3.7 shows that for any $k \leq n$ we have

$$\mathbf{I}_{[0, k]} H^n = H^k.$$

We obtain a progressive process $H \in \mathbb{L}_{\text{loc}}^2(B)$ uniquely determined by the requirement

$$\mathbf{I}_{[0, n]} H = H^n, \quad \forall n \in \mathbb{N}.$$

Note that

$$M_t = M_0 + \int_0^t H_s dB_s, \quad \forall t \geq 0.$$

In particular (M_t) is continuous.

□

Remark 6.3.10. (a) Under certain additional assumptions on M , one can give an explicit description of H^M via the so called Clark-Ocone formula. A most suggestive description of this formula involves the so called Malliavin calculus. For more details I refer to [145, Thm.1.27] or [58, Thm. 4.1].

(b) The martingale representation theorem has applications in financial mathematics; see e.g. [193, Sec. 5.2, 5.3].

□

6.4. Exercises

Exercise 6.1. Suppose that (M_t) is a continuous positive local martingale such that $\mathbb{E}[M_0] < \infty$. Prove that the following are equivalent.

- (i) The process (M_t) is a martingale..
- (ii) $\mathbb{E}[M_t] = \mathbb{E}[M_0], \forall t$.

□

Exercise 6.2. Suppose that (M_t) is a continuous local martingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$ such that, for any stopping time, the family $(M_{T \wedge t})_{t \geq 0}$ is UI. Prove that (M_t) is a martingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$.

□

Exercise 6.3. Suppose that (B_t) is a Brownian motion and $(\mathcal{F}_t)$ is a B -admissible filtration. Let $\lambda > 0$.

- (i) Prove that

$$M_t = e^{\lambda t} B_t - \lambda \int_0^t e^{\lambda s} B_s ds$$

is an L^2 -integrable..

- (ii) Prove that if $\lambda < 0$ then M_t is UI.
- (iii) Compute the distribution of M_∞ .

□

Exercise 6.4. Let B_1, B_2 be two independent Brownian motions. Decide which of the following processes is also a Brownian motion.

$$X_1(t) = \int_0^t \sin(s) dB_1(s) + \int_0^t \cos(s) dB_1(s), \quad (6.4.1a)$$

$$X_1(s) = \int_0^t \sin(s) dB_1(s) + \int_0^t \cos(s) dB_2(s) \quad (6.4.1b)$$

$$X_3(t) = \int_0^t \sin B_1(s) dB_1(s) + \int_0^t \cos B_2(s) dB_2(s) \quad (6.4.1c)$$

□

Exercise 6.5. Fix a Brownian motion $\mathbf{B} = (B^1, \dots, B^n)$ on $\mathbb{R}^n$. Denote by $e\mathcal{D}$ the space of compactly supported smooth functions on $\mathbb{R}^n$. For any $f \in \mathcal{D}$ we set

$$M(f)_t := f(\mathbf{B}_t) - f(0) - \int_0^t Lf(\mathbf{B}_s) ds$$

where

$$Lf := \frac{1}{2} \sum_{i=1}^n f''_{x^i x^i}$$

Define the “carré du champ” operator

$$\Gamma : \mathcal{D} \times \mathcal{D} \rightarrow \mathcal{D}, \quad \Gamma(f, g) := L(fg) - (Lf)g - fLg.$$

- (i) Prove that for any $f \in \mathcal{D}$ the process is a continuous L^2 -martingale.
(ii) Let $f, g \in \mathcal{D}$. Prove that

$$\langle M(f), M(g) \rangle_t = 2 \int_0^t \Gamma(f, g)(B_s) ds$$

□

Exercise 6.6. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space and suppose that $\mathbb{Q}$ is equivalent to $\mathbb{P}$, i.e.,

$$\mathbb{P}[S] = 0 \iff \mathbb{Q}[S] = 0.$$

Suppose that $\mathcal{F}_t$ is a filtration satisfying the usual conditions. Set

$$\mathbb{P}_t = \mathbb{P}|_{\mathcal{F}_t}, \quad \mathbb{Q}_t = \mathbb{Q}|_{\mathcal{F}_t}, \quad \rho_t = \frac{d\mathbb{Q}_t}{d\mathbb{P}_t}.$$

In other words, ρ_t is a nonnegative $\mathcal{F}_t$ -measurable function such that

$$\forall S \in \mathcal{F}_\infty, \quad \forall F \in \mathcal{F}_t, \quad \mathbb{Q}[F] = \int_F \rho_t(\omega) \mathbb{P}[d\omega].$$

- (i) Prove that the family $(\rho_t)_{t \geq 0}$ is a UI martingale with respect to the probability measure $\mathbb{P}$. In particular, since $\mathcal{F}_t$ satisfies the usual conditions, ρ_t admits a modification as an R-martingale.
(ii) Prove that

$$\inf_{t > 0} \rho_t > -0 \quad \mathbb{P} - \text{a.s.}$$

- (iii) Suppose that the martingale ρ_t admits a continuous modification. Prove that if (X_t) is a continuous local martingale with respect to $\mathbb{P}$, it is also a continuous local martingale with respect to $\mathbb{Q}$.

□

Exercise 6.7. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space and suppose that $(X_t)_{t \geq 0}$ be a strictly positive local positive continuous local martingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$ that satisfies the usual conditions. Prove that there exists a unique continuous local martingale (M_t) such that

$$X_t = \text{EXP}(M)_t, \quad \forall t \geq 0.$$

Hint. Try $M_t = \log X_t + \int_0^t \frac{1}{X_s} dX_s$.

□

Exercise 6.8. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space and B_t a Brownian motion defined on this space. For $\mu \in \mathbb{R}$ consider the Brownian motion with a drift

$$B_t^\mu := B_t + \mu t$$

Set

$$M_t^\mu := \text{EXP}(\mu B)_t.$$

Prove that for any bounded Borel measurable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and any

$$0 \leq t_1 < t_2 < \dots < t_n < T$$

we have

$$\mathbb{E}[f(B_{t_1}^\mu, \dots, B_{t_n}^\mu)] = \mathbb{E}[f(B_{t_1}, \dots, B_{t_n}) M_T^\mu]$$

□

Exercise 6.9. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space and $\mathcal{F}_\bullet = (\mathcal{F}_t)_{t \geq 0}$ a filtration of sigma subalgebras satisfying the usual conditions.

A subset $A \subset [0, \infty) \times \Omega$ is called *evanescent* if the indicator I_A is indistinguishable from the zero process.

Denote by $\mathcal{F}_{\text{pred}}$ the sigma-subalgebra of $\mathcal{B}_{[0, \infty)} \otimes \mathcal{S}$ generated by the evanescent sets and the stochastic intervals $[[0, T]]$, where T is an $\mathcal{F}_\bullet$ -stopping time. We will refer to $\mathcal{F}_{\text{pred}}$ as the *predictable* sigma-subalgebra. A stochastic process

$$X : [0, \infty) \times \Omega \rightarrow \mathbb{R}$$

is called *predictable* if it is $e\mathcal{F}_{\text{pred}}$ -measurable.

- (i) Prove that $\mathcal{F}_{\text{pred}}$ is generated by the evanescent sets and the stochastic intervals of the form $[[0, t]]$, $t \in \mathbb{R}$.
- (ii) Prove that a predictable process is $\mathcal{F}_\bullet$ -adapted.
- (iii) Prove that any predictable process is progressive.

□

A few useful facts

A.1. The Gamma function

Definition A.1.1 (Gamma and Beta functions). The *Gamma function* is the function

$$\Gamma : (0, \infty) \rightarrow \mathbb{R}, \quad \Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt. \quad (\text{A.1.1})$$

The *Beta function* is the function of two positive variables

$$B(x, y) := \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}, \quad x, y > 0. \quad (\text{A.1.2})$$

□

We gather here a few basic facts about the Gamma and Beta functions used in the text. For proofs we refer to [129, Chap. 1] or [222, Chap. 12].

Proposition A.1.2. *The following hold.*

- (i) $\Gamma(1) = 1$.
- (ii) $\Gamma(x+1) = x\Gamma(x)$, $\forall x > 0$.
- (iii) For any $n = 1, 2, \dots$ we have

$$\Gamma(n) = (n-1)!. \quad (\text{A.1.3})$$

- (iv) $\Gamma(1/2) = \sqrt{\pi}$.
- (v) For any $x, y > 0$ we have Euler's formula

$$B(x, y) = \int_0^1 s^{x-1} (1-s)^{y-1} ds = \int_0^\infty \frac{u^{x-1}}{(1+u)^{x+y}} du. \quad (\text{A.1.4})$$

- (vi) For any $x \in (0, 1)$ we have

$$B(x, 1-x) = \Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin \pi x} \quad (\text{A.1.5})$$

□

The equality (iv) above reads

$$\sqrt{\pi} = \Gamma(1/2) = \int_0^\infty e^{-t} t^{-1/2} dt$$

$$(t = x^2, t^{-1/2} = x^{-1} dt = 2x dx)$$

$$= 2 \int_0^\infty e^{-x^2} dx = \int_{-\infty}^0 e^{-x^2} dx + \int_0^\infty e^{-x^2} dx = \int_{-\infty}^\infty e^{-x^2} dx.$$

If we make the change in variables $x = \frac{s}{\sqrt{2}}$ so that $x^2 = \frac{s^2}{2}$ and $dx = \frac{1}{\sqrt{2}} ds$, then we deduce

$$\sqrt{\pi} = \frac{1}{\sqrt{2}} \int_{-\infty}^\infty e^{-\frac{x^2}{2}} dx.$$

From this we obtain the fundamental equality

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty e^{-\frac{x^2}{2}} dx = 1. \quad (\text{A.1.6})$$

The function $\Gamma(x)$ grows very fast as $x \rightarrow \infty$. Its asymptotics is governed by the *Stirling's formula*

$$\Gamma(x+1) = x\Gamma(x) \sim \sqrt{2\pi x} \left(\frac{x}{e}\right)^x \text{ as } x \rightarrow \infty. \quad (\text{A.1.7})$$

Note that for $n \in \mathbb{N}$ the above estimate reads

$$n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \text{ as } n \rightarrow \infty. \quad (\text{A.1.8})$$

There are very sharp estimates for the ratio

$$q_n = \frac{n!}{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n}.$$

More precisely we have (see [75, II.9])

$$\frac{1}{12n+1} < \log q_n < \frac{1}{12n}. \quad (\text{A.1.9})$$

In other words

$$\log n! = n \log n - n + \frac{1}{2} \log n + \frac{1}{2} \log(2\pi) + O(n^{-1}), \text{ as } n \rightarrow \infty.$$

A.2. Basic invariants of frequently used probability distributions

$$X \sim \text{Bin}(n, p) \iff \mathbb{P}[X = k] = \binom{n}{k} p^k q^{n-k}, \quad k = 0, 1, \dots, n, \quad q = 1 - p.$$

$$\text{Ber}(p) \sim \text{Bin}(1, p).$$

$$X \sim \text{NegBin}(k, p) \iff \mathbb{P}[X = n] = \binom{n-1}{k-1} p^k q^{n-k}, \quad n = k, k+1, \dots$$

$$\text{Geom}(p) \sim \text{NegBin}(1, p).$$

$$X \sim \text{HGeom}(w, b; n), \quad \mathbb{P}[X = k] = \frac{\binom{w}{k} \binom{b}{n-k}}{\binom{w+b}{n}}, \quad k = 0, 1, \dots, w.$$

$$X \sim \text{Poi}(\lambda), \quad \lambda > 0 \iff \mathbb{P}[X = n] = e^{-\lambda} \frac{\lambda^n}{n!}, \quad n = 0, 1, \dots$$

$$\begin{aligned}
X \sim \text{Unif}(a, b) &\iff \mathbb{P}_X = \frac{1}{b-a} \mathbf{I}_{[a,b]} dx. \\
X \sim \text{Exp}(\lambda), \lambda > 0 &\iff \mathbb{P}_X = \lambda e^{-\lambda x} \mathbf{I}_{[0,\infty)} dx \\
X \sim N(\mu, \sigma^2), \mu \in \mathbb{R}, \sigma > 0 &\iff \mathbb{P}_X = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx, x \in \mathbb{R}. \\
X \sim \text{Gamma}(\nu, \lambda) &\iff p_X(x) = \frac{\lambda^\nu}{\Gamma(\nu)} x^{\nu-1} e^{-\lambda x} \mathbf{I}_{[0,\infty)} dx \\
X \sim \text{Beta}(a, b) &\iff p_X = \frac{1}{B(a, b)} x^{a-1} (1-x)^{b-1} \mathbf{I}_{(0,1)} dx. \\
X \sim \text{Stud}_p &\iff p_X = \frac{1}{\sqrt{p\pi}} \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(1+x^2/p)^{(p+1)/2}} dx, x \in \mathbb{R}.
\end{aligned}$$

Name	Mean	Variance	pgf	mgf
Ber(p)	p	pq	$(q + ps)$	pe^t
Bin(n, p)	np	npq	$(q + ps)^n$	$p^n e^{nt}$
Geom(p)	$\frac{1}{p}$	$\frac{q}{p^2}$	$\frac{ps}{1-qs}$	$\frac{pe^t}{1-qe^t}$
NegBin(k, p)	$\frac{k}{p}$	$\frac{kq}{p^2}$	$\left(\frac{ps}{1-qs}\right)^k$	$\left(\frac{pe^t}{1-qe^t}\right)^k$
Poi(λ)	λ	λ	$e^{\lambda(s-1)}$	$e^{\lambda(e^t-1)}$
HGeom($w, b; n$)	$\frac{w}{w+b} \cdot n$	*	*	*
Unif(a, b)	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$	NA	$\frac{e^{tb}-e^{ta}}{tb-ta}$
Exp(λ)	λ^{-1}	λ^{-2}	NA	$\frac{\lambda}{\lambda-t}$
$N(\mu, \sigma^2)$	μ	σ^2	NA	$\exp\left(\frac{\sigma^2}{2}t^2 + \mu t\right)$
Gamma(ν, λ)	$\frac{\nu}{\lambda}$	$\frac{\nu}{\lambda^2}$	NA	$\left(\frac{\lambda}{\lambda-t}\right)^\nu$
Beta(a, b)	$\frac{a}{a+b}$	$\frac{ab}{(a+b)^2(a+b+1)}$	NA	$1 + \sum_{k=1}^{\infty} \frac{a^{(k)}}{(a+b)^{(k)}} \frac{t^k}{k!}$
Stud $_p$	0, $p > 1$	$\frac{p}{p-2}, p > 2$	NA	NA

Above, $a^{(k)}$ denotes the ascending *Pochhammer symbol*

$$a^{(k)} = a(a+1) \cdots a(a+k-1).$$

The descending *Pochhammer symbol* or descending factorial is

$$a_{(k)} = a(a-1) \cdots a(a-k+1).$$

A.3. A glimpse at R

This section is merely an invitation to programming in R. It is not meant as a serious guide to learning R. It mainly lists a few basic tricks that will get the curious reader started and cover many with simple probability simulations I used in my classes.

First, here is how you install R on your computers.

For Mac users

<https://cran.r-project.org/bin/macosx/>

For Windows users

<https://cran.r-project.org/bin/windows/base/>

Next, install **R Studio** (the Desktop version). This is a very convenient interface for using R.

<https://www.rstudio.com/products/RStudio/>

(Install first R and then R Studio.) You can also access RStudio and R in the cloud

<https://www.rollapp.com/app/rstudio>

The GitHub¹ has a repository of many R programs (or R scripts) that you can use as models.

The reader familiar with the basics of programming will have no problems learning the basics of R. This section is addressed to such a reader.

We list some of the commands and objects most frequently used in probability and we have included several examples to help the reader get started. R-Studio comes with links to various freely available web sources for R-programming. A commercial source that I find very useful is “*The Book of R*”, [49]. I often ask **Google** how to do this or that in R and I receive many satisfactory solutions. The advent of AI, made the job of converting a pseudocode into a genuine code a lot quicker, though not error free.

Example A.3.1 (Operations with vectors). The workhorse of R is the object called *vector*. An n -dimensional vector is essentially an element in $\mathbb{R}^n$. An n -dimensional vector in R can be more general in the sense that its entries need not be just numbers. They can be quantities of the same type.

To generate in R the vector $(1, 2, 4.5)$ and then naming it x use the command

```
x<-c(1, 2, 4.5)
```

To see what the vector x is type

```
x
```

To see what the k -th entry of x is use the command

```
x[k]
```

The command

```
x[j:k]
```

will generate all the entries of x from the j -th up to and including the k -th. If you want to add an entry to x , say you want to generate the longer vector $y = (1, 2, 4, 5, 2)$, use the command

```
y<-c(x, 2)
```

If you want to remove the 3-rd entry from y use the command

```
y[-3]
```

To remove all the entries equal to 2 or 5 from y use the command

```
setdiff(y, c(2, 5))
```

Describing long vectors entry by entry can be time consuming. The process of describing vectors can be accelerated if the entries of the vector x are subject to patterns.

For example, the vector of length 22 with all entries equal to the same number, say 1.5, can be generated using the command

¹<https://github.com/>


```
rep(1.5, 22)
```

To generate the vector listing in increasing order all the integers between -2 and 10 (included) use the command

```
(-2):10
```

To generate the vector named x consisting of 25 equidistant numbers starting at 1 and ending at 7 use the command

```
x<-seq(from=1, to=7, length.out=25)
```

To add all the entries of a vector $x = (x_1, \dots, x_n)$ use the command

```
sum(x)
```

To add all the natural numbers from 50 to 200 use the command

```
sum(50:200)
```

The result is 18,875.

You can sort the entries of a vector, if they are numerical. For example

```
> z<-c(1, 4, 3)
> sort(z)
[1] 1 3 4
```

A very convenient feature of working with vectors in R is that the basic algebraic operations involving numbers extend to vectors, component wise. For example, if z is the above vector, and $y = (1, 8, 9)$, then the command y/z returns $(1/1, 8/4, 9/3) = (1, 2, 3)$, while z^2 returns $(1, 16, 9)$ $\square$

Example A.3.2 (Logical operators). These are operators whose output is a TRUE or FALSE or a vector whose entries are TRUE/FALSE.

For example, the command $2 < 5$ returns TRUE. On the other hand if x is the vector $(2, 3, 7, 8)$, then the command $x < 5$ return

```
TRUE, TRUE, FALSE, FALSE.
```

In R, the logicals TRUE/FALSE also have arithmetic meaning,

$$\text{TRUE} = 1, \text{FALSE} = 0.$$

The output of $x < 5$ is a vector whose entries are TRUE/FALSE. To see how many of the entries of x are < 5 use the command

```
sum(x<5)
```

Above $x < 5$ is interpreted as a vector with 0/1-entries. When we add them we count how many are equal to 1 or, equivalently, how many of the entries of x are < 5 .

The R language also has two very convenient logical operators `any` and `all`. When we apply `any` to a vector with TRUE/FALSE entries it returns TRUE if at least one of the entries of v are TRUE and returns FALSE otherwise. When we apply `all` to a vector v with TRUE/FALSE entries it returns TRUE if *all* of the entries of v are TRUE and returns FALSE otherwise. $\square$

Example A.3.3 (Functions in R). One can define and work with functions in R. For example, to define the function

$$f(q) = 1 + 6q + 10q^2(1 - q)^4$$

use the command

```
f<-function(q) (1+4*q+10*q^2)*(1-q)^4
```

To find the value of f at $q = 0.73$ use the command

```
f(0.73)
```

To display the values of f at all the points

0, 0.01, 0.02, 0.03, ..., 0.15, 0.16

use the command

```
x<-seq(from=0, to=0.16, by=0.01)
f(x)
```

To plot the values of f over 100 equidistant points in the interval $[2, 7]$ use the command

```
x<-seq(from=2, to=7, length.out=100)
y<-f(x)
plot(x,y, type="l")
```

Equivalently, there is the simple command `curve(-)` that allows drawing multiple graphs in the same coordinate system. For example, the code

```
curve(x^2, from = -5, to = 5, ylim=c(-10,10))
curve(x^3, from = -5, to = 5, ylim=c(-10,10), col = "red", add = TRUE)
grid(nx = 10, ny = 10, col = "blue", lty = "dotted", lwd = 1)
```

produces the graph in Figure A.1. Above `col` stands for “color”. When this option is used different graphs are depicted in different colors. The optional `grid` command produces a grid of desired density and color.

Here is how we define in R the indicator function of the unit disc in the plane

$$I_D(x, y) = \begin{cases} 1, & x^2 + y^2 \leq 1, \\ 0, & x^2 + y^2 > 1. \end{cases}$$

```
indicator<-function(x,y) if(1 >= x^2+y^2) 1 else 0
```

Another possible code that generates this indicator function is

```
indicator<-function(x,y) as.integer(x^2+y^2<= 1)
```

Above, the command `as.integer` converts TRUE/FALSE to the integer 1/0. □

Example A.3.4 (Samples with replacement). For example, to sample *with replacement* 7 balls from a bin containing balls *labeled* 1 through 23 use the R command

```
sample(1:23, 7, replace=TRUE)
```

The result is a 7-dimensional vector whose entries consists of 7 numbers sampled with replacement from the set $\{1, \dots, 23\}$. Similarly, to simulate rolling a fair die 137 times use the command

```
sample(1:6, 137, replace=TRUE)
```




Figure A.1. The graphs of x^2 and x^3 in the same Cartesian plane.

Example A.3.5 (Rolling a die). Let us show how to simulate rolling a die a number n of times and then count how many times we get 6. Suppose $n = 20$. We indicate this using the command

```
n<-20
```

We now roll the die n times and store the results in a vector x

```
x<-sample(1:6, n, replace=TRUE)
```

Next we test which of the entries of x are equal to 6 and store the results of these 20 tests in a vector y

```
y<-x==6
```

The entries of y are *True* or *False*, depending on whether the corresponding entry of x was equal to 6 or not. To find how many entries of y are *T* use the command

```
sum(y)
```

The result is equal to the number of 6s we got during the string of 20 rolls of a fair die.

We can visualize data. Suppose we roll a die a large number $N = 1200$ of times. For each $1 \leq k \leq N$ we denote by $z(k)$ the fraction of the first k rolls when we rolled a 6. For $k \rightarrow \infty$ the Law of Large Numbers states that this frequency should approach $\frac{1}{6}$. The vector z can be generated in R using the commands

```
N<-12000
```

```
x<-sample(1:6, N, replace=TRUE)
```

```
z<-cumsum(x==6)/(1:N)
```

Above, `cumsum` stands for “cumulative sum”. The input of this operator is a numerical vector $x = (x_1, \dots, x_n)$. The output is a numerical vector s of the same dimension, with $s_k = s_1 = \dots + s_k$. We can visualize the fluctuations of $z(k)$ around the expected value $\frac{1}{6}$ using the R code

```
plot(1:N, z, type="l", xlab="# of rolls",
     ylab="average number 6-s")
abline(h=1/6,col="red")
```

Figure A.2 depicts the output.




Figure A.2. Rolling a die

□

Example A.3.6 (Samples without replacement). To sample without replacement 7 balls from an urn containing balls labeled 1 through 23 use the R command

```
sample(1:23, 7)
```

The number of possible samples above is $(27)_7$ and to compute it use the R command

```
prod(21:27)
```

□

Example A.3.7 (Permutations). To sample a random permutation of 7 objects use the R command

```
sample(1:7, 7)
```

To sample 10 random permutations of 7 objects use the R command

```
for (i in 1:10 ) print(sample(1:7, 7))
```

To compute $7!$ in R use the command

```
factorial(7)
```

□

Example A.3.8 (Combinations). Sampling random m -element subsets out of an n -element set possible is possible in R. For example, to sample 4 random subsets with 2 elements out of a 7-element set possible the following command

```
replicate(4, sort( sample(1:7, 2) ))
```

The sampled sets will appear as columns. To compute $\binom{52}{5}$ in R use the command

```
choose(52, 5)
```

□

Example A.3.9 (Custom discrete distribution). We can produce custom discrete random variables in R.

Suppose that we want to simulate a discrete random variable X whose values, sorted in increasing order, are

$$x_1 = 0.1, \quad x_2 = 0.2, \quad x_3 = 0.3, \quad x_4 = 0.7.$$

The corresponding probabilities are

$$p_1 = 1/3, \quad p_2 = 1/6, \quad p_3 = 1/4, \quad p_4 = 1/4.$$

The R-commands below describe how to compute the mean and the variance of X and how to sample X . In R, the symbol `#` indicates a comment. It is only for the programmer/user benefit. Anything following a `#` is not treated by R as a command.

```
X<-c(0.1,0.2,0.3,0.7)
# stores the values of X in
increasing order.

prob<-c(1/3,1/6,1/4,1/4) # stores the probabilities.
sum(prob)
# This is a test. If the result is 1, t
# then prob is a pmf.

m<-sum(X*prob)
# computes the mean of X and stores in m.
v<-sum((X^2)*prob) -m^2
# computes the variance of X and stores it in v.

sample(X,15, replace=TRUE, prob)
# produces 15 random samples of X.

cumsum(prob)

# computes the values of the cdf of X at x_1,x_2,...
```

□

Example A.3.10 (Useful discrete distributions). The standard discrete distributions are implemented in R.

The distribution	The R command
The binomial distribution $\text{Bin}(n, p)$	<code>binom(n,p)</code>
The geometric distribution $\text{Geom}(p)$	<code>geom(p)</code>
The negative binomial distribution $\text{NegBin}(k, p)$	<code>nbinom(k,p)</code>
The Poisson distribution $\text{Poi}(\lambda)$	<code>pois(lambda)</code>

The R library however uses rather different conventions

- (i) The geometric distribution in R is slightly different from the one described in this book. In R, the range of $\text{Geom}(p)$ variable T is $\{0, 1, \dots\}$ and its pmf is $\mathbb{P}[T = n] = p(1-p)^n$. In this book, a geometric random variable has range $\{1, 2, \dots\}$ and its pmf is $\mathbb{P}[T = n] = p(1-p)^{n-1}$; see Example A.3.12.
- (ii) In R the equality `nbinom(k, p) = n` represents the number of *failures* until we register the k -th success; see Example A.3.13.

The above commands by themselves mean nothing if they are not accompanied by one of the prefixes

- d produces the density or `pmf`.
- p produces the `cdf`.
- r produces random `samples`.
- q produces `quantiles`.

□

You can learn more details using R's help function. The examples below describe some concrete situations.

Example A.3.11 (Binomial). For example, suppose that $X \sim \text{Bin}(10, 0.2)$, i.e., X is the number of successes in a sequence of 10 independent Bernoulli trials with success probability 0.2.

To find the probability $\mathbb{P}(X = 3)$ use the R command

```
dbinom(3, 10, 0.2)
```

If $F_X(x) = \mathbb{P}(X \leq x)$ is the cdf of X , then you can compute $F_X(4)$ using the R command

```
pbinom(4, 10, 0.2)
```

To generate 253 random samples of X use the command

```
rbinom(253, 10, 0.2)
```

To find the 0.8-quantile of X use the R command

```
qbinom(0.8, 10, 0.2)
```

□

Example A.3.12 (Geometric). Suppose now that $T \sim \text{Geom}(0.2)$ is the waiting time until the first success in a sequence of independent Bernoulli trials with success probability $p = 0.2$.

To find the probability $\mathbb{P}(T = 3)$ use the command

```
dgeom(3-1, 0.2)
```

To find the probability $\mathbb{P}(T \leq 4)$ use the command

```
pgeom(4-1, 0.2)
```

To generate 253 random samples of T use the command

```
1+rgeom(253, 0.2)
```

To find the 0.8-quantile of T use the R command

```
qgeom(0.8, 0.2)+1
```

Example A.3.13 (Negative Binomial). Suppose that $T \sim \text{NegBin}(8, 0.2)$ is the waiting time for the first 8 successes in a string of Bernoulli trials with success probability.

To find the probability $\mathbb{P}(T = 12)$ use the R command

```
dnbinom(12-8, 8, 0.2)
```

You can compute $\mathbb{P}(T \leq 14)$ using the R command

```
pnbinom(14-8, 8, 0.2)
```

To generate 253 random samples of T use the command

```
8+rnbinom(253, 8, 0.2)
```

To find the 0.8-quantile of T use the R command

```
8+qnbinom(0.8, 8, 0.2)
```

□

Example A.3.14 (Poisson). Suppose that $X \sim \text{Poi}(0.2)$ is a Poisson random variable with parameter $\lambda = 0.2$.

To find the probability $\mathbb{P}(X = 3)$ use the command

```
dpois(3, 0.2)
```

To find the probability $\mathbb{P}(X \leq 4)$ use the command

```
ppois(4, 0.2)
```

To generate 253 random samples of X use the command

```
rpois(253, 0.2)
```

To find the 0.8-quantile of X use the R command

```
qpois(0.8, 0.2)
```

□

Example A.3.15 (Continuous distributions in R). The continuous distributions $\text{Unif}(a, b)$, $\exp_\lambda$ and $N(\mu, \sigma^2)$ can be simulated in R by invoking

```
unif(min=a, max=b)
```

```
exp(rate=lambda)
```

```
norm(mean=mu, sd=sigma)
```

where `sd`:=standard deviation.

To invoke the standard normal random variable you could use the shorter command

```
norm
```

□

As in the case of discrete distributions, we utilize these commands with the prefixes *d*–, *p*–, *q*– and *r*– that have the same meaning as in R-Session A.3.10. Thus *d*– will generate the pdf, *p*– the cdf, *r*– generates a random sample, and *q*– produces quantiles.

Example A.3.16. Here are some concrete examples. To find the probability density of $\exp_3$ at $x = 1.7$ use the command

```
dexp(1.7, 3)
```

To find the probability density of $N(\mu = 5, \sigma^2 = 7)$ at $x = 2.6$ use the command

```
dnorm(2.6, 5, sqrt(7))
```

To produce 1000 samples from $\text{Unif}(3, 13)$ use the command

```
runif(1000, 3, 13)
```

□

Example A.3.17 (Gambler's ruin). Consider two players the first with fortune $\$a$, and the second with fortune $\$b$. Set $N := a + b$. They flip a fair coin. Heads, player 1 gets a dollar from player 2, Tails, player 1 gives a dollar to player 2. The game ends when one of them is ruined. One can simulate this in R using the code

```
r<-function(a,N){
  t<-0
  x<-a
  v<-c(0,N)
  while(all(v!=x)){
    f<-sample(0:1,1, replace=TRUE)
    x<-x+(2*f-1)
    t<-t+1
  }
  y<-c(x,t)
  y
}
```

The output is a two-dimensional vector. Its first entry is the fortune of the first player at the end of the game, while the second entry is duration of the game, i.e., the number of coin flips until one of them is ruined.

To compute the winning probability of the first player and the expected duration of a game we can use the Law of Large Numbers and run a large number G of games

```
empiric_r<-function(G,a,N){
  P<-c()
  T<-c()
```



```

for(i in 1:G){
  P<-c(P,r(a,N)[1])
  T<-c(T,r(a,N)[2])
}
c(sum(P==N)/G,sum(T)/G)
}

```

For example, if we want to run a number $G = 1200$ of games with the first player's initial fortune $a = 8$ and the combined fortune of the two players is $N = 15$ use the command

```
empiric_r(1200,8,15)
```

The output is a two-dimensional vector. Its first entry describes the fraction of the G games won by the first player, and the second entry is the average duration of these G games.




Figure A.3. A zig-zag illustrating the ruin problem. In this picture the first player wins and the second player is ruined.

One can also visualize a game. The code below produces a vector whose entries describe the evolution of the fortune of the first player.

```

rgr<-function(a,N){
  x<-a
  z<-c(a)
  v<-c(0,N)
  while(all(v!=x)){
    f<-sample(0:1,1,replace=TRUE)
    x<-x+(2*f-1)
    z<-c(z,x)
  }
  z
}

```

For given values of N and a say, $N = 25$, $a = 12$, one can visualize the evolution of the fortune of the first player using the code below. Its output is a graph similar to the one in Figure A.3.

```

N<-25
a<-12
u<-rgr(a,N)
l<-length(u)-1

```

```
plot(0:1, u, type="l", xlab="# of flips",
     ylab="the fortune of the first player", ylim=c(0,N))
abline(h=c(0,N), col=c("red", "red"))
```

□

Example A.3.18 (Buffon's needle problem). The R program below uses the Buffon needle problem (see Exercise 1.30) to find an approximation of π .

```
L<-0.7 # L is the length of the needle. It is <1.
N<-1000000 # N is the number of times we throw the needle.
f<-0
#the next loop simulates the tossing of
#N random needles and computes
# the number f of times they intersect a line

for (i in 1:N){
  y<-runif(1, min=-1/2,max=1/2) #this locates
  # the center of the needle
  t<-runif(1, min=-pi/2,max=pi/2)#this determines
  #the inclination of the needle
  if ( abs(y)< 0.5*L*cos(t) ) f<-f+1 }
#f/N is the empirical frequency
"the aproximate value of pi is"; (N/f)*2*L
```

□

Example A.3.19 (Monte Carlo). The R-command lines below implement the Monte Carlo strategy for computing a double integral over the unit square

```
# Monte Carlo integration of the function f(x,y)
#over the rectangle [a,b] x[c,d]
# First we describe the function
f<- function(x,y) sin(x*y)
# Next, we describe the region of integration [a,b]x[c,d]
a=0
b=1
c=0
d=1
# Finally, we decide the number N of sample points in
# the region of integration
N=100000
#S will store the integral
S=0
for (i in 1:N){
  x<- runif(1,a,b) #we sample a point uniformly in [a,b]
  y<- runif(1,c,d) #we sample a point uniformly in [c,d]
  S<-S+f(x[1],y[1])
}
```

```
'the integral is'; (b-a)*(d-c)*S/N
```

The next code describes a Monte-Carlo computation of the area of the unit circle.

```
nsim<-1000000#nsim is the number of simulations
x<-runif(nsim,-1,1)#we choose nsim uniform samples
#in the interval (-1,1) on the x axis
y<-runif(nsim,-1,1)#we choose nsim uniform samples
#in the interval (-1,1) on the y axis
area<-4*sum(x^2+y^2<1)/nsim
"the area of the unit circle is very likely"; area
```

□

Example A.3.20. Suppose that we have a probability distribution `prob` on the alphabet $\{1, 2, \dots, L\}$. One experiment consists of sampling the alphabet according to the distribution `prob` until we first observe the given word (or pattern) `patt`. The following R-routine performs m such experiments and returns an m -dimensional vector `f` whose components are the cumulative means of the waiting times

$$f_k = \frac{1}{k} \sum_{j=1}^k T_j, \quad k = 1, \dots, m,$$

where T_j is the time to observe the pattern in the j -th experiment.

```
Tpattern<-function(patt, prob, m, L){
  k<-length(patt)
  T<-c()
  for (i in 1:m){
    x<-sample(1:L,k,replace=TRUE, prob)
    n<-k
    while ( all(x[(n-k+1):n]==patt)==0 ){
      x<-c(x, sample(1:L,1,replace=TRUE, prob) )
      n<-n+1
    }
    T<-c(T,n)
  }
  f<-cumsum(T)/(1:m)
  f
}
```

If `prob` is the uniform distribution use the faster routine

```
Tpatt_unif<-function(patt, m, L){
  k<-length(patt)
  T<-c()
  for (i in 1:m){
    x<-sample(1:L,k,replace=TRUE)
    n<-k
```

```

while ( all(x[(n-k+1):n]==patt)==0 ){
  x<-c(x, sample(1:L,1,replace=TRUE) )
  n<-n+1
}
T<-c(T,n)
}
f<-cumsum(T) / (1:m)
f
}

```

In the uniform case, the expected waiting time to observe the pattern `patt` can be determined using routine below that relies on the identity (3.1.12) in Example 3.1.33.

```

tau<-function(patt,L){
  n<-length(patt)
  m<-n-1
  t<-2^n
  for (i in 1:m){
    j<-n-i
    k<-i+1
    t<-t+ all(patt[1:j]==patt[k:n])*L^(n-i)
  }
  t
}

```

Example A.3.21 (Monty Hall Problem). This is a classical and publicly debated problem about a gameshow.

A contestant is told that behind one of the three closed doors there is a car he can win if he opens the correct door.

The contestant picks one door, does not open it. The gameshow host then opens another door he knows does to hide the award car. At this moment the contestant has two choices: either open the door he has already picked, or switch, and open the last remaining closed door. The Monty Hall problem asks which is the strategy that has a higher probability of guessing the winning door.

It turns out that the second option, switch and open the last remaining closed door has a winning probability $\frac{2}{3}$ while the first option has a winning probability $\frac{1}{3}$. The R-script below simulates a sequence of N independent games in which the players use the same strategy, stick with the initial choice, or switch. The comments in the script ought to make this more precise.

```

# We simulate the Monty Hall problem,
# Car is behind door 3.
# The contestant chooses randomly door x in 1,2,3
# MH(x) is the door indicated by the gameshow host.

MH<-function(x){
  if(x==3) sample(1:2,1) else setdiff(c(1:2), c(x))[1]
}

# N is the number of trials.

```

```

#MHP0(N)  simulates N Monty Hall  experiments
# in which the contestant does not change his initial choice
# The output MHP0(N)  is the number of winning guesses

MHP0<-function(N) {
  k<-0
  for (i in 1:N){
    doors<-c(1:3)
    x<-sample(1:3,1)
    k<- k+as.integer(x==3)
  }
  k
}

# N is the number of trials.
#MHP(N)  simulates N Monty Hall  experiments
# in which the contestant changes his opinion
#after choosing to a door.
# The output  MHP(N) is the number of winning guesses

MHP<-function(N) {
  k<-0
  for (i in 1:N){
    doors<-c(1:3)
    x<-sample(1:3,1)
    r<- c(x, MH(x))
    doors<-setdiff(doors,r)
    k<-k+as.integer(doors[1]==3) }
  k
}

```

□

Example A.3.22 (Approximating the Brownian motion). To generate the graph of a sample of the random zig-zag $W^{100}(t)$ defined by (2.5.2) use the command

```

# set.seed(1213)

#uncomment the above command  for reproducibility

x<-10*cumsum(2*rbinom(100,1, 0.5)-1)
plot(x, type='l')

```

Here is the R code that produced Figure 3.5.

```

#set.seed(1213)

#uncomment the above command  for reproducibility

```

```
#generate 100 independent samples
# of the Rademacher variable

x=2*rbinom(100,1, 0.5)-1

#prepare for reflection at t=60

r=c(rep(1,60), rep(-1,40))

#y will describe the reflected walk

y=x*r

#x approximates the Brownian motion
x=10*cumsum(x)

# y approximates the reflected Brownian motion

y=10*cumsum(y)

x=c(0,x)
y=c(0,y)
t=c(0:100)

plot(t,x, type='l', col="black", ylim=c(-70,120))
lines(c(61:101),c(y[61:101]),col="red", lty="dashed", ylim=c(-70,120))
points(c(61),c(y[61]), col="red" )
```

□

Bibliography

- [1] D. Aldous: *Exchangeability and related topics*, École d'été de Probabilités des Saint Fleur XIII-1983, p. 2-199, Lect. Notes in Math vol. 1117, Springer Verlag, 1985.
- [2] D. Aldous: *Probability Theory*, Course notes, Spring 2017.
https://www.stat.berkeley.edu/~aldous/205B/chewi_notes.pdf
- [3] J. Aldrich: *But you have to remember P. J. Daniell of Sheffield*, Electronic Journal for History of Probability and Statistics, Dec. 2007.
<https://www.emis.de/journals/JEHPS/Decembre2007/Aldrich.pdf>
- [4] V. I. Arnold, A. Avez: *Ergodic Problems of Classical Mechanics*, Addison Wesley, 1968.
- [5] V. I. Arnold: *Stochastic and deterministic characteristics of orbits in chaotically looking dynamical systems*, Trans. Moscow Math. Soc., **70**(2009), 31-69.
- [6] R.B. Ash: *Probability and Measure Theory*, (with contributions from C. Doléans-Dade), 2nd Edition, Academic Press, 2000.
- [7] S. Asmussen: *Applied Probability and Queues*, 2nd Edition, Stoch. Modelling and Appl. Probab., vol. 51, Springer Verlag, 2003.
- [8] K.B. Athreya, P. E. Ney: *Branching Processes*, Springer Verlag, 1972.
- [9] P. Baldi: *Stochastic Calculus. An Introduction Through Theory and Exercises*, Universitext, Springer Verlag, 2017.
- [10] S. Banach: *Über die Baire'sche Kategorie gewisser Functionenmengen*, Studia Mathematica, 3(1931), 174-179.
- [11] P. Bamberg, S. Sternberg: *A Course in Mathematics for Students of Physics*, vol. 2, Cambridge University Press, 1990.
- [12] V. Barbu: *Differential Equations*, Springer Undergraduate Mathematics Series, Springer Verlag, 2016.
- [13] R.F. Bass: *Real Analysis for Graduate Students*, Version 5, 2024.
- [14] M. Benaïm, T. Hurth: *Markov Chains on Metric Spaces. A Short Course*, Springer Verlag, 2022.
- [15] R. N. Bhattacharya, E. C. Waymire: *Stochastic Processes with Applications*, SIAM, 2009.
- [16] R. N. Bhattacharya, E. C. Waymire: *A Basic Course in Probability Theory*, 2nd Edition, Springer Verlag, 2016.
- [17] P. Billingsley: *Ergodic Theory and Information*, John Wiley & Sons, 1965.
- [18] P. Billingsley: *Convergence of Probability Measures*, 2nd Edition, John Wiley & Sons, 1999.
- [19] N.H. Bingham: *Fluctuation theory for the Ehrenfest Urn*, Adv. Appl. Prob. **23**(1991), 598-611.
- [20] D. Blackwell, D. Freedman: *The tail σ -field of a Markov chain and a theorem of Orey*, Ann. Math. Statist., **35**(1964), 1291-1295.
- [21] V.I. Bogachev: *Measure Theory. Vol. 1*, Springer Verlag, 2007.
- [22] A. N. Borodin: *Stochastic Processes*, Birkhäuser, 2017.

- [23] R. Bott: *On induced representations*, in volume *The Mathematical Heritage of Hermann Weyl*, Proc. Symp. Pure Math., vol. 48, Amer. Math. Soc., 1988
- [24] S. Boucheron, G. Lugosi, P. Massart: *Concentration Inequalities. A Nonasymptotic Theory of Independence*, Oxford University Press, 2013.
- [25] N. Bourbaki: *General Topology*, Part 2, Hermann, 1966
- [26] L. Breiman: *Probability*, SIAM, 1992.
- [27] P. Brémaud: *Markov Chains, Gibbs Fields, Monte Carlo Simulations and Queues*, Springer Verlag, 2nd Edition, 2020.
- [28] P. Brémaud: *Probability Theory and Stochastic Processes*, Springer Verlag, 2020.
- [29] H. Brezis: *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer Verlag, 2011.
- [30] J. Bricmont: *Making Sense of Statistical Mechanics*, Springer Verlag, 2022.
- [31] S. A. Broughton, K. Bryan: *Discrete Fourier Analysis and Wavelets. Applications to Signal and Image Processing*, Second Edition, John Wiley & Sons, 2018.
- [32] H. E. Buchanan, T. H. Hildebrandt: *Note on the convergence of sequences of functions of a certain type*, Ann. of Math., **9**(1908), 123-126.
- [33] J. T. Chang, D. Pollard: *Conditioning as Disintegration*, Statistica Neerlandica, **51**(1997), 287-317.
- [34] I. Chavel: *Eigenvalues in Riemann Geometry*, Academic Press, 1984.
- [35] J. Cheeger: *A lower bound for the smallest eigenvalue of the Laplacian*, in Gunning (ed.) *Problems in Analysis*, p. 199-205, Princeton University Press, 1970.
- [36] Y. S. Chow, H. Robbins, D. Siegel: *Great Expectations: The Theory of Optimal Stopping*, Houghton Mifflin Co., 1971.
- [37] Y. S. Chow, H. Teicher: *Probability Theory. Independence, Interchangeability, Martingales*, 3rd Edition, Springer Verlag, 1997.
- [38] K.L. Chung, J. L. Doob: *Fields, Optionality and Measurability*, Amer. J. Math., **87**(1965), 397-424.
- [39] K. L. Chung, F. AitSahlia: *Elementary probability theory: with stochastic processes and an introduction to mathematical finance*, Springer Verlag, 2003.
- [40] K.C. Chung, R.J. Williams: *Introduction to Stochastic Integration*, 2nd Edition, Birkhäuser, 1990.
- [41] V. Chvátal, D. Sankoff: *Longest common subsequences of two random sequences*, J. Appl. Prob. **12**(1975), 306-315.
- [42] E. Çinlar: *Introduction to Stochastic Processes*, Dover, 2014.
- [43] E. Çinlar: *Probability and Stochastics*, Graduate Texts in Math., vol. 261 Springer Verlag, 2011.
- [44] E.G. Coffman Jr., G. S. Lueker: *Probabilistic Analysis of Packing and Partitioning Algorithms*, John Wiley & Sons, 1991.
- [45] D. L. Cohn: *Measure Theory*, 2nd Edition, Birkhäuser, 2013.
- [46] I. P. Cornfeld, S. V. Fomin, Ya. G. Sinai: *Ergodic Theory*, Springer Verlag, 1982.
- [47] T.M. Cover, M. Thomas, J. A. Thomas: *Elements of Information Theory*, Wiley-Interscience, 2006.
- [48] P. J. Daniell: *Integrals in an infinite number of dimensions*, Ann. of Math. **20**(1919), 281-288.
- [49] T.M. Davies: *The Boof of R*, No Starch Press, 2015,
<https://nostarch.com/bookofr>
- [50] C. Dellacherie, P.-A. Meyer: *Probabilities and Potential*, vol. A, North Holland Mathematical Studies, vol. 29, Hermann Paris, 1978.
- [51] C. Dellacherie, P.-A. Meyer: *Probabilities and Potential*, vol. C, North Holland, 1988.
- [52] P. Diaconis: *Group Representations in Probability and Statistics*, Institute of Mathematical Statistics, 1988.
- [53] P. Diaconis: *The Markov chain Monte Carlo revolution*, Bull. Amer. Math. Soc., **46**(2009), 179-205.
- [54] P. Diaconis, R. Griffiths: *Exchangeable pairs of Bernoulli random variables, Krawtchouk polynomials, and Ehrenfest urns*, Aust. N. Z. J. Stat. **1**(2012), 81-101.
- [55] P. Diaconis, R. Griffiths: *An Introduction to multivariate Krawtchouk polynomials and their polynomials*, arXiv: 1309.0112, J. Stat. Plann. Inference, **154**(2014), 39-53.

- [56] P. Diaconis, B. Skyrms: *Ten Great Ideas About Chance*, Princeton University Press, 2018.
- [57] P. Diaconis, D. Stroock: *Geometric bounds for eigenvalues of Markov chains*, Ann. Appl. Prob., **1**(1991), 31-61.
- [58] G. Di Nunno, B. Øksendal, F. Proske: *Malliavin Calculus for Lévy Processes with Applications to Finance*, Springer Verlag, 2009.
- [59] W. Doeblin: *Exposé de la Théorie des Chaînes simples constantes de Markoff à un nombre fini d'États*, Rev. Math. de l'Union Interbalkanique, **2**(1938), 77-105. o
- [60] J. L. Doob: *The Brownian motion and stochastic equations*, Ann. of Math., **43**(1942), 351-369.
- [61] J. L. Doob: *Stochastic Processes*, John Wiley & Sons, 1953.
- [62] P. G. Doyle, J. L. Snell: *Random Walks and Electrical Networks*, MAA, 1984.
- [63] L. E. Dubins, L. J. Savage: *How to Gamble if you Must. Inequalities for Stochastic Processes*, Dover, 2014.
- [64] R. M. Dudley: *Real Analysis and Probability*, Cambridge University Press, 2004.
- [65] R. M. Dudley: *Uniform Central Limit theorems*, Cambridge University Press, 2014.
- [66] N. Dunford, J.T. Schwartz: *Linear Operators. Part I: General Theory*, John Wiley & Sons, 1957.
- [67] R. Durrett: *Stochastic Calculus: A Practical Guide*, CRC Press, 1996.
- [68] R. Durrett: *Probability. Theory and Examples*, 5th Edition, Cambridge University Press, 2019.
- [69] A. Dvoretzky, P. Erdős, S. Kakutani: *Nonincrease everywhere of the Brownian motion process*, 1961 Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. II pp. 103-116 Univ. California Press, Berkeley, Calif.
- [70] E. B. Dynkin, A. A. Yushkevich: *Markov Processes. Theorems and Problems*, Plenum Press, 1969.
- [71] Y. Eisner, B. Farkas, M. Haase, R. Nagel: *Operator Theoretic Aspects of Ergodic Theory*, Springer Verlag, 2015.
- [72] P. Erdős, A. Rényi: *On a classical problem of probability theory*, Magyar Tudományos Akadémia Matematikai Kutató Intézetének Közleményei, **6**(1961), 215-220,
- [73] G. Fayolle, V. A. Malyshev, M. V. Menshikov: *Topics in the Constructive Theory of Countable Markov Chains*, Cambridge University Press, 1995.
- [74] W. Feller: *The Kolmogorov-Smirnov theorems for empirical distributions*, Ann. Math. Statistics, **19**(1948), 177-189.
- [75] W. Feller: *An Introduction to Probability Theory and its Applications*, Volume 1, 3rd Edition, John Wiley & Sons, 1970.
- [76] W. Feller: *An Introduction to Probability Theory and its Applications*, Volume 2, 2nd Edition, John Wiley & Sons, 1970.
- [77] L. Floridi: *Information Theory. A Very Short Introduction*, Oxford University Press, 2010.
- [78] D. Foata, A. Fuchs: *Processus Stochastiques. Processus de Poisson, Chaînes de Markov et Martingales*, 2nd edition, Dunod, 1998.
- [79] T. Frankel: *The Geometry of Physics*, 3rd Edition, Cambridge University Press, 2011.
- [80] D. Freedman: *Brownian Motion and Diffusion*, Springer Verlag, 1983.
- [81] B. Fristedt, L. Gray: *A Modern Approach to Probability Theory*, Birkhäuser, 1997.
- [82] F. R. Gantmacher: *Theory of Matrices*, vol.2, AMS, Chelsea Publishing, 2000.
- [83] M. Gardner: *Time Travel and Other Mathematical Bewilderments*, W. H. Freeman & Co., 1988.
- [84] A. Garsia: *A simple proof of E. Hopf's maximal ergodic theorem*, J. Math. Mech., **14**(1965), 381-382.
- [85] A. Garsia: *Topics in Almost Everywhere Convergence*, Markham Publishing Co., 1970
- [86] I. M. Gelfand, N. Ya. Vilenkin: *Generalized Functions. Volume 4. Applications of Harmonic Analysis*, Academic Press, 1964.
- [87] D. Gilbarg, N. S. Trudinger: *Elliptic Partial Differential Equations of Second Order*, 2nd Edition, Springer Verlag, 2001.
- [88] J. P. Gilbert, F. Mosteller: *Recognizing the maximum of a sequence*, J. Amer. Stat. Assoc., **61**(1966), 35-73.
- [89] E. Giné, R. Nickl: *Mathematical Foundations of Infinite Dimensional Statistical Models*, Cambridge University Press, 2016.

- [90] J. Gleick: *The Information. A History. A Theory. A Flood*, Pantheon Books, 2011.
- [91] B. V. Gnedenko, A. N. Kolmogorov: *Limit Distributions for Sums of Independent Random Variables*, Addison Wesley, 1968.
- [92] A. Grigoryan: *Introduction to Analysis on Graphs*, University Lect. Series, Amer. Math. Soc., 2018.
- [93] G. R. Grimmett: *Probability on Graphs. Random Processes on Graphs and Lattices*, Cambridge University Press, 2011.
- [94] G. R. Grimmett, D. R. Stirzaker: *Probability and Stochastic Processes*, 4th Edition, Oxford University Press, 2020.
- [95] L. J. Guibas, A. M. Odlyzko: *String overlaps, pattern matching and nontransitive games*, J. Copmb. Th. Series A, **30**(1981), 183-208.
- [96] O. Häggström: *Finite Markov Chains and Algorithmic Applications*, Cambridge University Press, 2002.
- [97] M. Hairer: *Convergence of Markov Proceses*, Minicourse Univ. of Warwick 2010, <https://www.hairer.org/notes/Convergence.pdf>
- [98] P. Hall, C. C. Heyde: *Martingale Limit Theory and Its Applications*, Academic Press, 1980.
- [99] P. R. Halmos: *Lectures on Ergodic Theory*, Dover, 2017.
- [100] G.H. Hardy: *Divergent Series*, Oxford University Press, 1949.
- [101] T. E. Harris: *The Theory of Branching Processes*, Springer Verlag, 1963.
- [102] J. Hawkins: *Ergodic Dynamics. From Basic Theory to Applications*, Springer Verlag, 2021.
- [103] B. Hayes: *The first links in a Markov chain*, American Scientist, **101**(2013), No. 2, p. 92.
- [104] K. Itô: *Multiple Wiener integral*, J. Math. Soc. Japan, **3**(1951), 157-169.
<https://www.americanscientist.org/article/first-links-in-the-markov-chain>
- [105] M. Jacobsen: *Laplace and the origin of the Ornstein-Uhlenbeck process*, Bernoulli, **2**(1996), 271-286,
<https://www.jstor.org/stable/3318524>
- [106] R. Jarrow, P. Protter: *A short history of stochastic integration and mathematical finance: the early years, 1880-1970*, A Festschrift for Herman Rubin, Institute of Mathematical Statistics Lecture Notes – Monograph Series, **45**(2004), 75–91.
- [107] M. Jerrum, M. Sinclair: *Approximate Counting, Uniform Generation and Rapidly Mixing of Markov Chains*, Information and Computation, **82**(1989), 93-133.
- [108] M. Kac: *Random walk and the theory of Brownian motion*, Amer. Math. Monthly, **54**(1947), 369-391.
- [109] O. Kallenberg: *Foundations of Modern Probability*, 3rd Edition, Springer Verlag, 2021.
- [110] I. Karatzas, S.E. Shreve: *Brownian Motion and Stochastic Calculus*, 2nd Edition, Grad. Texts in Math., Springer Verlag, 1998.
- [111] S. Karlin, H. M. Taylor: *A First Course in Stochastic Processes*, 2nd Edition, Academic Press, 1975.
- [112] Y. Katznelson, B. Weiss: *A simple proof of some ergodic theorems*, Israel J. of Math, **4**(1982), 291-296.
- [113] J. G. Kemeny, J. L. Snell: *Finite Markov Chains*, with a new appendix “Generalization of a fundamental matrix”, Springer Verlag, 1983.
- [114] J. H. B. Kemperman: *The Passage Problem for a Stationary Markov Chain*, The University of Chicago Press, 1961.
- [115] H. Kesten, P. Ney, F. Spitzer: *The Galton-Watson process with mean one and finite variance*, Th. Prob. Appl., **11**(1966), 513-540.
- [116] J. M. Keynes: *A Treatise on Probability*, MacMillan and Co. London, 1921. Available at Project Guttenberg .
<http://www.gutenberg.org/ebooks/32625>
- [117] J.F.C. Kingman: *Uses of exchangeability*, Ann. Prob., **6**(1978), 183-197.
- [118] A. Klenke: *Probability Theory. A Comprehensive Course*, Universitext, Springer Verlag, 2008.
- [119] A. N. Kolmogorov: *Foundations of the Theory of Probability*, Chelsea 1950.
- [120] A. N. Kolmogorov: *The theory of transmission of information*, the volume *Selected Works of A. N. Kolmogorov. Volume III. Information Theory and the Theory of Algorithms*, p. 6-33, Kluwer Academic Publishers, 1993

- [121] E. Kowalski: *An Introduction to Expander Graphs*, Société Mathématiques de France, 2019.
- [122] U. Krengel: *Ergodic Theorems. With a supplement by Antoine Brunel*, Walter de Gruyter, 1985.
- [123] N. V. Krylov: *A few comments on a result of A. Novikov and Girsanov's theorem*, Stochastics, **91**(2019), 1186-1189.
- [124] L. Kuipers, H. Niederreiter: *Uniform Distribution of Sequences*, John Wiley & Sons, 1974. Dover reprint 2006.
- [125] H. Kunita, S. Watanabe: *On square integrable martingales*, Nagoya Math. J., **30**(1967), 209-245.
- [126] K. Kuratowski, A. Mostowski: *Set Theory. With an Introduction To Descriptive Set Theory*, North Hollan Publishing Co., 1976.
- [127] S. Lang: *Linear Algebra*, 3rd Edition, Springer Verlag, 1987.
- [128] G. F. Lawler: *Introduction to Stochastic Processes*, 2nd Edition, Chapman & Hall, 2006.
- [129] N. N. Lebedev: *Special Functions and Their Applications*, Dover, 1972.
- [130] M. Ledoux, M. Talagrand: *Probability in Banach Spaces*, Springer Verlag, 1991.
- [131] J.-F. Le Gall: *Measure Theory, Probability and Stochastic Processes*, Springer Verlag, 2022.
<https://www.math.u-psud.fr/~jfllegall/IPPA2.pdf>
- [132] J.-F. Le Gall: *Brownian Motion, Martingales, and Stochastic Calculus*, Graduate Texts in Math., vol. 274, Springer Verlag 2016.
- [133] D. A. Levin, Y. Perez, E. L. Wilmer: *Markov Chains and Mixing Times*, Amer. Math. Soc., 2009.
- [134] P. Lévy: *Théorie de l'Addition des Variables Aléatoires*, Gauthier-Villars, 1937.
- [135] P. Lévy: *Processus Stochastiques et Mouvement Brownien*, Gauthier Villars, 1965.
- [136] S.-Y. R. Li: *A martingale approach to the study of occurrence of sequence patterns in repeated experiments*, Ann. Prob. **8**(1980), 1171-1176.
- [137] M. Loève: *Probability Theory*, vol I, 4th Edition, Graduate Texts in Math. no.45, Springer Verlag, 1977.
- [138] T. Lindvall: *Lectures on the Coupling Method*, John Wiley & Sons, 1992.
- [139] L. H. Loomis: *An Introduction to Abstract Harmonic Analysis*, Dover, 2011.
- [140] A. Lubotzky: *Expander graphs in pure and applied mathematics*, Bull. A.M.S., **49**(2012), p. 113-162.
- [141] T. Lyons: *A simple criterion of transience of a reversible Markov chain*, Ann. Prob., **11**(1983), 393-402.
- [142] T. Lyons: *Strong Laws of Large Numbers for weakly correlated random variables*, Mich. Math. J., **35**, 353-359.
- [143] T. Lyons, Y. Peres: *Probability on Trees and Networks*, Cambridge University Press, 2017.
- [144] D. J. MacKay: *Information Theory, Inference and Learning Algorithms*, Cambridge University Press, 18th printing, 2017.
- [145] P. Malliavin, A. Thalmaier: *Stochastic Calculus of Variations in Mathematical Finance*, Springer Finance, Springer Verlag, 2005.
- [146] R. Mansuy: *The origins of the word "martingale"* Electronic Journal for History of Probability and Statistics, vol. 5, Fasc. 1, (2009), 1-9.
<http://www.jehps.net/juin2009.html>
- [147] J. Matoušek: *Lectures on Discrete Geometry*, Graduate Texts in Math. no. 212, Springer Verlag, 2002.
- [148] S. Mazurkewicz: *Sur les fonctions non dérivables*, Studia Mathematica, **3**(1931), 92-94.
- [149] M. McCaffrey: *Markov Chains: A Random Walk Through Particles, Cryptography, Websites and Card Shuffling*, Senior Thesis, University of Notre Dame, 2017,
https://academicweb.nd.edu/~lnicolae/Thesis_v3.pdf.
- [150] H. P. McKean: *Stochastic Integrals*, Academic Press, 1969.
- [151] P. A. Meyer: *Probability and Potentials*, Blaisdell Publishing Vo., 1966
- [152] M. Mitzenmacher, E. Upfal: *Probability and Computing. Randomized Algorithms and Probability Analysis*, 7th printing, Cambridge University Press, 2013.
- [153] M. Mohri, A. Rostamizadeh, A. Talwalkar: *Foundations of Machine Learning*, The MIT Press, 2012.
- [154] P. Mörters, U. Peres: *Brownian Motion*, Cambridge University Press, 2010.
- [155] C. St. J. A. Nash-Williams: *Random walks and electric currents in networks*, Proc. Cambridge. Phil. Soc., **55**(1959), 181-194.

- [156] J. Neveu: *Mathematical Foundations of the Calculus of Probability*, Holden Day Inc., 1965.
- [157] D. J. Newmann, L. Shepp: *The double dixie-cup problem*, Amer. Math. Monthly, **67**(1960), 58-61.
- [158] L. I. Nicolaescu: *Introduction to Real Analysis*, World Scientific, 2020.
- [159] L. I. Nicolaescu: *Lectures on the Geometry of Manifolds*, 3rd Edition, World Scientific, 2021.
- [160] J. R. Norris: *Markov Chains*, Cambridge University Press, 1997.
- [161] B. Oksendal: *Stochastic Differential Equations*, 6th Edition, Springer Verlag, 2007.
- [162] S. Olekser-Taylor, D. Schmidt: *Limit profile for the Bernoulli-Laplace urn*,
<https://arxiv.org/pdf/2409.07900>
- [163] R. A. Olshen: *The coincidence of measure algebras under an exchangeable probability*, Wahrscheinlichkeitstheorie Verw. Geb., **18**(1971), 153-158
- [164] R. E. A. C. Paley, N. Wiener, A. Zygmund, *Note on random functions*, Math. Z., 1933.
- [165] J. L. Palacios: *Fluctuation theory for the Ehrenfest urn via electric networks*, Adv. Appl. Prob., **25**(1993), 472-476.
- [166] K. R. Parthasarathy: *Probability Measures on Metric Spaces*, Academic Press, 1967.
- [167] A. Pascucci: *Probability II. Stochastic Calculus*, Springer Verlag, 2024.
- [168] G. Peskir, A. Shiryaev: *Optimal Stopping and Free-Boundary Problems*, Birkhäuser, 2006.
- [169] V. V. Petrov: *Limit Theorems of Probability Theory. Sequences of Independent Random Variables*, Oxford University Press, 1995.
- [170] I. Pinelis: *Martingales converging in probability and not a.s.*, MathOverflow,
<https://mathoverflow.net/a/410350/20302>
- [171] D. Pollard: *Convergence of Stochastic Processes*, Springer Verlag, 1984.
- [172] D. Pollard: *Empirical Processes: Theory and Applications*, NSF-CBMS Regional Conference Series in Probability and Statistics, vol. 2, 1990
- [173] M. Pollicott, M. Yuri: *Dynamical Systems and Ergodic Theory*, Cambridge University Press, 1998.
- [174] P. E. Protter: *Stochastic Integration and Differential Equations*, 2nd Edition, Corrected 3rd Printing Springer Verlag, 2005.
- [175] S. I. Resnick: *Adventures in Stochastic processes*, Birkhäuser, 2002.
- [176] S. I. Resnick: *Extreme Values, Regular Variation, and Point Processes*, Springer Verlag, 2008.
- [177] D. Revuz, M. Yor: *Continuous Martingales and Brownian Motion*, 3rd Edition, Corrected 3rd printing, Springer Verlag, 2005.
- [178] F. Riesz: *Sur la théorie ergodique*, Comment. Math. Helv. **17**(1944), 221-239.
- [179] R. T. Rockefellar: *Convex Analysis*, Princeton University Press, 1997.
- [180] L. C. G. Rogers, D. Williams: *Diffusions, Markov Processes and Martingales. Volume 1. Foundations*, Cambridge University Press, 2000.
- [181] L.C.G. Rogers, D. Williams: *Diffusions, Markov Processes and Martingales. Volume 2. Itô Calculus*, Cambridge University Press, 2000.
- [182] G.-C. Rota: *Indiscrete Thoughts*, Birkhäuser, 2008.
- [183] W. Rudin: *Fourier Analysis on Groups*, John Wiley & Sons, 1962.
- [184] E. Schechter: *Handbook of Analysis and Its Foundations*, Academic Press, 1997.
- [185] R. L. Schilling, L. Partzch: *Brownian Motion. An Introduction to Stochastic Processes*, DeGruyter, 2012.
- [186] R. L. Schilling: *Measure, Integral, Probability & Processes*, Amazon, 2021.
- [187] K. Schmüdgen: *The Moment Problem*, Springer Verlag, 2017.
- [188] D. Serre: *Matrices. Theory and Applications*, 2nd Edition, Grad. Texts Math., vol.216, Springer Verlag, 2010.
- [189] S. Shalev-Shwartz, S. Ben-David: *Understanding Machine Learning. From Theory to Algorithms*, Cambridge University Press, 2014.
- [190] A. N. Shiryaev: *Optimal Stopping Rules*, Springer Verlag, 1978, 2008.
- [191] A. N. Shiryaev: *Probability-I*, 3rd Edition, Springer Verlag, 2016.

- [192] A. N. Shiryaev: *Probability-2*, 3rd Edition, Springer Verlag, 2016.
- [193] S. E. Shreve: *Stochastic Calculus for Finance II. Continuous Time Models*, Springer Verlag, 2004.
- [194] B. Simon: *Convexity: An Analytic Viewpoint*, Cambridge University Press, 2011.
- [195] A. Sinclair: *Algorithms for Random Generation and Counting: A Markov Chain Approach*, Progress in Theoretical Com. Sci., Springer Verlag, 1993.
- [196] P. M. Soardi: *Potential Theory of Infinite Networks*, Lect. Notes. Math., vol. 1590, Springer Verlag, 1994.
- [197] R. Stanley: *Enumerative Combinatorics. vol.1*, 2nd Edition, Cambridge University Press, 2012.
- [198] J.M. Steele: *Probability Theory and Combinatorial Optimization*, CBMS-NSF Regional Conf. Series in Appl. Math., SIAM, 1997.
- [199] J. M. Steele: *Stochastic Calculus and Financial Applications*, Springer Verlag, 2001.
- [200] H. Steinhaus: *Sur les distances des points dans les ensembles de mesure positive*, Fund. Mathematica, **1**(1920), 93-104.
<https://eudml.org/doc/212590>
- [201] J. M. Stoyanov: *Counterexamples in Probability*, 3rd Edition, Dover, 2013.
- [202] D. W. Strook: *Elements of Stochastic Calculus and Analysis*, CRM Short Course, Springer Verlag, 2018.
- [203] D. W. Strook: *Gaussian Measures in Finite and Infinite Dimensions*, Springer Verlag, 2023.
- [204] Y. Suhov, M. Kelbert: *Probability and Statistics by Example: II Markov Chains: A Primer in Random Processes and Their Applications*, Cambridge University Press, 2008.
- [205] L. Takács: *On an urn problem of Paul and Tatiana Ehrenfest*, Math. Proc. Camb. Phil. Soc., **86**(1979), 127-130.
- [206] M. Talagrand: *Upper and Lower Bounds for Stochastic Processes. Modern Methods and Classical Problems*, Springer Verlag, 2014.
- [207] M. Taylor: *Measure Theory and Integration*, Grad. Studies in Math., Amer Math. Soc., 2006.
- [208] C. B. Thomas: *Representation Theory of Finite and Lie Groups*, World Scientific, 2004.
- [209] H. Thorisson: *Coupling, Stationarity and Regeneration*, Springer Verlag, 2000.
- [210] H. F. Trotter: *An Elementary Proof of the Central Limit Theorem*, Arch. der Math., **10**(1959), 226-234.
- [211] R. van Handel: *Stochastic Calculus, Filtering, and Stochastic Control*, 2007 Course Notes.
<https://web.math.princeton.edu/~rvan/acm217/ACM217.pdf>,
- [212] R. van Handel: *Probability in High Dimensions*, APC 550, Lecture Notes, Princeton University, 2016.
<https://web.math.princeton.edu/~rvan/APC550.pdf>
- [213] A. W. van der Vaart, J. A. Wellner: *Weak Convergence and Empirical Processes. With Applications to Statistics*, Springer Verlag, 2nd Edition, 2023.
- [214] V. N. Vapnik, A. Ja. Cervonenkis: *On the uniform convergence of relative frequencies to their probabilities*, Theor. Probab. Appl., **16**(1971), 264-240.
- [215] V. N. Vapnik: *The Nature of Statistical Learning Theory*, 2nd Edition, Springer Verlag, 2000.
- [216] R. S. Varadhan: *Probability*, Courant Lect. Notes in Math., Amer. Math. Soc., 2001.
- [217] M. Viana, K. Oliveira: *Foundations of Ergodic Theory*, Cambridge University Press, 2016.
- [218] R. von Mises: *Probability, Statistics and Truth*, 2nd Edition, Dover, 1981.
- [219] M. J. Wainright: *High Dimensional Statistics. A Non-Asymptotic Point of View*, Cambridge University Press, 2019.
- [220] H. Wendland: *Scattered Data Approximation*, Cambridge University Press, 2005.
- [221] H. Weyl: *Über die Gleichverteilung von Zahlen mod Eins*, Math. Ann., **77**(1916), 313-352.
- [222] E. T. Whittaker, G. N. Watson: *A Course in Modern Analysis*, 4th Edition, Cambridge University Press, 1950.
- [223] H. S. Wilf: *Generatingfunctionology*, Academic Press, 1994
- [224] D. Williams: *Probability with Martingales*, Cambridge University Press, 1991.

Index

- $\mathcal{M}_{\text{loc}}^0$, 608
- $(n)_k$, v
- A^c , v
- $B(a, b)$, 140
- $B(x, y)$, 625
- $B_k(x)$, 134, 391
- $B_{a,b}(x)$, 140
- D_A , 415
- D_Γ , 347
- $D_n[f]$, 10
- $F_\# \mu$, 14
- H_x , 415
- L -function, 344
- $L^p(\Omega, \mathcal{S}, \mu)$, 37
- $N(\mu, \sigma^2)$, 62
- N_x , 417
- PG_X , 139
- R -function, 344
- T_x^k , 417
- T_A , 415
- T_Γ , 347
- T_x , 415
- $X \sim Y$, 41
- $X \sim \mu$, 41
- $X \stackrel{d}{=} Y$, 41
- $X_\bullet^T$, 279
- X_T , 278, 348
- $[M]$, 592
- $[f]_n$, 10
- $\#S$, v
- $\text{Ber}(p)$, 14
- $\text{Bin}(n, p)$, 15, 52
- $\text{Couple}(\mu_i, i \in I)$, 459
- $\text{EXP}(\lambda M)$, 615
- $\text{Elem}(\Omega, \mathcal{S})$, 9, 30
- Ent_2 , 161
- $\text{Gamma}(\nu, \lambda)$, 64
- $\Gamma(x)$, 625
- $\text{Geom}(p)$, 53
- $\text{HGeom}(w, b; n)$, 55
- $\text{HGeom}(w, b; n)$, 423
- $\Lambda(\mathbb{C})$, 4
- $\text{Markov}(\mathfrak{X}, \mu, Q)$, 400
- $\text{Meas}(\mathbf{X})$, 166
- $\text{NegBin}(k, p)$, 54
- $\Phi(x)$, 63
- Φ_X , 180
- $\text{Poi}(\lambda)$, 56
- $\text{Poi}_\nu(\lambda t)$, 257
- $\text{Prob}(\Omega, \mathcal{S})$, 12
- $\text{Prob}(\mathbf{X})$, 166
- $\text{Probs}(\mathbf{X})$, 166
- $\text{Proc}(\mathcal{F}_\bullet)$, 344
- $\text{Proc}(\mathcal{F}_{\text{prog}})$, 345
- $\text{Proc}_{\text{prog}}$, 345
- $\mathcal{H}_{\text{eff}}(x_+, S_-)$, 468
- Stud_p , 267
- $\text{Unif}(a, b)$, 62
- $\|f\|_p$, 37
- a.s., 13
- $\mathbb{E}[X]$, 41
- $\mathbb{E}[X \mid \mathcal{F}]$, 90
- $\mathbb{E}[X \mid F]$, 88
- $\mathbb{E}[Y \mid X]$, 90
- $\mathbb{E}[Y \mid X = x]$, 91
- $\mathbb{E}_{\mathbb{P}}[X]$, 41
- $\mathbb{G}(\sigma^2)$, 202
- $\mathbf{\Gamma}_{\mu, \sigma^2}$, 62
- $\mathbb{H}_\varphi$, 47
- $\mathbb{I}_n$, v
- $\mathbb{L}^2$, 555
- $\mathbb{L}^2(0, T)$, 555
- $\mathbb{L}^2(\mathcal{F}_\bullet)$, 555
- $\mathbb{L}_c^2$, 555
- $\mathbb{L}^{p, \text{loc}}$, 560
- $\mathbb{M}_X(t)$, 49
- $\mathbb{P}[S \mid \mathcal{F}]$, 94, 111
- T-filtration
 - see filtration, 270
- $\beta_{a,b}(x)$, 66

- γ_{μ, σ^2} , 62
- $\bigvee_{i \in I} \mathcal{S}_i$, 3
- λ_n , v
- ω_n , vi
- $\mathcal{C}(X_\bullet)$, 275
- σ_{n-1} , vi
- $\chi(r, \mathfrak{X})$, 306
- $\chi^2(n)$, 65
- $\text{Cov}[X, Y]$, 48
- $\text{dens}(\mathcal{F})$, 217
- $\dim_{V_C}(\mathcal{F})$, 217
- $\mathcal{B}_T$, v
- $\mathcal{B}_X$, 3
- $\mathcal{C}_T$, 125
- $\mathcal{E}_{\text{pred}}$, 555
- $\mathcal{E}_{\text{pred}}(0, T)$, 556
- $\mathcal{F} \perp\!\!\!\perp \mathfrak{g}\mathcal{H}$, 106
- $\mathcal{F}_T$, 280, 346
- $\mathcal{F}_{\text{pred}}$, 624
- $\mathcal{F}_{\text{prog}}$, 345
- $\mathcal{F}_{t+}$, 345
- $\mathcal{L}^0(\Omega, \mathcal{S})$, 8
- $\mathcal{L}^0(\Omega, \mathcal{S})_*$, 67
- $\mathcal{L}^0(\mathcal{S})$, 8
- $\mathcal{L}^1(\Omega, \mathcal{S}, \mu)$, 31
- $\mathcal{L}^\infty(\Omega, \mathcal{S})$, 8
- $\mathcal{L}^p$, 567
- $\mathcal{L}^{p, \text{loc}}$, 567
- $\mathcal{L}^0_+(\Omega, \mathcal{S})$, 8
- $\mathcal{L}^1_+(\Omega, \mathcal{S}, \mu)$, 31
- $\mathcal{M}^c$, 608
- $\mathcal{M}^{c, \text{loc}}$, 570
- $\mathcal{N}_\mu$, 15
- $\mathcal{O}_T$, 526
- $\mathcal{P}_x$, 412
- $\mathcal{SM}$, 601
- $\mathcal{S}|_X$, 4
- $\mathcal{S}^\mu$, 15
- $\mathcal{S}_1 \vee \mathcal{S}_2$, 3
- $\mathcal{S}_\infty$, 337
- $\mathcal{V}$, 572
- ess sup, 37
- $\frac{d\nu}{d\mu}$, 36
- $\mathfrak{S}_n$, v
- i.i.d., 127
- $\perp\!\!\!\perp Y$, 22
- i.o., 79
- λ -system, 4
- $\langle M, N \rangle_t$, 600
- $\langle M \rangle$, 276
- $\lceil x \rceil$, v
- $\lfloor x \rfloor$, v
- $\mu * \nu$, 74
- $\mu_0 \otimes \mu_1$, 67
- $\nu \ll \mu$, 36
- π -system, 4
- π^x , 426
- $\rho[X, Y]$, 49
- σ -additive, 12
- σ -algebra, 2
 - Borel, 3
 - complete, 15
 - completion of a, 15
 - trace of a, 4
- $\sigma(F)$, 3
- $\sigma(X_i, i \in I)$, 10
- $\sigma[X]$, 47
- $\xrightarrow{\text{a.s.}}$, 79
- $\xrightarrow{d}$, 167
- $\xrightarrow{P}$, 81
- 2^X , v
- 2^T_0 , 124
- 2^X_0 , v
- $\text{Var}[X]$, 47
- $|S|$, v
- φ -entropy, 47
- $\hat{X}_T$, 314
- $\{F \in S\}$, 3
- $]]S, T]]$, 279
- $b\mathbb{L}^2(0, T)$, 557
- $c\mathbb{L}^2(0, T)$, 555
- $c\mathbb{L}^p(0, T)$, 555
- d_v , 455
- f^+ , 9
- f^- , 9
- $g_\nu(x; \lambda)$, 64
- $u \sim v$, 433
- $x_0 \leftrightarrow x_1$, 409
- $x_0 \rightarrow x_1$, 409
- Abel's trick, 157
- absolutely continuous, 36
- absorbing state, 410
- acceptance matrix, 503
- acceptance-rejection, 244
- accessible states, 409
- activation function, 503
- AEP, 163
- algebra, 2
 - Bernoulli, 3
- alphabet, 160
 - entropy of the, 161
- André's reflection trick, 28, 134, 355
- aperiodic, 412
- arcsine distribution, 66
- Arnold's cat map, 524, 545, 547
- atom, 306
- baker's transform, 550
- ballot problem, 283
- Banach's principle, 538
- Banch space, 37
- Bernoulli
 - algebra, 3
 - random variable, 52
 - shift, 526
 - trial, 52
- Bernoulli-Laplace urn, 406, 423
- Beta distribution, 66, 119, 142, 344
- Beta function, 66, 343, 625
 - incomplete, 140
- bin packing, 291

- binomial distribution, 15
- binomial inversion, 59, 134
- birth-and-death chain, 407
- bit, 161
- Boltzmann machine, 509
- Borel
 - σ -algebra, 3
 - set, 3
 - subset, 3
- Borel-Cantelli, *see also* lemma
- boundary operator, 462
- bounded difference, 292
- bounded variation, 571
- bracket
 - of local martingales, 600
- branching process, 272, 312, 333, 408
- Brownian bridge, 381, 383
- Brownian event, 228
- Brownian motion, 232, 235, 555
 - Markov property, 348
 - multidimensional, 364
 - planar, 362
 - quadratic variation, 239
 - reflection principle, 355, 358
 - started at 0, 238
 - strong Markov property, 353, 355, 376
- Brwonina motion
 - quadratic variation, 564
- Cèsaro convergent, 155, 542
- Cèsaro means, 155, 537
- Cayley graph, 487
- cdf, 19, 20, 40
- chain
 - energy of, 463
- chains, 462
- chamber, 3, 88
- character, 488, 546
- characteristic function, 180
- Cheeger constant, 500
- Chernoff
 - bound, 197
 - method, 197, 289
- closed set, 410
 - irreducible, 410
- coboundary operator, 465
- cochain, 464
- code
 - binary, 164
 - instantaneous, 164
 - Shannon, 165
 - uniquely decodable, 164
- communication class, 410
- compensator, 275, 601
- conditional
 - expectation, 89, 90, 132
 - independence, 106
 - index, 131
 - probability, 25, 94, 111
- conditional distribution, 112
- conductance, 461, 500
- consistent family, 123
- convenient descriptions, 556
 - compatible, 556
 - nodes of, 556
- convergence
 - LP , 83
 - almost sure, 79
 - in p -mean, 83
 - in distribution, 166
 - in law, 166
 - in probability, 81, 309
- convex function, 46, 198
 - conjugate of, 199
- convolution, 74
- correlation, 49
- coupling, 458, 519
 - event, 460
 - optimal, 459, 519
 - time, 457
- coupon collector, 54, 132, 253, 320, 520
- covariance, 48
- covariance form, 225
- covariance kernel, 226, 348
- coverting time, 514
- Cramér-Wold device, 263, 264
- cummulative distribution function, 19
- cumulant, 197
- current, 462
- cutting
 - see* electric network, 473
- cylinder, 125
- début time, 347, 415, 435, 440, 444
- derangement, 61
- derangements problem, 61
- detailed balance equation, 503, 504
- detailed balance equations, 422, 496
- dipole, 109, 375, 377, 472
- Dirac measure, 14
- Dirichlet form, 496
- disintegration, 115
 - kernel, 115
- distribution
 - Bernoulli, 14
 - Beta, 66, 119, 142, 344
 - binomial, 15
 - chi-squared, 65
 - compound Poisson, 257
 - Erlang, 64
 - exponential, 71, 76, 252
 - Gamma, 64, 142
 - geometric, 53, 70, 252
 - Gumbel, 253
 - hypergeometric, 55, 56, 423
 - negative binomial, 53, 54
 - Poisson, 77
 - Student, 267
- distribution function, 19
- Doebelin condition, 521
- Doléans exponential, 616
- Doob
 - conditions, 280, 287
 - decomposition, 275, 391, 392

- downcrossing, 298
- effective
 - conductance, 468
 - resistance, 468
- Ehrenfest urn, 406, 413, 423, 479, 490, 495, 522
- electric network, 461
 - current, 462
 - cutting, 473
 - potential, 464
 - shorting, 473
 - source, 462
 - voltage, 464
- empirical
 - distribution, 207
 - process, 212
- empirical gap, 119
- empirical risk, 219
- energy, 463
- entropy, 47, 161, 261, 520
 - differential, 202
 - information, 161, 201
 - rate, 520
 - relative, 261, 521
 - Shannon, 161, 201
- equidistribution, 538
- ergodic, 529, 548
 - map, 529
- ess sup, 37
- Euler means, 458
- evaluation map, 122, 129, 238, 377
- evanescent, 624
- event, 13
 - almost sure, 13
 - exchangeable, 337
 - improbable, 13
 - permutable, 337
- excess risk, 219
- excessive function, 431
- exchangeable, 336
 - event, 337, 528
 - sequence, 336, 343, 525, 528, 539
- expectation, 41
- exponential martingale, 364
- extinction
 - event, 301, 312
 - probability, 301
- falling factorial, v
- Fatou's lemma, 34
- filtration, 270
 - complete, 345
 - admissible, 352, 584
 - non-anticipating, 352, 575
 - right-continuous, 345
 - usual, 346
- finite variation
 - function, 571
 - process, 572
- flow
 - Kirchhoff, 469
- formula
 - Bayes', 29
 - Duhamel, 590
 - Fourier inversion, 488
 - Gaussian regression, 266
 - Itô, 576, 613
 - Stirling, 254, 420, 421
 - Stirling's, 626
 - Viète, 255
 - Wald, 133, 317, 318
- Fourier transform, 180, 488
- function
 - Beta, 625
 - convex, 46
 - elementary, 9
 - Gamma, 625
 - nonnegative definite, 259
 - positive definite, 259
 - strictly convex, 46
 - Young, 263, 307
- Galton-Watson process, 272, 300, 312
- gambler's ruin, 319, 406, 420, 636
- game, 439
 - reward function of a , 439
 - value of a , 439
- Gamma function, 63, 64, 625
- Gaussian
 - random variables, 63
 - Hilbert space, 229
 - measure, 63, 224, 264
 - covariance form, 225
 - process, 226
 - centered, 226
 - random function, 228
 - vector, 225, 264, 265
 - white noise, 229, 397, 553
- gaussian
 - measure
 - centered, 225
- Gibbs
 - distribution, 507
 - sampler, 506
- graph, 274
 - locally finite, 274
 - path in, 434
 - length of, 434
 - random walk on, 274
- Haar
 - basis, 545
 - functions, 545, 550
- Hamming distance, 292, 491
- harmonic function, 274, 432, 467
- Hermite polynomials, 136, 137, 259, 364, 584
- hitting time, 432
- hitting time, 278, 347, 415, 432, 435, 468
- HMC, 400
 - Laplacian of, 431
 - reversible, 422
 - time reversed, 422
- hypothesis class, 218
 - PAC learnable, 220

- independence
 - conditional, 106
- independency, 21
- independent
 - events, 22
 - families, 22
 - random variables, 22
- indicator function, vi
- inequality
 - Hölder, 199
 - Azuma, 288, 290, 292
 - Bonferroni, 58
 - motivic, 58
 - Cauchy-Schwartz, 86
 - Chebyshev, 48
 - Doob's L^p , 330, 369
 - Doob's maximal, 329–331, 349, 369, 562, 596
 - Doob's upcrossing, 298, 300, 370
 - Gibbs, 161, 201
 - Hölder, 37
 - Hoeffding, 203, 288, 293
 - information, 201
 - Jensen, 46, 161
 - Kolmogorov's maximal, 148
 - Kunita-Watanabe, 600, 606
 - Markov, 33
 - McDiarmid, 216, 292, 294, 295
 - Mills ratio, 64, 137, 237
 - Minkowski, 37
 - one-sided Chebyshev, 48
- infinitely divisible
 - distribution, 256
 - random variable, 256
- integrable, 31
- integral, 31
 - Daniell-Stone, 33, 39
 - Lebesgue, 31
- integration by parts, 72
- invariant
 - measure, 422
 - distribution, 422, 426
 - function, 526
 - set, 526
- irreducible
 - HMC, 410
 - set, 410
- Ising potential, 508
- Itô
 - calculus, 576
 - formula, 576, 613
 - integral, 555, 568
 - isometry, 557
 - process, 575, 598
- joint probability distribution, 73
- kernel, 107
 - disintegration, 115
 - Markovian, 107
 - probability, 107
 - pullback, 108
 - push-forward by, 108
- Kirchhoff current, 466
 - potential of, 466
- Kirchhoff's
 - first law, 463
 - second law, 466
- Kirchhoff's laws, 463
- KL-divergence, 200, 521
- Kolmogorov sequence, 529
- Kolmogorov sequence
 - mixing, 543
- Koopman operator, 531
- Kullback-Leibler divergence, 200, 261, 521
- L-process, 345
- Lévy's
 - equivalence, 153, 301
 - forgery theorem, 362
 - inequality, 243
 - lois à trois variables, 360
 - martingale, 432
- Lévy-Prokhorov metric, 175
- Langevin equation, 590
- Laplace transform, 140, 250
- Laplacian, 432, 496
- law
 - of rare events, 61
 - of total probability, 26, 117
 - conditional, 26, 451
- Law of Large Numbers, 153, 294, 453, 539
- lazy chain, 458, 504
- Lebesgue
 - integral, 31
 - measurable, 69
 - measure, 19, 69
- lemma
 - 'sooner-rather-than-later', 285, 319, 417
 - Borel-Cantelli, 79, 80, 155, 238, 301, 350, 562, 579
 - first, 80, 151
 - second, 80, 151, 390, 392
 - Fatou, 35, 175, 176, 300, 309, 310, 315, 335, 370
 - Fekete, 86, 291
 - Hoeffding, 204, 289
 - Kronecker, 392
 - Kronecker's, 155
 - maximal, 536
 - Sauer, 217
 - Scheffé's, 310
- likelihood, 29
- likelihood ratio, 332
- Lindeberg condition, 194
- local martingale, 570
 - bracket, 600
 - quadratic variation of a, 592
- localizing sequence, 567, 570
- log-normal distribution, 139
- logistic map, 550
- longest common subsequence problem, 85
- Lusin space, *see also* space
- Lyapunov function, 431
 - coercive, 436
- Lyapunov's condition, 196

- map
 - measurable, 5
- marginal, 73
- Markov chain, 400
 - path space, 539
 - aperiodic, 413, 455, 458, 543
 - homogeneous, 400
 - initial distribution, 400
 - irreducible, 410
 - null recurrent, 426
 - path space, 403
 - positively recurrent, 426, 437, 453, 540
 - recurrent, 418, 436, 481, 539
 - reversible, 422, 460, 461, 486
 - state space, 400
 - transient, 418, 433, 481
- Markov property, 400, 404, 416, 424
 - strong, 415, 417–419, 429, 454
- martingale, 270–273, 275, 277, 279, 281, 284, 285, 288, 291, 301, 318, 320, 321, 331, 332, 362, 431, 432
 - L^p -bounded, 331
 - L^p , 331
 - L^p -bounded, 598, 602
 - backwards, 334
 - bounded L^p , 331
 - closed, 271, 290
 - component, 275
 - continuous, 368, 562
 - continuous local, 570
 - De Moivre, 272, 320
 - discrete time, 270
 - Doob, 271, 290
 - exponential, 364, 584
 - Levy's, 432
 - quadratic variation, 276
- matrix
 - primitive, 485
 - stochastic, 484
- mean, 41
- measurable
 - map, 5
 - set, 2
 - space, 2
 - isomorphism, 5
- measure, 12
 - Borel, 15
 - σ -finite, 12
 - atom of, 306
 - Dirac, 14
 - finite, 12
 - inner regular, 126
 - Lebesgue, 19
 - Lebesgue-Stieltjes, 71
 - outer regular, 126
 - probability, 12
 - pushforward of a, 14
 - Radon, 126, 173
 - regular, 126
 - subprobability, 166
 - uniform, 14, 15
- measure preserving, 523
- measured space, 13
- median, 21, 136, 243
 - interval, 21
- memoryless property, 71
- metric
 - Lévy-Prokhorov, 175
 - space, 166
 - variation, 456
- Metropolis
 - algorithm, 503
- mgf, 49
- Mills ratio, 64, 137, 237
- mixing, 542, 543
 - weakly, 547
- mixture, 74, 108, 116, 337
- moment generating function, 49
- monotone class, 10
- Monte-Carlo method, 160
- motion
 - Brownian, 235
 - Brownian standard, 232
 - pre-Brownian, 224
- Nedoma's pathology, 4
- negligible, 15
- noisy dynamical system, 409
- nonnegative definite function, 259
- null recurrent, 426
- Ohm's law, 466
- optimal gambling strategy, 323
- optimal stopping, 98, 439
- orbit, 526, 538
- order statistics, 143, 144, 254, 382
- orientation, 462
- Orlicz space, 263
- Ornstein-Uhlenbeck operator, 137
- paradox
 - waiting time, 77
- partition, 3
 - chamber of, 3
- path
 - in a graph, 434
- path space, 525
- perceptron, 218
- period, 412
- persistent state, 416
- pgf, 43, 51, 139, 394
- pmf, 41
- Pochhammer symbol, 627
- Poincaré phenomenon, 206, 262
- Poisson approximation, 60
- Poisson process, 76
- poissonization, 60
- Polya's urn, 133, 273, 342
- population risk, 219
- positive definite function, 259
- positively recurrent, 426
- posterior, 29
- predictable, 275, 624
- predictor, 93
- premeasure, 16

- σ -finite, 17
- principle
 - Dirichlet, 471
 - inclusion-exclusion, 57
 - Raleigh, 472, 473
 - reflection, 355
 - Thompson, 471
- prior, 29
- probability
 - density, 62
 - generating function, 43, 51, 139
 - measure, 12
 - Euclidean, 39
 - space, 13
 - filtered, 270
- probability distribution
 - continuous, 62
 - joint, 73
- problem
 - ballot, 27, 283
 - Banach's matchbox, 133
 - bin packing, 291
 - birthday, 135, 253
 - Buffon, 136, 638
 - coupon collector, 54, 132, 253, 320
 - derangements, 61
 - gambler's ruin, 319
 - longest common subsequence, 85, 290
 - Monty Hall, 640
 - occupancy, 253, 293
 - optimal stopping, 439
 - Polya's urn, 133, 273, 342
 - secretary, 101, 448
- process
 - independent increments, 363
 - T -sample, 348
 - branching, 272, 312, 333
 - distribution of, 224
 - empirical, 212
 - exchangeable, 336
 - finite variation, 572
 - Galton-Watson, *see also* Galton-Watson process
 - Gaussian, *see also* Gaussian
 - Itô, 598
 - Itô, 575
 - L -, 345
 - measurable, 344
 - Ornstein-Uhlenbeck, 590
 - Poisson, 76, 145, 257, 319, 363, 397, 511
 - predictable, 275, 277, 391
 - elementary, 555
 - progressive, 344, 555, 603
 - R -, 345
 - renewal, 79
 - separable, 368, 370
 - stochastic, 120
 - stopped, 279, 348
- product formula, 25
- projective family, 123, 125
- pushforward, 14, 15
- quadratic variation, 239, 276, 396, 584, 592, 598
- optional, 388
- predictable, 388
- quantile, 20, 208, 519
 - coupling, 519
- quasi-invariant
 - function, 527
 - set, 527
- quasi-mixing, 548
- R -filtration, 346
- R -process, 345
- Rademacher
 - complexity, 216
 - function, 550
 - random variable, 27, 152, 214, 221, 295, 391
 - symmetrization, 214
- Rademacher complexity
 - empirical, 295
- random
 - measure, 107, 337
 - process, 120
 - variable, 13
 - walk, 27, 271, 274, 305, 391, 393, 401, 407, 413, 420, 423
 - on groups, 487
 - standard, 27
- random function, 122
- random variable, 13
 - exponential, 64
 - Gamma, 64
 - Bernoulli, 52
 - binomial, 52
 - cdf of, 40
 - discrete, 41
 - probability mass function, 41
 - distribution of, 15, 40
 - expectation, 41
 - exponential, 71, 382
 - finite, 13
 - Gaussian, 62
 - geometric, 53
 - hypergeometric, 55
 - Kolmogorov-Smirnov, 381
 - law of, 40
 - mean, 41
 - negative binomial, 54
 - normal, 62
 - Poisson, 56
 - probability distribution of, 40
 - quantile, 21
 - Raleigh, 253
 - standard normal, 63
 - subgaussian, 202
 - uniform, 62
- random vector, 73
 - probability distribution of, 73
- recurrence class, 418
- recurrent state, 416
- Reflection Principle, 355
- regression curve, 91
- regular version, 111, 112
- resistance, 461
- return time, 347, 415, 436, 468

- reversible, 422, 460, 495
- reward function, 439
- sample range, 119
- sample space, 13
- SDE, 588, 616
- secretary problem, 101
- semimartingale, 601, 610
- separating collection, 167
- shift, 525
 - Bernoulli, 526
- shorting, *see* electric network 473
- sieve, 60, 61
- sigma-algebra, 2
- Skorokhod
 - inequality, 243
- SLE, 486, 487, 489, 496
- SLLN, 147, 153, 158, 207, 210, 212, 333, 341, 349, 454
- source, 462
- space
 - Lusin, 112, 113, 116, 125
 - measurable, 2
 - measured, 13
 - metric, 112, 166
 - Polish, 113, 167, 458
- stable sequence, 248
- standard deviation, 47
- state
 - accessible, 409
 - aperiodic, 412
 - period of, 412
 - persistent, 416
 - recurrent, 416
 - transient, 416
- stationary
 - distribution, 422
 - measure, 422
- stationary sequence, 525
- Stieltjes
 - measure, 20
- Stirling's formula, 420, 421, 626
- stochastic differential, 575
 - noisy part, 576
- stochastic differential equation, 588
- stochastic integral, 605
 - discrete, 277, 392
- stochastic matrix, 401
- stochastic process, 120, 270, 344
 - distribution, 224
 - distribution of, 122, 377
 - indistinguishable, 232, 275
 - measurable, 344
 - path, 229
 - progressive, 344
 - stochastically equivalent, 232
 - version of, 232
- stopped process, 279
- stopping time, 98, 278, 279, 284, 285, 415
 - Doob, 280, 281, 287, 315
 - optimal, 99
 - reducing, 316, 373
- stopping times, 98
- strictly convex function, 46
- strong Markov property, 353
- subadditivity, 86
- subgaussian, *see also* random variable
- submartingale, 270, 275, 277, 279, 281, 298, 363
 - backwards, 334
 - discrete time, 270
- superadditivity, 85
- superharmonic function, 432
- supermartingale, 270, 324, 363, 389
 - backwards, 334
 - discrete time, 270
- survival function, 71
- tail
 - algebra, 24, 338, 527
 - events, 24
- tail-algebra, 529
- temporal average, 532
- tent map, 525, 545, 550
 - mixing, 545
- theorem
 - Lévy's characterization, 618
 - Monotone Convergence, 45
 - L^p -martingale convergence, 331
 - Alexandrov, 17, 126
 - asymptotic equipartition property, 162
 - Backwards Martingale Convergence, 336, 530
 - Birkhoff's ergodic, 158, 534
 - Blumenthal's 0-1 law, 350
 - Bochner, 184
 - Bounded Convergence, 87, 169, 310
 - Carathéodory Extension, 18, 126
 - Cayley-Hamilton, 486
 - central limit, 188
 - début, 347
 - Daniell-Stone, 34
 - de Finetti, 338, 342
 - de la Vallée-Poussin, 308
 - Dominated Convergence, 34, 39, 82, 86, 87, 90, 168, 171, 180, 181, 183, 186, 305, 309, 324, 332, 354
 - dominated convergence, 557
 - Donsker's invariance, 378
 - Doob's regularization, 369
 - Dunford-Pettis, 310
 - Dynkin, 10, 90
 - Dynkin's $\pi - \lambda$, 4, 6, 23, 24, 68
 - Egorov, 303
 - ergodic, 455, 538
 - Fubini-Tonelli, 67, 92, 307, 365, 555
 - Glivenko-Cantelli, 209, 381
 - Helly selection, 176, 178, 256
 - Hewitt-Savage 0-1 law, 341
 - Ionescu-Tulcea, 128
 - Kolmogorov continuity, 233
 - Kolmogorov existence, 52, 125, 402
 - Kolmogorov one series, 148, 151, 153, 229, 332
 - Kolmogorov's continuity, 554
 - Kolmogorov's zero-one, 25, 80, 148, 312, 341, 529, 539
 - Kolmogorov-Smirnov, 211
 - Kolmogorov-Smirnov, 387
 - Krein-Milman, 530

- Kuratowski, 113
- Lévy's 0-1, 312, 540
- Lévy's continuity, 185, 301
- Lévy's equivalence, 153
- Lévy's forgery, 362
- Lévy's equivalence, 301
- Lebesgue differentiation, 558
- Lebesgue-Vitali, 309
- Lindeberg, 194
- Lindenstrauss-Johnson, 206
- mapping, 167, 175, 383
- martingale representation, 621
- mean ergodic, 532
- Monotone Class, 10, 67, 90, 108, 111, 374, 403, 621
- Monotone Convergence, 32, 34, 35, 68, 80, 93, 108, 167, 332, 370, 375
- Optional Sampling, 281, 284, 287, 316, 318, 320, 321, 330
- optional sampling, 446
- Optional Stopping, 279, 305, 315
- Perron-Frobenius, 485
- portmanteau, 169, 171, 173, 378
- Radon-Nikodym, 36, 94
- Raleigh, 472
- Riesz Representation, 39, 114
- Skhorkhod representation, 378
- Skhorokhod representation, 376
- Skorokhod representation, 167
- Slutsky, 172, 223, 260, 383
- Strong Law of Large Numbers, 154, 333, 341, 349, 539
- submartingale convergence, 300, 301, 305, 433
- support, 357, 361
- Tikhonov's compactness, 127
- UI Optional Sampling, 314, 323
- Wald's formula, 317
- weak law of large numbers, 157
- Weyl's equidistribution, 541
- tight, 177, 178, 256
- time
 - hitting, 278
 - optional, 346
 - stopping, 278, 346
- top-to-random, 520
- total variation, 571
- trail, 497
- transience class, 418
- transient state, 416
- transition matrix, 400
 - n -th step, 401
 - locally finite, 430
- tree, 482
 - radially symmetric, 482
- UI, 305, 309, 311, 314, 315, 317, 318, 320, 322, 331
- unbiased estimator, 118
- uniform absolute continuity, 306
- uniform integrability, 305
- unimodality, 59
- union bound, 59
- upcrossing, 298
 - number, 298
- usual augmentation, 346
- usual conditions, 346, 350, 352, 369
- vague convergence, 166
- Vapnik-Chervonenkis, *see also* VC
- variance, 47
- variation distance, 455
- VC
 - family, 217
 - dimension, 217
- waiting time, 77
- Wald-Bellman equation, 440
- walk, 409
- weak convergence, 166
- weakly mixing, 547
- weight function, 14
- Wiener
 - integral, 231, 268, 397, 553
 - measure, 238, 377
 - process, 232
- WLLN, 157
- Young function, 263, 307
- zero-one
 - algebra, 25, 529, 540
 - event, 25, 529
- zig-zag, 27