

HOMEWORK 11

DUE: THURS, MAY 1

Turn in problems 1 and 4

1. Suppose that

$$\alpha : G \rightarrow G$$

is an inner automorphism (conjugation by an element of G). Show that the induced map

$$\alpha_* : BG \rightarrow BG$$

is homotopic to the identity.

2. Show that for complex line bundles L_i over a space X , $c_1(L_1 \otimes L_2) = c_1(L_1) + c_1(L_2)$. (Hint - consider the “universal example”: the external tensor product $L_{univ} \otimes L_{univ}$ over $\mathbb{C}P^\infty \times \mathbb{C}P^\infty$).

Most functors on the category of complex vector spaces extend to constructions in the category of complex vector bundles over a space X . For instance, given complex vector bundles V and W , there exist vector bundles $V \otimes W$ and $\text{Hom}(V, W)$. The fibers over a point $x \in X$ are given by

$$\begin{aligned}(V \otimes W)_x &= V_x \otimes_{\mathbb{C}} W_x \\ \text{Hom}(V, W)_x &= \text{Hom}_{\mathbb{C}}(V_x, W_x).\end{aligned}$$

These constructions are easily produced locally using a trivializing cover. Functoriality can then be used to give transition functions.

3. Suppose that L is a complex line bundle on a paracompact space. Let \bar{L} denote the *conjugate bundle*, where the fibers are given the conjugate action of \mathbb{C} .

- (a) Show that there is a bundle isomorphism $\bar{L} \cong \text{Hom}(L, \mathbb{C})$, where \mathbb{C} denotes the trivial line bundle.
(b) Conclude that there is a bundle isomorphism $L \otimes \bar{L} \cong \mathbb{C}$.
(c) Deduce that $c_1(\bar{L}) = -c_1(L)$.

4. Let \mathcal{L} be the restriction of the universal line bundle on $\mathbb{C}P^\infty$ to $\mathbb{C}P^n$.

(a) Show that the tangent bundle $T\mathbb{C}P^n$ to $\mathbb{C}P^n$ can be identified with the bundle $\text{Hom}(\mathcal{L}, \mathcal{L}^\perp)$. Here, \mathcal{L}^\perp is the perpendicular bundle of dimension n over $\mathbb{C}P^n$, whose fiber over a line L in \mathbb{C}^{n+1} is the perpendicular space L^\perp .

- (b) Use the axioms of Chern classes (i.e. the Cartan formula) to deduce that

$$c_i(T\mathbb{C}P^n) = (-1)^i \binom{n+1}{i} x^i$$

where $x \in H^2(\mathbb{C}P^n)$ is the generator given by $c_1(\mathcal{L})$. Hint: show that there is an isomorphism

$$TCP^n \oplus \mathbb{C} \cong TCP^n \oplus (\overline{\mathcal{L}} \otimes \mathcal{L}) \cong \text{Hom}(\mathcal{L}, \mathbb{C}^{n+1}).$$